

Assignment 1

Parker DeBruyne - V00837207

2023-05-21

1. (4 points) In November of each year, a walk-in clinic allows people to walk in to get a flu shot. Let X be the number of people who come to the clinic for a flu shot on a randomly selected day (in November). Suppose X has the following distribution:

| | | | | | | |
|------------|-----|-----|-----|-----|-----|-----|
| X | 0 | 1 | 2 | 3 | 4 | 5 |
| $P(X = x)$ | 0.3 | 0.1 | 0.2 | 0.1 | 0.1 | 0.2 |

If at least 3 people walk in for a flu shot on a particular day, what is the probability that there are 4 or fewer who walk in for a flu shot that day?

$$\begin{aligned}P(X \leq 4 | X \geq 3) &= \frac{P((X \leq 4) \cap (X \geq 3))}{P(X \geq 3)} \\&= \frac{0.2}{0.4} \\&= 0.5\end{aligned}$$

2. (6 points) Daily sales at a gas station are thought to be independent of one another with daily mean \$5000 and standard deviation \$700. Approximate the probability that the average daily sales over one year (i.e. 365 days) is greater than \$5,100.

With $n \geq 30$ it is appropriate to use the Normal distribution:

$$X \sim Normal(\mu = 5000, \sigma = 700)$$

$$P(X > 5100) = 1 - P(X \leq 5100)$$

$$= 1 - P\left(\frac{X - \mu}{\sigma} \leq \frac{5100 - 5000}{700}\right) = 1 - P(Z \leq 0.142857162)$$

$$\simeq 1 - P(Z \leq 0.14)$$

$$= 1 - 0.5517$$

$$= 0.4483$$

3. (10 points) Suppose that diseased trees are distributed randomly and uniformly throughout a large forest with an average of λ per acre. Let X denote the number of diseased trees in a randomly chosen one-acre forest plot where the range of X is, $X = \{0, 1, 2, \dots\}$.

- (a) We will use the Poisson distribution with mean λ to model X . Write down the probability mass function (p.m.f.) for X . Why would this distribution be suitable for modelling the number of diseased trees in a randomly chosen one-acre plot of forest?

The Poisson probability mass function is given as follows:

$$f(x) = \frac{\lambda^x e^{-\lambda}}{x!}$$

The poisson distribution models the random occurrence of events in any fixed interval, including time and space. In this case, we know the average number of diseased trees λ per one-acre interval, and that each interval is independent.

- (b) Suppose that we observe the number of diseased trees on n randomly chosen one-acre parcels, X_1, X_2, \dots, X_n . The random variables X_1, X_2, \dots, X_n can be assumed to be independent. Write down the JOINT probability mass function for X_1, X_2, \dots, X_n . Simplify this expression which is a function of λ and the X 's.
- s. Use this joint distribution for the remainder of Question 1.

$$\begin{aligned} P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n) &= \left(\frac{\lambda^{x_1} e^{-\lambda}}{x_1!}\right) \left(\frac{\lambda^{x_2} e^{-\lambda}}{x_2!}\right) \dots \left(\frac{\lambda^{x_n} e^{-\lambda}}{x_n!}\right) \\ &= \frac{\lambda^{\sum_{i=1}^n x_i} e^{-n\lambda}}{\prod_{i=1}^n x_i!} \end{aligned}$$

- (c) We are going to use the Method of Maximum Likelihood to estimate λ . Write down the Likelihood function $L(\lambda)$.

The Likelihood function omits multiplicative constants to simplify and focus on our desired parameter λ

$$\begin{aligned} L(\lambda) &= c p(x_i, \lambda), c = \prod_{i=1}^n x_i! \\ &= L(\lambda) = e^{-n\lambda} \lambda^{\sum_{i=1}^n x_i} \end{aligned}$$

- (d) Write down the Log-likelihood function $l(\lambda)$.

$$l(\lambda) = -n\lambda + \left(\sum_{i=1}^n x_i\right) \ln \lambda$$

- (e) Write down the Score Function $S(\lambda)$.

$$S(\lambda) = l'(\lambda) = -n + \frac{\sum_{i=1}^n x_i}{\lambda}$$

- (f) Derive the maximum likelihood estimate of λ .

The maximum likelihood is when the previous equality is set to zero and we solve for λ

$$0 = -n + \frac{\sum_{i=1}^n x_i}{\lambda}$$

Solving, we get:

$$\lambda = \frac{1}{n} \sum_{i=1}^n x_i$$

- (g) Write down the Information Function $I(\lambda)$.

$$I(\lambda) = -l''(\lambda) = -\left(\frac{-\sum_{i=1}^n x_i}{\lambda^2}\right) = \frac{\sum_{i=1}^n x_i}{\lambda^2}$$

- (h) Use the second derivative test to show that you have found a maximum.

$$l''(\lambda) = \frac{-\sum_{i=1}^n x_i}{\lambda^2}$$

$l''(\lambda)$ is always negative as n goes to ∞ implies that we have found a maximum.

- (i) Suppose that the numbers of diseased trees observed in $n = 4$ randomly chosen one-acre parcels were: 5, 8, 9, 2. Compute the maximum likelihood estimate of λ using this data.

$$\lambda = \frac{1}{4} \sum_{i=1}^4 x_i = \frac{1}{4}(5 + 8 + 9 + 2) = \frac{1}{4}(24)$$

$$\lambda = 6$$