

Strategic coexistence theory for evolutionary games

Sang Woo Park^{1,2,*}

1 School of Biological Sciences, Seoul National University, Seoul, Korea

2 Institute for Data Innovation in Science, Seoul National University, Seoul, Korea

*Corresponding author: sangwoopark@snu.ac.kr

Abstract

Evolutionary game theory and ecological coexistence theory both seek to predict the outcome of competition between biological entities, be they strategies or species, but the two fields have relied on largely separate approaches. Replicator equations provide a foundation for predicting the outcome of strategic competition, yet they do not explicitly decompose the mechanisms underlying these outcomes. Inspired by modern coexistence theory from community ecology, we develop Strategic Coexistence Theory (SCT), which translates payoff-based invasion conditions into strategic niche differences and fitness differences between any two competing strategies described by classic replicator dynamics. SCT recovers the classic classification of two-strategy games, distinguishing competitive exclusion, coexistence, and priority effects within a shared niche–fitness difference space. Applying SCT to five mechanisms for the evolution of cooperation further reveals that these mechanisms promote cooperation through distinct dynamical routes: kin selection, network reciprocity, and group selection primarily reduce fitness differences, whereas direct and indirect reciprocity destabilize competition and generate priority effects. Finally, applying SCT to microbial public-goods competition shows that nonlinear microbial growth can both stabilize and equalize competition between cooperators and defectors, allowing coexistence. Together, these results show that SCT provides a common language for comparing coexistence mechanisms across evolutionary games.

Significance statement

Evolutionary game theory provides a powerful framework for predicting the outcome of competition between strategies, including the dominance of one strategy, coexistence of multiple strategies, and bistability. However, a common framework for comparing how different mechanisms promote strategy coexistence remains elusive, even though stable polymorphisms, such as the coexistence of cooperators and defectors, are widely observed across biological systems. To address this gap, we developed Strategic Coexistence Theory (SCT), which extends Modern Coexistence Theory from community ecology to evolutionary games. Applying SCT to classic games, mechanisms of cooperation, and microbial public-goods competition shows

that seemingly similar evolutionary outcomes can arise through distinct mechanisms. This approach links ecological theory with replicator dynamics, providing a complementary tool for analyzing evolutionary games.

Introduction

Understanding the mechanisms that promote the evolutionary stability of competing strategies is a fundamental aim in evolutionary biology (Lewontin, 1961; Slobodkin, 1964; Smith and Price, 1973; Taylor and Jonker, 1978). To this end, evolutionary game theory (EGT) and replicator dynamics have provided key foundations, allowing us to predict the outcome of strategic competition through payoff inequalities, invasion criteria, and stability analysis (Schuster and Sigmund, 1983; Hofbauer and Sigmund, 1998; Cressman, 2003; Cressman and Tao, 2014). Recently, there has been growing interest in characterizing similarities between replicator dynamics and the dynamics of competing species (Gokhale and Traulsen, 2025; Tarnita and Traulsen, 2025; Allesina, 2026), building on the classical mathematical equivalence between replicator equations and generalized Lotka–Volterra models (Hofbauer, 1981; Page and Nowak, 2002). In particular, Tarnita and Traulsen (2025) recently argued that a closer synthesis between ecology and evolutionary game theory can provide new insights into ecological and evolutionary assembly, competition, and coexistence. This renewed perspective between ecological and evolutionary dynamics suggests that evolutionary games may be interpreted not only as payoff-based competition but also as problems of strategic coexistence.

Interestingly, even a simple two-strategy game can yield heterogeneous outcomes that mirror outcomes in community ecology (Hauert, 2002; Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025): the dominance of a single strategy (prisoner’s dilemma and harmony games), coexistence (hawk–dove games), and bistability (stag-hunt games). Such differences in conditions that promote evolutionary stability and dominance of a single strategy and those that permit strategy coexistence indicate that EGT can be understood from the perspective of coexistence theory (Motro, 1991; Doebeli et al., 2004; Hauert and Doebeli, 2004; Doebeli and Hauert, 2005; Gore et al., 2009; Souza et al., 2009; Archetti and Scheuring, 2011; Steidinger and Bever, 2014; Tarnita and Traulsen, 2025). In fact, stable polymorphisms of evolutionary strategies are observed in microbial and other biological systems (Gore et al., 2009; Cordero et al., 2012; Kocher et al., 2018; Pollak et al., 2021; Lin et al., 2024), but coexistence-theoretic perspective has often remained implicit. Recasting game dynamics as a coexistence problem immediately raises the following question: when two strategies coexist, does the corresponding mechanism stabilize or equalize competition (Chesson, 2000)? While EGT allows us to predict whether two strategies can coexist or not, a common framework for comparing coexistence mechanisms between competing strategies remains elusive.

Modern coexistence theory (MCT) from community ecology presents a particu-

larly useful framework for quantifying coexistence mechanisms (Chesson, 2000; Adler et al., 2007; Godoy et al., 2014; Godoy and Levine, 2014; Kraft et al., 2015; Adler et al., 2026). MCT predicts that coexistence of two competing entities, be they species or strategies, requires stabilizing niche differences to overcome average fitness differences (Chesson, 2000; Adler et al., 2007; Kraft et al., 2015). In community ecology, stabilizing niche differences represent differences in factors that limit the growth of competing species, causing each species to limit itself more than it limits its competitor (Chesson, 2000). In EGT, a strategy’s niche can be interpreted as strategic environments defined by the composition of competing strategies and the resulting interactions (Taylor and Jonker, 1978; Cressman and Tao, 2014; Tarnita and Traulsen, 2025). Thus, stabilizing niche differences arise when each strategy is favored by a different strategic environment, such that the spread of each strategy creates conditions that limit its own relative growth and favor the recovery of its competitor. In contrast, fitness differences represent differences in the ability of each entity to compete under the limiting environments created by its competitor (Chesson, 2000). This differs from payoff differences in EGT, which depend on the underlying strategy frequency and therefore combine both stabilizing effects caused by changes in the strategic environment and residual fitness differences that favor one strategy over another (Taylor and Jonker, 1978; Cressman and Tao, 2014; Tarnita and Traulsen, 2025).

So far, MCT has been primarily utilized for understanding species coexistence, but recent work showed that MCT can be extended beyond species competition, especially for studying pathogen strain competition, allowing one to directly quantify niche and fitness differences of competing entities, be they species or strains (Sieben et al., 2022; Park et al., 2024, 2026). Inspired by this effort, we present Strategic Coexistence Theory (SCT) for predicting coexistence and exclusion of competing strategies in EGT by decomposing invasion growth rates into strategic niche and fitness differences. We begin by introducing SCT, which can be applied to any generic system of EGT described by replicator equations. We show that SCT can recover outcomes from classic, two-strategy games, providing a reinterpretation of classic results from coexistence-theoretic perspective. We then apply SCT to comparing five mechanisms for the evolution of cooperation, revealing that direct and indirect reciprocity destabilize competition between cooperators and defectors, thus pushing the system towards priority effect regime. Finally, we consider a case study of microbial cooperation, where nonlinear microbial growth can both stabilize and equalize competition between cooperation and defection, allowing coexistence of strategies. Together, these examples show that SCT provides a common, ecological language for comparing the mechanisms underlying strategic evolution.

Strategic coexistence theory

Here, we present a brief derivation of strategic coexistence theory (SCT) (Figure 1). To do so, we begin with a classic replicator equation, which allows us to model changes in the frequency of competing strategies based on their payoffs (Schuster and Sigmund, 1983; Hofbauer and Sigmund, 1998; Cressman, 2003; Cressman and Tao, 2014):

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (1)$$

where x_i represents the frequency of strategy i , $\pi_i(\mathbf{x})$ represents the payoff of strategy i for a given strategy distribution $\mathbf{x}$, and $\bar{\pi}(\mathbf{x}) = \sum_i \pi_i(\mathbf{x})x_i$ represents the average payoff. Since strategy frequencies must sum to 1 (i.e., $\sum_i x_i = 1$), a single-strategy equilibrium dominated by strategy i can be described by a unit vector $\mathbf{e}_i$, where all elements are equal to zero except for the i -th element. We use these single-strategy equilibria to define niche and fitness differences between two competing strategies i and j .

Because payoffs in many games can be negative, payoff ratios can be difficult to interpret. Thus, we work on a positive multiplicative scale by exponentiating the payoff terms:

$$W_i(\mathbf{x}) = \exp(\pi_i(\mathbf{x})). \quad (2)$$

This exponential transformation is convenient because it preserves the payoff ordering and does not alter the invasion conditions. Therefore, this allows us to compare strategic environments through ratios, as in modern coexistence theory. For convenience, we refer to $W_i(\mathbf{x})$ as the multiplicative payoff of strategy i .

First, we define the niche overlap ρ between strategies i and j as the ratio between the geometric mean of each strategy's multiplicative payoff in the strategic environment that it creates when common, $\sqrt{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}$, and the geometric mean of each strategy's multiplicative payoff in the environment generated by its competitor, $\sqrt{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}$. The corresponding niche difference is then given by

$$1 - \rho = 1 - \sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}}. \quad (3)$$

When $1 - \rho > 0$, the geometric mean of the multiplicative payoffs across the two strategies is greater in competitor-generated monomorphic environments than in self-generated monomorphic environments. In other words, intraspecific competition is stronger than interspecific competition at the boundary. When $\rho > 1$, we interpret the resulting negative niche difference, $1 - \rho < 0$, as a destabilizing effect corresponding to positive frequency dependence, meaning that the strategy that is already common is favored.

Second, we define the fitness difference as the ratio between the geometric means of the multiplicative payoff of each strategy across the two single-strategy environ-




Figure 1: **A schematic diagram summarizing strategic coexistence theory.** Strategic coexistence theory decomposes pairwise replicator dynamics into stabilizing and equalizing mechanisms. Mutual invasion conditions can be rewritten on a multiplicative payoff scale, allowing us to derive strategic niche difference $1 - \rho$ and fitness difference κ_j/κ_i based on multiplicative payoff ratios. Then, the inequality $\rho < \kappa_j/\kappa_i < 1/\rho$ determines mutual invasion. We note that the exact values of niche and fitness differences are sensitive to payoff scaling, while dynamical outcomes are still preserved.

ments:

$$\frac{\kappa_j}{\kappa_i} = \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}. \quad (4)$$

In other words, strategy j has an average competitive advantage over strategy i ($\kappa_j/\kappa_i > 1$) when it performs better, on average, across the environments generated by itself and its competitor.

Together, for any two strategies described by the classic replicator equation, mutual invasion requires stabilizing niche differences to overcome average fitness differences. Specifically, the condition for mutual invasion can be rewritten as (Chesson, 2000):

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (5)$$

In simple systems with monotonic interior dynamics, these boundary invasion conditions also determine the global dynamical outcome, allowing competition to be classified into coexistence, competitive-exclusion, and priority-effect regimes. However, MCT and SCT are based on invasion conditions at the boundaries and do not generally characterize nonlinear interior dynamics. Mutual invasion still guarantees at least one stable internal equilibrium, but it does not determine the number or stability of additional internal equilibria. Likewise, boundary conditions associated

with competitive exclusion or priority effects do not rule out stable internal equilibria. As we show later, nonlinear dynamics can allow stable coexistence even when a system lies in the boundary-defined competitive-exclusion regime. Therefore, when interior dynamics are nonmonotonic, we use these classifications to describe boundary invasibility rather than complete dynamical outcomes.

Despite this limitation, the mutual-invasion inequality provides a useful perspective for comparing how different mechanisms alter boundary invasibility. Notably, the inequality can be applied to any pairwise comparison between two strategies described by the classic replicator equation because ρ and κ_j/κ_i can be calculated directly from ratios of multiplicative payoffs. This provides a common framework for comparing qualitative changes in boundary invasibility across different games. Detailed derivations are provided in Supplementary Text S1.

We note that the absolute magnitudes of niche and fitness differences are sensitive to payoff scaling because they are defined on an exponentiated payoff scale. Therefore, while SCT allows us to compare different games on shared axes, we cannot compare the exact values of niche and fitness differences across different games. Nonetheless, as we show in Supplementary Text S2, the qualitative outcome of strategic competition under SCT is preserved because payoff rescaling preserves payoff orderings. Therefore, we focus on comparing qualitative outcomes from SCT, including directions of change in niche and fitness difference and resulting dynamical regimes, and refrain from making any comparisons of magnitude of niche and fitness difference values.

Classic two-strategy games

As an illustrative example, we begin with classic two-strategy games (Hauert, 2002; Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025), which can be described by the following payoff matrix (Figure 2A):

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (6)$$

Here, the first and second rows represent strategies 1 and 2, respectively, and the columns represent the strategy of the interacting opponent. Writing x_1 to denote the frequency of strategy 1, the expected payoffs are

$$\pi_1(\mathbf{x}) = Rx_1 + S(1 - x_1), \quad (7)$$

$$\pi_2(\mathbf{x}) = Tx_1 + P(1 - x_1). \quad (8)$$

Thus, the dynamics can be written as

$$\frac{dx_1}{dt} = x_1(1 - x_1) [(R - T)x_1 + (S - P)(1 - x_1)]. \quad (9)$$

The signs of $T - R$ and $S - P$ determine whether each strategy can invade the other when rare. Specifically, strategy 2 can invade strategy 1 when $T > R$, whereas

197 strategy 1 can invade strategy 2 when $S > P$. These two invasion conditions partition
 198 the game space into four regimes (Figure 2B): prisoner’s dilemma ($T > R > P >$
 199 S), hawk–dove game ($T > R, S > P$), stag–hunt game ($R > T, P > S$), and
 200 harmony game ($R > T, S > P$). In standard EGT, these outcomes are obtained by
 201 analyzing the stability of equilibrium conditions. For simple games, this analysis is
 202 straightforward.




Figure 2: **Coexistence-theoretic interpretation of classic two-strategy games.** (A) General payoff matrix for a two-strategy game, where rows represent the focal strategy and columns represent the strategy of the interacting opponent. (B) Classification of four classic games based on the payoff differences $T - R$ and $S - P$. Prisoner’s dilemma and harmony games lead to the dominance of one strategy; hawk–dove games allow coexistence; and stag–hunt games result in bistability. (C) Mapping of the same four games onto niche–fitness difference space. Hawk–dove games correspond to the coexistence regime, where stabilizing niche differences overcome fitness differences. Prisoner’s dilemma and harmony games correspond to competitive-exclusion regimes, whereas stag–hunt games correspond to the priority-effect regime. Negative values of $1 - \rho$ indicate positive frequency dependence and destabilization. $R > P$ is assumed throughout, but this does not affect coexistence outcome in any way.

203 A coexistence-theoretic approach recovers the same classification but provides
 204 a different interpretation (Figure 2C). For a two-strategy game determined by this
 205 payoff matrix, the niche difference and fitness difference are given by the following
 206 set of equations (Supplementary Text S3):

$$1 - \rho = 1 - \exp\left(\frac{R + P - S - T}{2}\right), \quad (10)$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{T + P - R - S}{2}\right). \quad (11)$$


As shown above, mutual invasion requires $\rho < \kappa_2/\kappa_1 < 1/\rho$, which is equivalent to $T > R$ and $S > P$. This corresponds to the hawk–dove game, in which each strategy can invade its competitor when rare and the two strategies coexist. The remaining games also have natural interpretations in niche–fitness difference space. Prisoner’s dilemma and harmony games represent competitive exclusion regimes, where one strategy has a fitness advantage that exceeds niche differences. In contrast, stag–hunt games represent a priority-effect regime, where neither strategy can invade the other and the outcome depends on initial conditions. Therefore, coexistence theory not only allows us to recover the classic results from two-strategy games but also provides new interpretations for these results in terms of niche and fitness differences. Building on this, we next apply this framework to comparing core mechanisms of cooperation.

Comparisons of mechanisms for the evolution of cooperation

Understanding mechanisms that promote the evolution of cooperation is a central goal in evolutionary biology (West et al., 2007; Fletcher and Doebeli, 2008; Clutton-Brock, 2009; Kingma et al., 2014; Rakoff-Nahoum et al., 2016; Hilbe et al., 2018; Apicella and Silk, 2019; West et al., 2021; Lee et al., 2022; Palmer and Foster, 2022; Michel-Mata et al., 2024). Here, we compare five classic mechanisms for the evolution of cooperation using SCT (Nowak, 2006) and ask how they differ from the perspective of SCT.

Specifically, we consider the following mechanisms: kin selection, direct reciprocity, indirect reciprocity, network reciprocity, and group selection (Figure 3A). Kin selection favors cooperation when two interacting individuals are genetic relatives, where the genetic relatedness r must be greater than the cost-to-benefit ratio: $r > c/b$ (Hamilton, 1964). Direct reciprocity favors cooperation when current cooperation may promote future cooperation by the same individual through repeated encounters (Trivers, 1971; Axelrod and Hamilton, 1981). In this case, the probability of repeated interaction (i.e., “next round”) must be greater than the cost-to-benefit ratio: $w > c/b$. Indirect reciprocity favors cooperation when current cooperation may promote future cooperation by different individuals through reputation (Nowak and Sigmund, 1998; Wedekind and Milinski, 2000; Nowak and Sigmund, 2005). Thus, the social acquaintanceship (i.e., the probability of knowing someone’s reputation) must be greater than the cost-to-benefit ratio: $q > c/b$. Network reciprocity favors cooperation via clustering of cooperators, requiring the benefit-to-cost ratio to be greater than the average number of neighbors: $b/c > k$ (Nowak and May, 1992; Lieberman et al., 2005). Finally, group selection favors cooperation via multilevel selection, where groups with more cooperators exhibit faster growth at the group level even if defectors reproduce faster at the individual level (Traulsen and Nowak, 2006). This requires $b/c > 1 + (n/m)$, where n is the maximum group size and m is the number of groups. All of these mechanisms can be described by a simple

247 2×2 payoff matrix (Nowak, 2006), from which the corresponding niche and fitness
 248 differences can be computed using Eq. (10) and Eq. (11). Details on niche and fitness
 249 difference calculations are presented in Supplementary Text S4.



250 First, the prisoner's dilemma provides a baseline scenario, where defection is

evolutionarily stable (Figure 3A). From the perspective of SCT, there is complete niche overlap, and therefore defection competitively excludes cooperation (Figure 3B). Building on this, SCT shows that five mechanisms promote evolutionary stability of cooperation via different dynamical routes (Figure 3B). For example, kin selection allows the fitness of cooperation to increase as genetic relatedness r increases (Figure 3B). However, it does not affect the niche difference between cooperation and defection. Similarly, network reciprocity and group selection modulate the fitness of cooperation without causing any changes in strategic niches. In contrast, direct and indirect reciprocity destabilize competition while increasing the fitness of cooperation, thereby pushing the competition outcome towards a priority effect regime. Equivalently, these two mechanisms promote bistability of defection and cooperation.

SCT further reveals another difference between direct and indirect reciprocity. Direct reciprocity remains within the priority-effect regime as w reaches 1, whereas indirect reciprocity reaches the boundary between the priority-effect and competitive-exclusion regimes at $q = 1$ (Supplementary Text S4).

We note that none of these mechanisms allow coexistence of cooperation and defection. In many biological systems, such coexistence is often observed (Gore et al., 2009; Cordero et al., 2012; Kocher et al., 2018; Pollak et al., 2021; Lin et al., 2024). Therefore, we move on to the final example, applying SCT to microbial cooperation, where nonlinear benefits and environmental context can promote coexistence of cooperators and defectors.

Can stochasticity prevent coexistence?

So far, we have focused on simple deterministic games. However, many ecological and evolutionary processes are intrinsically stochastic. Recently, Wang et al. (2025) showed that environmental fluctuations can promote coexistence in games that would otherwise lead to competitive exclusion in a deterministic environment. From the perspective of SCT, this suggests that stochasticity can stabilize competition by promoting the recovery of strategies when rare. Interestingly, Wang et al. (2025) also showed that stochasticity can generate bistable coexistence and multiple internal equilibria. Thus, although stochasticity can stabilize competition near the boundaries, it can also generate destabilizing frequency dependence in the interior. This immediately raises the question of whether stochasticity can generate sufficiently strong positive frequency dependence to destabilize competition, thus pushing the system out of the coexistence regime. Here, we use SCT to separate these effects and clarify whether stochasticity stabilizes or destabilizes strategic competition (Supplementary Text S5).

Following Wang et al. (2025), we consider a 2×2 game in the presence of environmental noise. The stochastic payoff matrix is given by

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix} + \frac{\zeta}{\sqrt{s}} \begin{pmatrix} \sigma_R & \sigma_S \\ \sigma_T & \sigma_P \end{pmatrix}, \quad (12)$$

where the first matrix represents deterministic payoffs, ζ represents a noise term with zero mean and unit variance, and the second matrix determines the intensity of environmental stochasticity. Assuming weak selection $s \ll 1$, the evolution of two competing strategies in a large population can be described by the following equation:

$$\frac{dx}{dt} = sx(1-x) \left(\pi_C - \pi_D + \left(\frac{1}{2} - x \right) (\sigma_C - \sigma_D)^2 \right), \quad (13)$$

where $\pi_C = Rx + S(1-x)$ and $\pi_D = Tx + P(1-x)$ and $\sigma_C = \sigma_R x + \sigma_S(1-x)$ and $\sigma_D = \sigma_T x + \sigma_P(1-x)$. After rescaling time as $\tau = st$, the invasion growth rates of each strategy when rare can be written as:

$$r_C = S - P + \frac{1}{2}(\sigma_S - \sigma_P)^2, \quad (14)$$

$$r_D = T - R + \frac{1}{2}(\sigma_T - \sigma_R)^2. \quad (15)$$

This shows that invasion growth rates under environmental stochasticity will be greater than or equal to their deterministic counterparts, meaning that environmental stochasticity at the boundaries is intrinsically stabilizing.

Then, an extended SCT decomposition shows that niche overlap and fitness difference can be written as:

$$\rho = \exp \left(\frac{R + P - S - T}{2} - \frac{(\sigma_R - \sigma_T)^2 + (\sigma_S - \sigma_P)^2}{4} \right) \quad (16)$$

$$\frac{\kappa_D}{\kappa_C} = \exp \left(\frac{T + P - R - S}{2} + \frac{(\sigma_R - \sigma_T)^2 - (\sigma_S - \sigma_P)^2}{4} \right) \quad (17)$$

Writing $\hat{\rho}$ and $\hat{\kappa}_D/\hat{\kappa}_C$ to denote niche overlap and fitness difference in the absence of stochasticity and taking logs of both sides for convenience, we have:

$$\Delta \log \rho = \log \rho - \log \hat{\rho} = -\frac{(\sigma_R - \sigma_T)^2 + (\sigma_S - \sigma_P)^2}{4} \quad (18)$$

$$\Delta \log \frac{\kappa_D}{\kappa_C} = \log \frac{\kappa_D}{\kappa_C} - \log \frac{\hat{\kappa}_D}{\hat{\kappa}_C} = \frac{(\sigma_R - \sigma_T)^2 - (\sigma_S - \sigma_P)^2}{4}. \quad (19)$$

Therefore, stochasticity reduces niche overlap and increases niche difference. In contrast, stochasticity can either amplify or reduce fitness differences depending on which strategy's invasion rate is more strongly affected by environmental fluctuations. Notably, changes in fitness difference driven by environmental stochasticity will always be bounded by those in niche overlap:

$$\Delta \log \rho \leq \Delta \log \frac{\kappa_D}{\kappa_C} \leq -\Delta \log \rho \quad (20)$$

This result implies that environmental stochasticity cannot generate sufficient fitness difference to push the system out of the coexistence regime.




Figure 4: **Effects of environmental stochasticity on strategic competition.** (A) Assumed deterministic payoff structures across four scenarios. Environmental stochasticity is assumed to be proportional to these payoffs, scaled by a term k . (B) Arrows indicate the direction of changes in dynamical regime as the amount of stochasticity k increases. (C) Bifurcation diagrams under each scenario as a function of k . Solid and dashed lines represent stable and unstable equilibria, respectively.

312 to be proportional to deterministic payoffs (Wang et al., 2025):

$$\begin{pmatrix} \sigma_R & \sigma_S \\ \sigma_T & \sigma_P \end{pmatrix} = k \begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (21)$$

313 Following previous case studies, we consider four deterministic payoff structures (Fig-
 314 ure 4A): Prisoner’s Dilemma (PD), Hawk-Dove Game (HD), Stag-Hunt Game (SH)
 315 and Harmony Game (HG). Across all these examples, stochasticity eventually pushes
 316 the system inside the mutual invasion regime (Figure 4B). Notably, changes in dy-
 317 namical regimes predicted by SCT accurately capture bifurcations occurring at the
 318 boundaries (Figure 4C): at sufficiently large k , both boundary equilibria are unstable
 319 in all four games. Notably, for the SH game, the system begins from the priority
 320 effect regime and enters the competitive exclusion regime before eventually entering
 321 the mutual invasion regime (Figure 4B). Consequently, $x = 1$ (pure cooperation)
 322 becomes unstable first, and then $x = 0$ (pure defection) also becomes unstable.

323 This example also illustrates an important limitation of the original MCT as well
 324 as its extension, SCT. Since MCT and SCT solely rely on boundary conditions, they
 325 cannot predict the presence of multiple internal equilibria or their stability. In the SH
 326 example, there is an unstable internal equilibrium across all ranges of k we consider
 327 (Figure 4C). When the system enters the competitive-exclusion regime, an additional
 328 stable internal equilibrium appears, allowing coexistence. Furthermore, when mutual
 329 invasion is allowed, two stable coexistence equilibria are present. Therefore, while
 330 MCT and SCT accurately capture the behavior at the boundary via invasibility,
 331 they do not fully characterize the interior dynamics or all conditions under which
 332 coexistence is possible.

333 Evolution of cooperation in multi-player, nonlinear 334 public goodsgames

335 Previous examples focused on evolutionary games based on two-player interactions
 336 but real world behaviors, especially cooperation, depend on interactions between
 337 multiple players. In particular, cooperators can contribute to shared public goods,
 338 from which other individuals can benefit. As our final study, we consider a N-player
 339 public-goods game to investigate the effects of multiplayer interaction on strategy
 340 competition. In doing so, we also consider the effect of nonlinear public goods and
 341 ask when nonlinearity can stabilize competition.

342 To model the evolution of cooperation under multiplayer interactions, we begin
 343 by assuming that every individual receives the same amount of benefit $\beta(i)$ which
 344 increases monotonically with the number of cooperators i , and that cooperation
 345 incurs a constant cost $c > 0$. Assuming that each individual interacts with $N - 1$
 346 other selected individuals, the average payoff for cooperators $\pi_C(x)$ and defectors
 347 $\pi_D(x)$ depends on the probability $f_j(x)$ of finding j cooperators in the group, given

348 x proportion of cooperators in the population (Archetti and Scheuring, 2012):

$$\pi_C(x) = \sum_{j=0}^{N-1} f_j(x) \beta(j+1) - c \quad (22)$$

$$\pi_D(x) = \sum_{j=0}^{N-1} f_j(x) \beta(j) \quad (23)$$

349 Assuming random, homogeneous mixing in a large population, the probability $f_j(x)$
350 follows a binomial distribution:

$$\pi_C(x) = \sum_{j=0}^{N-1} \binom{N-1}{j} x^j (1-x)^{N-1-j} \beta(j+1) - c \quad (24)$$

$$\pi_D(x) = \sum_{j=0}^{N-1} \binom{N-1}{j} x^j (1-x)^{N-1-j} \beta(j) \quad (25)$$

351 The resulting replicator equation corresponds to:

$$\frac{dx}{dt} = x(1-x)G(x), \quad (26)$$

352 where $G(x) = \pi_C(x) - \pi_D(x)$ represents the payoff difference.

353 While this game consists of interactions between multiple individuals, from the
354 perspective of SCT, there are still two strategies competing. Thus, we can compute
355 niche overlap ρ and fitness difference κ_C/κ_D using standard SCT, which in turn can
356 be simplified in terms of payoff differences at the boundary:

$$\rho = \exp\left(\frac{\pi_C(1) + \pi_D(0) - \pi_C(0) - \pi_D(1)}{2}\right) = \exp\left(\frac{G(1) - G(0)}{2}\right) \quad (27)$$

$$\frac{\kappa_D}{\kappa_C} = \exp\left(\frac{\pi_D(0) + \pi_D(1) - \pi_C(1) - \pi_C(0)}{2}\right) = \exp\left(-\frac{G(1) + G(0)}{2}\right). \quad (28)$$

357 Here, the payoff values at the boundary correspond to:

$$G(0) = \beta(1) - \beta(0) - c \quad (29)$$

$$G(1) = \beta(N) - \beta(N-1) - c \quad (30)$$

358 Finally, mutual invasion at the boundary requires:

$$\rho < \frac{\kappa_D}{\kappa_C} < \frac{1}{\rho}. \quad (31)$$

359 This expression can be equivalently written as:

$$G(1) < 0 < G(0) \iff \beta(N) - \beta(N-1) < c < \beta(1) - \beta(0). \quad (32)$$

This formulation shows that mutual invasion is possible only when the marginal benefit generated by the first cooperator exceeds that generated by the N th cooperator, with the cost of cooperation lying between these two marginal benefits.

We can now draw three conclusions from this observation. First, when the benefit of public good $\beta(i)$ increases linearly with i (thus $\beta(1) - \beta(0) = \beta(N) - \beta(N - 1)$), then there is complete niche overlap because $G(1) = G(0)$. In this case, cooperators competitively exclude defectors when the marginal benefit $\beta(1) - \beta(0)$ is greater than cost c , and defectors competitively exclude cooperators otherwise. Stable coexistence becomes impossible. Second, when the benefit of public good $\beta(i)$ is a concave, increasing function of i (i.e., $d\beta/di > 0$ and $d^2\beta/di^2 < 0$), we have $\beta(1) - \beta(0) > \beta(N) - \beta(N - 1)$. This means that nonlinearity can generate niche difference and allow coexistence, provided that the cost c lies between the inequality. This generalizes the result of Gore et al. (2009), showing that the effect of concave nonlinearity on coexistence is robust to multiplayer extension (see Supplementary Text S1 and Figure S1 for detailed analyses). Third, when the benefit of public good $\beta(i)$ is a convex, increasing function of i (i.e., $d\beta/di > 0$ and $d^2\beta/di^2 > 0$), we have $\beta(1) - \beta(0) < \beta(N) - \beta(N - 1)$. In this case, nonlinearity can allow bistability of two boundary conditions and generate priority effect.

We note that this multi-player game can also permit

Discussion

Standard evolutionary game theory and replicator equations provide powerful tools for predicting the outcome of strategy competition, including strategic dominance, coexistence, and bistability (Motro, 1991; Doebeli et al., 2004; Hauert and Doebeli, 2004; Doebeli and Hauert, 2005; Gore et al., 2009; Souza et al., 2009; Archetti and Scheuring, 2011; Steidinger and Bever, 2014; Tarnita and Traulsen, 2025). However, these outcomes are typically understood by analyzing payoff differences within individual games, which makes it difficult to compare coexistence mechanisms across models. Building on Chesson's Modern Coexistence Theory from community ecology (Chesson, 2000), we developed a strategic coexistence theory (SCT) that allows us to decompose strategy competition in terms of stabilizing niche differences and fitness differences. This decomposition provides a unifying framework for comparing different games from the perspective of coexistence theory, allowing us to identify stabilizing and equalizing mechanisms for strategy competition.

To demonstrate the utility of SCT, we considered three case studies. First, the analysis of classic two-strategy games showed that SCT recovers the original classification for these classic games, highlighting the similarities between evolutionary games and community ecology (Hauert, 2002; Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025). Then, we compared different mechanisms for the evolution of cooperation, which showed that each mechanism promotes cooperation via different dynamical routes (Nowak, 2006). Kin selection, network reciprocity, and

group selection primarily affect the fitness difference between cooperation and defection, whereas direct and indirect reciprocity can destabilize competition and generate priority effects. Finally, our analysis of microbial cooperation showed that coexistence between cooperators and defectors requires more than reducing the fitness cost of cooperation: nonlinear benefits can generate stabilizing niche differences while also reducing fitness asymmetry (Gore et al., 2009). Taken together, these examples show that SCT can provide new insights into ecological underpinnings of strategy competition.

There are several limitations to SCT. First, similar to MCT from community ecology, SCT only allows for comparisons of pairwise strategies and therefore cannot determine the outcomes of games with more than two strategies (Chesson, 2000). Second, the exact values of strategic niche and fitness differences are sensitive to payoff scaling. SCT therefore supports quantitative comparisons only when models share a justified payoff normalization. Without such normalization, comparisons are restricted to qualitative changes in dynamical regimes, directions, and boundary crossings. Third, we focused on classic replicator equations, but Tarnita and Traulsen (2025) recently pointed out that these classic formulations implicitly ignore differences in intrinsic growth rates. In such cases, replicator equations may fail to predict the outcome of competition (Tarnita and Traulsen, 2025), and a full Lotka-Volterra model is needed to describe the abundances, rather than frequencies, of strategies. Thus, SCT inherits the same assumptions as the classic replicator framework: it is most appropriate when strategy competition can be described in terms of relative frequencies alone (Tarnita and Traulsen, 2025). When strategies differ intrinsically in growth, MCT may be required instead to tease apart mechanisms of strategy coexistence. Finally, we have focused on simple examples to illustrate the utility of SCT, but many games involve additional complexities, including population structure (Nowak et al., 2010), stochasticity (Wang et al., 2025; Bodin et al., 2026), and explicit ecological feedback (Tarnita and Traulsen, 2025). Extending SCT to these settings will be necessary to determine how broadly stabilizing and equalizing mechanisms can be compared across evolutionary games.

In conclusion, SCT provides a bridge between evolutionary game theory and community ecology, allowing strategy competition to be understood from the perspective of coexistence theory. Therefore, rather than replacing existing EGT or replicator-dynamics frameworks, SCT provides a complementary tool for teasing apart the coexistence mechanisms underlying evolutionary games. More broadly, our study builds on recent efforts to extend MCT beyond species competition (Park et al., 2024, 2026), providing a foundation for building unifying coexistence theory for competing entities, including species, pathogen strains, and behavioral strategies.

438 **Acknowledgement**

439 We thank Jonathan Levine for the helpful discussion. S.W.P. was supported by
440 the New Faculty Startup Fund from Seoul National University, the National Re-
441 search Foundation of Korea (NRF) grant funded by the Korea government (MSIT)
442 (RS-2026-25474574), and the Global-LAMP Program of the National Research Foun-
443 dation of Korea (NRF) grant funded by the Ministry of Education (No. RS-2023-
444 00301976).

445 **Competing interests**

446 The authors declare no competing interest.

447 **Data availability**

448 All data and code are stored in a publicly available GitHub repository (<https://github.com/parklab-snu/coexistence-strategy>).
449

450 Supplementary Text

451 S1 Derivation of the strategic coexistence theory

452 To derive strategic coexistence theory (SCT), we begin with replicator equations:

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S1})$$

453 where x_i represents the frequency of strategy i , π_i represents the payoff of strategy
 454 i , and $\bar{\pi}$ represents the average payoff. Then, the mutual invasion condition for two
 455 competing strategies i and j can be written as

$$\pi_j(\mathbf{e}_j) < \pi_i(\mathbf{e}_j), \quad (\text{S2})$$

$$\pi_i(\mathbf{e}_i) < \pi_j(\mathbf{e}_i). \quad (\text{S3})$$

456 These inequalities will be preserved when we exponentiate the payoffs, $W_i(\mathbf{x}) =$
 457 $\exp(\pi_i(\mathbf{x}))$:

$$W_j(\mathbf{e}_j) < W_i(\mathbf{e}_j), \quad (\text{S4})$$

$$W_i(\mathbf{e}_i) < W_j(\mathbf{e}_i). \quad (\text{S5})$$

458 Rearranging, we obtain the following inequality:

$$\frac{W_i(\mathbf{e}_i)}{W_j(\mathbf{e}_i)} < 1 < \frac{W_i(\mathbf{e}_j)}{W_j(\mathbf{e}_j)}. \quad (\text{S6})$$

459 By multiplying $\sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}$ on both sides, we obtain:

$$\sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}} < \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}} < \sqrt{\frac{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}} \quad (\text{S7})$$

460 Thus, by defining niche overlap ρ and fitness difference κ_j/κ_i as follows,

$$\rho = \sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}}, \quad (\text{S8})$$

$$\frac{\kappa_j}{\kappa_i} = \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}, \quad (\text{S9})$$

461 we obtain the following inequality that defines conditions for mutual invasion and
 462 long-term coexistence of two competing strategies:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (\text{S10})$$

463 S2 Niche and fitness difference payoff scaling

464 Consider a classic replicator equation:

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S11})$$

465 where x_i represents the frequency of strategy i , π_i represents the payoff of strategy
466 i , and $\bar{\pi}$ represents the average payoff. In this case, multiplying the payoffs by a
467 constant $s > 0$ only changes time and does not affect the overall dynamics:

$$\frac{dx_i}{dt} = x_i(\pi'_i(\mathbf{x}) - \bar{\pi}'(\mathbf{x})) \quad (\text{S12})$$

$$= sx_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S13})$$

468 where $\pi'_i(\mathbf{x}) = s\pi_i(\mathbf{x})$ is the scaled payoff. Then, the multiplicative payoff corresponds
469 to $W'_i(\mathbf{x}) = \exp(s\pi_i) = W_i(\mathbf{x})^s$.

470 If we try to calculate niche overlap and fitness difference based on the scaled
471 payoffs, we get:

$$\rho' = \sqrt{\frac{W'_i(\mathbf{e}_i)W'_j(\mathbf{e}_j)}{W'_j(\mathbf{e}_i)W'_i(\mathbf{e}_j)}}, \quad (\text{S14})$$

$$= \left(\sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}} \right)^s, \quad (\text{S15})$$

$$= \rho^s \quad (\text{S16})$$

$$\frac{\kappa'_j}{\kappa'_i} = \sqrt{\frac{W'_j(\mathbf{e}_i)W'_j(\mathbf{e}_j)}{W'_i(\mathbf{e}_i)W'_i(\mathbf{e}_j)}}, \quad (\text{S17})$$

$$= \left(\sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}} \right)^s, \quad (\text{S18})$$

$$= \left(\frac{\kappa_j}{\kappa_i} \right)^s \quad (\text{S19})$$

472 This implies that the exact values of niche and fitness difference are sensitive to
473 payoff scaling and are not intrinsic properties of a replicator system.

474 Nonetheless, the inequality for mutual invasion condition still holds under payoff
475 scaling:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho} \iff \rho^s < \left(\frac{\kappa_j}{\kappa_i} \right)^s < \frac{1}{\rho^s}. \quad (\text{S20})$$

476 Furthermore, boundary conditions for complete niche overlap ($\rho = \rho' = 1$) and com-
477 plete niche difference ($\rho = \rho' = 0$) are preserved. Moreover, the direction of changes
478 in fitness and niche difference is also preserved under payoff scaling. Specifically, for

parameter θ , we have:

$$\frac{d\rho'}{d\theta} = s\rho^{s-1} \frac{d\rho}{d\theta} \quad (\text{S21})$$

$$\frac{d}{d\theta} \left(\frac{\kappa'_j}{\kappa'_i} \right) = s \left(\frac{\kappa_j}{\kappa_i} \right)^{s-1} \frac{d}{d\theta} \left(\frac{\kappa_j}{\kappa_i} \right). \quad (\text{S22})$$

Since ρ and κ_j/κ_i are always positive, directional sensitivities of ρ and κ_j/κ_i to parameter θ are preserved. Similarly, directional sensitivities of $1 - \rho$ to θ are preserved:

$$\frac{d(1 - \rho')}{d\theta} = s\rho^{s-1} \frac{d(1 - \rho)}{d\theta} \quad (\text{S23})$$

Therefore, we focus on comparisons of qualitative outcomes from SCT and refrain from making any comparisons of magnitude of niche and fitness difference values.

S3 Applications to classic two-strategy games

Consider a two-strategy game described by the following payoff matrix:

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (\text{S24})$$

There are four classic games that we can define from this simple payoff matrix: prisoner's dilemma ($T > R > P > S$), hawk-dove game ($T > R, S > P$), stag-hunt game ($R > T, P > S$), and harmony game ($R > T, S > P$). Here, we show that each game corresponds to a distinct region in the niche-fitness difference space under SCT.

To do so, we first begin by deriving expressions for niche overlap ρ and fitness difference κ_j/κ_i . Writing x_1 to denote the frequency of strategy 1, the expected payoff for each strategy corresponds to:

$$\pi_1(\mathbf{x}) = Rx_1 + S(1 - x_1), \quad (\text{S25})$$

$$\pi_2(\mathbf{x}) = Tx_1 + P(1 - x_1). \quad (\text{S26})$$

Then, multiplicative payoffs for each strategy when rare can be written as:

$$W_1(1) = \exp(R) \quad (\text{S27})$$

$$W_1(0) = \exp(S) \quad (\text{S28})$$

$$W_2(1) = \exp(T) \quad (\text{S29})$$

$$W_2(0) = \exp(P) \quad (\text{S30})$$

Therefore, we have:

$$\rho = \exp \left(\frac{R + P - S - T}{2} \right), \quad (\text{S31})$$

$$\frac{\kappa_2}{\kappa_1} = \exp \left(\frac{T + P - R - S}{2} \right). \quad (\text{S32})$$

496 As before, SCT allows us to predict the mutual invasion of two strategy using the
 497 following inequality:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (\text{S33})$$

498 Then, simple algebra reveals that each condition in mutual invasion inequality
 499 can be equivalently described by the payoff inequality:

$$\rho < \frac{\kappa_2}{\kappa_1} \iff R < T \quad (\text{S34})$$

$$\rho > \frac{\kappa_2}{\kappa_1} \iff R > T \quad (\text{S35})$$

$$\frac{\kappa_2}{\kappa_1} < \frac{1}{\rho} \iff P < S \quad (\text{S36})$$

$$\frac{\kappa_2}{\kappa_1} > \frac{1}{\rho} \iff P > S \quad (\text{S37})$$

500 These inequalities then allow us to classify outcomes of two-strategy games from the
 501 perspective of SCT. Note that the inequality between R and P does not affect the
 502 outcome.

503 S4 Applications to mechanisms for the evolution of coopera- 504 tion

505 We consider five mechanisms for the evolution of cooperation, all of which can be
 506 represented as a simple 2×2 payoff matrix (Nowak, 2006). Here, we go through each
 507 mechanism and derive measures for niche and fitness differences.

508 **Prisoner's dilemma.** Here, we consider a special case of Prisoner's dilemma
 509 which can be described by the following payoff matrix:

$$\begin{pmatrix} b - c & -c \\ b & 0 \end{pmatrix}, \quad (\text{S38})$$

510 where b represents the benefit for the recipient and c represents cost for the donor.
 511 In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0 \quad (\text{S39})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(c). \quad (\text{S40})$$

512 Therefore, as long as the cost is greater than zero, prisoner's dilemma always results
 513 in the competitive exclusion of cooperation.

514 **Kin selection.** Kin selection can be described by the following payoff matrix:

$$\begin{pmatrix} (b - c)(1 + r) & br - c \\ b - rc & 0 \end{pmatrix}, \quad (\text{S41})$$

where r represents the genetic relatedness. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0 \quad (\text{S42})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(c - br). \quad (\text{S43})$$

Therefore, defection will exclude cooperation when $c > br$. Equivalently, cooperation will be favored when $r > c/b$, which recovers the classic result.

Direct reciprocity. Direct reciprocity can be described by the following payoff matrix:

$$\begin{pmatrix} (b-c)/(1-w) & -c \\ b & 0 \end{pmatrix}, \quad (\text{S44})$$

where w represents the probability of next round. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 1 - \exp\left(\frac{(b-c)/(1-w) + c - b}{2}\right) \quad (\text{S45})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{b + c - (b-c)/(1-w)}{2}\right). \quad (\text{S46})$$

Therefore, defection will exclude cooperation when

$$\rho < \frac{\kappa_2}{\kappa_1} \quad (\text{S47})$$

$$\iff \exp\left(\frac{(b-c)/(1-w) + c - b}{2}\right) < \exp\left(\frac{b + c - (b-c)/(1-w)}{2}\right) \quad (\text{S48})$$

$$\iff bw < c \quad (\text{S49})$$

However, note that the following inequality holds whenever $c > 0$:

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1}. \quad (\text{S50})$$

Therefore, when $w > c/b$, we will have

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1} < \rho, \quad (\text{S51})$$

which implies priority effect (equivalently, bistability). Even as $w \rightarrow 1^-$, this inequality holds and therefore, cooperation cannot competitively exclude defection.

Indirect reciprocity. Indirect reciprocity can be described by the following payoff matrix:

$$\begin{pmatrix} b-c & -c(1-q) \\ b(1-q) & 0 \end{pmatrix}, \quad (\text{S52})$$

530 where q represents social acquaintanceship. In this case, the niche and fitness differ-
 531 ences correspond to:

$$1 - \rho = 1 - \exp\left(\frac{(b - c)q}{2}\right), \quad (\text{S53})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{2c - (b + c)q}{2}\right). \quad (\text{S54})$$

532 In this case, the condition for the dominance of defection (and therefore competitive
 533 exclusion of cooperation) can be written as:

$$\rho < \frac{\kappa_2}{\kappa_1} \quad (\text{S55})$$

$$\iff bq < c \quad (\text{S56})$$

534 Moreover, the following inequality always holds as long as $c > 0$.

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1}, \quad (\text{S57})$$

535 which implies that coexistence between cooperation and defection is not possible.
 536 Therefore, when $q < c/b$, then defection will competitively exclude cooperation.
 537 Otherwise, the system will enter the priority-effect regime.

538 Interestingly, when $q \rightarrow 1$, we have

$$\rho = \exp\left(\frac{b - c}{2}\right) \quad (\text{S58})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{c - b}{2}\right) \quad (\text{S59})$$

$$\frac{1}{\rho} = \exp\left(\frac{c - b}{2}\right) \quad (\text{S60})$$

$$(\text{S61})$$

539 In other words, the following inequality will hold:

$$\frac{1}{\rho} = \frac{\kappa_2}{\kappa_1} < \rho \quad (\text{S62})$$

540 **Network reciprocity.** Network reciprocity can be described by the following
 541 payoff matrix:

$$\begin{pmatrix} b - c & H - c \\ b - H & 0 \end{pmatrix}, \quad (\text{S63})$$

542 where $H = [(b - c)k - 2c]/[(k + 1)(k - 2)]$ and k represents the number of neighbors.
 543 In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0, \quad (\text{S64})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(-H + c), \quad (\text{S65})$$

544 which implies that there is complete niche overlap and that fitness difference decreases
 545 with H . Since H is a monotonically decreasing function of k when $k > 2$ and $b/c \geq 2$,
 546 fitness difference will increase with k . In particular, cooperation will competitively
 547 exclude defection when $H > c$, meaning that:

$$b/c > k. \quad (\text{S66})$$

548 **Group selection.** Group selection can be described by the following payoff
 549 matrix:

$$\begin{pmatrix} (b-c)(m+n) & (b-c)m - cn \\ bn & 0 \end{pmatrix}, \quad (\text{S67})$$

550 where m represents the number of groups and n represents the group size. In this
 551 case, the niche and fitness differences correspond to:

$$1 - \rho = 0, \quad (\text{S68})$$

$$\frac{\kappa_2}{\kappa_1} = \exp((c-b)m + cn), \quad (\text{S69})$$

552 which implies that there is complete niche overlap. The fitness difference will increase
 553 with n but decrease with m , provided that $b > c$. In particular, cooperation will
 554 competitively exclude defection under the following condition:

$$1 + \frac{n}{m} < \frac{b}{c}. \quad (\text{S70})$$

555 **S5 Applications to evolutionary games with envi-** 556 **ronmental stochasticity**

557 Consider a competition of two strategies under stochastic payoffs:

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix} + \frac{\zeta}{\sqrt{s}} \begin{pmatrix} \sigma_R & \sigma_S \\ \sigma_T & \sigma_P \end{pmatrix}, \quad (\text{S71})$$

558 Assuming weak selection $s \ll 1$, the evolution of two competing strategies in a large
 559 population can be described by the following equation:

$$\frac{dx}{dt} = sx(1-x) \left(\pi_C - \pi_D + \left(\frac{1}{2} - x \right) (\sigma_C - \sigma_D)^2 \right), \quad (\text{S72})$$

560 To simplify, we write

$$\frac{dx}{dt} = sx(1-x)G(x), \quad (\text{S73})$$

561 where

$$G(x) = \left(\pi_C - \pi_D + \left(\frac{1}{2} - x \right) (\sigma_C - \sigma_D)^2 \right). \quad (\text{S74})$$

Thus, the sign of $G(x)$ at boundary conditions determines whether a strategy can invade when the competing strategy is at equilibrium.

Note that we cannot immediately apply the original SCT presented in the main text here because $G(x)$ does not uniquely separate into two payoff measures. Instead, we can still utilize the invasion growth rate when rare and follow the same procedure. In particular, the mutual invasion condition at the monomorphic boundary can be written as:

$$0 < G(0) \tag{S75}$$

$$0 < -G(1) \tag{S76}$$

Exponentiating both, we can combine them into a single inequality:

$$\exp(G(1)) < 1 < \exp(G(0)) \tag{S77}$$

Dividing both sides by $\exp((G(1) + G(0))/2)$, we get:

$$\exp\left(\frac{G(1) - G(0)}{2}\right) < \exp\left(-\frac{G(1) + G(0)}{2}\right) < \exp\left(\frac{G(0) - G(1)}{2}\right), \tag{S78}$$

which gives us expressions for niche overlap and fitness difference:

$$\rho = \exp\left(\frac{G(1) - G(0)}{2}\right) \tag{S79}$$

$$\frac{\kappa_D}{\kappa_C} = \exp\left(-\frac{G(1) + G(0)}{2}\right) \tag{S80}$$

Then, by substituting expressions for $G(0)$ and $G(1)$, we obtain the following expressions for niche and fitness difference:

$$1 - \rho = 1 - \exp\left(\frac{R + P - S - T}{2} - \frac{(\sigma_R - \sigma_T)^2 + (\sigma_S - \sigma_P)^2}{4}\right) \tag{S81}$$

$$\frac{\kappa_D}{\kappa_C} = \exp\left(\frac{T + P - R - S}{2} + \frac{(\sigma_R - \sigma_T)^2 - (\sigma_S - \sigma_P)^2}{4}\right) \tag{S82}$$

Now, to study the interior behavior of the system, we assume that the amount of stochasticity is proportional to deterministic payoffs (Wang et al., 2025):

$$\begin{pmatrix} \sigma_R & \sigma_S \\ \sigma_T & \sigma_P \end{pmatrix} = k \begin{pmatrix} R & S \\ T & P \end{pmatrix}. \tag{S83}$$

576 Then, the expression for $G(x)$ can be written as:

$$G(x) = ((R - T)x + (S - P)(1 - x)) \times \quad (\text{S84})$$

$$\left(1 + k^2 \left(\frac{1}{2} - x\right) ((R - T)x + (S - P)(1 - x))\right) \quad (\text{S85})$$

$$= ((R + P - T - S)x + (S - P)) \times \quad (\text{S86})$$

$$\left(1 + k^2 \left(\frac{1}{2} - x\right) ((R + P - T - S)x + (S - P))\right) \quad (\text{S87})$$

$$= k^2(R + P - T - S)^2 (x - x^*) \left(K + \left(\frac{1}{2} - x\right) (x - x^*)\right), \quad (\text{S88})$$

577 where $x^* = (P - S)/(R + P - T - S)$ and $K = 1/(k^2(R + P - T - S))$ for $k > 0$
 578 and $R + P - T - S \neq 0$. Therefore, this system has an internal equilibrium $x = x^*$
 579 when $0 < (P - S)/(R + P - T - S) < 1$.

580 The system can also have up to two additional equilibria when $K + \left(\frac{1}{2} - x\right) (x -$
 581 $x^*) = 0$ has positive, real roots and when these roots fall between 0 and 1. In
 582 particular, the quadratic equation can be written as:

$$x^2 - \left(\frac{1}{2} + x^*\right) x + \frac{x^*}{2} - K = 0. \quad (\text{S89})$$

583 Thus, the two solutions of the quadratic equation correspond to:

$$x = \frac{x^* + \frac{1}{2} \pm \sqrt{\left(\frac{1}{2} - x^*\right)^2 + 4K}}{2}. \quad (\text{S90})$$

584 These solutions are real when:

$$\left(\frac{1}{2} - x^*\right)^2 + 4K > 0. \quad (\text{S91})$$

585 When these solutions fall between 0 and 1, they correspond to additional internal
 586 equilibria.

587 **S6 Microbial cooperation and public goods**

588 Here, we consider a simple, phenomenological model of competition between cooper-
 589 ating and cheating yeast in a cooperative sucrose-metabolism system, where invertase
 590 production by cooperators allows cheaters to utilize glucose, a public good released
 591 through sucrose hydrolysis (Carlson and Botstein, 1982; Dickinson, 1998; Gore et al.,
 592 2009). Specifically, invertase production by cooperators incurs a cost c and gener-
 593 ates a total benefit of 1 that is captured privately with efficiency ϵ . Assuming a

linear relationship between glucose availability and microbial growth, the payoffs for cooperators π_C and cheaters (defectors) π_D can then be written as

$$\pi_C(f) = \epsilon + f(1 - \epsilon) - c, \quad (\text{S92})$$

$$\pi_D(f) = f(1 - \epsilon), \quad (\text{S93})$$

where the use of glucose depends on the fraction of cooperators f . The dynamics of cooperators and defectors are described by the following replicator equation:

$$\frac{df}{dt} = f(1 - f) [\pi_C(f) - \pi_D(f)] \quad (\text{S94})$$

$$= f(1 - f)(\epsilon - c) \quad (\text{S95})$$

For this linear growth model, the payoff difference is fixed to $\epsilon - c$ independent of f (Figure S1A). Therefore, cooperation is favored when efficiency is greater than the cost of cooperation, $\epsilon > c$, whereas defection is favored when the cost is greater than efficiency, $c > \epsilon$ (Gore et al., 2009).

Our framework allows us to reinterpret this result from the perspective of coexistence theory. In particular, the two strategies have no niche difference, corresponding to complete niche overlap, $\rho = 1$. In contrast, the fitness difference is determined by the balance between private efficiency and the cost of cooperation:

$$\frac{\kappa_D}{\kappa_C} = \exp(c - \epsilon). \quad (\text{S96})$$

Therefore, this model always predicts competitive exclusion, except in the neutral case $c = \epsilon$. The outcome is determined entirely by the fitness difference between cooperators and defectors, rather than by stabilizing niche differences.

In real biological systems, the effect of glucose on microbial growth follows a nonlinear relationship:

$$\pi_C(f) = [\epsilon + f(1 - \epsilon)]^\alpha - c, \quad (\text{S97})$$

$$\pi_D(f) = [f(1 - \epsilon)]^\alpha, \quad (\text{S98})$$

where $\alpha = 0.15$ reflects the experimental relationship between glucose and microbial growth (Gore et al., 2009). Then, the dynamics of cooperators and defectors are described by the following replicator equation:

$$\frac{df}{dt} = f(1 - f) [\{\epsilon + f(1 - \epsilon)\}^\alpha - c - \{f(1 - \epsilon)\}^\alpha]. \quad (\text{S99})$$

Under this nonlinear assumption, the payoff difference decreases with the cooperator fraction f , which can promote the coexistence of cooperators and defectors (Figure S1A): cooperators have a competitive advantage when rare, whereas defectors have a competitive advantage when cooperators are abundant.

618 From a coexistence-theoretic perspective, this means that nonlinear growth can
619 generate sufficient stabilizing niche difference to overcome the fitness difference. In
620 particular, the niche and fitness differences for this model correspond to:

$$1 - \rho = 1 - \exp\left(\frac{1 - \epsilon^\alpha - (1 - \epsilon)^\alpha}{2}\right), \quad (\text{S100})$$

$$\frac{\kappa_D}{\kappa_C} = \exp\left(c + \frac{(1 - \epsilon)^\alpha - 1 - \epsilon^\alpha}{2}\right). \quad (\text{S101})$$

621 These expressions clarify the separate roles of cost, efficiency, and nonlinearity. For
622 example, the cost of cooperation does not affect niche difference and only affects
623 fitness difference. Instead, niche difference is generated by the concave relationship
624 between glucose availability and microbial growth, which increases from 0 to $1 - \exp(-1/2)$
625 as α goes from 1 to 0 (Figure S1B,C). In contrast, as $\alpha \rightarrow 0^+$, the fitness
626 difference approaches

$$\log(\kappa_D/\kappa_C) \rightarrow c - \frac{1}{2}, \quad (\text{S102})$$

627 which no longer depends on the private capture efficiency ϵ (Figure S1C). Thus,
628 strong concavity not only generates stabilizing niche differences, but also acts as an
629 equalizing mechanism (Figure S1C). Such stabilizing mechanisms are key ingredients
630 for coexistence of cooperators and defectors.

Supplementary Figure




Figure S1: **Effects of nonlinear microbial growth on coexistence of cooperation and defection.** (A) Payoff differences between cooperation and defection for linear and nonlinear models of microbial growth. In panel A, we assumed $\epsilon = 0.4$, $c = 0.3$, and $\alpha = 0.15$ for illustrative purposes. (B) Effect of nonlinearity α on niche difference $1 - \rho$. Each line assumes a different value of efficiency ϵ . Note that lines for $\epsilon = 0.6$ and $\epsilon = 0.4$ are nearly overlapping and are indistinguishable. (C) Effect of nonlinearity α on both niche difference $1 - \rho$ and fitness difference $\log(\kappa_D/\kappa_C)$. The gray shaded area represents the coexistence regime. The square represents the condition when $\alpha \rightarrow 0^+$. The circles represent the condition when $\alpha = 0.15$, which correspond to the experimental relationship between glucose and microbial growth (Gore et al., 2009). In panels B and C, we assumed $c = 0.3$, while allowing ϵ and α to vary.

References

- Adler, P. B., M. Detto, S. P. Ellner, T. L. Gibbs, Z. J. Gold, S. J. Leonard, J. I. Levine, A. E. Schiffer, C. Song, M. Stemkovski, et al. (2026). What have we learned from empirical applications of modern coexistence theory? *Ecology letters* 29(6), e70404.
- Adler, P. B., J. HilleRisLambers, and J. M. Levine (2007). A niche for neutrality. *Ecology letters* 10(2), 95–104.
- Allesina, S. (2026). Global stability of ecological and evolutionary dynamics via equivalence. *Proceedings of the National Academy of Sciences* 123(13), e2534915123.
- Apicella, C. L. and J. B. Silk (2019). The evolution of human cooperation. *Current Biology* 29(11), R447–R450.
- Archetti, M. and I. Scheuring (2011). Coexistence of cooperation and defection in public goods games. *Evolution* 65(4), 1140–1148.
- Archetti, M. and I. Scheuring (2012). Game theory of public goods in one-shot social dilemmas without assortment. *Journal of theoretical biology* 299, 9–20.
- Axelrod, R. and W. D. Hamilton (1981). The evolution of cooperation. *science* 211(4489), 1390–1396.
- Bodin, F., G. Wang, and J. B. Plotkin (2026). Eco-evolutionary games in noisy environments. *bioRxiv*, 2026–05.
- Carlson, M. and D. Botstein (1982). Two differentially regulated mRNAs with different 5' ends encode secreted and intracellular forms of yeast invertase. *Cell* 28(1), 145–154.
- Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual review of Ecology and Systematics* 31(1), 343–366.
- Clutton-Brock, T. (2009). Cooperation between non-kin in animal societies. *Nature* 462(7269), 51–57.
- Cooney, D. B. (2020). Analysis of multilevel replicator dynamics for general two-strategy social dilemma. *Bulletin of Mathematical Biology* 82(6), 66.
- Cordero, O. X., L.-A. Ventouras, E. F. DeLong, and M. F. Polz (2012). Public good dynamics drive evolution of iron acquisition strategies in natural bacterioplankton populations. *Proceedings of the National Academy of Sciences* 109(49), 20059–20064.

- 665 Cressman, R. (2003). *Evolutionary dynamics and extensive form games*, Volume 5.
666 MIT Press.
- 667 Cressman, R. and Y. Tao (2014). The replicator equation and other game dynamics.
668 *Proceedings of the National Academy of Sciences* 111(supplement_3), 10810–10817.
- 669 Dickinson, J. R. (1998). *Metabolism and molecular physiology of Saccharomyces*
670 *cerevisiae*. Taylor & Francis.
- 671 Doebeli, M. and C. Hauert (2005). Models of cooperation based on the Prisoner’s
672 Dilemma and the Snowdrift game. *Ecology letters* 8(7), 748–766.
- 673 Doebeli, M., C. Hauert, and T. Killingback (2004). The evolutionary origin of coop-
674 erators and defectors. *science* 306(5697), 859–862.
- 675 Fletcher, J. A. and M. Doebeli (2008). A simple and general explanation for the evolu-
676 tion of altruism. *Proceedings of the Royal Society B: Biological Sciences* 276(1654),
677 13.
- 678 Godoy, O., N. J. Kraft, and J. M. Levine (2014). Phylogenetic relatedness and the
679 determinants of competitive outcomes. *Ecology letters* 17(7), 836–844.
- 680 Godoy, O. and J. M. Levine (2014). Phenology effects on invasion success: insights
681 from coupling field experiments to coexistence theory. *Ecology* 95(3), 726–736.
- 682 Gokhale, C. S. and A. Traulsen (2025). Higher-order equivalence of Lotka-Volterra
683 and replicator dynamics. *bioRxiv*, 2025–03.
- 684 Gore, J., H. Youk, and A. Van Oudenaarden (2009). Snowdrift game dynamics and
685 facultative cheating in yeast. *Nature* 459(7244), 253–256.
- 686 Hamilton, W. D. (1964). The genetical evolution of social behaviour. II. *Journal of*
687 *theoretical biology* 7(1), 17–52.
- 688 Hauert, C. (2002). Effects of space in 2×2 games. *International Journal of Bifur-*
689 *cation and Chaos* 12(07), 1531–1548.
- 690 Hauert, C. and M. Doebeli (2004). Spatial structure often inhibits the evolution of
691 cooperation in the snowdrift game. *Nature* 428(6983), 643–646.
- 692 Hilbe, C., Š. Šimsa, K. Chatterjee, and M. A. Nowak (2018). Evolution of cooperation
693 in stochastic games. *Nature* 559(7713), 246–249.
- 694 Hofbauer, J. (1981). On the occurrence of limit cycles in the volterra-lotka equation.
695 *Nonlinear Analysis: Theory, Methods & Applications* 5(9), 1003–1007.
- 696 Hofbauer, J. and K. Sigmund (1998). *Evolutionary games and population dynamics*.
697 Cambridge university press.

- Kingma, S. A., P. Santema, M. Taborsky, and J. Komdeur (2014). Group augmentation and the evolution of cooperation. *Trends in ecology & evolution* 29(8), 476–484.
- Kocher, S. D., R. Mallarino, B. E. Rubin, D. W. Yu, H. E. Hoekstra, and N. E. Pierce (2018). The genetic basis of a social polymorphism in halictid bees. *Nature Communications* 9(1), 4338.
- Kraft, N. J., O. Godoy, and J. M. Levine (2015). Plant functional traits and the multidimensional nature of species coexistence. *Proceedings of the National Academy of Sciences* 112(3), 797–802.
- Lee, I. P. A., O. T. Eldakar, J. P. Gogarten, and C. P. Andam (2022). Bacterial cooperation through horizontal gene transfer. *Trends in Ecology & Evolution* 37(3), 223–232.
- Lewontin, R. C. (1961). Evolution and the theory of games. *Journal of Theoretical Biology* 1(3), 382–403.
- Lieberman, E., C. Hauert, and M. A. Nowak (2005). Evolutionary dynamics on graphs. *Nature* 433(7023), 312–316.
- Lin, H., D. Wang, Q. Wang, J. Mao, Y. Bai, and J. Qu (2024). Interspecific competition prevents the proliferation of social cheaters in an unstructured environment. *The ISME Journal* 18(1), wrad038.
- Michel-Mata, S., M. Kawakatsu, J. Sartini, T. A. Kessinger, J. B. Plotkin, and C. E. Tarnita (2024). The evolution of private reputations in information-abundant landscapes. *Nature* 634(8035), 883–889.
- Motro, U. (1991). Co-operation and defection: playing the field and the ESS. *Journal of Theoretical Biology* 151(2), 145–154.
- Nowak, M. A. (2006). Five rules for the evolution of cooperation. *science* 314(5805), 1560–1563.
- Nowak, M. A. and R. M. May (1992). Evolutionary games and spatial chaos. *nature* 359(6398), 826–829.
- Nowak, M. A. and K. Sigmund (1998). Evolution of indirect reciprocity by image scoring. *Nature* 393(6685), 573–577.
- Nowak, M. A. and K. Sigmund (2005). Evolution of indirect reciprocity. *Nature* 437(7063), 1291–1298.
- Nowak, M. A., C. E. Tarnita, and T. Antal (2010). Evolutionary dynamics in structured populations. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365(1537), 19.

- 733 Page, K. M. and M. A. Nowak (2002). Unifying evolutionary dynamics. *Journal of*
734 *theoretical biology* 219(1), 93–98.
- 735 Palmer, J. D. and K. R. Foster (2022). Bacterial species rarely work together. *Sci-*
736 *ence* 376(6593), 581–582.
- 737 Park, S. W., S. Cobey, C. J. E. Metcalf, J. M. Levine, and B. T. Grenfell (2024).
738 Predicting pathogen mutual invasibility and co-circulation. *Science* 386(6718),
739 175–179.
- 740 Park, S. W., J. Levine, and B. Grenfell (2026). Unifying coexistence theory for
741 ecological communities and pathogen strain competition. *bioRxiv*, 2026–05.
- 742 Pollak, S., M. Gralka, Y. Sato, J. Schwartzman, L. Lu, and O. X. Cordero (2021).
743 Public good exploitation in natural bacterioplankton communities. *Science Ad-*
744 *vances* 7(31), eabi4717.
- 745 Rakoff-Nahoum, S., K. R. Foster, and L. E. Comstock (2016). The evolution of
746 cooperation within the gut microbiota. *Nature* 533(7602), 255–259.
- 747 Schuster, P. and K. Sigmund (1983). Replicator dynamics. *Journal of theoretical*
748 *biology* 100(3), 533–538.
- 749 Sieben, A. J., J. R. Mihaljevic, and L. G. Shoemaker (2022). Quantifying mechanisms
750 of coexistence in disease ecology. *Ecology* 103(12), e3819.
- 751 Slobodkin, L. B. (1964). The strategy of evolution. *American scientist* 52(3), 342–
752 357.
- 753 Smith, J. M. and G. R. Price (1973). The logic of animal conflict. *Nature* 246(5427),
754 15–18.
- 755 Souza, M. O., J. M. Pacheco, and F. C. Santos (2009). Evolution of cooperation
756 under N-person snowdrift games. *Journal of Theoretical Biology* 260(4), 581–588.
- 757 Steidinger, B. S. and J. D. Bever (2014). The coexistence of hosts with different
758 abilities to discriminate against cheater partners: an evolutionary game-theory
759 approach. *The American Naturalist* 183(6), 762–770.
- 760 Szabó, G. and G. Fath (2007). Evolutionary games on graphs. *Physics reports* 446(4-
761 6), 97–216.
- 762 Tarnita, C. E. and A. Traulsen (2025). Reconciling ecology and evolutionary game
763 theory or “When not to think cooperation”. *Proceedings of the National Academy*
764 *of Sciences* 122(14), e2413847122.
- 765 Taylor, P. D. and L. B. Jonker (1978). Evolutionary stable strategies and game
766 dynamics. *Mathematical biosciences* 40(1-2), 145–156.

- 767 Traulsen, A. and M. A. Nowak (2006). Evolution of cooperation by multilevel selec-
768 tion. *Proceedings of the National Academy of Sciences* 103(29), 10952–10955.
- 769 Trivers, R. L. (1971). The evolution of reciprocal altruism. *The Quarterly review of*
770 *biology* 46(1), 35–57.
- 771 Wang, G., Q. Su, L. Wang, and J. B. Plotkin (2025). Evolution of social behaviors
772 in noisy environments. *arXiv*.
- 773 Wedekind, C. and M. Milinski (2000). Cooperation through image scoring in humans.
774 *Science* 288(5467), 850–852.
- 775 West, S. A., G. A. Cooper, M. B. Ghou, and A. S. Griffin (2021). Ten recent insights
776 for our understanding of cooperation. *Nature ecology & evolution* 5(4), 419–430.
- 777 West, S. A., A. S. Griffin, and A. Gardner (2007). Evolutionary explanations for
778 cooperation. *Current biology* 17(16), R661–R672.