

Strategic coexistence theory for evolutionary games

Sang Woo Park^{1,2,*}

1 School of Biological Sciences, Seoul National University, Seoul, Korea

2 Institute for Data Innovation in Science, Seoul National University, Seoul, Korea

*Corresponding author: sangwoopark@snu.ac.kr

Abstract

Evolutionary game theory and ecological coexistence theory both seek to predict the outcome of competition between biological entities, be they strategies or species, but the two fields have relied on largely separate approaches. Replicator equations provide a foundation for predicting the outcome of strategic competition, yet they do not explicitly decompose the mechanisms underlying these outcomes. Inspired by modern coexistence theory from community ecology, we develop Strategic Coexistence Theory (SCT), which translates payoff-based invasion conditions into strategic niche differences and fitness differences between any two competing strategies described by classic replicator dynamics. SCT recovers the classic classification of two-strategy games, distinguishing competitive exclusion, coexistence, and priority effects within a shared niche–fitness difference space. Applying SCT to five mechanisms for the evolution of cooperation further reveals that these mechanisms promote cooperation through distinct dynamical routes: kin selection, network reciprocity, and group selection primarily reduce fitness differences, whereas direct and indirect reciprocity destabilize competition and generate priority effects. Finally, applying SCT to microbial public-goods competition shows that nonlinear microbial growth can both stabilize and equalize competition between cooperators and defectors, allowing coexistence. Together, these results show that SCT provides a common language for comparing coexistence mechanisms across evolutionary games.

Significance statement

Evolutionary game theory provides a powerful framework for predicting the outcome of competition between strategies, including the dominance of one strategy, coexistence of multiple strategies, and bistability. However, a common framework for comparing how different mechanisms promote strategy coexistence remains elusive, even though stable polymorphisms, such as the coexistence of cooperators and defectors, are widely observed across biological systems. To address this gap, we developed Strategic Coexistence Theory (SCT), which extends Modern Coexistence Theory from community ecology to evolutionary games. Applying SCT to classic games, mechanisms of cooperation, and microbial public-goods competition shows

that seemingly similar evolutionary outcomes can arise through distinct mechanisms. This approach links ecological theory with replicator dynamics, providing a complementary tool for analyzing evolutionary games.

Introduction

Understanding the mechanisms that promote the evolutionary stability of competing strategies is a fundamental aim in evolutionary biology (Lewontin, 1961; Slobodkin, 1964; Smith and Price, 1973; Taylor and Jonker, 1978). To this end, evolutionary game theory (EGT) and replicator dynamics have provided key foundations, allowing us to predict the outcome of strategic competition through payoff inequalities, invasion criteria, and stability analysis (Schuster and Sigmund, 1983; Hofbauer and Sigmund, 1998; Cressman, 2003; Cressman and Tao, 2014). Recently, there has been growing interest in characterizing similarities between replicator dynamics and the dynamics of competing species (Gokhale and Traulsen, 2025; Tarnita and Traulsen, 2025; Allesina, 2026), building on the classical mathematical equivalence between replicator equations and generalized Lotka–Volterra models (Hofbauer, 1981; Page and Nowak, 2002). In particular, Tarnita and Traulsen (2025) recently argued that a closer synthesis between ecology and evolutionary game theory can provide new insights into ecological and evolutionary assembly, competition, and coexistence. This renewed perspective between ecological and evolutionary dynamics suggests that evolutionary games may be interpreted not only as payoff-based competition but also as problems of strategic coexistence.

Interestingly, even a simple two-strategy game can yield heterogeneous outcomes that mirror outcomes in community ecology (Hauert, 2002; Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025): the dominance of a single strategy (prisoner’s dilemma and harmony games), coexistence (hawk–dove games), and bistability (stag-hunt games). Such differences in conditions that promote evolutionary stability and dominance of a single strategy and those that permit strategy coexistence indicate that EGT can be understood from the perspective of coexistence theory (Motro, 1991; Doebeli et al., 2004; Hauert and Doebeli, 2004; Doebeli and Hauert, 2005; Gore et al., 2009; Souza et al., 2009; Archetti and Scheuring, 2011; Steidinger and Bever, 2014; Tarnita and Traulsen, 2025). In fact, stable polymorphisms of evolutionary strategies, such as coexistence of cooperators and defectors, are observed in microbial and other biological systems (Gore et al., 2009; Cordero et al., 2012; Kocher et al., 2018; Pollak et al., 2021; Lin et al., 2024), but coexistence-theoretic perspective has often remained implicit. Recasting game dynamics as a coexistence problem immediately raises the following question: when two strategies coexist, does the corresponding mechanism stabilize or equalize competition (Chesson, 2000)? While EGT allows us to predict whether two strategies can coexist or not, a common framework for comparing coexistence mechanisms between competing strategies remains elusive.

Modern coexistence theory (MCT) from community ecology presents a particularly useful framework for quantifying coexistence mechanisms (Chesson, 2000; Adler et al., 2007; Godoy et al., 2014; Godoy and Levine, 2014; Kraft et al., 2015; Adler et al., 2026). MCT predicts that coexistence of two competing entities, be they species or strategies, requires stabilizing niche differences to overcome average fitness differences (Chesson, 2000; Adler et al., 2007; Kraft et al., 2015). In community ecology, stabilizing niche differences represent differences in factors that limit the growth of competing species, causing each species to limit itself more than it limits its competitor (Chesson, 2000). In EGT, a strategy’s niche can be interpreted as strategic environments defined by the composition of competing strategies and the resulting interactions (Taylor and Jonker, 1978; Cressman and Tao, 2014; Tarnita and Traulsen, 2025). Thus, stabilizing niche differences arise when each strategy is favored by a different strategic environment, such that the spread of each strategy creates conditions that limit its own relative growth and favor the recovery of its competitor. In contrast, fitness differences represent differences in the ability of each entity to compete under the limiting environments created by its competitor (Chesson, 2000). This differs from payoff differences in EGT, which depend on the underlying strategy frequency and therefore combine both stabilizing effects caused by changes in the strategic environment and residual fitness differences that favor one strategy over another (Taylor and Jonker, 1978; Cressman and Tao, 2014; Tarnita and Traulsen, 2025).

So far, MCT has been primarily utilized for understanding species coexistence, but recent work showed that MCT can be extended beyond species competition, especially for studying pathogen strain competition, allowing one to directly quantify niche and fitness differences of competing entities, be they species or strains (Sieben et al., 2022; Park et al., 2024, 2026). Inspired by this effort, we present Strategic Coexistence Theory (SCT) for predicting coexistence and exclusion of competing strategies in EGT by decomposing invasion growth rates into strategic niche and fitness differences. We begin by introducing SCT, which can be applied to any generic system of EGT described by replicator equations. We show that SCT can recover outcomes from classic, two-strategy games, providing a reinterpretation of classic results from coexistence-theoretic perspective. We then apply SCT to comparing five mechanisms for the evolution of cooperation, revealing that direct and indirect reciprocity destabilize competition between cooperators and defectors, thus pushing the system towards priority effect regime. Finally, we consider a case study of microbial cooperation, where nonlinear microbial growth can both stabilize and equalize competition between cooperation and defection, allowing coexistence of strategies. Together, these examples show that SCT provides a common, ecological language for comparing the mechanisms underlying strategic evolution.

Strategic coexistence theory

Here, we present a brief derivation of strategic coexistence theory (SCT) (Figure 1). To do so, we begin with a classic replicator equation, which allows us to model changes in the frequency of competing strategies based on their payoffs (Schuster and Sigmund, 1983; Hofbauer and Sigmund, 1998; Cressman, 2003; Cressman and Tao, 2014):

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (1)$$

where x_i represents the frequency of strategy i , $\pi_i(\mathbf{x})$ represents the payoff of strategy i for a given strategy distribution $\mathbf{x}$, and $\bar{\pi}(\mathbf{x}) = \sum_i \pi_i(\mathbf{x})x_i$ represents the average payoff. Since strategy frequencies must sum to 1 (i.e., $\sum_i x_i = 1$), a single-strategy equilibrium dominated by strategy i can be described by a unit vector $\mathbf{e}_i$, where all elements are equal to zero except for the i -th element. We use these single-strategy equilibria to define niche and fitness differences between two competing strategies i and j .

Because payoffs in many games can be negative, payoff ratios can be difficult to interpret. Thus, we work on a positive multiplicative scale by exponentiating the payoff terms:

$$W_i(\mathbf{x}) = \exp(\pi_i(\mathbf{x})). \quad (2)$$

This exponential transformation is convenient because it preserves the payoff ordering and does not alter the invasion conditions. Therefore, this allows us to compare strategic environments through ratios, as in modern coexistence theory. For convenience, we refer to $W_i(\mathbf{x})$ as the multiplicative payoff of strategy i .

First, we define the niche overlap ρ between strategies i and j as the ratio between the geometric mean of each strategy's multiplicative payoff in the strategic environment that it creates when common, $\sqrt{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}$, and the geometric mean of each strategy's multiplicative payoff in the environment generated by its competitor, $\sqrt{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}$. The corresponding niche difference is then given by

$$1 - \rho = 1 - \sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}}. \quad (3)$$

When $1 - \rho > 0$, each strategy performs better, on average, in the environment generated by its competitor than in the environment generated by itself. In other words, intraspecific competition is stronger than interspecific competition. Second, we define the fitness difference as the ratio between the geometric means of the multiplicative payoff of each strategy across the two single-strategy environments:

$$\frac{\kappa_j}{\kappa_i} = \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}. \quad (4)$$

In other words, strategy j has an average competitive advantage over strategy i ($\kappa_j/\kappa_i > 1$) when it performs better, on average, across the environments generated by itself and its competitor.




Figure 1: **A schematic diagram summarizing strategic coexistence theory.** Strategic coexistence theory decomposes pairwise replicator dynamics into stabilizing and equalizing mechanisms. Mutual invasion conditions can be rewritten on a multiplicative payoff scale, allowing us to derive strategic niche difference $1 - \rho$ and fitness difference κ_j/κ_i based on multiplicative payoff ratios. Then, the inequality $\rho < \kappa_j/\kappa_i < 1/\rho$ determines mutual invasion. We note that the exact values of niche and fitness differences are sensitive to payoff scaling, while dynamical outcomes are still preserved.

150 Together, for any two strategies described by the classic replicator equation, mu-
 151 tual invasion requires stabilizing niche differences to overcome average fitness differ-
 152 ences. Specifically, the condition for mutual invasion can be rewritten as (Chesson,
 153 2000):

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (5)$$

154 This inequality applies to any pairwise comparison between two strategies described
 155 by the classic replicator equation because ρ and κ_j/κ_i can be calculated directly from
 156 ratios of multiplicative payoffs. This decomposition not only allows us to understand
 157 how different mechanisms contribute to the coexistence or exclusion of competing
 158 strategies but also provides shared axes for comparing outcomes of different games.
 159 Detailed derivations are provided in Supplementary Text S1.

160 We note that the absolute magnitudes of niche and fitness differences are sen-
 161 sitive to payoff scaling because they are defined on an exponentiated payoff scale.
 162 Therefore, while SCT allows us to compare different games on shared axes, we can-
 163 not compare the exact values of niche and fitness differences across different games.
 164 Nonetheless, as we show in Supplementary Text S2, the qualitative outcome of strate-
 165 gic competition under SCT is preserved because the exponential transformation does
 166 not alter invasion conditions. Therefore, we focus on comparing qualitative outcomes

167 from SCT, including directions of change in niche and fitness difference and resulting
 168 dynamical regimes, and refrain from making any comparisons of magnitude of niche
 169 and fitness difference values.

170 **Classic two-strategy games**

171 As an illustrative example, we begin with classic two-strategy games (Hauert, 2002;
 172 Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025), which can be
 173 described by the following payoff matrix (Figure 2A):

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (6)$$

174 Here, the first and second rows represent strategies 1 and 2, respectively, and the
 175 columns represent the strategy of the interacting opponent. Writing x_1 to denote
 176 the frequency of strategy 1, the expected payoffs are

$$\pi_1(\mathbf{x}) = Rx_1 + S(1 - x_1), \quad (7)$$

$$\pi_2(\mathbf{x}) = Tx_1 + P(1 - x_1). \quad (8)$$

177 Thus, the dynamics can be written as

$$\frac{dx_1}{dt} = x_1(1 - x_1) [(R - T)x_1 + (S - P)(1 - x_1)]. \quad (9)$$

178 The signs of $T - R$ and $S - P$ determine whether each strategy can invade the
 179 other when rare. Specifically, strategy 2 can invade strategy 1 when $T > R$, whereas
 180 strategy 1 can invade strategy 2 when $S > P$. These two invasion conditions partition
 181 the game space into four regimes (Figure 2B): prisoner’s dilemma ($T > R > P >$
 182 S), hawk–dove game ($T > R, S > P$), stag–hunt game ($R > T, P > S$), and
 183 harmony game ($R > T, S > P$). In standard EGT, these outcomes are obtained by
 184 analyzing the stability of equilibrium conditions. For simple games, this analysis is
 185 straightforward.

186 A coexistence-theoretic approach recovers the same classification but provides
 187 a different interpretation (Figure 2C). For a two-strategy game determined by this
 188 payoff matrix, the niche difference and fitness difference are given by the following
 189 set of equations (Supplementary Text S3):

$$1 - \rho = 1 - \exp\left(\frac{R + P - S - T}{2}\right), \quad (10)$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{T + P - R - S}{2}\right). \quad (11)$$

190 As shown above, mutual invasion requires $\rho < \kappa_2/\kappa_1 < 1/\rho$, which is equivalent to
 191 $T > R$ and $S > P$. This corresponds to the hawk–dove game, in which each strategy




Figure 2: **Coexistence-theoretic interpretation of classic two-strategy games.** (A) General payoff matrix for a two-strategy game, where rows represent the focal strategy and columns represent the strategy of the interacting opponent. (B) Classification of four classic games based on the payoff differences $T - R$ and $S - P$. Prisoner’s dilemma and harmony games lead to the dominance of one strategy; hawk–dove games allow coexistence; and stag–hunt games result in bistability. (C) Mapping of the same four games onto niche–fitness difference space. Hawk–dove games correspond to the coexistence regime, where stabilizing niche differences overcome fitness differences. Prisoner’s dilemma and harmony games correspond to competitive-exclusion regimes, whereas stag–hunt games correspond to the priority-effect regime. $R > P$ is assumed throughout, but this does not affect coexistence outcome in any way.

192 can invade its competitor when rare and the two strategies coexist. The remaining
 193 games also have natural interpretations in niche–fitness difference space. Prisoner’s
 194 dilemma and harmony games represent competitive exclusion regimes, where one
 195 strategy has a fitness advantage that exceeds niche differences. In contrast, stag–hunt
 196 games represent a priority-effect regime, where neither strategy can invade the other
 197 and the outcome depends on initial conditions. Therefore, coexistence theory not
 198 only allows us to recover the classic results from two-strategy games but also provides
 199 new interpretations for these results in terms of niche and fitness differences. Building
 200 on this, we next apply this framework to comparing core mechanisms of cooperation.

201 Comparisons of mechanisms for the evolution of co- 202 operation

203 Even though cooperative behavior can be found across a wide range of biological
 204 systems, a simple prisoner’s dilemma shows that cooperation is not an evolutionarily

stable strategy (Axelrod and Hamilton, 1981; May, 1987). Thus, understanding mechanisms that promote the evolution of cooperation has remained a central goal in evolutionary biology (West et al., 2007; Fletcher and Doebeli, 2008; Clutton-Brock, 2009; Kingma et al., 2014; Rakoff-Nahoum et al., 2016; Hilbe et al., 2018; Apicella and Silk, 2019; West et al., 2021; Lee et al., 2022; Palmer and Foster, 2022; Michel-Mata et al., 2024). Here, we compare five classic mechanisms for the evolution of cooperation using SCT (Nowak, 2006) and ask how they differ from the perspective of SCT.

Specifically, we consider the following mechanisms: kin selection, direct reciprocity, indirect reciprocity, network reciprocity, and group selection (Figure 3A). Kin selection favors cooperation when two interacting individuals are genetic relatives, where the genetic relatedness r must be greater than the cost-to-benefit ratio: $r > c/b$ (Hamilton, 1964). Direct reciprocity favors cooperation when current cooperation may promote future cooperation by the same individual through repeated encounters (Trivers, 1971; Axelrod and Hamilton, 1981). In this case, the probability of repeated interaction (i.e., “next round”) must be greater than the cost-to-benefit ratio: $w > c/b$. Indirect reciprocity favors cooperation when current cooperation may promote future cooperation by different individuals through reputation (Nowak and Sigmund, 1998; Wedekind and Milinski, 2000; Nowak and Sigmund, 2005). Thus, the social acquaintanceship (i.e., the probability of knowing someone’s reputation) must be greater than the cost-to-benefit ratio: $q > c/b$. Network reciprocity favors cooperation via clustering of cooperators, requiring the benefit-to-cost ratio to be greater than the average number of neighbors: $b/c > k$ (Nowak and May, 1992; Lieberman et al., 2005). Finally, group selection favors cooperation via multilevel selection, where groups with more cooperators exhibit faster growth at the group level even if defectors reproduce faster at the individual level (Traulsen and Nowak, 2006). This requires $b/c > 1 + (n/m)$, where n is the maximum group size and m is the number of groups. All of these mechanisms can be described by a simple 2×2 payoff matrix (Nowak, 2006), from which the corresponding niche and fitness differences can be computed using Eq. (10) and Eq. (11). Details on niche and fitness difference calculations are presented in Supplementary Text S4.

First, the prisoner’s dilemma provides a baseline scenario, where defection is evolutionarily stable (Figure 3A). From the perspective of SCT, there is complete niche overlap, and therefore defection competitively excludes cooperation (Figure 3B). Building on this, SCT shows that five mechanisms promote evolutionary stability of cooperation via different dynamical routes (Figure 3B). For example, kin selection allows the fitness of cooperation to increase as genetic relatedness r increases (Figure 3B). However, it does not affect the niche difference between cooperation and defection. Similarly, network reciprocity and group selection modulate the fitness of cooperation without causing any changes in strategic niches. In contrast, direct and indirect reciprocity destabilize competition while increasing the fitness of cooperation, thereby pushing the competition outcome towards a priority effect regime. Equivalently, these two mechanisms promote bistability of defection and cooperation.




Figure 3: **Qualitative comparisons of five mechanisms for the evolution of cooperation using SCT.** (A) Payoff matrices for the baseline scenario (prisoner's dilemma) and five mechanisms for the evolution of cooperation (Nowak, 2006). Prisoner's dilemma is characterized by the benefit for the recipient b and the cost for the donor c . Network reciprocity is parameterized based on $H = [(b-c)k - 2c]/[(k+1)(k-2)]$. Parameters for all other mechanisms are described in panel B. (B) Changes in niche and fitness differences driven by each mechanism. Each arrow represents the direction of change as we increase the corresponding parameter, indicated in figure legends. Note that kin selection, network reciprocity, and group selection all result in changes in fitness difference without causing any changes in niche difference. Therefore, we indicate the direction of change for network reciprocity and group selection separately to avoid overlapping arrows. For simplicity, we focus on the effect of number of groups m for group selection; increasing the number of maximum group size n has the opposite effect (Supplementary Text S4). Across all models, we assume $b = 3$ and $c = 1$. Because SCT decompositions depend on positive payoff scaling, exact values of niche and fitness difference (e.g., arrow lengths, slopes, and distances from regime boundaries) should not be compared across mechanisms.

248 SCT further reveals another difference between direct and indirect reciprocity.
 249 In particular, direct reciprocity is unable to generate sufficient fitness difference to
 250 competitively exclude defection even when the probability of next round w reaches
 251 1 (Supplementary Text S4). In contrast, indirect reciprocity is able to generate

sufficient fitness difference to push the system towards the boundary between priority-effect regime and competitive-exclusion regime as social acquaintanceship q reaches 1 (Supplementary Text S4).

We note that none of these mechanisms allow coexistence of cooperation and defection. In many biological systems, such coexistence is often observed (Gore et al., 2009; Cordero et al., 2012; Kocher et al., 2018; Pollak et al., 2021; Lin et al., 2024). Therefore, we move on to the final example, applying SCT to microbial cooperation, where nonlinear benefits and environmental context can promote coexistence of cooperators and defectors.

Microbial cooperation and public goods

Here, we consider a simple, phenomenological model of competition between cooperating and cheating yeast in a cooperative sucrose-metabolism system, where invertase production by cooperators allows cheaters to utilize glucose, a public good released through sucrose hydrolysis (Carlson and Botstein, 1982; Dickinson, 1998; Gore et al., 2009). Specifically, invertase production by cooperators incurs a cost c and generates a total benefit of 1 that is captured privately with efficiency ϵ . Assuming a linear relationship between glucose availability and microbial growth, the payoffs for cooperators π_C and cheaters (defectors) π_D can then be written as

$$\pi_C(f) = \epsilon + f(1 - \epsilon) - c, \quad (12)$$

$$\pi_D(f) = f(1 - \epsilon), \quad (13)$$

where the use of glucose depends on the fraction of cooperators f . The dynamics of cooperators and defectors are described by the following replicator equation:

$$\frac{df}{dt} = f(1 - f) [\pi_C(f) - \pi_D(f)] \quad (14)$$

$$= f(1 - f)(\epsilon - c) \quad (15)$$

For this linear growth model, the payoff difference is fixed to $\epsilon - c$ independent of f (Figure 4A). Therefore, cooperation is favored when efficiency is greater than the cost of cooperation, $\epsilon > c$, whereas defection is favored when the cost is greater than efficiency, $c > \epsilon$ (Gore et al., 2009).

Our framework allows us to reinterpret this result from the perspective of coexistence theory. In particular, the two strategies have no niche difference, corresponding to complete niche overlap, $\rho = 1$. In contrast, the fitness difference is determined by the balance between private efficiency and the cost of cooperation:

$$\frac{\kappa_D}{\kappa_C} = \exp(c - \epsilon). \quad (16)$$

Therefore, this model always predicts competitive exclusion, except in the neutral case $c = \epsilon$. The outcome is determined entirely by the fitness difference between




Figure 4: **Effects of nonlinear microbial growth on coexistence of cooperation and defection.** (A) Payoff differences between cooperation and defection for linear and nonlinear models of microbial growth. In panel A, we assumed $\epsilon = 0.4$, $c = 0.3$, and $\alpha = 0.15$ for illustrative purposes. (B) Effect of nonlinearity α on niche difference $1 - \rho$. Each line assumes a different value of efficiency ϵ . Note that lines for $\epsilon = 0.6$ and $\epsilon = 0.4$ are nearly overlapping and are indistinguishable. (C) Effect of nonlinearity α on both niche difference $1 - \rho$ and fitness difference $\log(\kappa_D/\kappa_C)$. The gray shaded area represents the coexistence regime. The square represents the condition when $\alpha \rightarrow 0^+$. The circles represent the condition when $\alpha = 0.15$, which correspond to the experimental relationship between glucose and microbial growth (Gore et al., 2009). In panels B and C, we assumed $c = 0.3$, while allowing ϵ and α to vary.

cooperators and defectors, rather than by stabilizing niche differences. This example resembles the prisoner's dilemma discussed in the previous example (Figure 3).

In real biological systems, the effect of glucose on microbial growth follows a nonlinear relationship:

$$\pi_C(f) = [\epsilon + f(1 - \epsilon)]^\alpha - c, \quad (17)$$

$$\pi_D(f) = [f(1 - \epsilon)]^\alpha, \quad (18)$$

where $\alpha = 0.15$ reflects the experimental relationship between glucose and microbial growth (Gore et al., 2009). Then, the dynamics of cooperators and defectors are described by the following replicator equation:

$$\frac{df}{dt} = f(1 - f) [\{\epsilon + f(1 - \epsilon)\}^\alpha - c - \{f(1 - \epsilon)\}^\alpha]. \quad (19)$$

Under this nonlinear assumption, the payoff difference decreases with the cooperator fraction f , which can promote the coexistence of cooperators and defectors (Figure 4A): cooperators have a competitive advantage when rare, whereas defectors have a competitive advantage when cooperators are abundant.

From a coexistence-theoretic perspective, this means that nonlinear growth can generate sufficient stabilizing niche difference to overcome the fitness difference. In particular, the niche and fitness differences for this model correspond to:

$$1 - \rho = 1 - \exp\left(\frac{1 - \epsilon^\alpha - (1 - \epsilon)^\alpha}{2}\right), \quad (20)$$

$$\frac{\kappa_D}{\kappa_C} = \exp\left(c + \frac{(1 - \epsilon)^\alpha - 1 - \epsilon^\alpha}{2}\right). \quad (21)$$

These expressions clarify the separate roles of cost, efficiency, and nonlinearity. For example, the cost of cooperation does not affect niche difference and only affects fitness difference. Instead, niche difference is generated by the concave relationship between glucose availability and microbial growth, which increases from 0 to $1 - \exp(-1/2)$ as α goes from 1 to 0 (Figure 4B,C). In contrast, as $\alpha \rightarrow 0^+$, the fitness difference approaches

$$\log(\kappa_D/\kappa_C) \rightarrow c - \frac{1}{2}, \quad (22)$$

which no longer depends on the private capture efficiency ϵ (Figure 4C). Thus, strong concavity not only generates stabilizing niche differences, but also acts as an equalizing mechanism (Figure 4C). Such stabilizing mechanisms are key ingredients for coexistence of cooperators and defectors.

Discussion

Standard evolutionary game theory and replicator equations provide powerful tools for predicting the outcome of strategy competition, including strategic dominance, coexistence, and bistability (Motro, 1991; Doebeli et al., 2004; Hauert and Doebeli, 2004; Doebeli and Hauert, 2005; Gore et al., 2009; Souza et al., 2009; Archetti and Scheuring, 2011; Steidinger and Bever, 2014; Tarnita and Traulsen, 2025). However, these outcomes are typically understood by analyzing payoff differences within individual games, which makes it difficult to compare coexistence mechanisms across models. Building on Chesson's Modern Coexistence Theory from community ecology (Chesson, 2000), we developed a strategic coexistence theory (SCT) that allows us to decompose strategy competition in terms of stabilizing niche differences and fitness differences. This decomposition provides a unifying framework for comparing different games from the perspective of coexistence theory, allowing us to identify stabilizing and equalizing mechanisms for strategy competition.

To demonstrate the utility of SCT, we considered three case studies. First, the analysis of classic two-strategy games showed that SCT recovers the original classification for these classic games, highlighting the similarities between evolutionary games and community ecology (Hauert, 2002; Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025). Then, we compared different mechanisms for the evolution of cooperation, which showed that each mechanism promotes cooperation via

different dynamical routes (Nowak, 2006). Kin selection, network reciprocity, and group selection primarily affect the fitness difference between cooperation and defection, whereas direct and indirect reciprocity can destabilize competition and generate priority effects. Finally, our analysis of microbial cooperation showed that coexistence between cooperators and defectors requires more than reducing the fitness cost of cooperation: nonlinear benefits can generate stabilizing niche differences while also reducing fitness asymmetry (Gore et al., 2009). Taken together, these examples show that SCT can provide new insights into ecological underpinnings of strategy competition.

There are several limitations to SCT. First, similar to MCT from community ecology, SCT only allows for comparisons of pairwise strategies and therefore cannot determine the outcomes of games with more than two strategies (Chesson, 2000). Second, the exact values of strategic niche and fitness differences are sensitive to payoff scaling. SCT therefore supports quantitative comparisons only when models share a justified payoff normalization. Without such normalization, comparisons are restricted to qualitative changes in dynamical regimes, directions, and boundary crossings. Second, we focused on classic replicator equations, but Tarnita and Traulsen (2025) recently pointed out that these classic formulations implicitly ignore differences in intrinsic growth rates. In such cases, replicator equations may fail to predict the outcome of competition (Tarnita and Traulsen, 2025), and a full Lotka-Volterra model is needed to describe the abundances, rather than frequencies, of strategies. Thus, SCT inherits the same assumptions as the classic replicator framework: it is most appropriate when strategy competition can be described in terms of relative frequencies alone (Tarnita and Traulsen, 2025). When strategies differ intrinsically in growth, MCT may be required instead to tease apart mechanisms of strategy coexistence. Finally, we have focused on simple examples to illustrate the utility of SCT, but many games involve additional complexities, including population structure (Nowak et al., 2010), stochasticity (Wang et al., 2025; Bodin et al., 2026), and explicit ecological feedback (Tarnita and Traulsen, 2025). Extending SCT to these settings will be necessary to determine how broadly stabilizing and equalizing mechanisms can be compared across evolutionary games.

In conclusion, SCT provides a bridge between evolutionary game theory and community ecology, allowing strategy competition to be understood from the perspective of coexistence theory. Therefore, rather than replacing existing EGT or replicator-dynamics frameworks, SCT provides a complementary tool for teasing apart the coexistence mechanisms underlying evolutionary games. More broadly, our study builds on recent efforts to extend MCT beyond species competition (Park et al., 2024, 2026), providing a foundation for building unifying coexistence theory for competing entities, including species, pathogen strains, and behavioral strategies.

365 **Acknowledgement**

366 We thank Jonathan Levine for the helpful discussion. S.W.P. was supported by
367 the New Faculty Startup Fund from Seoul National University, the National Re-
368 search Foundation of Korea (NRF) grant funded by the Korea government (MSIT)
369 (RS-2026-25474574), and the Global-LAMP Program of the National Research Foun-
370 dation of Korea (NRF) grant funded by the Ministry of Education (No. RS-2023-
371 00301976).

372 **Competing interests**

373 The authors declare no competing interest.

374 **Data availability**

375 All data and code are stored in a publicly available GitHub repository ([https:](https://github.com/parklab-snu/coexistence-strategy)
376 [//github.com/parklab-snu/coexistence-strategy](https://github.com/parklab-snu/coexistence-strategy)).

Supplementary Text

S1 Derivation of the strategic coexistence theory

To derive strategic coexistence theory (SCT), we begin with replicator equations:

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S1})$$

where x_i represents the frequency of strategy i , π_i represents the payoff of strategy i , and $\bar{\pi}$ represents the average payoff. Then, the mutual invasion condition for two competing strategies i and j can be written as

$$\pi_j(\mathbf{e}_j) < \pi_i(\mathbf{e}_j), \quad (\text{S2})$$

$$\pi_i(\mathbf{e}_i) < \pi_j(\mathbf{e}_i). \quad (\text{S3})$$

These inequalities will be preserved when we exponentiate the payoffs, $W_i(\mathbf{x}) = \exp(\pi_i(\mathbf{x}))$:

$$W_j(\mathbf{e}_j) < W_i(\mathbf{e}_j), \quad (\text{S4})$$

$$W_i(\mathbf{e}_i) < W_j(\mathbf{e}_i). \quad (\text{S5})$$

Rearranging, we obtain the following inequality:

$$\frac{W_i(\mathbf{e}_i)}{W_j(\mathbf{e}_i)} < 1 < \frac{W_i(\mathbf{e}_j)}{W_j(\mathbf{e}_j)}. \quad (\text{S6})$$

By multiplying $\sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}$ on both sides, we obtain:

$$\sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}} < \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}} < \sqrt{\frac{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}} \quad (\text{S7})$$

Thus, by defining niche overlap ρ and fitness difference κ_j/κ_i as follows,

$$\rho = \sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}}, \quad (\text{S8})$$

$$\frac{\kappa_j}{\kappa_i} = \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}, \quad (\text{S9})$$

we obtain the following inequality that defines conditions for mutual invasion and long-term coexistence of two competing strategies:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (\text{S10})$$

390 S2 Niche and fitness difference payoff scaling

391 Consider a classic replicator equation:

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S11})$$

392 where x_i represents the frequency of strategy i , π_i represents the payoff of strategy
 393 i , and $\bar{\pi}$ represents the average payoff. In this case, multiplying the payoffs by a
 394 constant $s > 0$ only changes time and does not affect the overall dynamics:

$$\frac{dx_i}{dt} = x_i(\pi'_i(\mathbf{x}) - \bar{\pi}'(\mathbf{x})) \quad (\text{S12})$$

$$= sx_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S13})$$

395 where $\pi'_i = s\pi$ is the scaled payoff. Then, the multiplicative payoff corresponds to
 396 $W'_i = \exp(s\pi_i) = W_i^s$.

397 If we try to calculate niche overlap and fitness difference based on the scaled
 398 payoffs, we get:

$$\rho' = \sqrt{\frac{W'_i(\mathbf{e}_i)W'_j(\mathbf{e}_j)}{W'_j(\mathbf{e}_i)W'_i(\mathbf{e}_j)}}, \quad (\text{S14})$$

$$= \left(\sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}} \right)^s, \quad (\text{S15})$$

$$= \rho^s \quad (\text{S16})$$

$$\frac{\kappa'_j}{\kappa'_i} = \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}, \quad (\text{S17})$$

$$= \left(\sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}} \right)^s, \quad (\text{S18})$$

$$= \left(\frac{\kappa_j}{\kappa_i} \right)^s \quad (\text{S19})$$

399 This implies that the exact values of niche and fitness difference are sensitive to
 400 payoff scaling and are not intrinsic properties of a replicator system.

401 Nonetheless, the inequality for mutual invasion condition still holds under payoff
 402 scaling:

$$\rho' < \frac{\kappa'_j}{\kappa'_i} < \frac{1}{\rho'}. \quad (\text{S20})$$

403 Furthermore, boundary conditions for complete niche overlap ($\rho = \rho' = 1$) and com-
 404 plete niche difference ($\rho = \rho' = 0$) are preserved. Moreover, the direction of changes

in fitness and niche difference is also preserved under payoff scaling. Specifically, for parameter θ , we have:

$$\frac{d\rho'}{d\theta} = s\rho^{s-1}\frac{d\rho}{d\theta} \quad (\text{S21})$$

$$\frac{d}{d\theta} \left(\frac{\kappa'_j}{\kappa'_i} \right) = s \left(\frac{\kappa_j}{\kappa_i} \right)^{s-1} \frac{d}{d\theta} \left(\frac{\kappa_j}{\kappa_i} \right). \quad (\text{S22})$$

Since ρ and κ_j/κ_i are always positive, directional sensitivities of ρ and κ_j/κ_i to parameter θ are preserved. Similarly, directional sensitivities of $1 - \rho$ to θ are preserved. Therefore, we focus on comparisons of qualitative outcomes from SCT and refrain from making any comparisons of magnitude of niche and fitness difference values.

S3 Applications to classic two-strategy games

Consider a two-strategy game described by the following payoff matrix:

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (\text{S23})$$

There are four classic games that we can define from this simple payoff matrix: prisoner's dilemma ($T > R > P > S$), hawk-dove game ($T > R, S > P$), stag-hunt game ($R > T, P > S$), and harmony game ($R > T, S > P$). Here, we show that each game corresponds to a distinct region in the niche-fitness difference space under SCT.

To do so, we first begin by deriving expressions for niche overlap ρ and fitness difference κ_j/κ_i . Writing x_1 to denote the frequency of strategy 1, the expected payoff for each strategy corresponds to:

$$\pi_1(\mathbf{x}) = Rx_1 + S(1 - x_1), \quad (\text{S24})$$

$$\pi_2(\mathbf{x}) = Tx_1 + P(1 - x_1). \quad (\text{S25})$$

Then, multiplicative payoffs for each strategy when rare can be written as:

$$W_1(1) = \exp(R) \quad (\text{S26})$$

$$W_1(0) = \exp(S) \quad (\text{S27})$$

$$W_2(1) = \exp(T) \quad (\text{S28})$$

$$W_2(0) = \exp(P) \quad (\text{S29})$$

Therefore, we have:

$$\rho = \exp \left(\frac{R + P - S - T}{2} \right), \quad (\text{S30})$$

$$\frac{\kappa_2}{\kappa_1} = \exp \left(\frac{T + P - R - S}{2} \right). \quad (\text{S31})$$

423 As before, SCT allows us to predict the mutual invasion of two strategy using the
 424 following inequality:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (\text{S32})$$

425 Then, simple algebra reveals that each condition in mutual invasion inequality
 426 can be equivalently described by the payoff inequality:

$$\rho < \frac{\kappa_2}{\kappa_1} \iff R < T \quad (\text{S33})$$

$$\rho > \frac{\kappa_2}{\kappa_1} \iff R > T \quad (\text{S34})$$

$$\frac{\kappa_2}{\kappa_1} < \frac{1}{\rho} \iff P < S \quad (\text{S35})$$

$$\frac{\kappa_2}{\kappa_1} > \frac{1}{\rho} \iff P > S \quad (\text{S36})$$

427 These inequalities then allow us to classify outcomes of two-strategy games from the
 428 perspective of SCT. Note that the inequality between R and P does not affect the
 429 outcome.

430 **S4 Applications to mechanisms for the evolution of coopera-** 431 **tion**

432 We consider five mechanisms for the evolution of cooperation, all of which can be
 433 represented as a simple 2×2 payoff matrix (Nowak, 2006). Here, we go through each
 434 mechanism and derive measures for niche and fitness differences.

435 **Prisoner's dilemma.** Here, we consider a special case of Prisoner's dilemma
 436 which can be described by the following payoff matrix:

$$\begin{pmatrix} b - c & -c \\ b & 0 \end{pmatrix}, \quad (\text{S37})$$

437 where b represents the benefit for the recipient and c represents cost for the donor.
 438 In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0 \quad (\text{S38})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(c). \quad (\text{S39})$$

439 Therefore, as long as the cost is greater than zero, prisoner's dilemma always results
 440 in the competitive exclusion of cooperation.

441 **Kin selection.** Kin selection can be described by the following payoff matrix:

$$\begin{pmatrix} (b - c)(1 + r) & br - c \\ b - rc & 0 \end{pmatrix}, \quad (\text{S40})$$

where r represents the genetic relatedness. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0 \quad (\text{S41})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(c - br). \quad (\text{S42})$$

Therefore, defection will exclude cooperation when $c > br$. Equivalently, cooperation will be favored when $r > c/b$, which recovers the classic result.

Direct reciprocity. Direct reciprocity can be described by the following payoff matrix:

$$\begin{pmatrix} (b-c)/(1-w) & -c \\ b & 0 \end{pmatrix}, \quad (\text{S43})$$

where w represents the probability of next round. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 1 - \exp\left(\frac{(b-c)/(1-w) + c - b}{2}\right) \quad (\text{S44})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{b + c - (b-c)/(1-w)}{2}\right). \quad (\text{S45})$$

Therefore, defection will exclude cooperation when

$$\rho < \frac{\kappa_2}{\kappa_1} \quad (\text{S46})$$

$$\iff \exp\left(\frac{(b-c)/(1-w) + c - b}{2}\right) < \exp\left(\frac{b + c - (b-c)/(1-w)}{2}\right) \quad (\text{S47})$$

$$\iff bw < c \quad (\text{S48})$$

However, note that the following inequality holds whenever $c > 0$:

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1}. \quad (\text{S49})$$

Therefore, when $w > c/b$, we will have

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1} < \rho, \quad (\text{S50})$$

which implies priority effect (equivalently, bistability). Even as $w \rightarrow 1^-$, this inequality holds and therefore, cooperation cannot competitively exclude defection.

Indirect reciprocity. Indirect reciprocity can be described by the following payoff matrix:

$$\begin{pmatrix} b-c & -c(1-q) \\ b(1-q) & 0 \end{pmatrix}, \quad (\text{S51})$$

457 where q represents social acquaintanceship. In this case, the niche and fitness differ-
 458 ences correspond to:

$$1 - \rho = 1 - \exp\left(\frac{(b - c)q}{2}\right), \quad (\text{S52})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{2c - (b + c)q}{2}\right). \quad (\text{S53})$$

459 In this case, the condition for the dominance of defection (and therefore competitive
 460 exclusion of cooperation) can be written as:

$$\rho < \frac{\kappa_2}{\kappa_1} \quad (\text{S54})$$

$$\iff bq < c \quad (\text{S55})$$

461 Moreover, the following inequality always holds as long as $c > 0$.

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1}, \quad (\text{S56})$$

462 which implies that coexistence between cooperation and defection is not possible.
 463 Therefore, when $q < c/b$, then defection will competitively exclude cooperation.
 464 Otherwise, the system will enter the priority-effect regime.

465 Interestingly, when $q \rightarrow 1$, we have

$$\rho = \exp\left(\frac{b - c}{2}\right) \quad (\text{S57})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{c - b}{2}\right) \quad (\text{S58})$$

$$\frac{1}{\rho} = \exp\left(\frac{c - b}{2}\right) \quad (\text{S59})$$

$$(\text{S60})$$

466 In other words, the following inequality will hold:

$$\frac{1}{\rho} = \frac{\kappa_2}{\kappa_1} < \rho \quad (\text{S61})$$

467 **Network reciprocity.** Network reciprocity can be described by the following
 468 payoff matrix:

$$\begin{pmatrix} b - c & H - c \\ b - H & 0 \end{pmatrix}, \quad (\text{S62})$$

469 where $H = [(b - c)k - 2c]/[(k + 1)(k - 2)]$ and k represents the number of neighbors.
 470 In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0, \quad (\text{S63})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(-H + c), \quad (\text{S64})$$

471 which implies that there is complete niche overlap and that fitness difference decreases
 472 with H . Since H is a monotonically decreasing function of k when $k > 2$ and $b/c \geq 2$,
 473 fitness difference will increase with k . In particular, cooperation will competitively
 474 exclude defection when $H > c$, meaning that:

$$b/c > k. \quad (\text{S65})$$

475 **Group selection.** Group selection can be described by the following payoff
 476 matrix:

$$\begin{pmatrix} (b-c)(m+n) & (b-c)m - cn \\ bn & 0 \end{pmatrix}, \quad (\text{S66})$$

477 where m represents the number of groups and n represents the group size. In this
 478 case, the niche and fitness differences correspond to:

$$1 - \rho = 0, \quad (\text{S67})$$

$$\frac{\kappa_2}{\kappa_1} = \exp((c-b)m + cn), \quad (\text{S68})$$

479 which implies that there is complete niche overlap. The fitness difference will increase
 480 with n but decrease with m , provided that $b > c$. In particular, cooperation will
 481 competitively exclude defection under the following condition:

$$1 + \frac{n}{m} < \frac{b}{c}. \quad (\text{S69})$$

482 References

- 483 Adler, P. B., M. Detto, S. P. Ellner, T. L. Gibbs, Z. J. Gold, S. J. Leonard, J. I.
 484 Levine, A. E. Schiffer, C. Song, M. Stemkovski, et al. (2026). What have we learned
 485 from empirical applications of modern coexistence theory? *Ecology letters* 29(6),
 486 e70404.
- 487 Adler, P. B., J. HilleRisLambers, and J. M. Levine (2007). A niche for neutrality.
 488 *Ecology letters* 10(2), 95–104.
- 489 Allesina, S. (2026). Global stability of ecological and evolutionary dynamics
 490 via equivalence. *Proceedings of the National Academy of Sciences* 123(13),
 491 e2534915123.
- 492 Apicella, C. L. and J. B. Silk (2019). The evolution of human cooperation. *Current*
 493 *Biology* 29(11), R447–R450.
- 494 Archetti, M. and I. Scheuring (2011). Coexistence of cooperation and defection in
 495 public goods games. *Evolution* 65(4), 1140–1148.
- 496 Axelrod, R. and W. D. Hamilton (1981). The evolution of cooperation. *sci-*
 497 *ence* 211(4489), 1390–1396.

498 Bodin, F., G. Wang, and J. B. Plotkin (2026). Eco-evolutionary games in noisy
499 environments. *bioRxiv*, 2026–05.

500 Carlson, M. and D. Botstein (1982). Two differentially regulated mRNAs with differ-
501 ent 5' ends encode secreted and intracellular forms of yeast invertase. *Cell* 28(1),
502 145–154.

503 Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual review*
504 *of Ecology and Systematics* 31(1), 343–366.

505 Clutton-Brock, T. (2009). Cooperation between non-kin in animal societies. *Na-*
506 *ture* 462(7269), 51–57.

507 Cooney, D. B. (2020). Analysis of multilevel replicator dynamics for general two-
508 strategy social dilemma. *Bulletin of Mathematical Biology* 82(6), 66.

509 Cordero, O. X., L.-A. Ventouras, E. F. DeLong, and M. F. Polz (2012). Public good
510 dynamics drive evolution of iron acquisition strategies in natural bacterioplankton
511 populations. *Proceedings of the National Academy of Sciences* 109(49), 20059–
512 20064.

513 Cressman, R. (2003). *Evolutionary dynamics and extensive form games*, Volume 5.
514 MIT Press.

515 Cressman, R. and Y. Tao (2014). The replicator equation and other game dynamics.
516 *Proceedings of the National Academy of Sciences* 111(supplement_3), 10810–10817.

517 Dickinson, J. R. (1998). *Metabolism and molecular physiology of Saccharomyces*
518 *cerevisiae*. Taylor & Francis.

519 Doebeli, M. and C. Hauert (2005). Models of cooperation based on the Prisoner’s
520 Dilemma and the Snowdrift game. *Ecology letters* 8(7), 748–766.

521 Doebeli, M., C. Hauert, and T. Killingback (2004). The evolutionary origin of coop-
522 erators and defectors. *science* 306(5697), 859–862.

523 Fletcher, J. A. and M. Doebeli (2008). A simple and general explanation for the evolu-
524 tion of altruism. *Proceedings of the Royal Society B: Biological Sciences* 276(1654),
525 13.

526 Godoy, O., N. J. Kraft, and J. M. Levine (2014). Phylogenetic relatedness and the
527 determinants of competitive outcomes. *Ecology letters* 17(7), 836–844.

528 Godoy, O. and J. M. Levine (2014). Phenology effects on invasion success: insights
529 from coupling field experiments to coexistence theory. *Ecology* 95(3), 726–736.

530 Gokhale, C. S. and A. Traulsen (2025). Higher-order equivalence of Lotka-Volterra
531 and replicator dynamics. *bioRxiv*, 2025–03.

532 Gore, J., H. Youk, and A. Van Oudenaarden (2009). Snowdrift game dynamics and
533 facultative cheating in yeast. *Nature* 459(7244), 253–256.

534 Hamilton, W. D. (1964). The genetical evolution of social behaviour. II. *Journal of*
535 *theoretical biology* 7(1), 17–52.

536 Hauert, C. (2002). Effects of space in 2×2 games. *International Journal of Bifur-*
537 *cation and Chaos* 12(07), 1531–1548.

538 Hauert, C. and M. Doebeli (2004). Spatial structure often inhibits the evolution of
539 cooperation in the snowdrift game. *Nature* 428(6983), 643–646.

540 Hilbe, C., Š. Šimsa, K. Chatterjee, and M. A. Nowak (2018). Evolution of cooperation
541 in stochastic games. *Nature* 559(7713), 246–249.

542 Hofbauer, J. (1981). On the occurrence of limit cycles in the volterra-lotka equation.
543 *Nonlinear Analysis: Theory, Methods & Applications* 5(9), 1003–1007.

544 Hofbauer, J. and K. Sigmund (1998). *Evolutionary games and population dynamics*.
545 Cambridge university press.

546 Kingma, S. A., P. Santema, M. Taborsky, and J. Komdeur (2014). Group augmen-
547 tation and the evolution of cooperation. *Trends in ecology & evolution* 29(8),
548 476–484.

549 Kocher, S. D., R. Mallarino, B. E. Rubin, D. W. Yu, H. E. Hoekstra, and N. E.
550 Pierce (2018). The genetic basis of a social polymorphism in halictid bees. *Nature*
551 *Communications* 9(1), 4338.

552 Kraft, N. J., O. Godoy, and J. M. Levine (2015). Plant functional traits and the mul-
553 tidimensional nature of species coexistence. *Proceedings of the National Academy*
554 *of Sciences* 112(3), 797–802.

555 Lee, I. P. A., O. T. Eldakar, J. P. Gogarten, and C. P. Andam (2022). Bacterial co-
556 operation through horizontal gene transfer. *Trends in Ecology & Evolution* 37(3),
557 223–232.

558 Lewontin, R. C. (1961). Evolution and the theory of games. *Journal of Theoretical*
559 *Biology* 1(3), 382–403.

560 Lieberman, E., C. Hauert, and M. A. Nowak (2005). Evolutionary dynamics on
561 graphs. *Nature* 433(7023), 312–316.

562 Lin, H., D. Wang, Q. Wang, J. Mao, Y. Bai, and J. Qu (2024). Interspecific compe-
563 tition prevents the proliferation of social cheaters in an unstructured environment.
564 *The ISME Journal* 18(1), wrad038.

565 May, R. M. (1987). More evolution of cooperation. *Nature* 327(6117), 15–17.

Michel-Mata, S., M. Kawakatsu, J. Sartini, T. A. Kessinger, J. B. Plotkin, and C. E. Tarnita (2024). The evolution of private reputations in information-abundant landscapes. *Nature* 634(8035), 883–889.

Motro, U. (1991). Co-operation and defection: playing the field and the ESS. *Journal of Theoretical Biology* 151(2), 145–154.

Nowak, M. A. (2006). Five rules for the evolution of cooperation. *science* 314(5805), 1560–1563.

Nowak, M. A. and R. M. May (1992). Evolutionary games and spatial chaos. *nature* 359(6398), 826–829.

Nowak, M. A. and K. Sigmund (1998). Evolution of indirect reciprocity by image scoring. *Nature* 393(6685), 573–577.

Nowak, M. A. and K. Sigmund (2005). Evolution of indirect reciprocity. *Nature* 437(7063), 1291–1298.

Nowak, M. A., C. E. Tarnita, and T. Antal (2010). Evolutionary dynamics in structured populations. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365(1537), 19.

Page, K. M. and M. A. Nowak (2002). Unifying evolutionary dynamics. *Journal of theoretical biology* 219(1), 93–98.

Palmer, J. D. and K. R. Foster (2022). Bacterial species rarely work together. *Science* 376(6593), 581–582.

Park, S. W., S. Cobey, C. J. E. Metcalf, J. M. Levine, and B. T. Grenfell (2024). Predicting pathogen mutual invasibility and co-circulation. *Science* 386(6718), 175–179.

Park, S. W., J. Levine, and B. Grenfell (2026). Unifying coexistence theory for ecological communities and pathogen strain competition. *bioRxiv*, 2026–05.

Pollak, S., M. Gralka, Y. Sato, J. Schwartzman, L. Lu, and O. X. Cordero (2021). Public good exploitation in natural bacterioplankton communities. *Science Advances* 7(31), eabi4717.

Rakoff-Nahoum, S., K. R. Foster, and L. E. Comstock (2016). The evolution of cooperation within the gut microbiota. *Nature* 533(7602), 255–259.

Schuster, P. and K. Sigmund (1983). Replicator dynamics. *Journal of theoretical biology* 100(3), 533–538.

Sieben, A. J., J. R. Mihaljevic, and L. G. Shoemaker (2022). Quantifying mechanisms of coexistence in disease ecology. *Ecology* 103(12), e3819.

600 Slobodkin, L. B. (1964). The strategy of evolution. *American scientist* 52(3), 342–
601 357.

602 Smith, J. M. and G. R. Price (1973). The logic of animal conflict. *Nature* 246(5427),
603 15–18.

604 Souza, M. O., J. M. Pacheco, and F. C. Santos (2009). Evolution of cooperation
605 under N-person snowdrift games. *Journal of Theoretical Biology* 260(4), 581–588.

606 Steidinger, B. S. and J. D. Bever (2014). The coexistence of hosts with different
607 abilities to discriminate against cheater partners: an evolutionary game-theory
608 approach. *The American Naturalist* 183(6), 762–770.

609 Szabó, G. and G. Fath (2007). Evolutionary games on graphs. *Physics reports* 446(4–
610 6), 97–216.

611 Tarnita, C. E. and A. Traulsen (2025). Reconciling ecology and evolutionary game
612 theory or “When not to think cooperation”. *Proceedings of the National Academy
613 of Sciences* 122(14), e2413847122.

614 Taylor, P. D. and L. B. Jonker (1978). Evolutionary stable strategies and game
615 dynamics. *Mathematical biosciences* 40(1-2), 145–156.

616 Traulsen, A. and M. A. Nowak (2006). Evolution of cooperation by multilevel selec-
617 tion. *Proceedings of the National Academy of Sciences* 103(29), 10952–10955.

618 Trivers, R. L. (1971). The evolution of reciprocal altruism. *The Quarterly review of
619 biology* 46(1), 35–57.

620 Wang, G., Q. Su, L. Wang, and J. B. Plotkin (2025). Evolution of social behaviors
621 in noisy environments. *arXiv*.

622 Wedekind, C. and M. Milinski (2000). Cooperation through image scoring in humans.
623 *Science* 288(5467), 850–852.

624 West, S. A., G. A. Cooper, M. B. Ghoul, and A. S. Griffin (2021). Ten recent insights
625 for our understanding of cooperation. *Nature ecology & evolution* 5(4), 419–430.

626 West, S. A., A. S. Griffin, and A. Gardner (2007). Evolutionary explanations for
627 cooperation. *Current biology* 17(16), R661–R672.