

Strategic coexistence theory for evolutionary games

Sang Woo Park^{1,2,*}

1 School of Biological Sciences, Seoul National University, Seoul, Korea

2 Institute for Data Innovation in Science, Seoul National University, Seoul, Korea

*Corresponding author: sangwoopark@snu.ac.kr

Abstract

Evolutionary game theory and ecological coexistence theory both seek to predict the outcome of competition between biological entities, be they strategies or species, but the two fields have relied on largely separate approaches. Replicator equations provide a foundation for predicting the outcome of strategic competition, yet they do not explicitly decompose the mechanisms underlying these outcomes. Inspired by modern coexistence theory from community ecology, we develop Strategic Coexistence Theory (SCT), which translates payoff-based invasion conditions into strategic niche differences and fitness differences between any two competing strategies described by classic replicator dynamics. SCT recovers the classic classification of two-strategy games, distinguishing competitive exclusion, coexistence, and priority effects within a shared niche–fitness difference space. Applying SCT to five mechanisms for the evolution of cooperation further reveals that these mechanisms promote cooperation through distinct dynamical routes: kin selection, network reciprocity, and group selection primarily reduce fitness differences, whereas direct and indirect reciprocity destabilize competition and generate priority effects. Finally, applying SCT to microbial public-goods competition shows that nonlinear microbial growth can both stabilize and equalize competition between cooperators and defectors, allowing coexistence. Together, these results show that SCT provides a common language for comparing coexistence mechanisms across evolutionary games.

Significance statement

Evolutionary game theory provides a powerful framework for predicting the outcome of competition between strategies, including the dominance of one strategy, coexistence of multiple strategies, and bistability. However, a common framework for comparing how different mechanisms promote strategy coexistence remains elusive, even though stable polymorphisms, such as the coexistence of cooperators and defectors, are widely observed across biological systems. To address this gap, we developed Strategic Coexistence Theory (SCT), which extends Modern Coexistence Theory from community ecology to evolutionary games. Applying SCT to classic games, mechanisms of cooperation, and microbial public-goods competition shows

38 that seemingly similar evolutionary outcomes can arise through distinct mechanisms.
39 This approach links ecological theory with replicator dynamics, providing a comple-
40 mentary tool for analyzing evolutionary games.

41 Introduction

42 Understanding the mechanisms that promote the evolutionary stability of competing
43 strategies is a fundamental aim in evolutionary biology (Lewontin, 1961; Slobodkin,
44 1964; Smith and Price, 1973; Taylor and Jonker, 1978). To this end, evolutionary
45 game theory (EGT) and replicator dynamics have provided key foundations, allow-
46 ing us to predict the outcome of strategic competition through payoff inequalities,
47 invasion criteria, and stability analysis (Schuster and Sigmund, 1983; Hofbauer and
48 Sigmund, 1998; Cressman, 2003; Cressman and Tao, 2014). Recently, there has been
49 growing interest in characterizing similarities between replicator dynamics and the
50 dynamics of competing species (Gokhale and Traulsen, 2025; Tarnita and Traulsen,
51 2025; Allesina, 2026), building on the classical mathematical equivalence between
52 replicator equations and generalized Lotka–Volterra models (Hofbauer, 1981; Page
53 and Nowak, 2002). In particular, Tarnita and Traulsen (2025) recently argued that a
54 closer synthesis between ecology and evolutionary game theory can provide new in-
55 sights into ecological and evolutionary assembly, competition, and coexistence. This
56 renewed perspective between ecological and evolutionary dynamics suggests that evo-
57 lutionary games may be interpreted not only as payoff-based competition but also
58 as problems of strategic coexistence.

59 Interestingly, even a simple two-strategy game can yield heterogeneous outcomes
60 that mirror outcomes in community ecology (Hauert, 2002; Szabó and Fath, 2007;
61 Cooney, 2020; Tarnita and Traulsen, 2025): the dominance of a single strategy (pris-
62 oner’s dilemma and harmony games), coexistence (hawk–dove games), and bistability
63 (stag-hunt games). Such differences in conditions that promote evolutionary stability
64 and dominance of a single strategy and those that permit strategy coexistence indi-
65 cate that EGT can be understood from the perspective of coexistence theory (Motro,
66 1991; Doebeli et al., 2004; Hauert and Doebeli, 2004; Doebeli and Hauert, 2005; Gore
67 et al., 2009; Souza et al., 2009; Archetti and Scheuring, 2011; Steidinger and Bever,
68 2014; Tarnita and Traulsen, 2025). In fact, stable polymorphisms of evolutionary
69 strategies, such as coexistence of cooperators and defectors, are observed in micro-
70 bial and other biological systems (Gore et al., 2009; Cordero et al., 2012; Kocher
71 et al., 2018; Pollak et al., 2021; Lin et al., 2024), but coexistence-theoretic perspec-
72 tive has often remained implicit. Recasting game dynamics as a coexistence problem
73 immediately raises the following question: when two strategies coexist, does the
74 corresponding mechanism stabilize or equalize competition (Chesson, 2000)? While
75 EGT allows us to predict whether two strategies can coexist or not, a common frame-
76 work for comparing coexistence mechanisms between competing strategies remains
77 elusive.

Modern coexistence theory (MCT) from community ecology presents a particularly useful framework for quantifying coexistence mechanisms (Chesson, 2000; Adler et al., 2007; Godoy et al., 2014; Godoy and Levine, 2014; Kraft et al., 2015; Adler et al., 2026). MCT predicts that coexistence of two competing entities, be they species or strategies, requires stabilizing niche differences to overcome average fitness differences (Chesson, 2000; Adler et al., 2007; Kraft et al., 2015). In community ecology, stabilizing niche differences represent differences in factors that limit the growth of competing species, causing each species to limit itself more than it limits its competitor (Chesson, 2000). In EGT, a strategy’s niche can be interpreted as strategic environments defined by the composition of competing strategies and the resulting interactions (Taylor and Jonker, 1978; Cressman and Tao, 2014; Tarnita and Traulsen, 2025). Thus, stabilizing niche differences arise when each strategy is favored by a different strategic environment, such that the spread of each strategy creates conditions that limit its own relative growth and favor the recovery of its competitor. In contrast, fitness differences represent differences in the ability of each entity to compete under the limiting environments created by its competitor (Chesson, 2000). This differs from payoff differences in EGT, which depend on the underlying strategy frequency and therefore combine both stabilizing effects caused by changes in the strategic environment and residual fitness differences that favor one strategy over another (Taylor and Jonker, 1978; Cressman and Tao, 2014; Tarnita and Traulsen, 2025).

So far, MCT has been primarily utilized for understanding species coexistence, but recent work showed that MCT can be extended beyond species competition, especially for studying pathogen strain competition, allowing one to directly quantify niche and fitness differences of competing entities, be they species or strains (Sieben et al., 2022; Park et al., 2024, 2026). Inspired by this effort, we present Strategic Coexistence Theory (SCT) for predicting coexistence and exclusion of competing strategies in EGT by decomposing invasion growth rates into strategic niche and fitness differences. We begin by introducing SCT, which can be applied to any generic system of EGT described by replicator equations. We show that SCT can recover outcomes from classic, two-strategy games, providing a reinterpretation of classic results from coexistence-theoretic perspective. We then apply SCT to comparing five mechanisms for the evolution of cooperation, revealing that direct and indirect reciprocity destabilize competition between cooperators and defectors, thus pushing the system towards priority effect regime. Finally, we consider a case study of microbial cooperation, where nonlinear microbial growth can both stabilize and equalize competition between cooperation and defection, allowing coexistence of strategies. Together, these examples show that SCT provides a common, ecological language for comparing the mechanisms underlying strategic evolution.

Strategic coexistence theory

Here, we present a brief derivation of strategic coexistence theory (SCT) (Figure 1). To do so, we begin with a classic replicator equation, which allows us to model changes in the frequency of competing strategies based on their payoffs (Schuster and Sigmund, 1983; Hofbauer and Sigmund, 1998; Cressman, 2003; Cressman and Tao, 2014):

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (1)$$

where x_i represents the frequency of strategy i , $\pi_i(\mathbf{x})$ represents the payoff of strategy i for a given strategy distribution $\mathbf{x}$, and $\bar{\pi}(\mathbf{x}) = \sum_i \pi_i(\mathbf{x})x_i$ represents the average payoff. Since strategy frequencies must sum to 1 (i.e., $\sum_i x_i = 1$), a single-strategy equilibrium dominated by strategy i can be described by a unit vector $\mathbf{e}_i$, where all elements are equal to zero except for the i -th element. We use these single-strategy equilibria to define niche and fitness differences between two competing strategies i and j .

Because payoffs in many games can be negative, payoff ratios can be difficult to interpret. Thus, we work on a positive multiplicative scale by exponentiating the payoff terms:

$$W_i(\mathbf{x}) = \exp(\pi_i(\mathbf{x})). \quad (2)$$

This exponential transformation is convenient because it preserves the payoff ordering and does not alter the invasion conditions. Therefore, this allows us to compare strategic environments through ratios, as in modern coexistence theory. For convenience, we refer to $W_i(\mathbf{x})$ as the multiplicative payoff of strategy i .

First, we define the niche overlap ρ between strategies i and j as the ratio between the geometric mean of each strategy's multiplicative payoff in the strategic environment that it creates when common, $\sqrt{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}$, and the geometric mean of each strategy's multiplicative payoff in the environment generated by its competitor, $\sqrt{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}$. The corresponding niche difference is then given by

$$1 - \rho = 1 - \sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}}. \quad (3)$$

When $1 - \rho > 0$, each strategy performs better, on average, in the environment generated by its competitor than in the environment generated by itself. In other words, intraspecific competition is stronger than interspecific competition. Second, we define the fitness difference as the ratio between the geometric means of the multiplicative payoff of each strategy across the two single-strategy environments:

$$\frac{\kappa_j}{\kappa_i} = \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}. \quad (4)$$

In other words, strategy j has an average competitive advantage over strategy i ($\kappa_j/\kappa_i > 1$) when it performs better, on average, across the environments generated by itself and its competitor.




Figure 1: **A schematic diagram summarizing strategic coexistence theory.** Strategic coexistence theory decomposes pairwise replicator dynamics into stabilizing and equalizing mechanisms. Mutual invasion conditions can be rewritten on a multiplicative payoff scale, allowing us to derive strategic niche difference $1 - \rho$ and fitness difference κ_j/κ_i based on multiplicative payoff ratios. Then, the inequality $\rho < \kappa_j/\kappa_i < 1/\rho$ determines mutual invasion. We note that the exact values of niche and fitness differences are sensitive to payoff scaling, while dynamical outcomes are still preserved.

150 Together, for any two strategies described by the classic replicator equation, mu-
 151 tual invasion requires stabilizing niche differences to overcome average fitness differ-
 152 ences. Specifically, the condition for mutual invasion can be rewritten as (Chesson,
 153 2000):

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (5)$$

154 This inequality applies to any pairwise comparison between two strategies described
 155 by the classic replicator equation because ρ and κ_j/κ_i can be calculated directly from
 156 ratios of multiplicative payoffs. This decomposition not only allows us to understand
 157 how different mechanisms contribute to the coexistence or exclusion of competing
 158 strategies but also provides shared axes for comparing outcomes of different games.
 159 Detailed derivations are provided in Supplementary Text S1.

160 We note that the absolute magnitudes of niche and fitness differences are sen-
 161 sitive to payoff scaling because they are defined on an exponentiated payoff scale.
 162 Therefore, while SCT allows us to compare different games on shared axes, we can-
 163 not compare the exact values of niche and fitness differences across different games.
 164 Nonetheless, as we show in Supplementary Text S2, the qualitative outcome of strate-
 165 gic competition under SCT is preserved because payoff rescaling preserves payoff
 166 orderings. Therefore, we focus on comparing qualitative outcomes from SCT, in-

cluding directions of change in niche and fitness difference and resulting dynamical regimes, and refrain from making any comparisons of magnitude of niche and fitness difference values.

Classic two-strategy games

As an illustrative example, we begin with classic two-strategy games (Hauert, 2002; Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025), which can be described by the following payoff matrix (Figure 2A):

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (6)$$

Here, the first and second rows represent strategies 1 and 2, respectively, and the columns represent the strategy of the interacting opponent. Writing x_1 to denote the frequency of strategy 1, the expected payoffs are

$$\pi_1(\mathbf{x}) = Rx_1 + S(1 - x_1), \quad (7)$$

$$\pi_2(\mathbf{x}) = Tx_1 + P(1 - x_1). \quad (8)$$

Thus, the dynamics can be written as

$$\frac{dx_1}{dt} = x_1(1 - x_1) [(R - T)x_1 + (S - P)(1 - x_1)]. \quad (9)$$

The signs of $T - R$ and $S - P$ determine whether each strategy can invade the other when rare. Specifically, strategy 2 can invade strategy 1 when $T > R$, whereas strategy 1 can invade strategy 2 when $S > P$. These two invasion conditions partition the game space into four regimes (Figure 2B): prisoner’s dilemma ($T > R > P > S$), hawk–dove game ($T > R, S > P$), stag–hunt game ($R > T, P > S$), and harmony game ($R > T, S > P$). In standard EGT, these outcomes are obtained by analyzing the stability of equilibrium conditions. For simple games, this analysis is straightforward.

A coexistence-theoretic approach recovers the same classification but provides a different interpretation (Figure 2C). For a two-strategy game determined by this payoff matrix, the niche difference and fitness difference are given by the following set of equations (Supplementary Text S3):

$$1 - \rho = 1 - \exp\left(\frac{R + P - S - T}{2}\right), \quad (10)$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{T + P - R - S}{2}\right). \quad (11)$$

As shown above, mutual invasion requires $\rho < \kappa_2/\kappa_1 < 1/\rho$, which is equivalent to $T > R$ and $S > P$. This corresponds to the hawk–dove game, in which each strategy




Figure 2: **Coexistence-theoretic interpretation of classic two-strategy games.** (A) General payoff matrix for a two-strategy game, where rows represent the focal strategy and columns represent the strategy of the interacting opponent. (B) Classification of four classic games based on the payoff differences $T - R$ and $S - P$. Prisoner’s dilemma and harmony games lead to the dominance of one strategy; hawk–dove games allow coexistence; and stag–hunt games result in bistability. (C) Mapping of the same four games onto niche–fitness difference space. Hawk–dove games correspond to the coexistence regime, where stabilizing niche differences overcome fitness differences. Prisoner’s dilemma and harmony games correspond to competitive-exclusion regimes, whereas stag–hunt games correspond to the priority-effect regime. $R > P$ is assumed throughout, but this does not affect coexistence outcome in any way.

192 can invade its competitor when rare and the two strategies coexist. The remaining
 193 games also have natural interpretations in niche–fitness difference space. Prisoner’s
 194 dilemma and harmony games represent competitive exclusion regimes, where one
 195 strategy has a fitness advantage that exceeds niche differences. In contrast, stag–hunt
 196 games represent a priority-effect regime, where neither strategy can invade the other
 197 and the outcome depends on initial conditions. Therefore, coexistence theory not
 198 only allows us to recover the classic results from two-strategy games but also provides
 199 new interpretations for these results in terms of niche and fitness differences. Building
 200 on this, we next apply this framework to comparing core mechanisms of cooperation.

201 Comparisons of mechanisms for the evolution of co- 202 operation

203 Even though cooperative behavior can be found across a wide range of biological
 204 systems, a simple prisoner’s dilemma shows that cooperation is not an evolutionarily

stable strategy (Axelrod and Hamilton, 1981; May, 1987). Thus, understanding mechanisms that promote the evolution of cooperation has remained a central goal in evolutionary biology (West et al., 2007; Fletcher and Doebeli, 2008; Clutton-Brock, 2009; Kingma et al., 2014; Rakoff-Nahoum et al., 2016; Hilbe et al., 2018; Apicella and Silk, 2019; West et al., 2021; Lee et al., 2022; Palmer and Foster, 2022; Michel-Mata et al., 2024). Here, we compare five classic mechanisms for the evolution of cooperation using SCT (Nowak, 2006) and ask how they differ from the perspective of SCT.

Specifically, we consider the following mechanisms: kin selection, direct reciprocity, indirect reciprocity, network reciprocity, and group selection (Figure 3A). Kin selection favors cooperation when two interacting individuals are genetic relatives, where the genetic relatedness r must be greater than the cost-to-benefit ratio: $r > c/b$ (Hamilton, 1964). Direct reciprocity favors cooperation when current cooperation may promote future cooperation by the same individual through repeated encounters (Trivers, 1971; Axelrod and Hamilton, 1981). In this case, the probability of repeated interaction (i.e., “next round”) must be greater than the cost-to-benefit ratio: $w > c/b$. Indirect reciprocity favors cooperation when current cooperation may promote future cooperation by different individuals through reputation (Nowak and Sigmund, 1998; Wedekind and Milinski, 2000; Nowak and Sigmund, 2005). Thus, the social acquaintanceship (i.e., the probability of knowing someone’s reputation) must be greater than the cost-to-benefit ratio: $q > c/b$. Network reciprocity favors cooperation via clustering of cooperators, requiring the benefit-to-cost ratio to be greater than the average number of neighbors: $b/c > k$ (Nowak and May, 1992; Lieberman et al., 2005). Finally, group selection favors cooperation via multilevel selection, where groups with more cooperators exhibit faster growth at the group level even if defectors reproduce faster at the individual level (Traulsen and Nowak, 2006). This requires $b/c > 1 + (n/m)$, where n is the maximum group size and m is the number of groups. All of these mechanisms can be described by a simple 2×2 payoff matrix (Nowak, 2006), from which the corresponding niche and fitness differences can be computed using Eq. (10) and Eq. (11). Details on niche and fitness difference calculations are presented in Supplementary Text S4.

First, the prisoner’s dilemma provides a baseline scenario, where defection is evolutionarily stable (Figure 3A). From the perspective of SCT, there is complete niche overlap, and therefore defection competitively excludes cooperation (Figure 3B). Building on this, SCT shows that five mechanisms promote evolutionary stability of cooperation via different dynamical routes (Figure 3B). For example, kin selection allows the fitness of cooperation to increase as genetic relatedness r increases (Figure 3B). However, it does not affect the niche difference between cooperation and defection. Similarly, network reciprocity and group selection modulate the fitness of cooperation without causing any changes in strategic niches. In contrast, direct and indirect reciprocity destabilize competition while increasing the fitness of cooperation, thereby pushing the competition outcome towards a priority effect regime. Equivalently, these two mechanisms promote bistability of defection and cooperation.




Figure 3: **Qualitative comparisons of five mechanisms for the evolution of cooperation using SCT.** (A) Payoff matrices for the baseline scenario (prisoner's dilemma) and five mechanisms for the evolution of cooperation (Nowak, 2006). Prisoner's dilemma is characterized by the benefit for the recipient b and the cost for the donor c . Network reciprocity is parameterized based on $H = [(b-c)k - 2c]/[(k+1)(k-2)]$. Parameters for all other mechanisms are described in panel B. (B) Changes in niche and fitness differences driven by each mechanism. Each arrow represents the direction of change as we increase the corresponding parameter, indicated in figure legends. Note that kin selection, network reciprocity, and group selection all result in changes in fitness difference without causing any changes in niche difference. Therefore, we indicate the direction of change for network reciprocity and group selection separately to avoid overlapping arrows. For simplicity, we focus on the effect of number of groups m for group selection; increasing the number of maximum group size n has the opposite effect (Supplementary Text S4). Across all models, we assume $b = 3$ and $c = 1$. Because SCT decompositions depend on positive payoff scaling, exact values of niche and fitness difference (e.g., arrow lengths, slopes, and distances from regime boundaries) should not be compared across mechanisms.

248 SCT further reveals another difference between direct and indirect reciprocity.
 249 Direct reciprocity remains within the priority-effect regime as w reaches 1, whereas
 250 indirect reciprocity reaches the boundary between the priority-effect and competitive-
 251 exclusion regimes at $q = 1$ (Supplementary Text S4).

252 We note that none of these mechanisms allow coexistence of cooperation and de-
 253 fection. In many biological systems, such coexistence is often observed (Gore et al.,
 254 2009; Cordero et al., 2012; Kocher et al., 2018; Pollak et al., 2021; Lin et al., 2024).
 255 Therefore, we move on to the final example, applying SCT to microbial coopera-
 256 tion, where nonlinear benefits and environmental context can promote coexistence
 257 of cooperators and defectors.

258 Microbial cooperation and public goods

259 Here, we consider a simple, phenomenological model of competition between coop-
 260 erating and cheating yeast in a cooperative sucrose-metabolism system, where invertase
 261 production by cooperators allows cheaters to utilize glucose, a public good released
 262 through sucrose hydrolysis (Carlson and Botstein, 1982; Dickinson, 1998; Gore et al.,
 263 2009). Specifically, invertase production by cooperators incurs a cost c and gener-
 264 ates a total benefit of 1 that is captured privately with efficiency ϵ . Assuming a
 265 linear relationship between glucose availability and microbial growth, the payoffs for
 266 cooperators π_C and cheaters (defectors) π_D can then be written as

$$\pi_C(f) = \epsilon + f(1 - \epsilon) - c, \quad (12)$$

$$\pi_D(f) = f(1 - \epsilon), \quad (13)$$

267 where the use of glucose depends on the fraction of cooperators f . The dynamics of
 268 cooperators and defectors are described by the following replicator equation:

$$\frac{df}{dt} = f(1 - f) [\pi_C(f) - \pi_D(f)] \quad (14)$$

$$= f(1 - f)(\epsilon - c) \quad (15)$$

269 For this linear growth model, the payoff difference is fixed to $\epsilon - c$ independent of
 270 f (Figure 4A). Therefore, cooperation is favored when efficiency is greater than the
 271 cost of cooperation, $\epsilon > c$, whereas defection is favored when the cost is greater than
 272 efficiency, $c > \epsilon$ (Gore et al., 2009).

273 Our framework allows us to reinterpret this result from the perspective of coexis-
 274 tence theory. In particular, the two strategies have no niche difference, corresponding
 275 to complete niche overlap, $\rho = 1$. In contrast, the fitness difference is determined by
 276 the balance between private efficiency and the cost of cooperation:

$$\frac{\kappa_D}{\kappa_C} = \exp(c - \epsilon). \quad (16)$$

277 Therefore, this model always predicts competitive exclusion, except in the neutral
 278 case $c = \epsilon$. The outcome is determined entirely by the fitness difference between
 279 cooperators and defectors, rather than by stabilizing niche differences. This example
 280 resembles the prisoner's dilemma discussed in the previous example (Figure 3).




Figure 4: **Effects of nonlinear microbial growth on coexistence of cooperation and defection.** (A) Payoff differences between cooperation and defection for linear and nonlinear models of microbial growth. In panel A, we assumed $\epsilon = 0.4$, $c = 0.3$, and $\alpha = 0.15$ for illustrative purposes. (B) Effect of nonlinearity α on niche difference $1 - \rho$. Each line assumes a different value of efficiency ϵ . Note that lines for $\epsilon = 0.6$ and $\epsilon = 0.4$ are nearly overlapping and are indistinguishable. (C) Effect of nonlinearity α on both niche difference $1 - \rho$ and fitness difference $\log(\kappa_D/\kappa_C)$. The gray shaded area represents the coexistence regime. The square represents the condition when $\alpha \rightarrow 0^+$. The circles represent the condition when $\alpha = 0.15$, which correspond to the experimental relationship between glucose and microbial growth (Gore et al., 2009). In panels B and C, we assumed $c = 0.3$, while allowing ϵ and α to vary.

281 In real biological systems, the effect of glucose on microbial growth follows a
 282 nonlinear relationship:

$$\pi_C(f) = [\epsilon + f(1 - \epsilon)]^\alpha - c, \quad (17)$$

$$\pi_D(f) = [f(1 - \epsilon)]^\alpha, \quad (18)$$

283 where $\alpha = 0.15$ reflects the experimental relationship between glucose and microbial
 284 growth (Gore et al., 2009). Then, the dynamics of cooperators and defectors are
 285 described by the following replicator equation:

$$\frac{df}{dt} = f(1 - f) [\{\epsilon + f(1 - \epsilon)\}^\alpha - c - \{f(1 - \epsilon)\}^\alpha]. \quad (19)$$

286 Under this nonlinear assumption, the payoff difference decreases with the cooperator
 287 fraction f , which can promote the coexistence of cooperators and defectors (Figure
 288 4A): cooperators have a competitive advantage when rare, whereas defectors have a
 289 competitive advantage when cooperators are abundant.

290 From a coexistence-theoretic perspective, this means that nonlinear growth can
 291 generate sufficient stabilizing niche difference to overcome the fitness difference. In

particular, the niche and fitness differences for this model correspond to:

$$1 - \rho = 1 - \exp\left(\frac{1 - \epsilon^\alpha - (1 - \epsilon)^\alpha}{2}\right), \quad (20)$$

$$\frac{\kappa_D}{\kappa_C} = \exp\left(c + \frac{(1 - \epsilon)^\alpha - 1 - \epsilon^\alpha}{2}\right). \quad (21)$$

These expressions clarify the separate roles of cost, efficiency, and nonlinearity. For example, the cost of cooperation does not affect niche difference and only affects fitness difference. Instead, niche difference is generated by the concave relationship between glucose availability and microbial growth, which increases from 0 to $1 - \exp(-1/2)$ as α goes from 1 to 0 (Figure 4B,C). In contrast, as $\alpha \rightarrow 0^+$, the fitness difference approaches

$$\log(\kappa_D/\kappa_C) \rightarrow c - \frac{1}{2}, \quad (22)$$

which no longer depends on the private capture efficiency ϵ (Figure 4C). Thus, strong concavity not only generates stabilizing niche differences, but also acts as an equalizing mechanism (Figure 4C). Such stabilizing mechanisms are key ingredients for coexistence of cooperators and defectors.

Discussion

Standard evolutionary game theory and replicator equations provide powerful tools for predicting the outcome of strategy competition, including strategic dominance, coexistence, and bistability (Motro, 1991; Doebeli et al., 2004; Hauert and Doebeli, 2004; Doebeli and Hauert, 2005; Gore et al., 2009; Souza et al., 2009; Archetti and Scheuring, 2011; Steidinger and Bever, 2014; Tarnita and Traulsen, 2025). However, these outcomes are typically understood by analyzing payoff differences within individual games, which makes it difficult to compare coexistence mechanisms across models. Building on Chesson’s Modern Coexistence Theory from community ecology (Chesson, 2000), we developed a strategic coexistence theory (SCT) that allows us to decompose strategy competition in terms of stabilizing niche differences and fitness differences. This decomposition provides a unifying framework for comparing different games from the perspective of coexistence theory, allowing us to identify stabilizing and equalizing mechanisms for strategy competition.

To demonstrate the utility of SCT, we considered three case studies. First, the analysis of classic two-strategy games showed that SCT recovers the original classification for these classic games, highlighting the similarities between evolutionary games and community ecology (Hauert, 2002; Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025). Then, we compared different mechanisms for the evolution of cooperation, which showed that each mechanism promotes cooperation via different dynamical routes (Nowak, 2006). Kin selection, network reciprocity, and

group selection primarily affect the fitness difference between cooperation and defection, whereas direct and indirect reciprocity can destabilize competition and generate priority effects. Finally, our analysis of microbial cooperation showed that coexistence between cooperators and defectors requires more than reducing the fitness cost of cooperation: nonlinear benefits can generate stabilizing niche differences while also reducing fitness asymmetry (Gore et al., 2009). Taken together, these examples show that SCT can provide new insights into ecological underpinnings of strategy competition.

There are several limitations to SCT. First, similar to MCT from community ecology, SCT only allows for comparisons of pairwise strategies and therefore cannot determine the outcomes of games with more than two strategies (Chesson, 2000). Second, the exact values of strategic niche and fitness differences are sensitive to payoff scaling. SCT therefore supports quantitative comparisons only when models share a justified payoff normalization. Without such normalization, comparisons are restricted to qualitative changes in dynamical regimes, directions, and boundary crossings. Third, we focused on classic replicator equations, but Tarnita and Traulsen (2025) recently pointed out that these classic formulations implicitly ignore differences in intrinsic growth rates. In such cases, replicator equations may fail to predict the outcome of competition (Tarnita and Traulsen, 2025), and a full Lotka-Volterra model is needed to describe the abundances, rather than frequencies, of strategies. Thus, SCT inherits the same assumptions as the classic replicator framework: it is most appropriate when strategy competition can be described in terms of relative frequencies alone (Tarnita and Traulsen, 2025). When strategies differ intrinsically in growth, MCT may be required instead to tease apart mechanisms of strategy coexistence. Finally, we have focused on simple examples to illustrate the utility of SCT, but many games involve additional complexities, including population structure (Nowak et al., 2010), stochasticity (Wang et al., 2025; Bodin et al., 2026), and explicit ecological feedback (Tarnita and Traulsen, 2025). Extending SCT to these settings will be necessary to determine how broadly stabilizing and equalizing mechanisms can be compared across evolutionary games.

In conclusion, SCT provides a bridge between evolutionary game theory and community ecology, allowing strategy competition to be understood from the perspective of coexistence theory. Therefore, rather than replacing existing EGT or replicator-dynamics frameworks, SCT provides a complementary tool for teasing apart the coexistence mechanisms underlying evolutionary games. More broadly, our study builds on recent efforts to extend MCT beyond species competition (Park et al., 2024, 2026), providing a foundation for building unifying coexistence theory for competing entities, including species, pathogen strains, and behavioral strategies.

362 **Acknowledgement**

363 We thank Jonathan Levine for the helpful discussion. S.W.P. was supported by
364 the New Faculty Startup Fund from Seoul National University, the National Re-
365 search Foundation of Korea (NRF) grant funded by the Korea government (MSIT)
366 (RS-2026-25474574), and the Global-LAMP Program of the National Research Foun-
367 dation of Korea (NRF) grant funded by the Ministry of Education (No. RS-2023-
368 00301976).

369 **Competing interests**

370 The authors declare no competing interest.

371 **Data availability**

372 All data and code are stored in a publicly available GitHub repository (<https://github.com/parklab-snu/coexistence-strategy>).
373

Supplementary Text

S1 Derivation of the strategic coexistence theory

To derive strategic coexistence theory (SCT), we begin with replicator equations:

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S1})$$

where x_i represents the frequency of strategy i , π_i represents the payoff of strategy i , and $\bar{\pi}$ represents the average payoff. Then, the mutual invasion condition for two competing strategies i and j can be written as

$$\pi_j(\mathbf{e}_j) < \pi_i(\mathbf{e}_j), \quad (\text{S2})$$

$$\pi_i(\mathbf{e}_i) < \pi_j(\mathbf{e}_i). \quad (\text{S3})$$

These inequalities will be preserved when we exponentiate the payoffs, $W_i(\mathbf{x}) = \exp(\pi_i(\mathbf{x}))$:

$$W_j(\mathbf{e}_j) < W_i(\mathbf{e}_j), \quad (\text{S4})$$

$$W_i(\mathbf{e}_i) < W_j(\mathbf{e}_i). \quad (\text{S5})$$

Rearranging, we obtain the following inequality:

$$\frac{W_i(\mathbf{e}_i)}{W_j(\mathbf{e}_i)} < 1 < \frac{W_i(\mathbf{e}_j)}{W_j(\mathbf{e}_j)}. \quad (\text{S6})$$

By multiplying $\sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}$ on both sides, we obtain:

$$\sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}} < \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}} < \sqrt{\frac{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}} \quad (\text{S7})$$

Thus, by defining niche overlap ρ and fitness difference κ_j/κ_i as follows,

$$\rho = \sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}}, \quad (\text{S8})$$

$$\frac{\kappa_j}{\kappa_i} = \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}, \quad (\text{S9})$$

we obtain the following inequality that defines conditions for mutual invasion and long-term coexistence of two competing strategies:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (\text{S10})$$

387 S2 Niche and fitness difference payoff scaling

388 Consider a classic replicator equation:

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S11})$$

389 where x_i represents the frequency of strategy i , π_i represents the payoff of strategy
 390 i , and $\bar{\pi}$ represents the average payoff. In this case, multiplying the payoffs by a
 391 constant $s > 0$ only changes time and does not affect the overall dynamics:

$$\frac{dx_i}{dt} = x_i(\pi'_i(\mathbf{x}) - \bar{\pi}'(\mathbf{x})) \quad (\text{S12})$$

$$= sx_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S13})$$

392 where $\pi'_i(\mathbf{x}) = s\pi_i(\mathbf{x})$ is the scaled payoff. Then, the multiplicative payoff corresponds
 393 to $W'_i(\mathbf{x}) = \exp(s\pi_i) = W_i(\mathbf{x})^s$.

394 If we try to calculate niche overlap and fitness difference based on the scaled
 395 payoffs, we get:

$$\rho' = \sqrt{\frac{W'_i(\mathbf{e}_i)W'_j(\mathbf{e}_j)}{W'_j(\mathbf{e}_i)W'_i(\mathbf{e}_j)}}, \quad (\text{S14})$$

$$= \left(\sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}} \right)^s, \quad (\text{S15})$$

$$= \rho^s \quad (\text{S16})$$

$$\frac{\kappa'_j}{\kappa'_i} = \sqrt{\frac{W'_j(\mathbf{e}_i)W'_j(\mathbf{e}_j)}{W'_i(\mathbf{e}_i)W'_i(\mathbf{e}_j)}}, \quad (\text{S17})$$

$$= \left(\sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}} \right)^s, \quad (\text{S18})$$

$$= \left(\frac{\kappa_j}{\kappa_i} \right)^s \quad (\text{S19})$$

396 This implies that the exact values of niche and fitness difference are sensitive to
 397 payoff scaling and are not intrinsic properties of a replicator system.

398 Nonetheless, the inequality for mutual invasion condition still holds under payoff
 399 scaling:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho} \iff \rho^s < \left(\frac{\kappa_j}{\kappa_i} \right)^s < \frac{1}{\rho^s}. \quad (\text{S20})$$

400 Furthermore, boundary conditions for complete niche overlap ($\rho = \rho' = 1$) and com-
 401 plete niche difference ($\rho = \rho' = 0$) are preserved. Moreover, the direction of changes
 402 in fitness and niche difference is also preserved under payoff scaling. Specifically, for

parameter θ , we have:

$$\frac{d\rho'}{d\theta} = s\rho^{s-1} \frac{d\rho}{d\theta} \quad (\text{S21})$$

$$\frac{d}{d\theta} \left(\frac{\kappa'_j}{\kappa'_i} \right) = s \left(\frac{\kappa_j}{\kappa_i} \right)^{s-1} \frac{d}{d\theta} \left(\frac{\kappa_j}{\kappa_i} \right). \quad (\text{S22})$$

Since ρ and κ_j/κ_i are always positive, directional sensitivities of ρ and κ_j/κ_i to parameter θ are preserved. Similarly, directional sensitivities of $1 - \rho$ to θ are preserved:

$$\frac{d(1 - \rho')}{d\theta} = s\rho^{s-1} \frac{d(1 - \rho)}{d\theta} \quad (\text{S23})$$

Therefore, we focus on comparisons of qualitative outcomes from SCT and refrain from making any comparisons of magnitude of niche and fitness difference values.

S3 Applications to classic two-strategy games

Consider a two-strategy game described by the following payoff matrix:

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (\text{S24})$$

There are four classic games that we can define from this simple payoff matrix: prisoner's dilemma ($T > R > P > S$), hawk-dove game ($T > R, S > P$), stag-hunt game ($R > T, P > S$), and harmony game ($R > T, S > P$). Here, we show that each game corresponds to a distinct region in the niche-fitness difference space under SCT.

To do so, we first begin by deriving expressions for niche overlap ρ and fitness difference κ_j/κ_i . Writing x_1 to denote the frequency of strategy 1, the expected payoff for each strategy corresponds to:

$$\pi_1(\mathbf{x}) = Rx_1 + S(1 - x_1), \quad (\text{S25})$$

$$\pi_2(\mathbf{x}) = Tx_1 + P(1 - x_1). \quad (\text{S26})$$

Then, multiplicative payoffs for each strategy when rare can be written as:

$$W_1(1) = \exp(R) \quad (\text{S27})$$

$$W_1(0) = \exp(S) \quad (\text{S28})$$

$$W_2(1) = \exp(T) \quad (\text{S29})$$

$$W_2(0) = \exp(P) \quad (\text{S30})$$

Therefore, we have:

$$\rho = \exp \left(\frac{R + P - S - T}{2} \right), \quad (\text{S31})$$

$$\frac{\kappa_2}{\kappa_1} = \exp \left(\frac{T + P - R - S}{2} \right). \quad (\text{S32})$$

420 As before, SCT allows us to predict the mutual invasion of two strategy using the
 421 following inequality:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (\text{S33})$$

422 Then, simple algebra reveals that each condition in mutual invasion inequality
 423 can be equivalently described by the payoff inequality:

$$\rho < \frac{\kappa_2}{\kappa_1} \iff R < T \quad (\text{S34})$$

$$\rho > \frac{\kappa_2}{\kappa_1} \iff R > T \quad (\text{S35})$$

$$\frac{\kappa_2}{\kappa_1} < \frac{1}{\rho} \iff P < S \quad (\text{S36})$$

$$\frac{\kappa_2}{\kappa_1} > \frac{1}{\rho} \iff P > S \quad (\text{S37})$$

424 These inequalities then allow us to classify outcomes of two-strategy games from the
 425 perspective of SCT. Note that the inequality between R and P does not affect the
 426 outcome.

427 **S4 Applications to mechanisms for the evolution of coopera-** 428 **tion**

429 We consider five mechanisms for the evolution of cooperation, all of which can be
 430 represented as a simple 2×2 payoff matrix (Nowak, 2006). Here, we go through each
 431 mechanism and derive measures for niche and fitness differences.

432 **Prisoner's dilemma.** Here, we consider a special case of Prisoner's dilemma
 433 which can be described by the following payoff matrix:

$$\begin{pmatrix} b - c & -c \\ b & 0 \end{pmatrix}, \quad (\text{S38})$$

434 where b represents the benefit for the recipient and c represents cost for the donor.
 435 In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0 \quad (\text{S39})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(c). \quad (\text{S40})$$

436 Therefore, as long as the cost is greater than zero, prisoner's dilemma always results
 437 in the competitive exclusion of cooperation.

438 **Kin selection.** Kin selection can be described by the following payoff matrix:

$$\begin{pmatrix} (b - c)(1 + r) & br - c \\ b - rc & 0 \end{pmatrix}, \quad (\text{S41})$$

where r represents the genetic relatedness. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0 \quad (\text{S42})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(c - br). \quad (\text{S43})$$

Therefore, defection will exclude cooperation when $c > br$. Equivalently, cooperation will be favored when $r > c/b$, which recovers the classic result.

Direct reciprocity. Direct reciprocity can be described by the following payoff matrix:

$$\begin{pmatrix} (b-c)/(1-w) & -c \\ b & 0 \end{pmatrix}, \quad (\text{S44})$$

where w represents the probability of next round. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 1 - \exp\left(\frac{(b-c)/(1-w) + c - b}{2}\right) \quad (\text{S45})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{b + c - (b-c)/(1-w)}{2}\right). \quad (\text{S46})$$

Therefore, defection will exclude cooperation when

$$\rho < \frac{\kappa_2}{\kappa_1} \quad (\text{S47})$$

$$\iff \exp\left(\frac{(b-c)/(1-w) + c - b}{2}\right) < \exp\left(\frac{b + c - (b-c)/(1-w)}{2}\right) \quad (\text{S48})$$

$$\iff bw < c \quad (\text{S49})$$

However, note that the following inequality holds whenever $c > 0$:

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1}. \quad (\text{S50})$$

Therefore, when $w > c/b$, we will have

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1} < \rho, \quad (\text{S51})$$

which implies priority effect (equivalently, bistability). Even as $w \rightarrow 1^-$, this inequality holds and therefore, cooperation cannot competitively exclude defection.

Indirect reciprocity. Indirect reciprocity can be described by the following payoff matrix:

$$\begin{pmatrix} b-c & -c(1-q) \\ b(1-q) & 0 \end{pmatrix}, \quad (\text{S52})$$

where q represents social acquaintanceship. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 1 - \exp\left(\frac{(b - c)q}{2}\right), \quad (\text{S53})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{2c - (b + c)q}{2}\right). \quad (\text{S54})$$

In this case, the condition for the dominance of defection (and therefore competitive exclusion of cooperation) can be written as:

$$\rho < \frac{\kappa_2}{\kappa_1} \quad (\text{S55})$$

$$\iff bq < c \quad (\text{S56})$$

Moreover, the following inequality always holds as long as $c > 0$.

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1}, \quad (\text{S57})$$

which implies that coexistence between cooperation and defection is not possible. Therefore, when $q < c/b$, then defection will competitively exclude cooperation. Otherwise, the system will enter the priority-effect regime.

Interestingly, when $q \rightarrow 1$, we have

$$\rho = \exp\left(\frac{b - c}{2}\right) \quad (\text{S58})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{c - b}{2}\right) \quad (\text{S59})$$

$$\frac{1}{\rho} = \exp\left(\frac{c - b}{2}\right) \quad (\text{S60})$$

$$(\text{S61})$$

In other words, the following inequality will hold:

$$\frac{1}{\rho} = \frac{\kappa_2}{\kappa_1} < \rho \quad (\text{S62})$$

Network reciprocity. Network reciprocity can be described by the following payoff matrix:

$$\begin{pmatrix} b - c & H - c \\ b - H & 0 \end{pmatrix}, \quad (\text{S63})$$

where $H = [(b - c)k - 2c]/[(k + 1)(k - 2)]$ and k represents the number of neighbors. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0, \quad (\text{S64})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(-H + c), \quad (\text{S65})$$

468 which implies that there is complete niche overlap and that fitness difference decreases
 469 with H . Since H is a monotonically decreasing function of k when $k > 2$ and $b/c \geq 2$,
 470 fitness difference will increase with k . In particular, cooperation will competitively
 471 exclude defection when $H > c$, meaning that:

$$b/c > k. \quad (\text{S66})$$

472 **Group selection.** Group selection can be described by the following payoff
 473 matrix:

$$\begin{pmatrix} (b-c)(m+n) & (b-c)m - cn \\ bn & 0 \end{pmatrix}, \quad (\text{S67})$$

474 where m represents the number of groups and n represents the group size. In this
 475 case, the niche and fitness differences correspond to:

$$1 - \rho = 0, \quad (\text{S68})$$

$$\frac{\kappa_2}{\kappa_1} = \exp((c-b)m + cn), \quad (\text{S69})$$

476 which implies that there is complete niche overlap. The fitness difference will increase
 477 with n but decrease with m , provided that $b > c$. In particular, cooperation will
 478 competitively exclude defection under the following condition:

$$1 + \frac{n}{m} < \frac{b}{c}. \quad (\text{S70})$$

479 References

- 480 Adler, P. B., M. Detto, S. P. Ellner, T. L. Gibbs, Z. J. Gold, S. J. Leonard, J. I.
 481 Levine, A. E. Schiffer, C. Song, M. Stemkovski, et al. (2026). What have we learned
 482 from empirical applications of modern coexistence theory? *Ecology letters* 29(6),
 483 e70404.
- 484 Adler, P. B., J. HilleRisLambers, and J. M. Levine (2007). A niche for neutrality.
 485 *Ecology letters* 10(2), 95–104.
- 486 Allesina, S. (2026). Global stability of ecological and evolutionary dynamics
 487 via equivalence. *Proceedings of the National Academy of Sciences* 123(13),
 488 e2534915123.
- 489 Apicella, C. L. and J. B. Silk (2019). The evolution of human cooperation. *Current*
 490 *Biology* 29(11), R447–R450.
- 491 Archetti, M. and I. Scheuring (2011). Coexistence of cooperation and defection in
 492 public goods games. *Evolution* 65(4), 1140–1148.
- 493 Axelrod, R. and W. D. Hamilton (1981). The evolution of cooperation. *sci-*
 494 *ence* 211(4489), 1390–1396.

495 Bodin, F., G. Wang, and J. B. Plotkin (2026). Eco-evolutionary games in noisy
496 environments. *bioRxiv*, 2026–05.

497 Carlson, M. and D. Botstein (1982). Two differentially regulated mRNAs with differ-
498 ent 5' ends encode secreted and intracellular forms of yeast invertase. *Cell* 28(1),
499 145–154.

500 Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual review*
501 *of Ecology and Systematics* 31(1), 343–366.

502 Clutton-Brock, T. (2009). Cooperation between non-kin in animal societies. *Na-*
503 *ture* 462(7269), 51–57.

504 Cooney, D. B. (2020). Analysis of multilevel replicator dynamics for general two-
505 strategy social dilemma. *Bulletin of Mathematical Biology* 82(6), 66.

506 Cordero, O. X., L.-A. Ventouras, E. F. DeLong, and M. F. Polz (2012). Public good
507 dynamics drive evolution of iron acquisition strategies in natural bacterioplankton
508 populations. *Proceedings of the National Academy of Sciences* 109(49), 20059–
509 20064.

510 Cressman, R. (2003). *Evolutionary dynamics and extensive form games*, Volume 5.
511 MIT Press.

512 Cressman, R. and Y. Tao (2014). The replicator equation and other game dynamics.
513 *Proceedings of the National Academy of Sciences* 111(supplement_3), 10810–10817.

514 Dickinson, J. R. (1998). *Metabolism and molecular physiology of Saccharomyces*
515 *cerevisiae*. Taylor & Francis.

516 Doebeli, M. and C. Hauert (2005). Models of cooperation based on the Prisoner’s
517 Dilemma and the Snowdrift game. *Ecology letters* 8(7), 748–766.

518 Doebeli, M., C. Hauert, and T. Killingback (2004). The evolutionary origin of coop-
519 erators and defectors. *science* 306(5697), 859–862.

520 Fletcher, J. A. and M. Doebeli (2008). A simple and general explanation for the evolu-
521 tion of altruism. *Proceedings of the Royal Society B: Biological Sciences* 276(1654),
522 13.

523 Godoy, O., N. J. Kraft, and J. M. Levine (2014). Phylogenetic relatedness and the
524 determinants of competitive outcomes. *Ecology letters* 17(7), 836–844.

525 Godoy, O. and J. M. Levine (2014). Phenology effects on invasion success: insights
526 from coupling field experiments to coexistence theory. *Ecology* 95(3), 726–736.

527 Gokhale, C. S. and A. Traulsen (2025). Higher-order equivalence of Lotka-Volterra
528 and replicator dynamics. *bioRxiv*, 2025–03.

- 529 Gore, J., H. Youk, and A. Van Oudenaarden (2009). Snowdrift game dynamics and
530 facultative cheating in yeast. *Nature* 459(7244), 253–256.
- 531 Hamilton, W. D. (1964). The genetical evolution of social behaviour. II. *Journal of*
532 *theoretical biology* 7(1), 17–52.
- 533 Hauert, C. (2002). Effects of space in 2×2 games. *International Journal of Bifur-*
534 *cation and Chaos* 12(07), 1531–1548.
- 535 Hauert, C. and M. Doebeli (2004). Spatial structure often inhibits the evolution of
536 cooperation in the snowdrift game. *Nature* 428(6983), 643–646.
- 537 Hilbe, C., Š. Šimsa, K. Chatterjee, and M. A. Nowak (2018). Evolution of cooperation
538 in stochastic games. *Nature* 559(7713), 246–249.
- 539 Hofbauer, J. (1981). On the occurrence of limit cycles in the volterra-lotka equation.
540 *Nonlinear Analysis: Theory, Methods & Applications* 5(9), 1003–1007.
- 541 Hofbauer, J. and K. Sigmund (1998). *Evolutionary games and population dynamics*.
542 Cambridge university press.
- 543 Kingma, S. A., P. Santema, M. Taborsky, and J. Komdeur (2014). Group augmen-
544 tation and the evolution of cooperation. *Trends in ecology & evolution* 29(8),
545 476–484.
- 546 Kocher, S. D., R. Mallarino, B. E. Rubin, D. W. Yu, H. E. Hoekstra, and N. E.
547 Pierce (2018). The genetic basis of a social polymorphism in halictid bees. *Nature*
548 *Communications* 9(1), 4338.
- 549 Kraft, N. J., O. Godoy, and J. M. Levine (2015). Plant functional traits and the mul-
550 tidimensional nature of species coexistence. *Proceedings of the National Academy*
551 *of Sciences* 112(3), 797–802.
- 552 Lee, I. P. A., O. T. Eldakar, J. P. Gogarten, and C. P. Andam (2022). Bacterial co-
553 operation through horizontal gene transfer. *Trends in Ecology & Evolution* 37(3),
554 223–232.
- 555 Lewontin, R. C. (1961). Evolution and the theory of games. *Journal of Theoretical*
556 *Biology* 1(3), 382–403.
- 557 Lieberman, E., C. Hauert, and M. A. Nowak (2005). Evolutionary dynamics on
558 graphs. *Nature* 433(7023), 312–316.
- 559 Lin, H., D. Wang, Q. Wang, J. Mao, Y. Bai, and J. Qu (2024). Interspecific compe-
560 tition prevents the proliferation of social cheaters in an unstructured environment.
561 *The ISME Journal* 18(1), wrad038.
- 562 May, R. M. (1987). More evolution of cooperation. *Nature* 327(6117), 15–17.

Michel-Mata, S., M. Kawakatsu, J. Sartini, T. A. Kessinger, J. B. Plotkin, and C. E. Tarnita (2024). The evolution of private reputations in information-abundant landscapes. *Nature* 634(8035), 883–889.

Motro, U. (1991). Co-operation and defection: playing the field and the ESS. *Journal of Theoretical Biology* 151(2), 145–154.

Nowak, M. A. (2006). Five rules for the evolution of cooperation. *science* 314(5805), 1560–1563.

Nowak, M. A. and R. M. May (1992). Evolutionary games and spatial chaos. *nature* 359(6398), 826–829.

Nowak, M. A. and K. Sigmund (1998). Evolution of indirect reciprocity by image scoring. *Nature* 393(6685), 573–577.

Nowak, M. A. and K. Sigmund (2005). Evolution of indirect reciprocity. *Nature* 437(7063), 1291–1298.

Nowak, M. A., C. E. Tarnita, and T. Antal (2010). Evolutionary dynamics in structured populations. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365(1537), 19.

Page, K. M. and M. A. Nowak (2002). Unifying evolutionary dynamics. *Journal of theoretical biology* 219(1), 93–98.

Palmer, J. D. and K. R. Foster (2022). Bacterial species rarely work together. *Science* 376(6593), 581–582.

Park, S. W., S. Cobey, C. J. E. Metcalf, J. M. Levine, and B. T. Grenfell (2024). Predicting pathogen mutual invasibility and co-circulation. *Science* 386(6718), 175–179.

Park, S. W., J. Levine, and B. Grenfell (2026). Unifying coexistence theory for ecological communities and pathogen strain competition. *bioRxiv*, 2026–05.

Pollak, S., M. Gralka, Y. Sato, J. Schwartzman, L. Lu, and O. X. Cordero (2021). Public good exploitation in natural bacterioplankton communities. *Science Advances* 7(31), eabi4717.

Rakoff-Nahoum, S., K. R. Foster, and L. E. Comstock (2016). The evolution of cooperation within the gut microbiota. *Nature* 533(7602), 255–259.

Schuster, P. and K. Sigmund (1983). Replicator dynamics. *Journal of theoretical biology* 100(3), 533–538.

Sieben, A. J., J. R. Mihaljevic, and L. G. Shoemaker (2022). Quantifying mechanisms of coexistence in disease ecology. *Ecology* 103(12), e3819.

597 Slobodkin, L. B. (1964). The strategy of evolution. *American scientist* 52(3), 342–
598 357.

599 Smith, J. M. and G. R. Price (1973). The logic of animal conflict. *Nature* 246(5427),
600 15–18.

601 Souza, M. O., J. M. Pacheco, and F. C. Santos (2009). Evolution of cooperation
602 under N-person snowdrift games. *Journal of Theoretical Biology* 260(4), 581–588.

603 Steidinger, B. S. and J. D. Bever (2014). The coexistence of hosts with different
604 abilities to discriminate against cheater partners: an evolutionary game-theory
605 approach. *The American Naturalist* 183(6), 762–770.

606 Szabó, G. and G. Fath (2007). Evolutionary games on graphs. *Physics reports* 446(4-
607 6), 97–216.

608 Tarnita, C. E. and A. Traulsen (2025). Reconciling ecology and evolutionary game
609 theory or “When not to think cooperation”. *Proceedings of the National Academy*
610 *of Sciences* 122(14), e2413847122.

611 Taylor, P. D. and L. B. Jonker (1978). Evolutionary stable strategies and game
612 dynamics. *Mathematical biosciences* 40(1-2), 145–156.

613 Traulsen, A. and M. A. Nowak (2006). Evolution of cooperation by multilevel selec-
614 tion. *Proceedings of the National Academy of Sciences* 103(29), 10952–10955.

615 Trivers, R. L. (1971). The evolution of reciprocal altruism. *The Quarterly review of*
616 *biology* 46(1), 35–57.

617 Wang, G., Q. Su, L. Wang, and J. B. Plotkin (2025). Evolution of social behaviors
618 in noisy environments. *arXiv*.

619 Wedekind, C. and M. Milinski (2000). Cooperation through image scoring in humans.
620 *Science* 288(5467), 850–852.

621 West, S. A., G. A. Cooper, M. B. Ghoul, and A. S. Griffin (2021). Ten recent insights
622 for our understanding of cooperation. *Nature ecology & evolution* 5(4), 419–430.

623 West, S. A., A. S. Griffin, and A. Gardner (2007). Evolutionary explanations for
624 cooperation. *Current biology* 17(16), R661–R672.