

Strategic coexistence theory for evolutionary games

Sang Woo Park^{1,2,*}

1 School of Biological Sciences, Seoul National University, Seoul, Korea

2 Institute for Data Innovation in Science, Seoul National University, Seoul, Korea

*Corresponding author: sangwoopark@snu.ac.kr

Abstract

Evolutionary game theory and ecological coexistence theory both seek to predict the outcome of competition between biological entities, be they strategies or species, but the two fields have relied on largely separate approaches. Replicator equations provide a foundation for predicting the outcome of strategic competition, but comparing how different mechanisms alter competition across games remains challenging. Inspired by modern coexistence theory from community ecology, we develop Strategic Coexistence Theory (SCT), which translates payoff-based invasion conditions at monomorphic boundary conditions into strategic niche differences and fitness differences between any two competing strategies described by classic replicator dynamics. We show that SCT recovers the classic classification of two-strategy games and also allows us to compare mechanisms for the evolution of cooperation from different games on the shared niche-fitness difference space. In stochastic games, environmental stochasticity increases both boundary invasion growth rates, whereas in multiplayer public-goods games, nonlinear benefits can either stabilize or destabilize competition. These applications also show that the number or stability of internal equilibria generally cannot be predicted from invasion growth rates at monomorphic boundaries, illustrating a key limitation of characterizing competitive systems using boundary invasion rates alone. Together, these results show that SCT provides a coexistence-theoretic perspective for characterizing how different mechanisms alter strategy competition across games.

Significance statement

Evolutionary game theory provides a powerful framework for predicting the outcome of competition between strategies. However, comparing how different mechanisms contribute to different outcomes across games remains challenging. Motivated by Modern Coexistence Theory from community ecology, we developed Strategic Coexistence Theory (SCT), which decomposes invasion rates of strategies when rare into strategic niche and fitness differences. Applications to classic games, mechanisms of cooperation, environmental stochasticity, and multiplayer public-goods games place diverse models onto shared axes of niche and fitness differences. These applications

also point to a limitation to predicting coexistence from mutual invasion conditions at the boundary. This approach links ecological theory with replicator dynamics, providing a common language for comparing strategy competition across games.

Introduction

Understanding the mechanisms that promote the evolutionary stability of competing strategies is a fundamental aim in evolutionary biology (Lewontin, 1961; Slobodkin, 1964; Smith and Price, 1973; Taylor and Jonker, 1978). To this end, evolutionary game theory (EGT) and replicator dynamics have provided key foundations, allowing us to predict the outcome of strategic competition through payoff inequalities, invasion criteria, and stability analysis (Schuster and Sigmund, 1983; Hofbauer and Sigmund, 1998; Cressman, 2003; Cressman and Tao, 2014). Recently, there has been growing interest in characterizing similarities between replicator dynamics and the dynamics of competing species (Gokhale and Traulsen, 2025; Tarnita and Traulsen, 2025; Allesina, 2026), building on the classical mathematical equivalence between replicator equations and generalized Lotka–Volterra models (Hofbauer, 1981; Page and Nowak, 2002). In particular, Tarnita and Traulsen (2025) argued that EGT approaches for studying complex interactions may inform analyses of analogous ecological interactions, such as higher order and hierarchical interactions. Building on this renewed connection between ecological and evolutionary dynamics, we suggest that interpreting evolutionary games from the ecological perspective of strategy coexistence may enable comparisons of mechanisms in different games within a common framework.

Interestingly, even a simple two-strategy game can yield heterogeneous outcomes that mirror those in community ecology (Hauert, 2002; Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025): the dominance of a single strategy (prisoner’s dilemma and harmony games), coexistence (hawk–dove games), and bistability (stag–hunt games). Such differences in conditions that promote evolutionary stability and dominance of a single strategy and those that permit strategy coexistence indicate that EGT can be understood from the perspective of coexistence theory (Motro, 1991; Doebeli et al., 2004; Hauert and Doebeli, 2004; Doebeli and Hauert, 2005; Gore et al., 2009; Souza et al., 2009; Archetti and Scheuring, 2011; Steidinger and Bever, 2014; Tarnita and Traulsen, 2025). In fact, stable polymorphisms of evolutionary strategies are observed in microbial and other biological systems (Gore et al., 2009; Cordero et al., 2012; Kocher et al., 2018; Pollak et al., 2021; Lin et al., 2024), but a coexistence-theoretic perspective has often remained implicit. Recasting game dynamics as a coexistence problem immediately raises the following question: when two strategies coexist, does the corresponding mechanism stabilize or equalize competition (Chesson, 2000)? While EGT allows us to predict whether two strategies can coexist or not, a common framework for comparing how different mechanisms stabilize or equalize competition across different games remains elusive.

Modern coexistence theory (MCT) from community ecology presents a particularly useful framework for quantifying coexistence mechanisms (Chesson, 2000; Adler et al., 2007; Godoy et al., 2014; Godoy and Levine, 2014; Kraft et al., 2015; Adler et al., 2026). MCT predicts that coexistence of two competing entities, be they species or strategies, requires stabilizing niche differences to overcome average fitness differences (Chesson, 2000; Adler et al., 2007; Kraft et al., 2015). In community ecology, stabilizing niche differences represent differences in factors that limit the growth of competing species, causing each species to limit itself more than it limits its competitor (Chesson, 2000). In EGT, a strategy’s niche can be interpreted as strategic environments defined by the composition of competing strategies and the resulting interactions (Taylor and Jonker, 1978; Cressman and Tao, 2014; Tarnita and Traulsen, 2025). From the perspective of MCT, stabilizing niche differences arise when each strategy is favored under the monomorphic boundary condition created by its competitor than under the one created by itself, such that the spread of each strategy limits its own relative growth and favors the recovery of its competitor. In contrast, fitness differences represent differences in the ability of each entity to compete under the limiting environments created by its competitor (Chesson, 2000). This differs from payoff differences in EGT, which depend on the underlying strategy frequency and therefore combine both stabilizing effects caused by changes in the strategic environment and residual fitness differences that favor one strategy over another (Taylor and Jonker, 1978; Cressman and Tao, 2014; Tarnita and Traulsen, 2025).

So far, MCT has been primarily utilized for understanding species coexistence, but recent work showed that analogous framework can be derived for other biological systems, especially for studying pathogen strain competition, allowing one to directly quantify niche and fitness differences of competing entities, be they species or strains (Sieben et al., 2022; Park et al., 2024, 2026). Inspired by this effort, we present Strategic Coexistence Theory (SCT), which translates the invasion growth rates of competing strategies at monomorphic boundary conditions into strategic niche and fitness differences. In doing so, we apply SCT to four different case studies: classic two-strategy games, mechanisms for the evolution of cooperation, environmental stochasticity, and multiplayer public-goods games. For classic-game and stochasticity applications, we consider four different game types: Prisoner’s Dilemma, Hawk–Dove, Stag–Hunt, and Harmony. In the cooperation-mechanism application, prisoner’s dilemma serves as a common baseline for comparing five pairwise mechanisms. In contrast, the multiplayer public-goods application considers an explicit N -player game with nonlinearity effect on public goods.

Overall, these examples illustrate the utility of a coexistence-theoretic perspective for comparing mechanisms of strategy competition. However, since SCT is based on invasion growth rates at monomorphic boundary conditions, it does not provide a complete picture for the interior dynamics. Within this scope, SCT provides a common framework for linking coexistence theory in community ecology with evolutionary game theory.

Strategic coexistence theory

Here, we present a brief derivation of strategic coexistence theory (SCT) (Figure 1). To do so, we begin with a classic replicator equation, which allows us to model changes in the frequency of competing strategies based on their payoffs (Schuster and Sigmund, 1983; Hofbauer and Sigmund, 1998; Cressman, 2003; Cressman and Tao, 2014):

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (1)$$

where x_i represents the frequency of strategy i , $\pi_i(\mathbf{x})$ represents the payoff of strategy i for a given strategy distribution $\mathbf{x}$, and $\bar{\pi}(\mathbf{x}) = \sum_i \pi_i(\mathbf{x})x_i$ represents the average payoff. Since strategy frequencies must sum to 1 (i.e., $\sum_i x_i = 1$), a single-strategy equilibrium dominated by strategy i can be described by a unit vector $\mathbf{e}_i$, where all elements are equal to zero except for the i -th element. We use these single-strategy equilibria to define niche and fitness differences between two competing strategies i and j .

Because payoffs in many games can be negative, payoff ratios can be difficult to interpret. Thus, we work on a positive multiplicative scale by exponentiating the payoff terms:

$$W_i(\mathbf{x}) = \exp(\pi_i(\mathbf{x})). \quad (2)$$

This exponential transformation is convenient because it preserves the payoff ordering and does not alter the invasion conditions. Therefore, this allows us to compare strategic environments through ratios, as in modern coexistence theory. For convenience, we refer to $W_i(\mathbf{x})$ as the multiplicative payoff of strategy i .

First, we define the niche overlap ρ between strategies i and j as the ratio between the geometric mean of each strategy's multiplicative payoff in the strategic environment that it creates when common, $\sqrt{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}$, and the geometric mean of each strategy's multiplicative payoff in the environment generated by its competitor, $\sqrt{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}$. The corresponding niche difference is then given by

$$1 - \rho = 1 - \sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}}. \quad (3)$$

When $1 - \rho > 0$, the geometric mean of the multiplicative payoffs across the two strategies is greater in competitor-generated monomorphic environments than in self-generated monomorphic environments. In other words, intraspecific competition is stronger than interspecific competition at the boundary. When $\rho > 1$, we interpret the resulting negative niche difference, $1 - \rho < 0$, as a destabilizing effect corresponding to positive frequency dependence, meaning that the strategy that is already common is favored.

Second, we define the fitness difference as the ratio between the geometric means of the multiplicative payoff of each strategy across the two single-strategy environ-




Figure 1: **A schematic diagram summarizing strategic coexistence theory.** Strategic coexistence theory decomposes pairwise replicator dynamics into stabilizing and equalizing mechanisms. Mutual invasion conditions can be rewritten on a multiplicative payoff scale, allowing us to derive strategic niche difference $1 - \rho$ and fitness difference κ_j/κ_i based on multiplicative payoff ratios. Then, the inequality $\rho < \kappa_j/\kappa_i < 1/\rho$ determines mutual invasion. We note that the exact values of niche and fitness differences are sensitive to payoff scaling, while dynamical outcomes are still preserved.

ments:

$$\frac{\kappa_j}{\kappa_i} = \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}. \quad (4)$$

In other words, strategy j has an average competitive advantage over strategy i ($\kappa_j/\kappa_i > 1$) when it performs better, on average, across the environments generated by itself and its competitor.

We note that niche and fitness difference can be equivalently written in terms of the invasion growth rates at the monomorphic boundary conditions:

$$1 - \rho = 1 - \exp\left(-\frac{r_{i|j} + r_{j|i}}{2}\right), \quad (5)$$

$$\frac{\kappa_j}{\kappa_i} = \exp\left(\frac{r_{j|i} - r_{i|j}}{2}\right), \quad (6)$$

where $r_{i|j} = \pi_i(\mathbf{e}_j) - \pi_j(\mathbf{e}_j)$ and $r_{j|i} = \pi_j(\mathbf{e}_i) - \pi_i(\mathbf{e}_i)$ represent the growth rate of strategy i and j when strategy j and i are at monomorphic equilibrium, respectively. These growth rates also correspond to payoff differences. This parameterization is particularly useful because it allows us to compute niche and fitness differences even when the selection gradient governing replicator dynamics do not uniquely separate into two payoff values.

167 Together, for any two strategies described by the classic replicator equation, mu-
 168 tual invasion requires stabilizing niche differences to overcome average fitness differ-
 169 ences. Specifically, the condition for mutual invasion can be rewritten as (Chesson,
 170 2000):

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (7)$$

171 In simple systems with monotonic interior dynamics, these boundary invasion con-
 172 ditions also determine the global dynamical outcome, allowing competition to be
 173 classified into coexistence, competitive-exclusion, and priority-effect regimes. How-
 174 ever, MCT and SCT are based on invasion conditions at the boundaries and do not
 175 generally characterize nonlinear interior dynamics. Mutual invasion still guarantees
 176 at least one stable internal equilibrium, but it does not determine the number or
 177 stability of additional internal equilibria. Likewise, boundary conditions associated
 178 with competitive exclusion or priority effects do not rule out stable internal equilib-
 179 ria. As we show later, nonlinear dynamics can allow stable coexistence even when a
 180 system lies in the boundary-defined competitive-exclusion regime. Therefore, when
 181 interior dynamics are nonmonotonic, we use these classifications to describe bound-
 182 ary invasibility rather than complete dynamical outcomes.

183 Despite this limitation, the mutual-invasion inequality provides a useful perspec-
 184 tive for comparing how different mechanisms alter boundary invasibility. Notably,
 185 the inequality can be applied to any pairwise comparison between two strategies
 186 described by the classic replicator equation because ρ and κ_j/κ_i can be calculated
 187 directly from ratios of multiplicative payoffs. This provides a common framework
 188 for comparing qualitative changes in boundary invasibility across different games.
 189 Detailed derivations are provided in Supplementary Text S1.

190 We note that the absolute magnitudes of niche and fitness differences are sen-
 191 sitive to payoff scaling because they are defined on an exponentiated payoff scale.
 192 Therefore, while SCT allows us to compare different games on shared axes, we can-
 193 not compare the exact values of niche and fitness differences across different games.
 194 Nonetheless, as we show in Supplementary Text S2, the qualitative outcome of strate-
 195 gic competition under SCT is preserved because payoff rescaling preserves payoff
 196 orderings. Therefore, we focus on comparing qualitative outcomes from SCT, in-
 197 cluding directions of change in niche and fitness difference and resulting dynamical
 198 regimes, and refrain from making any comparisons of magnitude of niche and fitness
 199 difference values.

200 Classic two-strategy games

201 As an illustrative example, we begin with classic two-strategy games (Hauert, 2002;
 202 Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025), which can be

described by the following payoff matrix (Figure 2A):

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (8)$$

Here, the first and second rows represent strategies 1 and 2, respectively, and the columns represent the strategy of the interacting opponent. Writing x_1 to denote the frequency of strategy 1, the expected payoffs are

$$\pi_1(\mathbf{x}) = Rx_1 + S(1 - x_1), \quad (9)$$

$$\pi_2(\mathbf{x}) = Tx_1 + P(1 - x_1). \quad (10)$$

Thus, the dynamics can be written as

$$\frac{dx_1}{dt} = x_1(1 - x_1) [(R - T)x_1 + (S - P)(1 - x_1)]. \quad (11)$$

The signs of $T - R$ and $S - P$ determine whether each strategy can invade the other when rare. Specifically, strategy 2 can invade strategy 1 when $T > R$, whereas strategy 1 can invade strategy 2 when $S > P$. These two invasion conditions partition the game space into four regimes (Figure 2B): prisoner's dilemma ($T > R > P > S$), hawk-dove game ($T > R, S > P$), stag-hunt game ($R > T, P > S$), and harmony game ($R > T, S > P$). In standard EGT, these outcomes are obtained by analyzing the stability of equilibrium conditions. For simple games, this analysis is straightforward.

A coexistence-theoretic approach recovers the same classification (Figure 2C). For a two-strategy game determined by this payoff matrix, the niche difference and fitness difference are given by the following set of equations (Supplementary Text S3):

$$1 - \rho = 1 - \exp\left(\frac{R + P - S - T}{2}\right), \quad (12)$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{T + P - R - S}{2}\right). \quad (13)$$

As shown above, mutual invasion requires $\rho < \kappa_2/\kappa_1 < 1/\rho$, which is equivalent to $T > R$ and $S > P$. This corresponds to the hawk-dove game, in which each strategy can invade its competitor when rare and the two strategies coexist. The remaining games also have natural interpretations in niche-fitness difference space. Prisoner's dilemma and harmony games represent competitive exclusion regimes, where one strategy has a fitness advantage that exceeds niche differences. In contrast, stag-hunt games represent a priority-effect regime, where neither strategy can invade the other and the outcome depends on initial conditions. Therefore, coexistence theory not only allows us to recover the classic results from two-strategy games but also expresses these results in terms of niche and fitness differences. Building on this, we next apply this framework to comparing core mechanisms of cooperation.




Figure 2: **Coexistence-theoretic interpretation of classic two-strategy games.** (A) General payoff matrix for a two-strategy game, where rows represent the focal strategy and columns represent the strategy of the interacting opponent. (B) Classification of four classic games based on the payoff differences $T - R$ and $S - P$. Prisoner's dilemma and harmony games lead to the dominance of one strategy; hawk-dove games allow coexistence; and stag-hunt games result in bistability. (C) Mapping of the same four games onto niche-fitness difference space. Hawk-dove games correspond to the coexistence regime, where stabilizing niche differences overcome fitness differences. Prisoner's dilemma and harmony games correspond to competitive-exclusion regimes, whereas stag-hunt games correspond to the priority-effect regime. Negative values of $1 - \rho$ indicate positive frequency dependence and destabilization. $R > P$ is assumed throughout, but this does not affect coexistence outcome in any way.

Comparisons of mechanisms for the evolution of co-operation

Understanding mechanisms that promote the evolution of cooperation is a central goal in evolutionary biology (West et al., 2007; Fletcher and Doebeli, 2008; Clutton-Brock, 2009; Kingma et al., 2014; Rakoff-Nahoum et al., 2016; Hilbe et al., 2018; Apicella and Silk, 2019; West et al., 2021; Lee et al., 2022; Palmer and Foster, 2022; Michel-Mata et al., 2024). Here, we compare five classic mechanisms for the evolution of cooperation using SCT (Nowak, 2006) and ask how they differ from the perspective of SCT. Specifically, we consider the following mechanisms: kin selection, direct reciprocity, indirect reciprocity, network reciprocity, and group selection (Figure 3A). Following Nowak (2006), these five mechanisms are formulated as modifications of a pairwise Prisoner's Dilemma. We use the Prisoner's Dilemma as a common baseline for comparing these specific formulations in this case study, rather than as a general model of cooperation.

245 Kin selection favors cooperation when two interacting individuals are genetic rel-
 246 atives, where the genetic relatedness r must be greater than the cost-to-benefit ratio:
 247 $r > c/b$ (Hamilton, 1964). Direct reciprocity favors cooperation when current co-
 248 operation may promote future cooperation by the same individual through repeated
 249 encounters (Trivers, 1971; Axelrod and Hamilton, 1981). In this case, the probability
 250 of repeated interaction (i.e., “next round”) must be greater than the cost-to-benefit
 251 ratio: $w > c/b$. Indirect reciprocity favors cooperation when current cooperation
 252 may promote future cooperation by different individuals through reputation (Nowak
 253 and Sigmund, 1998; Wedekind and Milinski, 2000; Nowak and Sigmund, 2005). Thus,
 254 the social acquaintanceship (i.e., the probability of knowing someone’s reputation)
 255 must be greater than the cost-to-benefit ratio: $q > c/b$. Network reciprocity favors
 256 cooperation via clustering of cooperators, requiring the benefit-to-cost ratio to be
 257 greater than the average number of neighbors: $b/c > k$ (Nowak and May, 1992;
 258 Lieberman et al., 2005). Finally, group selection favors cooperation via multilevel
 259 selection, where groups with more cooperators exhibit faster growth at the group
 260 level even if defectors reproduce faster at the individual level (Traulsen and Nowak,
 261 2006). This requires $b/c > 1 + (n/m)$, where n is the maximum group size and
 262 m is the number of groups. All of these mechanisms can be described by a simple
 263 2×2 payoff matrix (Nowak, 2006), from which the corresponding niche and fitness
 264 differences can be computed using Eq. (12) and Eq. (13). Details on niche and fitness
 265 difference calculations are presented in Supplementary Text S4.

266 First, the prisoner’s dilemma provides a baseline scenario, where defection is
 267 evolutionarily stable (Figure 3A). From the perspective of SCT, there is complete
 268 niche overlap, and therefore defection competitively excludes cooperation (Figure
 269 3B). Building on this, SCT shows that five mechanisms promote evolutionary sta-
 270 bility of cooperation via different dynamical routes (Figure 3B). For example, kin
 271 selection allows the fitness of cooperation to increase as genetic relatedness r increases
 272 (Figure 3B). However, it does not affect the niche difference between cooperation and
 273 defection. Similarly, network reciprocity and group selection modulate the fitness of
 274 cooperation without causing any changes in strategic niches. In contrast, direct
 275 and indirect reciprocity destabilize competition while increasing the fitness of coop-
 276 eration, thereby pushing the competition outcome towards a priority effect regime.
 277 Equivalently, these two mechanisms promote bistability of defection and cooperation.

278 SCT further illustrates another difference between direct and indirect reciprocity.
 279 Direct reciprocity remains within the priority-effect regime as w reaches 1, whereas
 280 indirect reciprocity reaches the boundary between the priority-effect and competitive-
 281 exclusion regimes at $q = 1$ (Supplementary Text S4).

282 We note that none of these mechanisms allow coexistence of cooperation and
 283 defection. In many biological systems, such coexistence is often observed (Gore et al.,
 284 2009; Cordero et al., 2012; Kocher et al., 2018; Pollak et al., 2021; Lin et al., 2024).
 285 Therefore, we move on to the next example, where environmental stochasticity can
 286 promote coexistence of cooperators and defectors.

(A)

Prisoner's dilemma	Kin selection
$\begin{pmatrix} b-c & -c \\ b & 0 \end{pmatrix}$	$\begin{pmatrix} (b-c)(1+r) & br-c \\ b-rc & 0 \end{pmatrix}$
Direct reciprocity	Indirect reciprocity
$\begin{pmatrix} (b-c)/(1-w) & -c \\ b & 0 \end{pmatrix}$	$\begin{pmatrix} b-c & -c(1-q) \\ b(1-q) & 0 \end{pmatrix}$
Network reciprocity	Group selection
$\begin{pmatrix} b-c & H-c \\ b-H & 0 \end{pmatrix}$	$\begin{pmatrix} (b-c)(m+n) & (b-c)m-cn \\ bn & 0 \end{pmatrix}$

(B)




Figure 3: **Qualitative comparisons of five mechanisms for the evolution of cooperation using SCT.** (A) Payoff matrices for the baseline scenario (prisoner's dilemma) and five mechanisms for the evolution of cooperation (Nowak, 2006). Prisoner's dilemma is characterized by the benefit for the recipient b and the cost for the donor c . Network reciprocity is parameterized based on $H = [(b-c)k - 2c]/[(k+1)(k-2)]$. Parameters for all other mechanisms are described in panel B. (B) Changes in niche and fitness differences driven by each mechanism. Each arrow represents the direction of change as we increase the corresponding parameter, indicated in figure legends. Note that kin selection, network reciprocity, and group selection all result in changes in fitness difference without causing any changes in niche difference. Therefore, we indicate the direction of change for network reciprocity and group selection separately to avoid overlapping arrows. For simplicity, we focus on the effect of number of groups m for group selection; increasing the number of maximum group size n has the opposite effect (Supplementary Text S4). Across all models, we assume $b = 3$ and $c = 1$. Because SCT decompositions depend on positive payoff scaling, exact values of niche and fitness difference (e.g., arrow lengths, slopes, and distances from regime boundaries) should not be compared across mechanisms.

Can stochasticity prevent coexistence?

So far, we have focused on simple deterministic games. However, many ecological and evolutionary processes are intrinsically stochastic. Recently, Wang et al. (2025)

showed that environmental fluctuations can promote coexistence in games that would otherwise lead to competitive exclusion in a deterministic environment. From the perspective of SCT, this suggests that stochasticity can stabilize competition by promoting the recovery of strategies when rare. Interestingly, Wang et al. (2025) also showed that stochasticity can generate bistable coexistence and multiple internal equilibria. Thus, although stochasticity can stabilize competition near the boundaries, it can also generate destabilizing frequency dependence in the interior. This immediately raises the question of whether stochasticity can generate sufficiently strong positive frequency dependence to destabilize competition, thus pushing the system out of the coexistence regime. Here, we use SCT to separate these effects and clarify whether stochasticity stabilizes or destabilizes strategic competition (Supplementary Text S5). Although we refer to the two strategies as cooperation and defection for consistency, this analysis is not restricted to the Prisoner's Dilemma and can be applied to any 2×2 games.

Following Wang et al. (2025), we consider a 2×2 game in the presence of environmental noise. The stochastic payoff matrix is given by

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix} + \frac{\zeta}{\sqrt{s}} \begin{pmatrix} \sigma_R & \sigma_S \\ \sigma_T & \sigma_P \end{pmatrix}, \quad (14)$$

where the first matrix represents deterministic payoffs, ζ represents a noise term with zero mean and unit variance, and the second matrix determines the intensity of environmental stochasticity. Assuming weak selection $s \ll 1$, the evolution of two competing strategies in a large population can be described by the following equation:

$$\frac{dx}{dt} = sx(1-x) \left(\pi_C - \pi_D + \left(\frac{1}{2} - x \right) (\sigma_C - \sigma_D)^2 \right), \quad (15)$$

where $\pi_C = Rx + S(1-x)$ and $\pi_D = Tx + P(1-x)$ and $\sigma_C = \sigma_R x + \sigma_S(1-x)$ and $\sigma_D = \sigma_T x + \sigma_P(1-x)$. After rescaling time as $\tau = st$, the invasion growth rates of each strategy when rare can be written as:

$$r_C = S - P + \frac{1}{2}(\sigma_S - \sigma_P)^2, \quad (16)$$

$$r_D = T - R + \frac{1}{2}(\sigma_T - \sigma_R)^2. \quad (17)$$

This shows that invasion growth rates under environmental stochasticity will be greater than or equal to their deterministic counterparts, meaning that environmental stochasticity at the boundaries is intrinsically stabilizing.

Applying SCT, niche overlap and fitness difference for this system can be written as:

$$\rho = \exp \left(\frac{R + P - S - T}{2} - \frac{(\sigma_R - \sigma_T)^2 + (\sigma_S - \sigma_P)^2}{4} \right) \quad (18)$$

$$\frac{\kappa_D}{\kappa_C} = \exp \left(\frac{T + P - R - S}{2} + \frac{(\sigma_R - \sigma_T)^2 - (\sigma_S - \sigma_P)^2}{4} \right) \quad (19)$$

319 Writing $\hat{\rho}$ and $\hat{\kappa}_D/\hat{\kappa}_C$ to denote niche overlap and fitness difference in the absence of
 320 stochasticity and taking logs of both sides for convenience, we have:

$$\Delta \log \rho = \log \rho - \log \hat{\rho} = -\frac{(\sigma_R - \sigma_T)^2 + (\sigma_S - \sigma_P)^2}{4} \quad (20)$$

$$\Delta \log \frac{\kappa_D}{\kappa_C} = \log \frac{\kappa_D}{\kappa_C} - \log \frac{\hat{\kappa}_D}{\hat{\kappa}_C} = \frac{(\sigma_R - \sigma_T)^2 - (\sigma_S - \sigma_P)^2}{4}. \quad (21)$$

321 Therefore, stochasticity reduces niche overlap and increases niche difference. In con-
 322 trast, stochasticity can either amplify or reduce fitness differences depending on which
 323 strategy's invasion rate is more strongly affected by environmental fluctuations. No-
 324 tably, changes in fitness difference driven by environmental stochasticity will always
 325 be bounded by those in niche overlap:

$$\Delta \log \rho \leq \Delta \log \frac{\kappa_D}{\kappa_C} \leq -\Delta \log \rho \quad (22)$$

326 This result implies that environmental stochasticity cannot generate sufficient fitness
 327 difference to push the system out of the coexistence regime.

328 As an example, consider a scenario where the amount of stochasticity is assumed
 329 to be proportional to deterministic payoffs (Wang et al., 2025):

$$\begin{pmatrix} \sigma_R & \sigma_S \\ \sigma_T & \sigma_P \end{pmatrix} = k \begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (23)$$

330 Following previous case studies, we consider four deterministic payoff structures (Fig-
 331 ure 4A): Prisoner's Dilemma (PD), Hawk-Dove Game (HD), Stag-Hunt Game (SH)
 332 and Harmony Game (HG). Across all these examples, stochasticity eventually pushes
 333 the system inside the mutual invasion regime (Figure 4B). Notably, changes in dy-
 334 namical regimes predicted by SCT accurately capture bifurcations occurring at the
 335 boundaries (Figure 4C): at sufficiently large k , both boundary equilibria are unstable
 336 in all four games. Notably, for the SH game, the system begins from the priority
 337 effect regime and enters the competitive exclusion regime before eventually entering
 338 the mutual invasion regime (Figure 4B). Consequently, $x = 1$ (pure cooperation)
 339 becomes unstable first, and then $x = 0$ (pure defection) also becomes unstable.

340 This example also illustrates an important limitation shared by MCT and SCT.
 341 Since MCT and SCT solely rely on boundary conditions, they cannot predict the
 342 presence of multiple internal equilibria or their stability. In the SH example, there is
 343 an unstable internal equilibrium across all ranges of k we consider (Figure 4C). When
 344 the system enters the competitive-exclusion regime, an additional stable internal
 345 equilibrium appears, allowing coexistence. Furthermore, when mutual invasion is
 346 allowed, two stable coexistence equilibria are present. Therefore, while MCT and
 347 SCT accurately capture the behavior at the boundary via invasibility, they do not
 348 fully characterize the interior dynamics or all conditions under which coexistence is
 349 possible.




Figure 4: **Effects of environmental stochasticity on strategic competition.** (A) Assumed deterministic payoff structures across four scenarios. Environmental stochasticity is assumed to be proportional to these payoffs, scaled by a term k . (B) Arrows indicate the direction of changes in dynamical regime as the amount of stochasticity k increases. (C) Bifurcation diagrams under each scenario as a function of k . Solid and dashed lines represent stable and unstable equilibria, respectively.

350 Evolution of cooperation in multiplayer, nonlinear 351 public goods games

352 Previous case studies focused on evolutionary games based on pairwise interactions,
353 but real-world behaviors, especially cooperation, often involve interactions among

multiple players (Motro, 1991; Archetti and Scheuring, 2012). In particular, co-operators can contribute to shared public goods, from which other individuals can benefit. As our final study, we consider an N-player public-goods game to investigate the effects of multiplayer interaction on strategy competition. In doing so, we also consider the effect of nonlinear public goods and ask when nonlinearity can stabilize or destabilize competition.

To model the evolution of cooperation under multiplayer interactions, we begin by assuming that every individual receives the same amount of benefit $\beta(i)$ which increases monotonically with the number of cooperators i , and that cooperation incurs a constant cost $c > 0$. Assuming that each individual interacts with $N - 1$ other selected individuals, the average payoff for cooperators $\pi_C(x)$ and defectors $\pi_D(x)$ depends on the probability $f_j(x)$ of finding j cooperators in the group, given x proportion of cooperators in the population (Archetti and Scheuring, 2012):

$$\pi_C(x) = \sum_{j=0}^{N-1} f_j(x) \beta(j+1) - c \quad (24)$$

$$\pi_D(x) = \sum_{j=0}^{N-1} f_j(x) \beta(j) \quad (25)$$

Assuming random, homogeneous mixing in a large population, the probability $f_j(x)$ follows a binomial distribution:

$$\pi_C(x) = \sum_{j=0}^{N-1} \binom{N-1}{j} x^j (1-x)^{N-1-j} \beta(j+1) - c \quad (26)$$

$$\pi_D(x) = \sum_{j=0}^{N-1} \binom{N-1}{j} x^j (1-x)^{N-1-j} \beta(j) \quad (27)$$

The resulting replicator equation corresponds to:

$$\frac{dx}{dt} = x(1-x)G(x), \quad (28)$$

where $G(x) = \pi_C(x) - \pi_D(x)$ represents the payoff difference.

While this game consists of interactions between multiple individuals, from the perspective of SCT, there are still two strategies competing. Thus, we can compute niche overlap ρ and fitness difference κ_D/κ_C using standard SCT, which in turn can be simplified in terms of payoff differences at the boundary:

$$\rho = \exp \left(\frac{\pi_C(1) + \pi_D(0) - \pi_C(0) - \pi_D(1)}{2} \right) = \exp \left(\frac{G(1) - G(0)}{2} \right) \quad (29)$$

$$\frac{\kappa_D}{\kappa_C} = \exp \left(\frac{\pi_D(0) + \pi_D(1) - \pi_C(1) - \pi_C(0)}{2} \right) = \exp \left(-\frac{G(1) + G(0)}{2} \right). \quad (30)$$

375 Here, the payoff values at the boundary correspond to:

$$G(0) = \beta(1) - \beta(0) - c \quad (31)$$

$$G(1) = \beta(N) - \beta(N-1) - c \quad (32)$$

376 Finally, mutual invasion at the boundary requires:

$$\rho < \frac{\kappa_D}{\kappa_C} < \frac{1}{\rho}. \quad (33)$$

377 This expression can be equivalently written as:

$$G(1) < 0 < G(0) \iff \beta(N) - \beta(N-1) < c < \beta(1) - \beta(0). \quad (34)$$

378 This formulation shows that mutual invasion is possible only when the marginal ben-
 379 efit generated by the first cooperator exceeds that generated by the N th cooperator,
 380 with the cost of cooperation lying between these two marginal benefits.

381 We can now draw three conclusions from this observation. First, when the benefit
 382 of public good $\beta(i)$ increases linearly with i (thus $\beta(1) - \beta(0) = \beta(N) - \beta(N-1)$),
 383 then there is complete niche overlap because $G(1) = G(0)$. In this case, cooperators
 384 competitively exclude defectors when the marginal benefit $\beta(1) - \beta(0)$ is greater than
 385 the cost c , and defectors competitively exclude cooperators when the marginal benefit
 386 is smaller than the cost. Stable coexistence becomes impossible. Second, when the
 387 benefit of public good $\beta(i)$ is a concave, increasing function of i (i.e., $d\beta/di > 0$ and
 388 $d^2\beta/di^2 < 0$), we have $\beta(1) - \beta(0) > \beta(N) - \beta(N-1)$ (Motro, 1991). This means
 389 that nonlinearity can generate niche difference and allow coexistence when

$$\beta(N) - \beta(N-1) < c < \beta(1) - \beta(0) \quad (35)$$

390 This generalizes the result of Gore et al. (2009), showing that the effect of concave
 391 nonlinearity on coexistence is robust to multiplayer extension (see Supplementary
 392 Text S6 and Figure S1 for detailed analyses). Third, when the benefit of public
 393 good $\beta(i)$ is a convex, increasing function of i (i.e., $d\beta/di > 0$ and $d^2\beta/di^2 > 0$),
 394 we have $\beta(1) - \beta(0) < \beta(N) - \beta(N-1)$ (Motro, 1991). In this case, nonlinearity
 395 can allow bistability of two boundary monomorphic equilibria and generate priority
 396 effect when

$$\beta(1) - \beta(0) < c < \beta(N) - \beta(N-1). \quad (36)$$

397 Therefore, SCT shows that nonlinearity can either stabilize or destabilize strategic
 398 coexistence depending on the shape of nonlinearity. Accordingly, $G(x)$ decreases
 399 monotonically for concave benefits and increases monotonically for convex benefits,
 400 so these cases can have at most one internal equilibrium.

401 We note that this multiplayer game can also permit multiple internal equilibria.
 402 For example, Archetti and Scheuring (2012) considers a scenario where $\beta(t)$ follows a
 403 sigmoid function, illustrating that coexistence between cooperators and defectors is
 404 possible even when defectors have higher payoffs at both boundaries. In such cases,
 405 SCT will be able to capture invasibility at the boundary but will not be able to
 406 predict internal dynamics, as we show in the previous case study.

Discussion

Standard evolutionary game theory and replicator equations provide powerful tools for predicting the outcome of strategy competition, including strategic dominance, coexistence, and bistability (Motro, 1991; Doebeli et al., 2004; Hauert and Doebeli, 2004; Doebeli and Hauert, 2005; Gore et al., 2009; Souza et al., 2009; Archetti and Scheuring, 2011; Steidinger and Bever, 2014; Tarnita and Traulsen, 2025). However, these outcomes are typically understood by analyzing payoff differences within individual games, which makes it difficult to compare coexistence mechanisms across models. Building on Chesson’s Modern Coexistence Theory from community ecology (Chesson, 2000), we developed a strategic coexistence theory (SCT) that allows us to decompose strategy competition at the invasion boundaries in terms of stabilizing niche differences and fitness differences.

To demonstrate the utility of SCT, we considered four case studies. First, the analysis of classic two-strategy games showed that SCT recovers the original classification for these classic games, highlighting the similarities between evolutionary games and community ecology (Hauert, 2002; Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025). Second, we compared different mechanisms for the evolution of cooperation, which showed that each mechanism promotes cooperation via different dynamical routes (Nowak, 2006). Kin selection, network reciprocity, and group selection primarily affect the fitness difference between cooperation and defection, whereas direct and indirect reciprocity can destabilize competition and generate priority effects. Third, we considered the effect of environmental stochasticity on strategy competition, showing that stochasticity cannot destabilize competition. Finally, a case study of multiplayer, public-goods games showed that nonlinear public benefit can either stabilize or destabilize competition. The final two applications also revealed an important limitation of analyses based on invasion growth rates at monomorphic boundary conditions: they are unable to predict the number or stability of interior equilibria. We note that SCT is not intended to produce dynamical predictions that differ from standard analyses in EGT. Instead, SCT provides a common visualization and expository framework for comparing how different mechanisms alter boundary invasibility through changes in strategic niche and fitness differences.

There are several limitations to SCT. First, similar to MCT in community ecology, SCT only allows for comparisons of pairwise strategies and therefore cannot determine the outcomes of games with more than two strategies (Chesson, 2000). While SCT can be still applied to multi-player interactions, conclusions based on 2-player games may not be robust to multi-player extensions, which can introduce nonlinearity. Thus, our analysis of cooperation mechanisms should be interpreted as specific to the pairwise formulations considered here and as an illustration of SCT, rather than as general conclusions about multiplayer versions of these mechanisms.

Second, the exact values of strategic niche and fitness differences are sensitive to payoff scaling. SCT therefore supports quantitative comparisons only when models share a justified payoff normalization. Without such normalization, comparisons

are restricted to qualitative changes in dynamical regimes, directions, and boundary crossings.

Third, we focused on classic replicator equations, but Tarnita and Traulsen (2025) recently pointed out that these classic formulations implicitly ignore differences in intrinsic growth rates. In such cases, replicator equations may fail to predict the outcome of competition (Tarnita and Traulsen, 2025), and a full Lotka-Volterra model is needed to describe the abundances, rather than frequencies, of strategies. Thus, SCT inherits the same assumptions as the classic replicator framework: it is most appropriate when strategy competition can be described in terms of relative frequencies alone (Tarnita and Traulsen, 2025). When strategies differ intrinsically in growth, MCT may be required instead to tease apart mechanisms of strategy coexistence.

Finally, we have focused on simple examples to illustrate the utility of SCT, but many games involve additional complexities, including population structure (Nowak et al., 2010) and explicit ecological feedback (Weitz et al., 2016; Tarnita and Traulsen, 2025). Extending SCT to these settings will be necessary to determine how broadly stabilizing and equalizing mechanisms can be compared across evolutionary games.

While SCT was inspired by MCT, there are important differences between the two frameworks. In particular, MCT describes changes in species abundance, whereas SCT describes changes in strategy frequency. For example, even when growth rates of the abundances of both competitors are negative, the growth rate of the frequency of a superior competitor will still be positive; this implies that the invasibility in frequency and abundance space are fundamentally different. Moreover, MCT relies on a simple, Lotka-Volterra model, which permits at most one internal equilibrium, which allows us to predict coexistence from mutual invasion. However, SCT relies on replicator dynamics, which can permit multiple internal equilibria, meaning that criterion for mutual invasion and coexistence can differ. Therefore, SCT should be interpreted as a decomposition framework that is inspired by MCT, rather than a direct extension.

In conclusion, SCT provides a bridge between evolutionary game theory and community ecology, allowing strategy competition to be understood from the perspective of coexistence theory. Therefore, rather than replacing existing EGT or replicator-dynamics frameworks, SCT provides a complementary tool for teasing apart the coexistence mechanisms underlying evolutionary games. More broadly, our study builds on recent efforts to extend MCT beyond species competition (Park et al., 2024, 2026), providing a foundation for building unifying coexistence theory for competing entities, including species, pathogen strains, and behavioral strategies.

Acknowledgement

We thank Jonathan Levine for the helpful discussion. S.W.P. was supported by the New Faculty Startup Fund from Seoul National University, the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT)

489 (RS-2026-25474574), and the Global-LAMP Program of the National Research Foun-
490 dation of Korea (NRF) grant funded by the Ministry of Education (No. RS-2023-
491 00301976).

492 **Competing interests**

493 The authors declare no competing interest.

494 **Data availability**

495 All data and code are stored in a publicly available GitHub repository (<https://github.com/parklab-snu/coexistence-strategy>).
496

Supplementary Text

S1 Derivation of the strategic coexistence theory

To derive strategic coexistence theory (SCT), we begin with replicator equations:

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S1})$$

where x_i represents the frequency of strategy i , π_i represents the payoff of strategy i , and $\bar{\pi}$ represents the average payoff. Then, the mutual invasion condition for two competing strategies i and j can be written as

$$\pi_j(\mathbf{e}_j) < \pi_i(\mathbf{e}_j), \quad (\text{S2})$$

$$\pi_i(\mathbf{e}_i) < \pi_j(\mathbf{e}_i). \quad (\text{S3})$$

These inequalities will be preserved when we exponentiate the payoffs, $W_i(\mathbf{x}) = \exp(\pi_i(\mathbf{x}))$:

$$W_j(\mathbf{e}_j) < W_i(\mathbf{e}_j), \quad (\text{S4})$$

$$W_i(\mathbf{e}_i) < W_j(\mathbf{e}_i). \quad (\text{S5})$$

Rearranging, we obtain the following inequality:

$$\frac{W_i(\mathbf{e}_i)}{W_j(\mathbf{e}_i)} < 1 < \frac{W_i(\mathbf{e}_j)}{W_j(\mathbf{e}_j)}. \quad (\text{S6})$$

By multiplying $\sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}$ on both sides, we obtain:

$$\sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}} < \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}} < \sqrt{\frac{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}} \quad (\text{S7})$$

Thus, by defining niche overlap ρ and fitness difference κ_j/κ_i as follows,

$$\rho = \sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}}, \quad (\text{S8})$$

$$\frac{\kappa_j}{\kappa_i} = \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}, \quad (\text{S9})$$

we obtain the following inequality that defines conditions for mutual invasion:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (\text{S10})$$

509 Note that mutual invasion condition can be also written in terms of invasion
 510 growth rates at monomorphic equilibrium $r_{i|j}$ and $r_{j|i}$:

$$0 < r_{i|j}, \quad (\text{S11})$$

$$0 < r_{j|i}. \quad (\text{S12})$$

511 Analogous to the previous derivation, the mutual condition can be then rewritten
 512 as:

$$-\frac{r_{j|i} + r_{j|i}}{2} < \frac{r_{j|i} - r_{j|i}}{2} < \frac{r_{i|j} + r_{j|i}}{2}. \quad (\text{S13})$$

513 Exponentiating all sides preserves the inequality and keep them positive:

$$\exp\left(-\frac{r_{j|i} + r_{j|i}}{2}\right) < \exp\left(\frac{r_{j|i} - r_{j|i}}{2}\right) < \exp\left(\frac{r_{i|j} + r_{j|i}}{2}\right). \quad (\text{S14})$$

514 Thus, by defining niche overlap ρ and fitness difference κ_j/κ_i as follows,

$$\rho = \exp\left(-\frac{r_{j|i} + r_{j|i}}{2}\right), \quad (\text{S15})$$

$$\frac{\kappa_j}{\kappa_i} = \exp\left(\frac{r_{j|i} - r_{j|i}}{2}\right), \quad (\text{S16})$$

515 we obtain the following inequality that defines conditions for mutual invasion:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (\text{S17})$$

516 **S2 Niche and fitness difference payoff scaling**

517 Consider a classic replicator equation:

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S18})$$

518 where x_i represents the frequency of strategy i , π_i represents the payoff of strategy
 519 i , and $\bar{\pi}$ represents the average payoff. In this case, multiplying the payoffs by a
 520 constant $s > 0$ only changes time and does not affect the overall dynamics:

$$\frac{dx_i}{dt} = x_i(\pi'_i(\mathbf{x}) - \bar{\pi}'(\mathbf{x})) \quad (\text{S19})$$

$$= sx_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S20})$$

521 where $\pi'_i(\mathbf{x}) = s\pi_i(\mathbf{x})$ is the scaled payoff. Then, the multiplicative payoff corresponds
 522 to $W'_i(\mathbf{x}) = \exp(s\pi_i) = W_i(\mathbf{x})^s$.

523 If we try to calculate niche overlap and fitness difference based on the scaled
524 payoffs, we get:

$$\rho' = \sqrt{\frac{W'_i(\mathbf{e}_i)W'_j(\mathbf{e}_j)}{W'_j(\mathbf{e}_i)W'_i(\mathbf{e}_j)}}, \quad (\text{S21})$$

$$= \left(\sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}} \right)^s, \quad (\text{S22})$$

$$= \rho^s \quad (\text{S23})$$

$$\frac{\kappa'_j}{\kappa'_i} = \sqrt{\frac{W'_j(\mathbf{e}_i)W'_j(\mathbf{e}_j)}{W'_i(\mathbf{e}_i)W'_i(\mathbf{e}_j)}}, \quad (\text{S24})$$

$$= \left(\sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}} \right)^s, \quad (\text{S25})$$

$$= \left(\frac{\kappa_j}{\kappa_i} \right)^s \quad (\text{S26})$$

525 This implies that the exact values of niche and fitness difference are sensitive to
526 payoff scaling and are not intrinsic properties of a replicator system.

527 Nonetheless, the inequality for mutual invasion condition still holds under payoff
528 scaling:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho} \iff \rho^s < \left(\frac{\kappa_j}{\kappa_i} \right)^s < \frac{1}{\rho^s}. \quad (\text{S27})$$

529 Furthermore, boundary conditions for complete niche overlap ($\rho = \rho' = 1$) and com-
530 plete niche difference ($\rho = \rho' = 0$) are preserved. Moreover, the direction of changes
531 in fitness and niche difference is also preserved under payoff scaling. Specifically, for
532 parameter θ , we have:

$$\frac{d\rho'}{d\theta} = s\rho^{s-1} \frac{d\rho}{d\theta} \quad (\text{S28})$$

$$\frac{d}{d\theta} \left(\frac{\kappa'_j}{\kappa'_i} \right) = s \left(\frac{\kappa_j}{\kappa_i} \right)^{s-1} \frac{d}{d\theta} \left(\frac{\kappa_j}{\kappa_i} \right). \quad (\text{S29})$$

533 Since ρ and κ_j/κ_i are always positive, directional sensitivities of ρ and κ_j/κ_i to pa-
534 rameter θ are preserved. Similarly, directional sensitivities of $1 - \rho$ to θ are preserved:

$$\frac{d(1 - \rho')}{d\theta} = s\rho^{s-1} \frac{d(1 - \rho)}{d\theta} \quad (\text{S30})$$

535 Therefore, we focus on comparisons of qualitative outcomes from SCT and refrain
536 from making any comparisons of magnitude of niche and fitness difference values.

537 S3 Applications to classic two-strategy games

538 Consider a two-strategy game described by the following payoff matrix:

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (\text{S31})$$

539 There are four classic games that we can define from this simple payoff matrix:
 540 prisoner's dilemma ($T > R > P > S$), hawk-dove game ($T > R, S > P$), stag-hunt
 541 game ($R > T, P > S$), and harmony game ($R > T, S > P$). Here, we show that
 542 each game corresponds to a distinct region in the niche-fitness difference space under
 543 SCT.

544 To do so, we first begin by deriving expressions for niche overlap ρ and fitness
 545 difference κ_j/κ_i . Writing x_1 to denote the frequency of strategy 1, the expected
 546 payoff for each strategy corresponds to:

$$\pi_1(\mathbf{x}) = Rx_1 + S(1 - x_1), \quad (\text{S32})$$

$$\pi_2(\mathbf{x}) = Tx_1 + P(1 - x_1). \quad (\text{S33})$$

547 Then, multiplicative payoffs for each strategy when rare can be written as:

$$W_1(1) = \exp(R) \quad (\text{S34})$$

$$W_1(0) = \exp(S) \quad (\text{S35})$$

$$W_2(1) = \exp(T) \quad (\text{S36})$$

$$W_2(0) = \exp(P) \quad (\text{S37})$$

548 Therefore, we have:

$$\rho = \exp\left(\frac{R + P - S - T}{2}\right), \quad (\text{S38})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{T + P - R - S}{2}\right). \quad (\text{S39})$$

549 As before, the mutual-invasion condition can be expressed as:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (\text{S40})$$

550 Then, simple algebra reveals that each condition in mutual invasion inequality
 551 can be equivalently described by the payoff inequality:

$$\rho < \frac{\kappa_2}{\kappa_1} \iff R < T \quad (\text{S41})$$

$$\rho > \frac{\kappa_2}{\kappa_1} \iff R > T \quad (\text{S42})$$

$$\frac{\kappa_2}{\kappa_1} < \frac{1}{\rho} \iff P < S \quad (\text{S43})$$

$$\frac{\kappa_2}{\kappa_1} > \frac{1}{\rho} \iff P > S \quad (\text{S44})$$

552 These inequalities then allow us to classify outcomes of two-strategy games from the
 553 perspective of SCT. Note that the inequality between R and P does not affect the
 554 outcome.

555 **S4 Applications to mechanisms for the evolution of coopera-** 556 **tion**

557 We consider five mechanisms for the evolution of cooperation, all of which can be
 558 represented as a simple 2×2 payoff matrix (Nowak, 2006). Here, we go through each
 559 mechanism and derive measures for niche and fitness differences.

560 **Prisoner's dilemma.** Here, we consider a special case of Prisoner's dilemma
 561 which can be described by the following payoff matrix:

$$\begin{pmatrix} b - c & -c \\ b & 0 \end{pmatrix}, \quad (\text{S45})$$

562 where b represents the benefit for the recipient and c represents cost for the donor.
 563 In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0 \quad (\text{S46})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(c). \quad (\text{S47})$$

564 Therefore, as long as the cost is greater than zero, prisoner's dilemma always results
 565 in the competitive exclusion of cooperation.

566 **Kin selection.** Kin selection can be described by the following payoff matrix:

$$\begin{pmatrix} (b - c)(1 + r) & br - c \\ b - rc & 0 \end{pmatrix}, \quad (\text{S48})$$

567 where r represents the genetic relatedness. In this case, the niche and fitness differ-
 568 ences correspond to:

$$1 - \rho = 0 \quad (\text{S49})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(c - br). \quad (\text{S50})$$

569 Therefore, defection will exclude cooperation when $c > br$. Equivalently, cooperation
 570 will be favored when $r > c/b$, which recovers the classic result.

571 **Direct reciprocity.** Direct reciprocity can be described by the following payoff
 572 matrix:

$$\begin{pmatrix} (b - c)/(1 - w) & -c \\ b & 0 \end{pmatrix}, \quad (\text{S51})$$

573 where w represents the probability of next round. In this case, the niche and fitness
 574 differences correspond to:

$$1 - \rho = 1 - \exp\left(\frac{(b - c)/(1 - w) + c - b}{2}\right) \quad (\text{S52})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{b + c - (b - c)/(1 - w)}{2}\right). \quad (\text{S53})$$

575 Therefore, defection will exclude cooperation when

$$\rho < \frac{\kappa_2}{\kappa_1} \quad (\text{S54})$$

$$\iff \exp\left(\frac{(b - c)/(1 - w) + c - b}{2}\right) < \exp\left(\frac{b + c - (b - c)/(1 - w)}{2}\right) \quad (\text{S55})$$

$$\iff bw < c \quad (\text{S56})$$

576 However, note that the following inequality holds whenever $c > 0$:

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1}. \quad (\text{S57})$$

577 Therefore, when $w > c/b$, we will have

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1} < \rho, \quad (\text{S58})$$

578 which implies priority effect (equivalently, bistability). Even as $w \rightarrow 1^-$, this in-
 579 equality holds and therefore, cooperation cannot competitively exclude defection.

580 **Indirect reciprocity.** Indirect reciprocity can be described by the following
 581 payoff matrix:

$$\begin{pmatrix} b - c & -c(1 - q) \\ b(1 - q) & 0 \end{pmatrix}, \quad (\text{S59})$$

582 where q represents social acquaintanceship. In this case, the niche and fitness differ-
 583 ences correspond to:

$$1 - \rho = 1 - \exp\left(\frac{(b - c)q}{2}\right), \quad (\text{S60})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{2c - (b + c)q}{2}\right). \quad (\text{S61})$$

584 In this case, the condition for the dominance of defection (and therefore competitive
 585 exclusion of cooperation) can be written as:

$$\rho < \frac{\kappa_2}{\kappa_1} \quad (\text{S62})$$

$$\iff bq < c \quad (\text{S63})$$

Moreover, the following inequality always holds as long as $c > 0$.

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1}, \quad (\text{S64})$$

which implies that coexistence between cooperation and defection is not possible. Therefore, when $q < c/b$, then defection will competitively exclude cooperation. Otherwise, the system will enter the priority-effect regime.

Interestingly, when $q \rightarrow 1$, we have

$$\rho = \exp\left(\frac{b-c}{2}\right) \quad (\text{S65})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{c-b}{2}\right) \quad (\text{S66})$$

$$\frac{1}{\rho} = \exp\left(\frac{c-b}{2}\right) \quad (\text{S67})$$

$$(\text{S68})$$

In other words, the following inequality will hold:

$$\frac{1}{\rho} = \frac{\kappa_2}{\kappa_1} < \rho \quad (\text{S69})$$

Network reciprocity. Network reciprocity can be described by the following payoff matrix:

$$\begin{pmatrix} b-c & H-c \\ b-H & 0 \end{pmatrix}, \quad (\text{S70})$$

where $H = [(b-c)k - 2c]/[(k+1)(k-2)]$ and k represents the number of neighbors. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0, \quad (\text{S71})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(-H + c), \quad (\text{S72})$$

which implies that there is complete niche overlap and that fitness difference decreases with H . Since H is a monotonically decreasing function of k when $k > 2$ and $b/c \geq 2$, fitness difference will increase with k . In particular, cooperation will competitively exclude defection when $H > c$, meaning that:

$$b/c > k. \quad (\text{S73})$$

Group selection. Group selection can be described by the following payoff matrix:

$$\begin{pmatrix} (b-c)(m+n) & (b-c)m - cn \\ bn & 0 \end{pmatrix}, \quad (\text{S74})$$

where m represents the number of groups and n represents the group size. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0, \quad (\text{S75})$$

$$\frac{\kappa_2}{\kappa_1} = \exp((c - b)m + cn), \quad (\text{S76})$$

which implies that there is complete niche overlap. The fitness difference will increase with n but decrease with m , provided that $b > c$. In particular, cooperation will competitively exclude defection under the following condition:

$$1 + \frac{n}{m} < \frac{b}{c}. \quad (\text{S77})$$

S5 Applications to evolutionary games with environmental stochasticity

Consider a competition of two strategies under stochastic payoffs:

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix} + \frac{\zeta}{\sqrt{s}} \begin{pmatrix} \sigma_R & \sigma_S \\ \sigma_T & \sigma_P \end{pmatrix}, \quad (\text{S78})$$

Assuming weak selection $s \ll 1$, the evolution of two competing strategies in a large population can be described by the following equation:

$$\frac{dx}{dt} = sx(1 - x) \left(\pi_C - \pi_D + \left(\frac{1}{2} - x \right) (\sigma_C - \sigma_D)^2 \right), \quad (\text{S79})$$

To simplify, we write

$$\frac{dx}{dt} = sx(1 - x)G(x), \quad (\text{S80})$$

where

$$G(x) = \left(\pi_C - \pi_D + \left(\frac{1}{2} - x \right) (\sigma_C - \sigma_D)^2 \right). \quad (\text{S81})$$

Thus, the sign of $G(x)$ at boundary conditions determines whether a strategy can invade when the competing strategy is at equilibrium.

Since $G(x)$ does not uniquely separate into two payoff measures, we define niche and fitness difference based on the invasion growth rates $r_{C|D} = G(0)$ and $r_{D|C} = -G(1)$:

$$1 - \rho = -\exp\left(\frac{G(1) - G(0)}{2}\right) \quad (\text{S82})$$

$$\frac{\kappa_D}{\kappa_C} = \exp\left(-\frac{G(1) + G(0)}{2}\right) \quad (\text{S83})$$

619 Then, by substituting expressions for $G(0)$ and $G(1)$, we obtain the following expres-
 620 sions for niche and fitness difference:

$$1 - \rho = 1 - \exp \left(\frac{R + P - S - T}{2} - \frac{(\sigma_R - \sigma_T)^2 + (\sigma_S - \sigma_P)^2}{4} \right) \quad (\text{S84})$$

$$\frac{\kappa_D}{\kappa_C} = \exp \left(\frac{T + P - R - S}{2} + \frac{(\sigma_R - \sigma_T)^2 - (\sigma_S - \sigma_P)^2}{4} \right) \quad (\text{S85})$$

621 Now, to study the interior behavior of the system, we assume that the amount
 622 of stochasticity is proportional to deterministic payoffs (Wang et al., 2025):

$$\begin{pmatrix} \sigma_R & \sigma_S \\ \sigma_T & \sigma_P \end{pmatrix} = k \begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (\text{S86})$$

623 Then, the expression for $G(x)$ can be written as:

$$G(x) = ((R - T)x + (S - P)(1 - x)) \times \quad (\text{S87})$$

$$\left(1 + k^2 \left(\frac{1}{2} - x \right) ((R - T)x + (S - P)(1 - x)) \right) \quad (\text{S88})$$

$$= ((R + P - T - S)x + (S - P)) \times \quad (\text{S89})$$

$$\left(1 + k^2 \left(\frac{1}{2} - x \right) ((R + P - T - S)x + (S - P)) \right) \quad (\text{S90})$$

$$= k^2 (R + P - T - S)^2 (x - x^*) \left(K + \left(\frac{1}{2} - x \right) (x - x^*) \right), \quad (\text{S91})$$

624 where $x^* = (P - S)/(R + P - T - S)$ and $K = 1/(k^2(R + P - T - S))$ for $k > 0$
 625 and $R + P - T - S \neq 0$. Therefore, this system has an internal equilibrium $x = x^*$
 626 when $0 < (P - S)/(R + P - T - S) < 1$.

627 The system can also have up to two additional equilibria when $K + \left(\frac{1}{2} - x \right) (x -$
 628 $x^*) = 0$ has positive, real roots and when these roots fall between 0 and 1. In
 629 particular, the quadratic equation can be written as:

$$x^2 - \left(\frac{1}{2} + x^* \right) x + \frac{x^*}{2} - K = 0. \quad (\text{S92})$$

630 Thus, the two solutions of the quadratic equation correspond to:

$$x = \frac{x^* + \frac{1}{2} \pm \sqrt{\left(\frac{1}{2} - x^* \right)^2 + 4K}}{2}. \quad (\text{S93})$$

631 These solutions are real when:

$$\left(\frac{1}{2} - x^* \right)^2 + 4K > 0. \quad (\text{S94})$$

632 When these solutions fall between 0 and 1, they correspond to additional internal
 633 equilibria.

634 S6 Microbial cooperation and public goods

635 Here, we consider a simple, phenomenological model of competition between co-
 636 operating and cheating yeast in a cooperative sucrose-metabolism system, where
 637 invertase production by cooperators allows cheaters to utilize glucose, a public good
 638 released through sucrose hydrolysis (Carlson and Botstein, 1982; Dickinson, 1998;
 639 Gore et al., 2009). We note that this is a phenomenological approximation of a
 640 public-goods game as it does not explicitly consider interactions between multiple
 641 players (Archetti and Scheuring, 2011). Nonetheless, this example provides a simple
 642 illustration of how a concave benefit function can promote coexistence.

643 Invertase production by cooperators incurs a cost c and generates a total benefit
 644 of 1 that is captured privately with efficiency ϵ . Assuming a linear relationship
 645 between glucose availability and microbial growth, the payoffs for cooperators π_C
 646 and cheaters (defectors) π_D can then be written as

$$\pi_C(f) = \epsilon + f(1 - \epsilon) - c, \quad (\text{S95})$$

$$\pi_D(f) = f(1 - \epsilon), \quad (\text{S96})$$

647 where the use of glucose depends on the fraction of cooperators f . The dynamics of
 648 cooperators and defectors are described by the following replicator equation:

$$\frac{df}{dt} = f(1 - f) [\pi_C(f) - \pi_D(f)] \quad (\text{S97})$$

$$= f(1 - f)(\epsilon - c) \quad (\text{S98})$$

649 For this linear growth model, the payoff difference is fixed to $\epsilon - c$ independent of f
 650 (Figure S1A). Therefore, cooperation is favored when efficiency is greater than the
 651 cost of cooperation, $\epsilon > c$, whereas defection is favored when the cost is greater than
 652 efficiency, $c > \epsilon$ (Gore et al., 2009).

653 Our framework allows us to reinterpret this result from the perspective of coexis-
 654 tence theory. In particular, the two strategies have no niche difference, corresponding
 655 to complete niche overlap, $\rho = 1$. In contrast, the fitness difference is determined by
 656 the balance between private efficiency and the cost of cooperation:

$$\frac{\kappa_D}{\kappa_C} = \exp(c - \epsilon). \quad (\text{S99})$$

657 Therefore, this model always predicts competitive exclusion, except in the neutral
 658 case $c = \epsilon$. The outcome is determined entirely by the fitness difference between
 659 cooperators and defectors, rather than by stabilizing niche differences.

660 In real biological systems, the effect of glucose on microbial growth follows a
 661 nonlinear relationship:

$$\pi_C(f) = [\epsilon + f(1 - \epsilon)]^\alpha - c, \quad (\text{S100})$$

$$\pi_D(f) = [f(1 - \epsilon)]^\alpha, \quad (\text{S101})$$

662 where $\alpha = 0.15$ reflects the experimental relationship between glucose and microbial
 663 growth (Gore et al., 2009). Then, the dynamics of cooperators and defectors are
 664 described by the following replicator equation:

$$\frac{df}{dt} = f(1-f) [\{\epsilon + f(1-\epsilon)\}^\alpha - c - \{f(1-\epsilon)\}^\alpha]. \quad (\text{S102})$$

665 Under this nonlinear assumption, the payoff difference decreases with the cooperator
 666 fraction f , which can promote the coexistence of cooperators and defectors (Figure
 667 S1A): cooperators have a competitive advantage when rare, whereas defectors have a
 668 competitive advantage when cooperators are abundant. Because the payoff difference
 669 decreases monotonically with f , this model can have at most one internal equilib-
 670 rium. Thus, while it provides a simple illustration of how concavity can promote
 671 coexistence, it cannot capture the multiple internal equilibria that can arise under
 672 more general nonlinear benefit functions, such as sigmoid functions.

673 From a coexistence-theoretic perspective, this means that nonlinear growth can
 674 generate sufficient stabilizing niche difference to overcome the fitness difference. In
 675 particular, the niche and fitness differences for this model correspond to:

$$1 - \rho = 1 - \exp\left(\frac{1 - \epsilon^\alpha - (1 - \epsilon)^\alpha}{2}\right), \quad (\text{S103})$$

$$\frac{\kappa_D}{\kappa_C} = \exp\left(c + \frac{(1 - \epsilon)^\alpha - 1 - \epsilon^\alpha}{2}\right). \quad (\text{S104})$$

676 These expressions clarify the separate roles of cost, efficiency, and nonlinearity. For
 677 example, the cost of cooperation does not affect niche difference and only affects
 678 fitness difference. Instead, niche difference is generated by the concave relationship
 679 between glucose availability and microbial growth, which increases from 0 to $1 -$
 680 $\exp(-1/2)$ as α goes from 1 to 0 (Figure S1B,C). In contrast, as $\alpha \rightarrow 0^+$, the fitness
 681 difference approaches

$$\log(\kappa_D/\kappa_C) \rightarrow c - \frac{1}{2}, \quad (\text{S105})$$

682 which no longer depends on the private capture efficiency ϵ (Figure S1C). Thus,
 683 strong concavity not only generates stabilizing niche differences, but also acts as an
 684 equalizing mechanism (Figure S1C). Such stabilizing mechanisms are key ingredients
 685 for coexistence of competing strategies.

Supplementary Figure




Figure S1: **Effects of nonlinear microbial growth on coexistence of cooperation and defection.** (A) Payoff differences between cooperation and defection for linear and nonlinear models of microbial growth. In panel A, we assumed $\epsilon = 0.4$, $c = 0.3$, and $\alpha = 0.15$ for illustrative purposes. (B) Effect of nonlinearity α on niche difference $1 - \rho$. Each line assumes a different value of efficiency ϵ . Note that lines for $\epsilon = 0.6$ and $\epsilon = 0.4$ are nearly overlapping and are indistinguishable. (C) Effect of nonlinearity α on both niche difference $1 - \rho$ and fitness difference $\log(\kappa_D/\kappa_C)$. The gray shaded area represents the coexistence regime. The square represents the condition when $\alpha \rightarrow 0^+$. The circles represent the condition when $\alpha = 0.15$, which correspond to the experimental relationship between glucose and microbial growth (Gore et al., 2009). In panels B and C, we assumed $c = 0.3$, while allowing ϵ and α to vary.

References

- Adler, P. B., M. Detto, S. P. Ellner, T. L. Gibbs, Z. J. Gold, S. J. Leonard, J. I. Levine, A. E. Schiffer, C. Song, M. Stemkovski, et al. (2026). What have we learned from empirical applications of modern coexistence theory? *Ecology letters* 29(6), e70404.
- Adler, P. B., J. HilleRisLambers, and J. M. Levine (2007). A niche for neutrality. *Ecology letters* 10(2), 95–104.
- Allesina, S. (2026). Global stability of ecological and evolutionary dynamics via equivalence. *Proceedings of the National Academy of Sciences* 123(13), e2534915123.
- Apicella, C. L. and J. B. Silk (2019). The evolution of human cooperation. *Current Biology* 29(11), R447–R450.
- Archetti, M. and I. Scheuring (2011). Coexistence of cooperation and defection in public goods games. *Evolution* 65(4), 1140–1148.
- Archetti, M. and I. Scheuring (2012). Game theory of public goods in one-shot social dilemmas without assortment. *Journal of theoretical biology* 299, 9–20.
- Axelrod, R. and W. D. Hamilton (1981). The evolution of cooperation. *science* 211(4489), 1390–1396.
- Carlson, M. and D. Botstein (1982). Two differentially regulated mRNAs with different 5' ends encode secreted and intracellular forms of yeast invertase. *Cell* 28(1), 145–154.
- Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual review of Ecology and Systematics* 31(1), 343–366.
- Clutton-Brock, T. (2009). Cooperation between non-kin in animal societies. *Nature* 462(7269), 51–57.
- Cooney, D. B. (2020). Analysis of multilevel replicator dynamics for general two-strategy social dilemma. *Bulletin of Mathematical Biology* 82(6), 66.
- Cordero, O. X., L.-A. Ventouras, E. F. DeLong, and M. F. Polz (2012). Public good dynamics drive evolution of iron acquisition strategies in natural bacterioplankton populations. *Proceedings of the National Academy of Sciences* 109(49), 20059–20064.
- Cressman, R. (2003). *Evolutionary dynamics and extensive form games*, Volume 5. MIT Press.

- 720 Cressman, R. and Y. Tao (2014). The replicator equation and other game dynamics.
721 *Proceedings of the National Academy of Sciences* 111(supplement_3), 10810–10817.
- 722 Dickinson, J. R. (1998). *Metabolism and molecular physiology of Saccharomyces*
723 *cerevisiae*. Taylor & Francis.
- 724 Doebeli, M. and C. Hauert (2005). Models of cooperation based on the Prisoner’s
725 Dilemma and the Snowdrift game. *Ecology letters* 8(7), 748–766.
- 726 Doebeli, M., C. Hauert, and T. Killingback (2004). The evolutionary origin of coop-
727 erators and defectors. *science* 306(5697), 859–862.
- 728 Fletcher, J. A. and M. Doebeli (2008). A simple and general explanation for the evolu-
729 tion of altruism. *Proceedings of the Royal Society B: Biological Sciences* 276(1654),
730 13.
- 731 Godoy, O., N. J. Kraft, and J. M. Levine (2014). Phylogenetic relatedness and the
732 determinants of competitive outcomes. *Ecology letters* 17(7), 836–844.
- 733 Godoy, O. and J. M. Levine (2014). Phenology effects on invasion success: insights
734 from coupling field experiments to coexistence theory. *Ecology* 95(3), 726–736.
- 735 Gokhale, C. S. and A. Traulsen (2025). Higher-order equivalence of Lotka-Volterra
736 and replicator dynamics. *bioRxiv*, 2025–03.
- 737 Gore, J., H. Youk, and A. Van Oudenaarden (2009). Snowdrift game dynamics and
738 facultative cheating in yeast. *Nature* 459(7244), 253–256.
- 739 Hamilton, W. D. (1964). The genetical evolution of social behaviour. II. *Journal of*
740 *theoretical biology* 7(1), 17–52.
- 741 Hauert, C. (2002). Effects of space in 2×2 games. *International Journal of Bifur-*
742 *cation and Chaos* 12(07), 1531–1548.
- 743 Hauert, C. and M. Doebeli (2004). Spatial structure often inhibits the evolution of
744 cooperation in the snowdrift game. *Nature* 428(6983), 643–646.
- 745 Hilbe, C., Š. Šimsa, K. Chatterjee, and M. A. Nowak (2018). Evolution of cooperation
746 in stochastic games. *Nature* 559(7713), 246–249.
- 747 Hofbauer, J. (1981). On the occurrence of limit cycles in the volterra-lotka equation.
748 *Nonlinear Analysis: Theory, Methods & Applications* 5(9), 1003–1007.
- 749 Hofbauer, J. and K. Sigmund (1998). *Evolutionary games and population dynamics*.
750 Cambridge university press.
- 751 Kingma, S. A., P. Santema, M. Taborsky, and J. Komdeur (2014). Group augmen-
752 tation and the evolution of cooperation. *Trends in ecology & evolution* 29(8),
753 476–484.

754 Kocher, S. D., R. Mallarino, B. E. Rubin, D. W. Yu, H. E. Hoekstra, and N. E.
755 Pierce (2018). The genetic basis of a social polymorphism in halictid bees. *Nature*
756 *Communications* 9(1), 4338.

757 Kraft, N. J., O. Godoy, and J. M. Levine (2015). Plant functional traits and the mul-
758 tidimensional nature of species coexistence. *Proceedings of the National Academy*
759 *of Sciences* 112(3), 797–802.

760 Lee, I. P. A., O. T. Eldakar, J. P. Gogarten, and C. P. Andam (2022). Bacterial co-
761 operation through horizontal gene transfer. *Trends in Ecology & Evolution* 37(3),
762 223–232.

763 Lewontin, R. C. (1961). Evolution and the theory of games. *Journal of Theoretical*
764 *Biology* 1(3), 382–403.

765 Lieberman, E., C. Hauert, and M. A. Nowak (2005). Evolutionary dynamics on
766 graphs. *Nature* 433(7023), 312–316.

767 Lin, H., D. Wang, Q. Wang, J. Mao, Y. Bai, and J. Qu (2024). Interspecific compe-
768 tition prevents the proliferation of social cheaters in an unstructured environment.
769 *The ISME Journal* 18(1), wrad038.

770 Michel-Mata, S., M. Kawakatsu, J. Sartini, T. A. Kessinger, J. B. Plotkin, and C. E.
771 Tarnita (2024). The evolution of private reputations in information-abundant
772 landscapes. *Nature* 634(8035), 883–889.

773 Motro, U. (1991). Co-operation and defection: playing the field and the ESS. *Journal*
774 *of Theoretical Biology* 151(2), 145–154.

775 Nowak, M. A. (2006). Five rules for the evolution of cooperation. *science* 314(5805),
776 1560–1563.

777 Nowak, M. A. and R. M. May (1992). Evolutionary games and spatial chaos. *na-*
778 *ture* 359(6398), 826–829.

779 Nowak, M. A. and K. Sigmund (1998). Evolution of indirect reciprocity by image
780 scoring. *Nature* 393(6685), 573–577.

781 Nowak, M. A. and K. Sigmund (2005). Evolution of indirect reciprocity. *Na-*
782 *ture* 437(7063), 1291–1298.

783 Nowak, M. A., C. E. Tarnita, and T. Antal (2010). Evolutionary dynamics in struc-
784 tured populations. *Philosophical Transactions of the Royal Society B: Biological*
785 *Sciences* 365(1537), 19.

786 Page, K. M. and M. A. Nowak (2002). Unifying evolutionary dynamics. *Journal of*
787 *theoretical biology* 219(1), 93–98.

- 788 Palmer, J. D. and K. R. Foster (2022). Bacterial species rarely work together. *Sci-*
789 *ence* 376(6593), 581–582.
- 790 Park, S. W., S. Cobey, C. J. E. Metcalf, J. M. Levine, and B. T. Grenfell (2024).
791 Predicting pathogen mutual invasibility and co-circulation. *Science* 386(6718),
792 175–179.
- 793 Park, S. W., J. Levine, and B. Grenfell (2026). Unifying coexistence theory for
794 ecological communities and pathogen strain competition. *bioRxiv*, 2026–05.
- 795 Pollak, S., M. Gralka, Y. Sato, J. Schwartzman, L. Lu, and O. X. Cordero (2021).
796 Public good exploitation in natural bacterioplankton communities. *Science Ad-*
797 *vances* 7(31), eabi4717.
- 798 Rakoff-Nahoum, S., K. R. Foster, and L. E. Comstock (2016). The evolution of
799 cooperation within the gut microbiota. *Nature* 533(7602), 255–259.
- 800 Schuster, P. and K. Sigmund (1983). Replicator dynamics. *Journal of theoretical*
801 *biology* 100(3), 533–538.
- 802 Sieben, A. J., J. R. Mihaljevic, and L. G. Shoemaker (2022). Quantifying mechanisms
803 of coexistence in disease ecology. *Ecology* 103(12), e3819.
- 804 Slobodkin, L. B. (1964). The strategy of evolution. *American scientist* 52(3), 342–
805 357.
- 806 Smith, J. M. and G. R. Price (1973). The logic of animal conflict. *Nature* 246(5427),
807 15–18.
- 808 Souza, M. O., J. M. Pacheco, and F. C. Santos (2009). Evolution of cooperation
809 under N-person snowdrift games. *Journal of Theoretical Biology* 260(4), 581–588.
- 810 Steidinger, B. S. and J. D. Bever (2014). The coexistence of hosts with different
811 abilities to discriminate against cheater partners: an evolutionary game-theory
812 approach. *The American Naturalist* 183(6), 762–770.
- 813 Szabó, G. and G. Fath (2007). Evolutionary games on graphs. *Physics reports* 446(4-
814 6), 97–216.
- 815 Tarnita, C. E. and A. Traulsen (2025). Reconciling ecology and evolutionary game
816 theory or “When not to think cooperation”. *Proceedings of the National Academy*
817 *of Sciences* 122(14), e2413847122.
- 818 Taylor, P. D. and L. B. Jonker (1978). Evolutionary stable strategies and game
819 dynamics. *Mathematical biosciences* 40(1-2), 145–156.
- 820 Traulsen, A. and M. A. Nowak (2006). Evolution of cooperation by multilevel selec-
821 tion. *Proceedings of the National Academy of Sciences* 103(29), 10952–10955.

- 822 Trivers, R. L. (1971). The evolution of reciprocal altruism. *The Quarterly review of*
823 *biology* 46(1), 35–57.
- 824 Wang, G., Q. Su, L. Wang, and J. B. Plotkin (2025). Evolution of social behaviors
825 in noisy environments. *arXiv*.
- 826 Wedekind, C. and M. Milinski (2000). Cooperation through image scoring in humans.
827 *Science* 288(5467), 850–852.
- 828 Weitz, J. S., C. Eksin, K. Paarporn, S. P. Brown, and W. C. Ratcliff (2016). An
829 oscillating tragedy of the commons in replicator dynamics with game-environment
830 feedback. *Proceedings of the National Academy of Sciences* 113(47), E7518–E7525.
- 831 West, S. A., G. A. Cooper, M. B. Ghoul, and A. S. Griffin (2021). Ten recent insights
832 for our understanding of cooperation. *Nature ecology & evolution* 5(4), 419–430.
- 833 West, S. A., A. S. Griffin, and A. Gardner (2007). Evolutionary explanations for
834 cooperation. *Current biology* 17(16), R661–R672.