

Strategic coexistence theory for evolutionary games

Sang Woo Park^{1,*}

¹ School of Biological Sciences, Seoul National University, Seoul, Korea

*Corresponding author: sangwoopark@snu.ac.kr

Abstract

Evolutionary game theory and ecological coexistence theory both seek to predict the outcome of competition between biological entities, be they strategies or species, but the two fields have relied on largely separate approaches. Replicator equations provide a foundation for analyzing strategy competition and the stability of strategic equilibria, yet they do not explicitly separate the mechanisms that stabilize competition from those that equalize fitness differences between strategies. Here, we extend modern coexistence theory from community ecology to develop strategic coexistence theory (SCT), a framework for quantifying strategic niche and fitness differences between competing strategies. SCT recovers the classic classification of two-strategy games, distinguishing competitive exclusion, coexistence, and priority effects within a shared niche–fitness difference space. Applying SCT to five mechanisms for the evolution of cooperation further reveals that these mechanisms promote cooperation through distinct dynamical routes: kin selection, network reciprocity, and group selection primarily reduce fitness differences, whereas direct and indirect reciprocity destabilize competition and generate priority effects. Finally, applying SCT to microbial public-goods game shows that nonlinear microbial growth can both stabilize and equalize competition between cooperators and defectors, allowing coexistence. Together, these results show that SCT provides a complementary framework for comparing evolutionary games and teasing apart the coexistence mechanisms underlying strategy competition.

Introduction

Understanding the mechanisms that promote the evolutionary stability of competing strategies is a fundamental aim in evolutionary biology (Lewontin, 1961; Slobodkin, 1964; Smith and Price, 1973; Taylor and Jonker, 1978). To this end, evolutionary game theory (EGT) and replicator dynamics have provided key foundations, allowing us to predict the outcome of strategic competition through payoff inequalities, invasion criteria, and stability analysis (Schuster and Sigmund, 1983; Hofbauer and Sigmund, 1998; Cressman, 2003; Shalizi, 2009; Cressman and Tao, 2014). However, less attention has been given to similarities between replicator dynamics and community ecology until recently (Gokhale and Traulsen, 2025; Tarnita and Traulsen,

2025; Allesina, 2026), even though the classic replicator equations and the generalized Lotka-Volterra equation are mathematically equivalent (Hofbauer, 1981; Page and Nowak, 2002).

Interestingly, even a simple two-strategy game can yield heterogeneous outcomes that mirror outcomes in community ecology (Hauert, 2002; Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025): the dominance of a single strategy (prisoner’s dilemma and harmony games), coexistence (hawk–dove games), and bistability (stag-hunt games). Such differences in conditions that promote evolutionary stability and dominance of a single strategy and those that permit strategy coexistence indicate that EGT can be understood from the perspective of coexistence theory (Motro, 1991; Doebeli et al., 2004; Hauert and Doebeli, 2004; Doebeli and Hauert, 2005; Gore et al., 2009; Souza et al., 2009; Archetti and Scheuring, 2011; Steidinger and Bever, 2014; Tarnita and Traulsen, 2025). In fact, stable polymorphism of evolutionary strategies, such as coexistence of cooperators and defectors, are observed in microbial and other biological systems (Gore et al., 2009; Cordero et al., 2012; Kocher et al., 2018; Pollak et al., 2021; Lin et al., 2024), but coexistence-theoretic perspective has often remained implicit. Recasting game dynamics as a coexistence problem immediately raises the following question: when two strategies coexist, does the corresponding mechanism stabilize or equalize competition (Chesson, 2000)? While EGT allows us to predict whether two strategies can coexist or not, a framework for quantifying coexistence mechanisms between competing strategies remains elusive.

Modern coexistence theory (MCT) from community ecology presents a particularly useful framework for quantifying coexistence mechanisms (Chesson, 2000; Godoy et al., 2014; Godoy and Levine, 2014; Kraft et al., 2015; Adler et al., 2026). MCT predicts that coexistence of two competing entities, be they species or strategies, requires stabilizing niche differences to overcome average fitness differences. In community ecology, stabilizing niche differences represent differences in factors that limit the growth of competing species, causing each species to limit itself more than it limits its competitor (Chesson, 2000). In EGT, a strategy’s niche can be interpreted as strategic environments defined by the composition of competing strategies and the resulting interactions. Thus, stabilizing niche differences arise when each strategy is favored by a different strategic environment, such that the spread of each strategy creates conditions that limit its own relative growth and favor the recovery of its competitor. In contrast, fitness differences represent differences in the ability of each entity to compete under the limiting environments created by its competitor (Chesson, 2000). This differs from payoff differences in EGT, which depend on the underlying strategy frequency and therefore combine both stabilizing effects caused by changes in the strategic environment and residual fitness differences that favor one strategy over another.

So far, MCT has been primarily utilized for understanding species coexistence, but recent work showed that MCT can be extended beyond species competition, especially for studying pathogen strain competition, allowing one to directly quantify niche and fitness differences of competing entities, be they species or strains (Sieben

et al., 2022; Park et al., 2024, 2026). Inspired by this effort, we present a Strategic Coexistence Theory (SCT) for predicting coexistence and exclusion of competing strategies in EGT by quantifying strategic niche and fitness differences. We begin by introducing SCT, which can be applied to any generic system of EGT described by replicator equations. We show that SCT can recover outcomes from classic, two-strategy games, providing reinterpretation of classic results from coexistence-theoretic perspective. We then apply SCT to comparing five mechanisms for the evolution of cooperation, revealing that direct and indirect reciprocity destabilize competition between cooperation and defection, thus pushing the system towards priority effect regime. Finally, we consider a case study of microbial cooperation, where nonlinear microbial growth can both stabilize and equalize competition between cooperation and defection, allowing coexistence of strategies.

Strategic coexistence theory

Here, we present a brief derivation of strategic coexistence theory (SCT) (Figure 1). To do so, we begin with a classic replicator equation, which allows us to model changes in the frequency of competing strategies based on their payoffs (Schuster and Sigmund, 1983; Hofbauer and Sigmund, 1998; Cressman, 2003; Shalizi, 2009; Cressman and Tao, 2014):

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (1)$$

where x_i represents the frequency of strategy i , $\pi_i(\mathbf{x})$ represents the payoff of strategy i for a given strategy distribution $\mathbf{x}$, and $\bar{\pi}(\mathbf{x}) = \sum_i \pi_i(\mathbf{x})x_i$ represents the average payoff. Since strategy frequencies must sum to 1 (i.e., $\sum_i x_i = 1$), a single-strategy equilibrium dominated by strategy i can be described by a unit vector $\mathbf{e}_i$, where all elements are equal to zero except for the i -th element. We use these single-strategy equilibria to define niche and fitness differences between two competing strategies i and j .

Because payoffs in many games can be negative, payoff ratios can be difficult to interpret. Thus, we work on a positive multiplicative scale by exponentiating the payoff terms:

$$W_i(\mathbf{x}) = \exp(\pi_i(\mathbf{x})). \quad (2)$$

This exponential transformation is convenient because it preserves the payoff ordering and does not alter the invasion conditions. Therefore, this allows us to compare strategic environments through ratios, as in modern coexistence theory. For convenience, we refer to $W_i(\mathbf{x})$ as the multiplicative payoff of strategy i .

First, we define the niche overlap ρ between strategies i and j as the ratio between the geometric mean of each strategy's multiplicative payoff in the strategic environment that it creates when common, $\sqrt{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}$, and the geometric mean of




Figure 1: **A schematic diagram summarizing strategic coexistence theory.** Strategic coexistence theory decomposes pairwise replicator dynamics into stabilizing and equalizing mechanisms. Mutual invasion conditions can be rewritten on a multiplicative payoff scale, allowing us to derive strategic niche difference $1 - \rho$ and fitness difference κ_j/κ_i . Then, the inequality $\rho < \kappa_j/\kappa_i < 1/\rho$ determines mutual invasion.

each strategy's multiplicative payoff in the environment generated by its competitor, $\sqrt{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}$. The corresponding niche difference is then given by

$$1 - \rho = 1 - \sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}}. \quad (3)$$

When $1 - \rho > 0$, each strategy performs better, on average, in the environment generated by its competitor than in the environment generated by itself. In other words, intraspecific competition is stronger than interspecific competition. Second, we define the fitness difference as the ratio between the geometric means of the multiplicative payoff of each strategy across the two single-strategy environments:

$$\frac{\kappa_j}{\kappa_i} = \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}. \quad (4)$$

In other words, strategy j has an average competitive advantage over strategy i ($\kappa_j/\kappa_i > 1$) when it performs better, on average, across the environments generated by itself and its competitor.

Together, for any two strategies described by the classic replicator equation, mutual invasion requires stabilizing niche differences to overcome average fitness differences. Specifically, the condition for mutual invasion can be rewritten as (Chesson,

129 2000):

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (5)$$

130 This inequality applies to any games described by the classic replicator equation
 131 because ρ and κ_j/κ_i can be calculated directly from ratios of multiplicative pay-
 132 offs. This decomposition not only allows us to understand how different mechanisms
 133 contribute to the coexistence or exclusion of competing strategies but also provides
 134 shared axes for comparing different games.

135 We note that the absolute magnitudes of niche and fitness differences are sensitive
 136 to payoff scaling because they are defined on an exponentiated payoff scale. However,
 137 the qualitative outcome of strategic competition under SCT is preserved because the
 138 exponential transformation does not alter invasion conditions.

139 Classic two-strategy games

140 As an illustrative example, we begin with classic two-strategy games (Hauert, 2002;
 141 Szabó and Fath, 2007; Cooney, 2020; Tarnita and Traulsen, 2025), which can be
 142 described by the following payoff matrix (Figure 2A):

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (6)$$

143 Here, the first and second rows represent strategies 1 and 2, respectively, and the
 144 columns represent the strategy of the interacting opponent. Writing x_1 to denote
 145 the frequency of strategy 1, the expected payoffs are

$$\pi_1(\mathbf{x}) = Rx_1 + S(1 - x_1), \quad (7)$$

$$\pi_2(\mathbf{x}) = Tx_1 + P(1 - x_1). \quad (8)$$

146 Thus, the dynamics can be written as

$$\frac{dx_1}{dt} = x_1(1 - x_1) [(R - T)x_1 + (S - P)(1 - x_1)]. \quad (9)$$

147 The signs of $T - R$ and $S - P$ determine whether each strategy can invade the
 148 other when rare. Specifically, strategy 2 can invade strategy 1 when $T > R$, whereas
 149 strategy 1 can invade strategy 2 when $S > P$. These two invasion conditions partition
 150 the game space into four regimes (Figure 2B): prisoner’s dilemma ($T > R > P >$
 151 S), hawk–dove game ($T > R, S > P$), stag–hunt game ($R > T, P > S$), and
 152 harmony game ($R > T, S > P$). In standard EGT, these outcomes are obtained by
 153 analyzing the stability of equilibrium conditions. For simple games, this analysis is
 154 straightforward.

155 A coexistence-theoretic approach recovers the same classification but provides
 156 a different interpretation (Figure 2C). For a two-strategy game determined by this




Figure 2: **Coexistence-theoretic interpretation of classic two-strategy games.** (A) General payoff matrix for a two-strategy game, where rows represent the focal strategy and columns represent the strategy of the interacting opponent. (B) Classification of four classic games based on the payoff differences $T - R$ and $S - P$. Prisoner's dilemma and harmony games lead to the dominance of one strategy; hawk-dove games allow coexistence; and stag-hunt games result in bistability. (C) Mapping of the same four games onto niche-fitness difference space. Hawk-dove games correspond to the coexistence regime, where stabilizing niche differences overcome fitness differences. Prisoner's dilemma and harmony games correspond to competitive-exclusion regimes, whereas stag-hunt games correspond to the priority-effect regime. $R > P$ is assumed throughout, but this does not affect coexistence outcome in any way.

157 payoff matrix, the niche difference and fitness difference are given by

$$1 - \rho = 1 - \exp\left(\frac{R + P - S - T}{2}\right), \quad (10)$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{T + P - R - S}{2}\right). \quad (11)$$

158 As shown above, mutual invasion requires $\rho < \kappa_2/\kappa_1 < 1/\rho$, which is equivalent to
 159 $T > R$ and $S > P$. This corresponds to the hawk-dove game, in which each strategy
 160 can invade when rare and the two strategies coexist. The remaining games also
 161 have natural interpretations in niche-fitness difference space. Prisoner's dilemma
 162 and harmony games represent competitive exclusion regimes, where one strategy
 163 has a fitness advantage that exceeds niche differences. In contrast, stag-hunt games
 164 represent a priority-effect regime, where neither strategy can invade the other and
 165 the outcome depends on initial conditions. Therefore, coexistence theory not only
 166 allows us to recover the classic results from two-strategy games but also provides new
 167 interpretations for these results in terms of niche and fitness differences. Building on

168 this, we next apply this framework to comparing core mechanisms of cooperation.

169 Comparisons of mechanisms for the evolution of co- 170 operation

171 Even though cooperative behavior can be found across a wide range of biological
172 systems, a simple prisoner’s dilemma shows that cooperation is not an evolution-
173 arily stable strategy (May, 1987). Thus, many researchers have sought to identify
174 mechanisms that promote the evolution of cooperation (West et al., 2007). Here, we
175 compare five classic mechanisms for the evolution of cooperation using SCT and ask
176 how they differ from the perspective of strategy coexistence.

177 Specifically, we consider the following mechanisms: kin selection, direct reci-
178 procity, indirect reciprocity, network reciprocity, and group selection (Figure 3A).
179 Kin selection favors cooperation when two interacting individuals are genetic rela-
180 tives, where the genetic relatedness r must be greater than the cost-to-benefit ratio:
181 $r > c/b$ (Hamilton, 1964). Direct reciprocity favors cooperation when current co-
182 operation may promote future cooperation by the same individual through repeated
183 encounters (Trivers, 1971; Axelrod and Hamilton, 1981). In this case, the probability
184 of repeated interaction (i.e., “next round”) must be greater than the cost-to-benefit
185 ratio: $w > c/b$. Indirect reciprocity favors cooperation when current cooperation
186 may promote future cooperation by different individuals through reputation (Nowak
187 and Sigmund, 1998; Wedekind and Milinski, 2000; Nowak and Sigmund, 2005). Thus,
188 the social acquaintanceship (i.e., the probability of knowing someone’s reputation)
189 must be greater than the cost-to-benefit ratio: $q > c/b$. Network reciprocity favors
190 cooperation via clustering of cooperators, requiring the benefit-to-cost ratio to be
191 greater than the average number of neighbors: $b/c > k$ (Nowak and May, 1992;
192 Lieberman et al., 2005). Finally, group selection favors cooperation via multilevel
193 selection, where groups with more cooperators exhibit faster growth at the group
194 level even if defectors reproduce faster at the individual level (Traulsen and Nowak,
195 2006). This requires $b/c > 1 + (n/m)$, where n is the maximum group size and
196 m is the number of groups. All of these mechanisms can be described by a simple
197 2×2 payoff matrix (Nowak, 2006), from which the corresponding niche and fitness
198 differences can be computed using Eq. (10) and Eq. (11).

199 First, the prisoner’s dilemma provides a baseline scenario, where defection is
200 evolutionarily stable (Figure 3A). From the perspective of SCT, there is complete
201 niche overlap, and therefore defection competitively excludes cooperation (Figure
202 3B). Building on this, SCT shows that five mechanisms promote evolutionary sta-
203 bility of cooperation via different dynamical routes (Figure 3B). For example, kin
204 selection allows the fitness of cooperation to increase as genetic relatedness r increases
205 (Figure 3B). However, it does not affect the niche difference between cooperation and
206 defectors. Similarly, network reciprocity and group selection modulate the fitness of
207 cooperation without causing any changes in strategic niches. In contrast, direct




Figure 3: **Comparisons of five mechanisms for the evolution of cooperation using SCT.** (A) Payoff matrices for the baseline scenario (prisoner's dilemma) and five mechanisms for the evolution of cooperation (Nowak, 2006). Prisoner's dilemma is characterized by the benefit for the recipient b and the cost for the donor c . Network reciprocity is parameterized based on $H = [(b-c)k - 2c]/[(k+1)(k-2)]$. (B) Changes in niche and fitness differences driven by each mechanism. Each arrow represents the direction of change as we increase the corresponding parameter, indicated in figure legends. Note that kin selection, network reciprocity, and group selection all result in changes in fitness difference without causing any changes in niche difference. Therefore, we indicate the direction of change for network reciprocity and group selection separately to avoid overlapping arrows.

and indirect reciprocity destabilize competition while increasing the fitness of cooperation, thereby pushing the competition outcome towards a priority effect regime. Equivalently, these two mechanisms promote bistability of defection and cooperation. SCT further reveals another difference between direct and indirect reciprocity. In particular, direct reciprocity is unable to generate sufficient fitness difference to competitively exclude defection even when the probability of next round w reaches 1. In contrast, indirect reciprocity is able to generate sufficient fitness difference to push the system towards the boundary between priority-effect regime and competitive-exclusion regime as social acquaintanceship q reaches 1, indicating that an additional mechanism will be able to promote complete dominance of cooperation.

We note that none of these mechanisms allow coexistence of cooperation and

defection. Therefore, we move on to the final example, applying SCT to microbial cooperation, where nonlinear benefits and environmental context can promote coexistence of cooperators and defectors.

Microbial cooperation and public goods

Here, we consider a simple, phenomenological model of competition between cooperating and cheating yeast in a cooperative sucrose-metabolism system, where invertase production by cooperators allows cheaters to utilize glucose, a public good released through sucrose hydrolysis (Carlson and Botstein, 1982; Dickinson, 1998; Gore et al., 2009). Specifically, invertase production by cooperators incurs a cost c and generates a total benefit of 1 that is captured privately with efficiency ϵ . Assuming a linear relationship between glucose availability and microbial growth, the payoffs for cooperators π_C and cheaters (defectors) π_D can then be written as

$$\pi_C(f) = \epsilon + f(1 - \epsilon) - c, \quad (12)$$

$$\pi_D(f) = f(1 - \epsilon), \quad (13)$$

where the use of glucose depends on the fraction of cooperators f . The dynamics of cooperators and defectors are described by the following replicator equation:

$$\frac{df}{dt} = f(1 - f) [\pi_C(f) - \pi_D(f)] \quad (14)$$

$$= f(1 - f)(\epsilon - c) \quad (15)$$

For this linear growth model, the payoff difference is fixed to $\epsilon - c$ independent of f (Figure 4A). Therefore, cooperation is favored when efficiency is greater than the cost of cooperation, $\epsilon > c$, whereas defection is favored when the cost is greater than efficiency, $c > \epsilon$ (Gore et al., 2009).

Our framework allows us to reinterpret this result from the perspective of coexistence theory. In particular, the two strategies have no niche difference, corresponding to complete niche overlap, $\rho = 1$. In contrast, the fitness difference is determined by the balance between private efficiency and the cost of cooperation:

$$\frac{\kappa_D}{\kappa_C} = \exp(c - \epsilon). \quad (16)$$

Therefore, this model always predicts competitive exclusion, except in the neutral case $c = \epsilon$. The outcome is determined entirely by the fitness difference between cooperators and defectors, rather than by stabilizing niche differences. This example resembles the prisoner's dilemma discussed in the previous example (Figure 3).

In real biological systems, the effect of glucose on microbial growth can follow a nonlinear relationship:

$$\pi_C(f) = [\epsilon + f(1 - \epsilon)]^\alpha - c, \quad (17)$$

$$\pi_D(f) = [f(1 - \epsilon)]^\alpha, \quad (18)$$




Figure 4: **Effects of nonlinear microbial growth on coexistence of cooperation and defection.** (A) Payoff differences for linear and nonlinear models of microbial growth. In panel A, we assumed $\epsilon = 0.4$, $c = 0.3$, and $\alpha = 0.15$ for illustrative purposes. (B) Effect of nonlinearity α on niche difference $1 - \rho$. Each line assumes a different value of efficiency ϵ . Note that lines for $\epsilon = 0.6$ and $\epsilon = 0.4$ are completely overlapping. (C) Effect of nonlinearity α on both niche difference $1 - \rho$ and fitness difference $\log(\kappa_D/\kappa_C)$. The gray shaded area represents the coexistence regime. The square represents the condition when $\alpha \rightarrow 0^+$. The circles represent the condition when $\alpha = 0.15$, which correspond to the experimental relationship between glucose and microbial growth (Gore et al., 2009). In panels B and C, we assumed $c = 0.3$, while allowing ϵ and α to vary.

where $\alpha = 0.15$ reflects the experimental relationship between glucose and microbial growth (Gore et al., 2009). Then, the dynamics of cooperators and defectors are described by the following replicator equation:

$$\frac{df}{dt} = f(1-f) [\{\epsilon + f(1-\epsilon)\}^\alpha - c - \{f(1-\epsilon)\}^\alpha]. \quad (19)$$

Under this nonlinear assumption, the payoff difference decreases with the cooperator fraction f , which can promote the coexistence of cooperators and defectors (Figure 4A): cooperators have a competitive advantage when rare, whereas defectors have a competitive advantage when cooperators are abundant.

From a coexistence-theoretic perspective, this means that nonlinear growth can generate sufficient stabilizing niche difference to overcome the fitness difference. In particular, the niche and fitness differences for this model correspond to:

$$1 - \rho = 1 - \exp\left(\frac{1 - \epsilon^\alpha - (1 - \epsilon)^\alpha}{2}\right), \quad (20)$$

$$\frac{\kappa_D}{\kappa_C} = \exp\left(c + \frac{(1 - \epsilon)^\alpha - 1 - \epsilon^\alpha}{2}\right). \quad (21)$$

257 These expressions clarify the separate roles of cost, efficiency, and nonlinearity. For
 258 example, the cost of cooperation does not affect niche difference and only affects
 259 fitness difference. Instead, niche difference is generated by the concave relationship
 260 between glucose availability and microbial growth, which increases from 0 to $1 -$
 261 $\exp(-1/2)$ as α goes from 1 to 0 (Figure 4B,C). In contrast, as $\alpha \rightarrow 0^+$, the fitness
 262 difference approaches

$$\log(\kappa_D/\kappa_C) \rightarrow c - \frac{1}{2}, \quad (22)$$

263 which no longer depends on the private capture efficiency ϵ (Figure 4C). Thus, strong
 264 concavity not only generates stabilizing niche differences, but also acts as an equal-
 265 izing mechanism (Figure 4C).

266 Discussion

267 Standard evolutionary game theory and replicator equations provide powerful tools
 268 for predicting the outcome of strategy competition, including strategic dominance,
 269 coexistence, and bistability (Hofbauer and Sigmund, 1998; Cressman and Tao, 2014).
 270 However, these outcomes are typically understood by analyzing payoff differences
 271 within individual games, which makes it difficult to compare coexistence mechanisms
 272 across models. Building on Chesson’s Modern Coexistence Theory from community
 273 ecology (Chesson, 2000), we developed a strategic coexistence theory (SCT) that
 274 allows us to decompose strategy competition in terms of stabilizing niche differences
 275 and fitness differences. This decomposition provides a unifying framework for com-
 276 paring different games from the perspective of coexistence theory, allowing us to
 277 identify stabilizing and equalizing mechanisms for strategy competition.

278 To demonstrate the utility of SCT, we considered three case studies. First, the
 279 analysis of classic two-strategy games showed that SCT recovers the original clas-
 280 sification for these classic games, highlighting the similarities between evolutionary
 281 games and community ecology (Hauert, 2002; Szabó and Fath, 2007; Cooney, 2020;
 282 Tarnita and Traulsen, 2025). Then, we compared different mechanisms for the evo-
 283 lution of cooperation, which showed that each mechanism promotes cooperation via
 284 different dynamical routes (Nowak, 2006). Kin selection, network reciprocity, and
 285 group selection primarily affect the fitness difference between cooperation and defec-
 286 tion, whereas direct and indirect reciprocity can destabilize competition and generate
 287 priority effects. Finally, our analysis of microbial cooperation showed that coexis-
 288 tence between cooperators and defectors requires more than reducing the fitness cost
 289 of cooperation: nonlinear benefits can generate stabilizing niche differences while
 290 also reducing fitness asymmetry (Gore et al., 2009). Taken together, these examples
 291 show that SCT can provide new insights into ecological underpinnings of strategy
 292 competition.

293 There are several limitations to SCT. First, similar to MCT from community
 294 ecology, SCT only allows for comparisons of pairwise strategies and therefore cannot
 295 determine the outcomes of games with more than two strategies (Chesson, 2000).

296 Second, we focused on classic replicator equations, but Tarnita and Traulsen (2025)
 297 recently pointed out that these classic formulations implicitly ignore differences in
 298 intrinsic growth rates. In such cases, replicator equations may fail to predict the out-
 299 come of competition (Tarnita and Traulsen, 2025), and a full Lotka-Volterra model is
 300 needed to describe the abundances, rather than frequencies, of strategies. Thus, SCT
 301 inherits the same assumptions as the classic replicator framework: it is most appro-
 302 priate when strategy competition can be described in terms of relative frequencies
 303 alone. When strategies differ intrinsically in growth, MCT may be required instead
 304 to tease apart mechanisms of strategy coexistence. Finally, we have focused on simple
 305 examples to illustrate the utility of SCT, but many games involve additional com-
 306 plexities, including population structure (Nowak et al., 2010), stochasticity (Wang
 307 et al., 2025; Bodin et al., 2026), and explicit ecological feedback. Extending SCT to
 308 these settings will be necessary to determine how broadly stabilizing and equalizing
 309 mechanisms can be compared across evolutionary games.

310 In conclusion, SCT provides a bridge between evolutionary game theory and
 311 community ecology, allowing strategy competition to be understood from the per-
 312 spective of coexistence theory. Therefore, rather than replacing existing EGT or
 313 replicator-dynamics frameworks, SCT provides a complementary tool for teasing
 314 apart the coexistence mechanisms underlying evolutionary games. More broadly,
 315 our study builds on recent efforts to extend MCT beyond species competition (Park
 316 et al., 2024, 2026), providing a foundation for building unifying coexistence theory
 317 for competing entities, including species, pathogen strains, and behavioral strategies.

318 Supplementary Text

319 S1 Derivation of the strategic coexistence theory

320 To derive strategic coexistence theory (SCT), we begin with replicator equations:

$$\frac{dx_i}{dt} = x_i(\pi_i(\mathbf{x}) - \bar{\pi}(\mathbf{x})), \quad (\text{S1})$$

321 where x_i represents the frequency of strategy i , π_i represents the payoff of strategy
 322 i , and $\bar{\pi}$ represents the average payoff. Then, the mutual invasion condition for two
 323 competing strategies i and j can be written as

$$\pi_j(\mathbf{e}_j) < \pi_i(\mathbf{e}_j), \quad (\text{S2})$$

$$\pi_i(\mathbf{e}_i) < \pi_j(\mathbf{e}_i). \quad (\text{S3})$$

324 These inequalities will be preserved when we exponentiate the payoffs, $W_i(\mathbf{x}) =$
 325 $\exp(\pi_i(\mathbf{x}))$:

$$W_j(\mathbf{e}_j) < W_i(\mathbf{e}_j), \quad (\text{S4})$$

$$W_i(\mathbf{e}_i) < W_j(\mathbf{e}_i). \quad (\text{S5})$$

326 Rearranging, we obtain the following inequality:

$$\frac{W_i(\mathbf{e}_i)}{W_j(\mathbf{e}_i)} < 1 < \frac{W_i(\mathbf{e}_j)}{W_j(\mathbf{e}_j)}. \quad (\text{S6})$$

327 By multiplying $\sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}$ on both sides, we obtain:

$$\sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}} < \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}} < \sqrt{\frac{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}} \quad (\text{S7})$$

328 Thus, by defining niche overlap ρ and fitness difference κ_j/κ_i as follows,

$$\rho = \sqrt{\frac{W_i(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_j(\mathbf{e}_i)W_i(\mathbf{e}_j)}}, \quad (\text{S8})$$

$$\frac{\kappa_j}{\kappa_i} = \sqrt{\frac{W_j(\mathbf{e}_i)W_j(\mathbf{e}_j)}{W_i(\mathbf{e}_i)W_i(\mathbf{e}_j)}}, \quad (\text{S9})$$

329 we obtain the following inequality that defines conditions for mutual invasion and
 330 long-term coexistence of two competing strategies:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (\text{S10})$$

331 S2 Applications to classic two-strategy games

332 Consider a two-strategy game described by the following payoff matrix:

$$\begin{pmatrix} R & S \\ T & P \end{pmatrix}. \quad (\text{S11})$$

333 There are four classic games that we can define from this simple payoff matrix:
 334 prisoner's dilemma ($T > R > P > S$), hawk-dove game ($T > R, S > P$), stag-hunt
 335 game ($R > T, P > S$), and harmony game ($R > T, S > P$). Here, we show that
 336 each game corresponds to a distinct region in the niche-fitness difference space under
 337 SCT.

338 To do so, we first begin by deriving expressions for niche overlap ρ and fitness
 339 difference κ_j/κ_i . Writing x_1 to denote the frequency of strategy 1, the expected
 340 payoff for each strategy corresponds to:

$$\pi_1(\mathbf{x}) = Rx_1 + S(1 - x_1), \quad (\text{S12})$$

$$\pi_2(\mathbf{x}) = Tx_1 + P(1 - x_1). \quad (\text{S13})$$

341 Then, multiplicative payoffs for each strategy when rare can be written as:

$$W_1(1) = \exp(R) \quad (\text{S14})$$

$$W_1(0) = \exp(S) \quad (\text{S15})$$

$$W_2(1) = \exp(T) \quad (\text{S16})$$

$$W_2(0) = \exp(P) \quad (\text{S17})$$

342 Therefore, we have:

$$\rho = \exp\left(\frac{R + P - S - T}{2}\right), \quad (\text{S18})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{T + P - R - S}{2}\right). \quad (\text{S19})$$

343 As before, SCT allows us to predict the mutual invasion of two strategy using the
 344 following inequality:

$$\rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}. \quad (\text{S20})$$

345 Then, simple algebra reveals that each condition in mutual invasion inequality
 346 can be equivalently described by the payoff inequality:

$$\rho < \frac{\kappa_2}{\kappa_1} \iff R < T \quad (\text{S21})$$

$$\rho > \frac{\kappa_2}{\kappa_1} \iff R > T \quad (\text{S22})$$

$$\frac{\kappa_2}{\kappa_1} < \frac{1}{\rho} \iff P < S \quad (\text{S23})$$

$$\frac{\kappa_2}{\kappa_1} > \frac{1}{\rho} \iff P > S \quad (\text{S24})$$

347 These inequalities then allow us to classify outcomes of two-strategy games from the
 348 perspective of SCT. Note that the inequality between R and P does not affect the
 349 outcome.

350 **S3 Applications to mechanisms for the evolution of coopera-** 351 **tion**

352 We consider five mechanisms for the evolution of cooperation, all of which can be
 353 represented as a simple 2×2 payoff matrix (Nowak, 2006). Here, we go through each
 354 mechanism and derive measures for niche and fitness differences.

355 **Prisoner's dilemma.** Here, we consider a special case of Prisoner's dilemma
 356 which can be described by the following payoff matrix:

$$\begin{pmatrix} b - c & -c \\ b & 0 \end{pmatrix}, \quad (\text{S25})$$

357 where b represents the benefit for the recipient and c represents cost for the donor.
 358 In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0 \quad (\text{S26})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(c). \quad (\text{S27})$$

359 Therefore, as long as the cost is greater than zero, prisoner's dilemma always results
 360 in the competitive exclusion of cooperation.

361 **Kin selection.** Kin selection can be described by the following payoff matrix:

$$\begin{pmatrix} (b - c)(1 + r) & br - c \\ b - rc & 0 \end{pmatrix}, \quad (\text{S28})$$

362 where r represents the genetic relatedness. In this case, the niche and fitness differ-
 363 ences correspond to:

$$1 - \rho = 0 \quad (\text{S29})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(c - br). \quad (\text{S30})$$

364 Therefore, defection will exclude cooperation when $c > br$. Equivalently, cooperation
 365 will be favored when $r > c/b$, which recovers the classic result.

366 **Direct reciprocity.** Direct reciprocity can be described by the following payoff
 367 matrix:

$$\begin{pmatrix} (b - c)/(1 - w) & -c \\ b & 0 \end{pmatrix}, \quad (\text{S31})$$

where w represents the probability of next round. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 1 - \exp\left(\frac{(b - c)/(1 - w) + c - b}{2}\right) \quad (\text{S32})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{b + c - (b - c)/(1 - w)}{2}\right). \quad (\text{S33})$$

Therefore, defection will exclude cooperation when

$$\rho < \frac{\kappa_2}{\kappa_1} \quad (\text{S34})$$

$$\Leftrightarrow \exp\left(\frac{(b - c)/(1 - w) + c - b}{2}\right) < \exp\left(\frac{b + c - (b - c)/(1 - w)}{2}\right) \quad (\text{S35})$$

$$\Leftrightarrow bw < c \quad (\text{S36})$$

However, note that the following equality holds whenever $c > 0$:

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1}. \quad (\text{S37})$$

Therefore, when $w > c/b$, we will have

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1} < \rho, \quad (\text{S38})$$

which implies priority effect (equivalently, bistability). Even as $w \rightarrow 1^-$, this inequality holds and therefore, cooperation cannot competitively exclude defection.

Indirect reciprocity. Indirect reciprocity can be described by the following payoff matrix:

$$\begin{pmatrix} b - c & -c(1 - q) \\ b(1 - q) & 0 \end{pmatrix}, \quad (\text{S39})$$

where q represents social acquaintanceship. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 1 - \exp\left(\frac{(b - c)q}{2}\right), \quad (\text{S40})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{2c - (b + c)q}{2}\right). \quad (\text{S41})$$

In this case, the condition for the dominance of defection (and therefore competitive exclusion of cooperation) can be written as:

$$\rho < \frac{\kappa_2}{\kappa_1} \quad (\text{S42})$$

$$\Leftrightarrow bq < c \quad (\text{S43})$$

381 Moreover, the following inequality always holds as long as $c > 0$.

$$\frac{1}{\rho} < \frac{\kappa_2}{\kappa_1}, \quad (\text{S44})$$

382 which implies that coexistence between cooperation and defection is not possible.
 383 Therefore, when $q < c/b$, then defection will competitively exclude cooperation.
 384 Otherwise, the system will enter the priority-effect regime.

385 Interestingly, when $q \rightarrow 1$, we have

$$\rho = \exp\left(\frac{b-c}{2}\right) \quad (\text{S45})$$

$$\frac{\kappa_2}{\kappa_1} = \exp\left(\frac{c-b}{2}\right) \quad (\text{S46})$$

$$\frac{1}{\rho} = \exp\left(\frac{c-b}{2}\right) \quad (\text{S47})$$

$$(\text{S48})$$

386 In other words, the following inequality will hold:

$$\frac{1}{\rho} = \frac{\kappa_2}{\kappa_1} < \rho \quad (\text{S49})$$

387 **Network reciprocity.** Network reciprocity can be described by the following
 388 payoff matrix:

$$\begin{pmatrix} b-c & H-c \\ b-H & 0 \end{pmatrix}, \quad (\text{S50})$$

389 where $H = [(b-c)k - 2c]/[(k+1)(k-2)]$ and k represents the number of neighbors.
 390 In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0, \quad (\text{S51})$$

$$\frac{\kappa_2}{\kappa_1} = \exp(-H + c), \quad (\text{S52})$$

391 which implies that there is complete niche overlap and that fitness difference decreases
 392 with H . Since H is a monotonically decreasing function of k for $k > 2$, fitness
 393 difference will increase with k . In particular, cooperation will competitively exclude
 394 defection when $H > c$, meaning that:

$$b/c > k. \quad (\text{S53})$$

395 **Group selection.** Group selection can be described by the following payoff
 396 matrix:

$$\begin{pmatrix} (b-c)(m+n) & (b-c)m - cn \\ bn & 0 \end{pmatrix}, \quad (\text{S54})$$

where m represents the number of groups and n represents the group size. In this case, the niche and fitness differences correspond to:

$$1 - \rho = 0, \quad (\text{S55})$$

$$\frac{\kappa_2}{\kappa_1} = \exp((c - b)m + cn), \quad (\text{S56})$$

which implies that there is complete niche overlap. The fitness difference will increase with n but decrease with m , provided that $b > c$. In particular, cooperation will competitively exclude defection under the following condition:

$$1 + \frac{n}{m} < \frac{b}{c}. \quad (\text{S57})$$

References

- Adler, P. B., M. Detto, S. P. Ellner, T. L. Gibbs, Z. J. Gold, S. J. Leonard, J. I. Levine, A. E. Schiffer, C. Song, M. Stemkovski, et al. (2026). What have we learned from empirical applications of modern coexistence theory? *Ecology letters* 29(6), e70404.
- Allesina, S. (2026). Global stability of ecological and evolutionary dynamics via equivalence. *Proceedings of the National Academy of Sciences* 123(13), e2534915123.
- Archetti, M. and I. Scheuring (2011). Coexistence of cooperation and defection in public goods games. *Evolution* 65(4), 1140–1148.
- Axelrod, R. and W. D. Hamilton (1981). The evolution of cooperation. *science* 211(4489), 1390–1396.
- Bodin, F., G. Wang, and J. B. Plotkin (2026). Eco-evolutionary games in noisy environments. *bioRxiv*, 2026–05.
- Carlson, M. and D. Botstein (1982). Two differentially regulated mRNAs with different 5' ends encode secreted and intracellular forms of yeast invertase. *Cell* 28(1), 145–154.
- Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual review of Ecology and Systematics* 31(1), 343–366.
- Cooney, D. B. (2020). Analysis of multilevel replicator dynamics for general two-strategy social dilemma. *Bulletin of Mathematical Biology* 82(6), 66.
- Cordero, O. X., L.-A. Ventouras, E. F. DeLong, and M. F. Polz (2012). Public good dynamics drive evolution of iron acquisition strategies in natural bacterioplankton populations. *Proceedings of the National Academy of Sciences* 109(49), 20059–20064.

427 Cressman, R. (2003). *Evolutionary dynamics and extensive form games*, Volume 5.
428 MIT Press.

429 Cressman, R. and Y. Tao (2014). The replicator equation and other game dynamics.
430 *Proceedings of the National Academy of Sciences* 111(supplement_3), 10810–10817.

431 Dickinson, J. R. (1998). *Metabolism and molecular physiology of Saccharomyces*
432 *cerevisiae*. Taylor & Francis.

433 Doebeli, M. and C. Hauert (2005). Models of cooperation based on the Prisoner’s
434 Dilemma and the Snowdrift game. *Ecology letters* 8(7), 748–766.

435 Doebeli, M., C. Hauert, and T. Killingback (2004). The evolutionary origin of coop-
436 erators and defectors. *science* 306(5697), 859–862.

437 Godoy, O., N. J. Kraft, and J. M. Levine (2014). Phylogenetic relatedness and the
438 determinants of competitive outcomes. *Ecology letters* 17(7), 836–844.

439 Godoy, O. and J. M. Levine (2014). Phenology effects on invasion success: insights
440 from coupling field experiments to coexistence theory. *Ecology* 95(3), 726–736.

441 Gokhale, C. S. and A. Traulsen (2025). Higher-order equivalence of Lotka-Volterra
442 and replicator dynamics. *bioRxiv*, 2025–03.

443 Gore, J., H. Youk, and A. Van Oudenaarden (2009). Snowdrift game dynamics and
444 facultative cheating in yeast. *Nature* 459(7244), 253–256.

445 Hamilton, W. D. (1964). The genetical evolution of social behaviour. II. *Journal of*
446 *theoretical biology* 7(1), 17–52.

447 Hauert, C. (2002). Effects of space in 2×2 games. *International Journal of Bifur-*
448 *cation and Chaos* 12(07), 1531–1548.

449 Hauert, C. and M. Doebeli (2004). Spatial structure often inhibits the evolution of
450 cooperation in the snowdrift game. *Nature* 428(6983), 643–646.

451 Hofbauer, J. (1981). On the occurrence of limit cycles in the volterra-lotka equation.
452 *Nonlinear Analysis: Theory, Methods & Applications* 5(9), 1003–1007.

453 Hofbauer, J. and K. Sigmund (1998). *Evolutionary games and population dynamics*.
454 Cambridge university press.

455 Kocher, S. D., R. Mallarino, B. E. Rubin, D. W. Yu, H. E. Hoekstra, and N. E.
456 Pierce (2018). The genetic basis of a social polymorphism in halictid bees. *Nature*
457 *Communications* 9(1), 4338.

458 Kraft, N. J., O. Godoy, and J. M. Levine (2015). Plant functional traits and the mul-
459 tidimensional nature of species coexistence. *Proceedings of the National Academy*
460 *of Sciences* 112(3), 797–802.

- 461 Lewontin, R. C. (1961). Evolution and the theory of games. *Journal of Theoretical*
462 *Biology* 1(3), 382–403.
- 463 Lieberman, E., C. Hauert, and M. A. Nowak (2005). Evolutionary dynamics on
464 graphs. *Nature* 433(7023), 312–316.
- 465 Lin, H., D. Wang, Q. Wang, J. Mao, Y. Bai, and J. Qu (2024). Interspecific compe-
466 tition prevents the proliferation of social cheaters in an unstructured environment.
467 *The ISME Journal* 18(1), wrad038.
- 468 May, R. M. (1987). More evolution of cooperation. *Nature* 327(6117), 15–17.
- 469 Motro, U. (1991). Co-operation and defection: playing the field and the ESS. *Journal*
470 *of Theoretical Biology* 151(2), 145–154.
- 471 Nowak, M. A. (2006). Five rules for the evolution of cooperation. *science* 314(5805),
472 1560–1563.
- 473 Nowak, M. A. and R. M. May (1992). Evolutionary games and spatial chaos. *na-*
474 *ture* 359(6398), 826–829.
- 475 Nowak, M. A. and K. Sigmund (1998). Evolution of indirect reciprocity by image
476 scoring. *Nature* 393(6685), 573–577.
- 477 Nowak, M. A. and K. Sigmund (2005). Evolution of indirect reciprocity. *Na-*
478 *ture* 437(7063), 1291–1298.
- 479 Nowak, M. A., C. E. Tarnita, and T. Antal (2010). Evolutionary dynamics in struc-
480 tured populations. *Philosophical Transactions of the Royal Society B: Biological*
481 *Sciences* 365(1537), 19.
- 482 Page, K. M. and M. A. Nowak (2002). Unifying evolutionary dynamics. *Journal of*
483 *theoretical biology* 219(1), 93–98.
- 484 Park, S. W., S. Cobey, C. J. E. Metcalf, J. M. Levine, and B. T. Grenfell (2024).
485 Predicting pathogen mutual invasibility and co-circulation. *Science* 386(6718),
486 175–179.
- 487 Park, S. W., J. Levine, and B. Grenfell (2026). Unifying coexistence theory for
488 ecological communities and pathogen strain competition. *bioRxiv*, 2026–05.
- 489 Pollak, S., M. Gralka, Y. Sato, J. Schwartzman, L. Lu, and O. X. Cordero (2021).
490 Public good exploitation in natural bacterioplankton communities. *Science Ad-*
491 *vances* 7(31), eabi4717.
- 492 Schuster, P. and K. Sigmund (1983). Replicator dynamics. *Journal of theoretical*
493 *biology* 100(3), 533–538.

494 Shalizi, C. R. (2009). Dynamics of bayesian updating with dependent data and
495 misspecified models. *arXiv*.

496 Sieben, A. J., J. R. Mihaljevic, and L. G. Shoemaker (2022). Quantifying mechanisms
497 of coexistence in disease ecology. *Ecology* 103(12), e3819.

498 Slobodkin, L. B. (1964). The strategy of evolution. *American scientist* 52(3), 342–
499 357.

500 Smith, J. M. and G. R. Price (1973). The logic of animal conflict. *Nature* 246(5427),
501 15–18.

502 Souza, M. O., J. M. Pacheco, and F. C. Santos (2009). Evolution of cooperation
503 under N-person snowdrift games. *Journal of Theoretical Biology* 260(4), 581–588.

504 Steidinger, B. S. and J. D. Bever (2014). The coexistence of hosts with different
505 abilities to discriminate against cheater partners: an evolutionary game-theory
506 approach. *The American Naturalist* 183(6), 762–770.

507 Szabó, G. and G. Fath (2007). Evolutionary games on graphs. *Physics reports* 446(4-
508 6), 97–216.

509 Tarnita, C. E. and A. Traulsen (2025). Reconciling ecology and evolutionary game
510 theory or “When not to think cooperation”. *Proceedings of the National Academy
511 of Sciences* 122(14), e2413847122.

512 Taylor, P. D. and L. B. Jonker (1978). Evolutionary stable strategies and game
513 dynamics. *Mathematical biosciences* 40(1-2), 145–156.

514 Traulsen, A. and M. A. Nowak (2006). Evolution of cooperation by multilevel selec-
515 tion. *Proceedings of the National Academy of Sciences* 103(29), 10952–10955.

516 Trivers, R. L. (1971). The evolution of reciprocal altruism. *The Quarterly review of
517 biology* 46(1), 35–57.

518 Wang, G., Q. Su, L. Wang, and J. B. Plotkin (2025). Evolution of social behaviors
519 in noisy environments. *arXiv*.

520 Wedekind, C. and M. Milinski (2000). Cooperation through image scoring in humans.
521 *Science* 288(5467), 850–852.

522 West, S. A., A. S. Griffin, and A. Gardner (2007). Evolutionary explanations for
523 cooperation. *Current biology* 17(16), R661–R672.