

## ICCS310: Assignment 4

Possawat Sanorkam

possawat2017@hotmail.com

February 24, 2021

---

### 1: Eh? They Have The Same Cardinality?

---

Prove the following statements using rigorous mathematical reasoning:

(1)  $|[0, \frac{1}{2})| = |[0, 1)|$

*Proof:* We want to show that  $|[0, \frac{1}{2})| = |[0, 1)|$  by direct proof. By definition, let  $A$  and  $B$  be sets. Say  $A$  and  $B$  have the same cardinality (size), denoted by  $|A| = |B|$ , if there exists a bijection between them.

We want to show that there exist a function  $f$  such that  $f$  is a bijection. Let  $f : A \rightarrow B$ ,  $A = [0, \frac{1}{2})$ , and  $B = [0, 1)$ . Then, let  $f(x) = 2x$ . Let  $a \in A$  and  $a' \in A$ . From observation,  $f(a)$  is unique for any arbitrary  $a$ . We have that  $(\forall a \neq a')[f(a) \neq f(a')]$ , so  $f$  is injective. Let  $b \in B$ . From observation, every  $b$  can be obtain from some  $f(a)$ . We have that  $(\forall b)(\exists a)[f(a) = b]$ , so  $f$  is also surjective. According to the definition we stated before,  $f$  is a bijective function. Hence,  $|A| = |B|$ . Therefore,  $|[0, \frac{1}{2})| = |[0, 1)|$ .  $\square$

(2)  $|[0, 1)| = |(-1, 1)|$

*Proof:* We want to show that  $|[0, 1)| = |(-1, 1)|$  by direct proof. By definition, let  $A$  and  $B$  be sets. Say  $A$  and  $B$  have the same cardinality (size), denoted by  $|A| = |B|$ , if there exists a bijection between them. In addition,  $|A| = |B|$  if and only if  $|A| \leq |B|$  and  $|B| \leq |A|$ .

We want to show that there exist functions  $f$  that is injective and  $g$  that is also injective. Let  $f : A \rightarrow B$ ,  $g : B \rightarrow A$ ,  $A = [0, 1)$ , and  $B = (-1, 1)$ . Then, let  $f(x) = x$ . Let  $a \in A$  and  $a' \in A$ . From observation,  $f(a)$  is unique for any arbitrary  $a$ . We have that  $(\forall a \neq a')[f(a) \neq f(a')]$ , so  $f$  is injective. Then, let  $g(x) = \frac{(x+1)}{2}$ . Let  $b \in B$  and  $b' \in B$ . From observation,  $g(b)$  is unique for any arbitrary  $b$ . We have that  $(\forall b \neq b')[g(b) \neq g(b')]$ , so  $g$  is injective. Hence,  $|A| \leq |B|$  and  $|B| \leq |A|$  which implies that  $|A| = |B|$ . Therefore,  $|[0, 1)| = |(-1, 1)|$ .  $\square$

(3)  $|[0, 1)| = |\mathbb{R}|$

*Proof:* We want to show that  $|[0, 1)| = |\mathbb{R}|$  by direct proof. Besides, we have that  $|[0, 1)| = |(-1, 1)|$  which means we can show that  $|\mathbb{R}| = |(-1, 1)|$  instead. Say  $A$  and  $B$  have the same cardinality (size), denoted by  $|A| = |B|$ , if there exists a bijection between them. In addition,  $|A| = |B|$  if and only if  $|A| \leq |B|$  and  $|B| \leq |A|$ .

We want to show that there exist a function  $f$  such that  $f$  is a bijection. Let  $f : A \rightarrow \mathbb{R}$ , and  $A = (-1, 1)$ . Then, let  $f(x) = \frac{x}{1-x^2}$ . This function is continuous on domain  $A$  when  $x \neq 1$  and  $x \neq -1$ . Let  $a \in A$  and  $a' \in A$ . From observation,  $f(a)$  is unique for any arbitrary  $a$ . We have that  $(\forall a \neq a')[f(a) \neq f(a')]$ , so  $f$  is injective. Let  $b \in B$ . From observation, every  $b$  can be obtain from some  $f(a)$ . We have that  $(\forall b)(\exists a)[f(a) = b]$ , so  $f$  is also surjective. According to the definition we stated before,  $f$  is a bijective function. So,  $|A| = |\mathbb{R}|$  or  $|\mathbb{R}| = |(-1, 1)|$ . Hence,  $|\mathbb{R}| = |(-1, 1)|$  implies that  $|\mathbb{R}| = |[0, 1)|$  also. Therefore,  $|[0, 1)| = |\mathbb{R}|$ .  $\square$

---

### 2: The Power Set of A

---

(1) Prove that  $|2^A| = |\{0,1\}^A|$ .

*Proof:*

(2) Prove that  $|A| < |\{0,1\}^A|$  and conclude that  $|A| < |2^A|$ .

*Proof:*

### 3: Hamming Code

Consider applying the Hamming coding scheme to send 8 bits of data. This will require 4 parity bits, so an encoded code word in this scheme is 12 bits long.

(1) If the data bits are  $d_1, d_2, d_3, \dots, d_8$ , what is  $\beta_2$  in terms of  $d_i$ 's?

*Solution:*  $\beta_2 = d_1 \oplus d_3 \oplus d_4 \oplus d_6 \oplus d_7$

(2) Encode the following 8-bit data: 01101010.

*Solution:* Hamming Code =  $(\beta_1 \beta_2 d_1 \beta_4 d_2 d_3 d_4 \beta_8 d_5 d_6 d_7 d_8)$

Encode 01101010 by adding the parity bits as followed

$$\beta_1 = d_1 \oplus d_2 \oplus d_4 \oplus d_5 \oplus d_7 = 0 \oplus 1 \oplus 0 \oplus 1 \oplus 1 = 1 \quad (1)$$

$$\beta_2 = d_1 \oplus d_3 \oplus d_4 \oplus d_6 \oplus d_7 = 0 \oplus 1 \oplus 0 \oplus 0 \oplus 1 = 0 \quad (2)$$

$$\beta_4 = d_2 \oplus d_3 \oplus d_4 \oplus d_8 = 1 \oplus 1 \oplus 0 \oplus 0 = 0 \quad (3)$$

$$\beta_8 = d_5 \oplus d_6 \oplus d_7 \oplus d_8 = 1 \oplus 0 \oplus 1 \oplus 0 = 0 \quad (4)$$

Therefore, encoded bits are 100011001010

(3) Assuming that at most a single single bit flip, decide the following codewords (indicate also whether there was any error):

*Solution:* Hamming Code =  $(\beta_1 \beta_2 d_1 \beta_4 d_2 d_3 d_4 \beta_8 d_5 d_6 d_7 d_8)$

(i) 010011111000

So,  $\beta_1 = 0, \beta_2 = 1, \beta_4 = 0$ , and  $\beta_8 = 1$ .

$$\beta_1 \oplus d_1 \oplus d_2 \oplus d_4 \oplus d_5 \oplus d_7 = 0 \oplus 0 \oplus 1 \oplus 1 \oplus 1 \oplus 0 = 1 \quad (5)$$

$$\beta_2 \oplus d_1 \oplus d_3 \oplus d_4 \oplus d_6 \oplus d_7 = 1 \oplus 0 \oplus 1 \oplus 1 \oplus 0 \oplus 0 = 1 \quad (6)$$

$$\beta_4 \oplus d_2 \oplus d_3 \oplus d_4 \oplus d_8 = 0 \oplus 1 \oplus 1 \oplus 1 \oplus 0 = 1 \quad (7)$$

$$\beta_8 \oplus d_5 \oplus d_6 \oplus d_7 \oplus d_8 = 1 \oplus 1 \oplus 0 \oplus 0 \oplus 0 = 0 \quad (8)$$

Error Position is  $0111_2 = 7$ . Corrected Data is 010011011000.

(ii) 011101010010

So,  $\beta_1 = 0, \beta_2 = 1, \beta_4 = 1$ , and  $\beta_8 = 1$ .

$$\beta_1 \oplus d_1 \oplus d_2 \oplus d_4 \oplus d_5 \oplus d_7 = 0 \oplus 1 \oplus 0 \oplus 0 \oplus 0 \oplus 0 = 1 \quad (9)$$

$$\beta_2 \oplus d_1 \oplus d_3 \oplus d_4 \oplus d_6 \oplus d_7 = 1 \oplus 1 \oplus 1 \oplus 0 \oplus 0 \oplus 0 = 1 \quad (10)$$

$$\beta_4 \oplus d_2 \oplus d_3 \oplus d_4 \oplus d_8 = 1 \oplus 0 \oplus 1 \oplus 0 \oplus 0 = 0 \quad (11)$$

$$\beta_8 \oplus d_5 \oplus d_6 \oplus d_7 \oplus d_8 = 1 \oplus 0 \oplus 0 \oplus 1 \oplus 0 = 0 \quad (12)$$

Error Position is  $0011_2 = 3$ . Corrected Data is 010101010010.

---

#### 4: Same Number of 0s and 1s

---

Consider the language  $L = \{w \in \{0,1\}^* \mid w \text{ contains an equal number of 0s and 1s}\}$ . Show that  $L$  is (Turing) decidable by providing a TM that decides it (a medium-level detail is preferred).

*Proof:*

---

#### 5: Infinite DFA

---

Show that the following language is (Turing) decidable:

$$\text{IDFA} = \{\langle M \rangle \mid M \text{ is a DFA and } L(M) \text{ is an infinite language}\}.$$

*Proof:* We want to show that IDFA is decidable. So, we will construct a TM  $T$  that decides IDFA. For all DFAs  $M$ , we want  $T(\langle M \rangle)$  to accept if  $L(M)$  is infinite language, else it will reject.

---

#### 6: Lucky 9

---

(1) Let  $L_1 \subseteq \Sigma^*$  be defined as

$$L_1 = \begin{cases} \emptyset & \text{if } 2^{74207281} - 1 \text{ is prime} \\ \{99\} & \text{if } 2^{74207281} - 1 \text{ is not prime} \end{cases}$$

Prove that  $L_1$  is (Turing) decidable.

*Proof:*

(2) Let  $L_2 \subseteq \Sigma^*$  be defined as

$w \in L_2 \iff w$  appears somewhere (not necessarily consecutively) in the decimal expansion of  $\pi$

Prove that  $L_2$  is (Turing) decidable.

*Proof:*

---

#### 7: $\beta$ -reduction

---

(1)  $(\lambda z.z)(\lambda z.zz)(\lambda z.zy)$

*Solution:*

$$(\lambda z.z)(\lambda z.zz)(\lambda z.zy) \rightarrow_{\beta} (\lambda z.zz)(\lambda z.zy) \quad (13)$$

$$\rightarrow_{\beta} (\lambda z.zy)(\lambda z.zy) \quad (14)$$

$$\rightarrow_{\beta} (\lambda z.zy)y \quad (15)$$

$$\rightarrow_{\beta} yy \quad (16)$$

(2)  $((\lambda x.\lambda y.(xy))(\lambda y.y))w$

*Solution:*

$$(((\lambda x.\lambda y.(xy))(\lambda y.y))w) \rightarrow_{\beta} (\lambda y.(xy))w \quad (17)$$

$$\rightarrow_{\beta} xw \quad (18)$$

## 8: Fibonacci

Using the functions we have developed (e.g., `pred`, `if_then_else`, `mult`, `add`, etc.), write down an explicit  $\lambda$ -term `fib` such that  $\overline{fib} \ \bar{n} =_{\beta} \overline{f(n)}$ .

*Solution:*

## 9: Power Of 2

Implement a  $\lambda$ -term for the  $pow(n) = 2^n$

*Solution:*