

# ICCS310: Assignment 5

Possawat Sanorkam

possawat2017@hotmail.com

March 12, 2021

---

## 1: Reject TM

---

$$\text{REJECT}_{\text{TM}} = \{\langle M, x \rangle \mid M \text{ is a TM that rejects input } x\}$$

Show directly (i.e., without resorting to reduction) that  $\text{REJECT}_{\text{TM}}$  is undecidable.

*Proof:*

---

## 2: Accept vs. Reject

---

$$\text{REJECT}_{\text{TM}} = \{\langle M, x \rangle \mid M \text{ is a TM that rejects input } x\}$$

Show directly (i.e., without resorting to reduction) that  $\text{REJECT}_{\text{TM}}$  is undecidable.

*Proof:*

---

## 3: Reverse on TM

---

$$\text{REJECT}_{\text{TM}} = \{\langle M, x \rangle \mid M \text{ is a TM that rejects input } x\}$$

Show directly (i.e., without resorting to reduction) that  $\text{REJECT}_{\text{TM}}$  is undecidable.

*Proof:*

---

## 4: Undecidability

---

$$\text{REJECT}_{\text{TM}} = \{\langle M, x \rangle \mid M \text{ is a TM that rejects input } x\}$$

Show directly (i.e., without resorting to reduction) that  $\text{REJECT}_{\text{TM}}$  is undecidable.

*Proof:*

---

## 5: Total Is No Harder Than Finite

---

$$\text{REJECT}_{\text{TM}} = \{\langle M, x \rangle \mid M \text{ is a TM that rejects input } x\}$$

Show directly (i.e., without resorting to reduction) that  $\text{REJECT}_{\text{TM}}$  is undecidable.

*Proof:*

---

## 6: Finite Is No Harder Than Total

---

$$\text{REJECT}_{\text{TM}} = \{\langle M, x \rangle \mid M \text{ is a TM that rejects input } x\}$$

Show directly (i.e., without resorting to reduction) that  $\text{REJECT}_{\text{TM}}$  is undecidable.

*Proof:*

---

### 7: Extra: Undecidability of Nontrivial Properties

---

$$\text{REJECT}_{\text{TM}} = \{\langle M, x \rangle \mid M \text{ is a TM that rejects input } x\}$$

Show directly (i.e., without resorting to reduction) that  $\text{REJECT}_{\text{TM}}$  is undecidable.

*Proof:*