

# ICCS310: Assignment 5

Possawat Sanorkam

possawat2017@hotmail.com

March 12, 2021

---

## 1: Reject TM

---

$$\text{REJECT}_{\text{TM}} = \{\langle M, x \rangle \mid M \text{ is a TM that rejects input } x\}$$

Show directly (i.e., without resorting to reduction) that  $\text{REJECT}_{\text{TM}}$  is undecidable.

*Proof:*

---

## 2: Accept vs. Reject

---

$$\text{ACCEPT}_{\text{TM}} = \{\langle M, x \rangle \mid M \text{ is a TM that accepts input } x\}$$

(i) Prove that  $\text{ACCEPT}_{\text{TM}} \leq \text{REJECT}_{\text{TM}}$

*Proof:*

(ii) Prove that  $\text{REJECT}_{\text{TM}} \leq \text{ACCEPT}_{\text{TM}}$

*Proof:*

---

## 3: Reverse on TM

---

$$\text{T} = \{\langle M \rangle \mid M \text{ is a TM that accepts } \text{rev}(w) \text{ whenever it accepts } w\}$$

where  $\text{rev}(w)$  is the reverse of the string  $w$ . Show that  $\text{T}$  is undecidable.

*Proof:*

---

## 4: Undecidability

---

(i) Show that

$$\text{TOTAL} = \{\langle M \rangle \mid M \text{ is a Turing machine that halts on every input}\}$$

is undecidable

*Proof:*

(ii) Show that

$$\text{FINITE} = \{\langle M \rangle \mid M \text{ is a Turing machine and } L(M) \text{ is a finite set}\}$$

is undecidable

*Proof:*

(iii) Show that

$$\text{REGULAR} = \{\langle M \rangle \mid M \text{ is a Turing machine and } L(M) \text{ is regular}\}$$

where  $\text{rev}(w)$  is the reverse of the string  $w$ . Show that T is undecidable.

*Proof:*

---

### 5: Total Is No Harder Than Finite

---

Prove that

$$\text{TOTAL} \leq_T \text{FINITE}$$

is undecidable

*Proof:*

---

### 6: Finite Is No Harder Than Total

---

Prove that

$$\text{FINITE} \leq_T \text{TOTAL}$$

is undecidable

*Proof:*

---

### 7: Extra: Undecidability of Nontrivial Properties

---

*Proof:* It is non trivial. How to decide on it though?