

## SECTION 1

### MODULE 3 : Elementary Combinatorics

## Permutation

The different arrangements which can be made out of a given set of things, by taking some or all of them at a time are called their Permutations.

## Permutation

The different arrangements which can be made out of a given set of things, by taking some or all of them at a time are called their Permutations.

The number of permutations of  $n$  different things taken  $r \leq n$  at a time is denoted by

$$P(n, r) \quad \text{or} \quad {}^n P_r.$$

## Permutation

The different arrangements which can be made out of a given set of things, by taking some or all of them at a time are called their Permutations.

The number of permutations of  $n$  different things taken  $r \leq n$  at a time is denoted by

$$P(n, r) \quad \text{or} \quad {}^n P_r.$$

The value of

$$P(n, r) = \frac{n!}{(n-r)!}.$$

## Permutation

The different arrangements which can be made out of a given set of things, by taking some or all of them at a time are called their Permutations.

The number of permutations of  $n$  different things taken  $r \leq n$  at a time is denoted by

$$P(n, r) \quad \text{or} \quad {}^n P_r.$$

The value of

$$P(n, r) = \frac{n!}{(n-r)!}.$$

Note :

$$0! = 1$$

## Permutation

The different arrangements which can be made out of a given set of things, by taking some or all of them at a time are called their Permutations.

The number of permutations of  $n$  different things taken  $r \leq n$  at a time is denoted by

$$P(n, r) \quad \text{or} \quad {}^n P_r.$$

The value of

$$P(n, r) = \frac{n!}{(n-r)!}.$$

Note :

$$0! = 1$$

$$P(n, n) = n!$$

## Example

(1) If  ${}^nP_2 = 72$ , find the value of  $n$ .

## Example

(1) If  ${}^nP_2 = 72$ , find the value of  $n$ .

**Solution :** Here

$$\begin{aligned} {}^nP_2 = 72 &\implies \frac{n!}{(n-2)!} = 72 \\ &\implies \frac{n \cdot (n-1) \cdot (n-2)!}{(n-2)!} = 72 \\ &\implies n(n-1) = 72 \implies n^2 - n - 72 = 0 \\ &\implies (n-9)(n+8) = 72 \end{aligned}$$

Hence

$$n = 9.$$



(2) If  ${}^{2n+1}P_{n-1} : {}^{2n-1}P_n = 3 : 5$ , find the value of  $n$ .

# Elementary Combinatorics

(2) If  ${}^{2n+1}P_{n-1} : {}^{2n-1}P_n = 3 : 5$ , find the value of  $n$ .

**Solution :** Here

$${}^{2n+1}P_{n-1} = \frac{(2n+1)!}{(n+2)!} \quad \text{and} \quad {}^{2n-1}P_n = \frac{(2n-1)!}{(n-1)!}$$

$$\begin{aligned} {}^{2n+1}P_{n-1} : {}^{2n-1}P_n = 3 : 5 &\implies \frac{(2n+1)!}{(n+2)!} \times \frac{(n-1)!}{(2n-1)!} = \frac{3}{5} \\ &\implies \frac{(2n+1) \times (2n) \times (2n-1)! \times (n-1)!}{(n+2) \times (n+1) \times (n) \times (n-1)! \times (2n-1)!} = \frac{3}{5} \\ &\implies \frac{(2n+1)(2n)}{(n+2)(n+1)(n)} = \frac{3}{5} \\ &\implies \frac{4n+2}{n^2+3n+2} = \frac{3}{5} \\ &\implies 3n^2 - 11n - 4 = 0 \\ &\implies (n-4)(3n+1) = 0 \end{aligned}$$

Hence

$$n = 4.$$

## Combination

The different selection which can be made out of a given set of things, by taking some or all of them at a time irrespective of order are called their Combinations.

## Combination

The different selection which can be made out of a given set of things, by taking some or all of them at a time irrespective of order are called their Combinations.

The number of combination of  $n$  different things taken  $r \leq n$  at a time is denoted by

$$C(n, r) \quad \text{or} \quad {}^nC_r.$$

## Combination

The different selection which can be made out of a given set of things, by taking some or all of them at a time irrespective of order are called their Combinations.

The number of combination of  $n$  different things taken  $r \leq n$  at a time is denoted by

$$C(n, r) \quad \text{or} \quad {}^nC_r.$$

The value of

$$C(n, r) = \frac{n!}{r!(n-r)!}.$$

## Combination

The different selection which can be made out of a given set of things, by taking some or all of them at a time irrespective of order are called their Combinations.

The number of combination of  $n$  different things taken  $r \leq n$  at a time is denoted by

$$C(n, r) \quad \text{or} \quad {}^nC_r.$$

The value of

$$C(n, r) = \frac{n!}{r!(n-r)!}.$$

Note :

$$C(n, n) = C(n, 0) = 1$$

Properties of  $C(n, r)$  or  ${}^nC_r$ .

1. For  $0 \leq r \leq n$ ,

$${}^nC_r = {}^nC_{n-r}$$

Properties of  $C(n, r)$  or  ${}^nC_r$ .

1. For  $0 \leq r \leq n$ ,

$${}^nC_r = {}^nC_{n-r}$$

2. For  $0 \leq r \leq n$ ,

$${}^nC_r + {}^nC_{r-1} = {}^{n+1}C_r.$$



Properties of  $C(n, r)$  or  ${}^nC_r$ .

1. For  $0 \leq r \leq n$ ,

$${}^nC_r = {}^nC_{n-r}$$

2. For  $0 \leq r \leq n$ ,

$${}^nC_r + {}^nC_{r-1} = {}^{n+1}C_r.$$

3.

$${}^nC_x = {}^nC_y \implies x = y \text{ or } x + y = n.$$

## Example

(1) If  ${}^{18}C_r = {}^{18}C_{r+2}$ , then find the value of  ${}^rC_5$ .

**Solution :** Here  ${}^{18}C_r = {}^{18}C_{r+2}$ , then by property of  ${}^nC_r$  we have

## Example

(1) If  ${}^{18}C_r = {}^{18}C_{r+2}$ , then find the value of  ${}^rC_5$ .

**Solution :** Here  ${}^{18}C_r = {}^{18}C_{r+2}$ , then by property of  ${}^nC_r$  we have

$$r + (r + 2) = 18 \implies 2r + 2 = 18 \implies r = 8.$$

Now

## Example

(1) If  ${}^{18}C_r = {}^{18}C_{r+2}$ , then find the value of  ${}^rC_5$ .

**Solution :** Here  ${}^{18}C_r = {}^{18}C_{r+2}$ , then by property of  ${}^nC_r$  we have

$$r + (r + 2) = 18 \implies 2r + 2 = 18 \implies r = 8.$$

Now

$${}^rC_5 = {}^8C_5 = \frac{8!}{5! \times (8-5)!} = \frac{8 \times 7 \times 6 \times 5!}{5! \times 3 \times 2 \times 1} = 56.$$

(2) In how many ways 3 questions can be selected from 9 questions ?

**Solution :** The required number of ways is

## Example

(1) If  ${}^{18}C_r = {}^{18}C_{r+2}$ , then find the value of  ${}^rC_5$ .

**Solution :** Here  ${}^{18}C_r = {}^{18}C_{r+2}$ , then by property of  ${}^nC_r$  we have

$$r + (r + 2) = 18 \implies 2r + 2 = 18 \implies r = 8.$$

Now

$${}^rC_5 = {}^8C_5 = \frac{8!}{5! \times (8-5)!} = \frac{8 \times 7 \times 6 \times 5!}{5! \times 3 \times 2 \times 1} = 56.$$

(2) In how many ways 3 questions can be selected from 9 questions ?

**Solution :** The required number of ways is

$${}^9C_3 = \frac{9 \times 8 \times 7}{3 \times 2 \times 1} = 84.$$

(3) If  ${}^nC_x = 56$  and  ${}^nP_x = 336$ , find  $n$  and  $x$ .

**Solution :** Given  ${}^nC_x = 56$  and  ${}^nP_x = 336$ .

Therefore

$$\frac{n!}{x! \times (n-x)!} = 56 \quad \text{and} \quad \frac{n!}{(n-x)!} = 336.$$

So

(3) If  ${}^nC_x = 56$  and  ${}^nP_x = 336$ , find  $n$  and  $x$ .

**Solution :** Given  ${}^nC_x = 56$  and  ${}^nP_x = 336$ .

Therefore

$$\frac{n!}{x! \times (n-x)!} = 56 \quad \text{and} \quad \frac{n!}{(n-x)!} = 336.$$

So

$$\frac{336}{x!} = 56 \implies \frac{336}{56} = x! \implies x! = 6 \implies x = 3$$

Now

(3) If  ${}^nC_x = 56$  and  ${}^nP_x = 336$ , find  $n$  and  $x$ .

**Solution :** Given  ${}^nC_x = 56$  and  ${}^nP_x = 336$ .

Therefore

$$\frac{n!}{x! \times (n-x)!} = 56 \quad \text{and} \quad \frac{n!}{(n-x)!} = 336.$$

So

$$\frac{336}{x!} = 56 \implies \frac{336}{56} = x! \implies x! = 6 \implies x = 3$$

Now

$${}^nP_x = 336 \implies {}^nP_3 = 336.$$

which gives

$$\begin{aligned} \frac{n!}{(n-3)!} = 336 &\implies \frac{n \times (n-1) \times (n-2) \times (n-3)!}{(n-3)!} = 336 \\ &\implies n \times (n-1) \times (n-2) = 336 = 8 \times 7 \times 6 \end{aligned}$$

Hence  $n = 8$ .



## Mathematical Induction

- It is a technique by which one can prove mathematical statements involving positive integers.

## Mathematical Induction

- It is a technique by which one can prove mathematical statements involving positive integers.
- It reduces the proof to a finite number of steps and guarantee that there is no positive integer for which the statement fails.

## Mathematical Induction

- It is a technique by which one can prove mathematical statements involving positive integers.
- It reduces the proof to a finite number of steps and guarantee that there is no positive integer for which the statement fails.

### Principle of Mathematical Induction

## Mathematical Induction

- It is a technique by which one can prove mathematical statements involving positive integers.
- It reduces the proof to a finite number of steps and guarantee that there is no positive integer for which the statement fails.

### Principle of Mathematical Induction

Let  $S(n)$  be the statement involving positive integer  $n = 1, 2, 3, \dots$  such that

## Mathematical Induction

- It is a technique by which one can prove mathematical statements involving positive integers.
- It reduces the proof to a finite number of steps and guarantee that there is no positive integer for which the statement fails.

### Principle of Mathematical Induction

Let  $S(n)$  be the statement involving positive integer  $n = 1, 2, 3, \dots$  such that

(i)  $S(1)$  is true.

## Mathematical Induction

- It is a technique by which one can prove mathematical statements involving positive integers.
- It reduces the proof to a finite number of steps and guarantee that there is no positive integer for which the statement fails.

### Principle of Mathematical Induction

Let  $S(n)$  be the statement involving positive integer  $n = 1, 2, 3, \dots$  such that

- (i)  $S(1)$  is true.
- (ii)  $S(k+1)$  is true whenever  $S(k)$  is true.

## Mathematical Induction

- It is a technique by which one can prove mathematical statements involving positive integers.
- It reduces the proof to a finite number of steps and guarantee that there is no positive integer for which the statement fails.

### Principle of Mathematical Induction

Let  $S(n)$  be the statement involving positive integer  $n = 1, 2, 3, \dots$  such that

- (i)  $S(1)$  is true.
- (ii)  $S(k+1)$  is true whenever  $S(k)$  is true.

Then  $S(n)$  is true for all positive integer  $n$ .

## Example

(1) Show that

$$1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1)$$

where  $n \geq 1$  by mathematical induction.



## Example

(1) Show that

$$1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1)$$

where  $n \geq 1$  by mathematical induction.

**Solution :**

## Example

(1) Show that

$$1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1)$$

where  $n \geq 1$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : 1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1).$$

For  $n = 1$ ,

## Example

(1) Show that

$$1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1)$$

where  $n \geq 1$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : 1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1).$$

For  $n = 1$ , LHS =  $1^2 = 1$  and

## Example

(1) Show that

$$1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1)$$

where  $n \geq 1$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : 1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1).$$

For  $n = 1$ , LHS =  $1^2 = 1$  and RHS =  $\frac{1}{6}n(n+1)(2n+1) = \frac{1}{6} \times 1 \times (1+1)(2 \times 1 + 1) = 1$ .

## Example

(1) Show that

$$1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1)$$

where  $n \geq 1$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : 1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1).$$

For  $n = 1$ , LHS =  $1^2 = 1$  and RHS =  $\frac{1}{6}n(n+1)(2n+1) = \frac{1}{6} \times 1 \times (1+1)(2 \times 1 + 1) = 1$ .  
Therefore LHS=RHS.

## Example

(1) Show that

$$1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1)$$

where  $n \geq 1$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : 1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1).$$

For  $n = 1$ , LHS =  $1^2 = 1$  and RHS =  $\frac{1}{6}n(n+1)(2n+1) = \frac{1}{6} \times 1 \times (1+1)(2 \times 1 + 1) = 1$ .  
Therefore LHS=RHS. Hence  $S(1)$  is true.

## Example

(1) Show that

$$1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1)$$

where  $n \geq 1$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : 1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{1}{6}n(n+1)(2n+1).$$

For  $n = 1$ , LHS =  $1^2 = 1$  and RHS =  $\frac{1}{6}n(n+1)(2n+1) = \frac{1}{6} \times 1 \times (1+1)(2 \times 1 + 1) = 1$ .  
Therefore LHS=RHS. Hence  $S(1)$  is true.

Assume  $S(k)$  is true

$$1^2 + 2^2 + 3^2 + \cdots + k^2 = \frac{1}{6}k(k+1)(2k+1)$$

We show that  $S(k+1)$  is true

$$S(k+1) : 1^2 + 2^2 + 3^2 + \cdots + k^2 + (k+1)^2 = \frac{1}{6}(k+1)(k+2)(2k+3).$$

Now

$$\begin{aligned} 1^2 + 2^2 + 3^2 + \cdots + k^2 + (k+1)^2 &= (1^2 + 2^2 + 3^2 + \cdots + k^2) + (k+1)^2 \\ &= \frac{1}{6}k(k+1)(2k+1) + (k+1)^2 \\ &= (k+1) \left[ \frac{1}{6}k(2k+1) + (k+1) \right] \\ &= (k+1) \left[ \frac{1}{6}(2k^2 + 7k + 6) \right] \\ &= \frac{1}{6}(k+1)(k+2)(2k+3). \end{aligned}$$

Hence  $S(k+1)$  is true. So  $S(k+1)$  is true whenever  $S(k)$  is true. By Principal of Mathematical Induction  $S(n)$  is true for all positive integer  $n$ .



(2) Show that  $n^2 > 2n + 1$  where  $n \geq 3$  by mathematical induction.

(2) Show that  $n^2 > 2n + 1$  where  $n \geq 3$  by mathematical induction.

**Solution :**

(2) Show that  $n^2 > 2n + 1$  where  $n \geq 3$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : n^2 > 2n + 1, \text{ where } n \geq 3$$

For  $n = 3$ ,

(2) Show that  $n^2 > 2n + 1$  where  $n \geq 3$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : n^2 > 2n + 1, \text{ where } n \geq 3$$

For  $n = 3$ , LHS is  $3^2 = 9$  and

(2) Show that  $n^2 > 2n + 1$  where  $n \geq 3$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : n^2 > 2n + 1, \text{ where } n \geq 3$$

For  $n = 3$ , LHS is  $3^2 = 9$  and RHS is  $2n + 1 = 7$ .

(2) Show that  $n^2 > 2n + 1$  where  $n \geq 3$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : n^2 > 2n + 1, \text{ where } n \geq 3$$

For  $n = 3$ , LHS is  $3^2 = 9$  and RHS is  $2n + 1 = 7$ . Therefore  $\text{LHS} > \text{RHS}$ .

(2) Show that  $n^2 > 2n + 1$  where  $n \geq 3$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : n^2 > 2n + 1, \text{ where } n \geq 3$$

For  $n = 3$ , LHS is  $3^2 = 9$  and RHS is  $2n + 1 = 7$ . Therefore  $\text{LHS} > \text{RHS}$ . Hence  $S(3)$  is true.

(2) Show that  $n^2 > 2n + 1$  where  $n \geq 3$  by mathematical induction.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : n^2 > 2n + 1, \text{ where } n \geq 3$$

For  $n = 3$ , LHS is  $3^2 = 9$  and RHS is  $2n + 1 = 7$ . Therefore  $\text{LHS} > \text{RHS}$ . Hence  $S(3)$  is true.

Assume  $S(k)$  is true

$$k^2 > 2k + 1$$

Now we show that  $S(k + 1)$  is true

$$S(k + 1) : (k + 1)^2 > 2k + 3.$$



Now

$$\begin{aligned}(k+1)^2 &= k^2 + 2k + 1 = (k^2) + 2k + 1 \\ &> 2k + 1 + 2k + 1 \\ &> 2k + 1 + 1 + 1 \\ &= 2k + 3\end{aligned}$$

Hence  $S(k+1)$  is true.

So  $S(k+1)$  is true whenever  $S(k)$  is true. By Principal of Mathematical Induction  $S(n)$  is true for all positive integer  $n \geq 3$ .

(3) For every positive integer  $n$ , prove that  $7^n - 3^n$  is divisible by 4.

(3) For every positive integer  $n$ , prove that  $7^n - 3^n$  is divisible by 4.

**Solution :**

(3) For every positive integer  $n$ , prove that  $7^n - 3^n$  is divisible by 4.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : 7^n - 3^n \text{ is divisible by } 4.$$

(3) For every positive integer  $n$ , prove that  $7^n - 3^n$  is divisible by 4.

**Solution** : Let  $S(n)$  be the given statement

$$S(n) : 7^n - 3^n \text{ is divisible by } 4.$$

For  $n = 1$ ,

(3) For every positive integer  $n$ , prove that  $7^n - 3^n$  is divisible by 4.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : 7^n - 3^n \text{ is divisible by } 4.$$

For  $n = 1$ ,  $7^n - 3^n$  is  $7 - 3 = 4$  and it is divisible by 4.

(3) For every positive integer  $n$ , prove that  $7^n - 3^n$  is divisible by 4.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : 7^n - 3^n \text{ is divisible by } 4.$$

For  $n = 1$ ,  $7^n - 3^n$  is  $7 - 3 = 4$  and it is divisible by 4. Hence  $S(1)$  is true.

(3) For every positive integer  $n$ , prove that  $7^n - 3^n$  is divisible by 4.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : 7^n - 3^n \text{ is divisible by } 4.$$

For  $n = 1$ ,  $7^n - 3^n$  is  $7 - 3 = 4$  and it is divisible by 4. Hence  $S(1)$  is true.

Assume  $S(k)$  is true

$$7^k - 3^k \text{ is divisible by } 4.$$



(3) For every positive integer  $n$ , prove that  $7^n - 3^n$  is divisible by 4.

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : 7^n - 3^n \text{ is divisible by } 4.$$

For  $n = 1$ ,  $7^n - 3^n$  is  $7 - 3 = 4$  and it is divisible by 4. Hence  $S(1)$  is true.

Assume  $S(k)$  is true

$$7^k - 3^k \text{ is divisible by } 4.$$

Now we show that  $S(k+1)$  is true  $7^{k+1} - 3^{k+1}$  is divisible by 4.

Now

$$\begin{aligned}7^{k+1} - 3^{k+1} &= 7^k 7 - 3^k 3 \\&= 7^k 7 - 7^k 3 + 7^k 3 - 3^k 3 \\&= 7^k (7 - 3) + 3(7^k - 3^k) \\&= 7^k (4) + 3(7^k - 3^k)\end{aligned}$$

which is divisible by 4.

Hence  $S(k+1)$  is true.

So  $S(k+1)$  is true whenever  $S(k)$  is true. By Principal of Mathematical Induction  $S(n)$  is true for all positive integer  $n$ .

(4) Prove that  $(1+x)^n \geq (1+nx)$ , for all natural number  $n$ , where  $x > -1$ .

(4) Prove that  $(1+x)^n \geq (1+nx)$ , for all natural number  $n$ , where  $x > -1$ .

**Solution :**

(4) Prove that  $(1+x)^n \geq (1+nx)$ , for all natural number  $n$ , where  $x > -1$ .

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : (1+x)^n \geq (1+nx)$$

where  $x > -1$  For  $n = 1$ ,

(4) Prove that  $(1+x)^n \geq (1+nx)$ , for all natural number  $n$ , where  $x > -1$ .

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : (1+x)^n \geq (1+nx)$$

where  $x > -1$  For  $n = 1$ ,  $\text{RHS} = (1+x)^1 = 1+x$  and

(4) Prove that  $(1+x)^n \geq (1+nx)$ , for all natural number  $n$ , where  $x > -1$ .

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : (1+x)^n \geq (1+nx)$$

where  $x > -1$  For  $n = 1$ ,  $\text{RHS} = (1+x)^1 = 1+x$  and  $\text{LHS} = (1+nx) = (1+x)$ .

(4) Prove that  $(1+x)^n \geq (1+nx)$ , for all natural number  $n$ , where  $x > -1$ .

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : (1+x)^n \geq (1+nx)$$

where  $x > -1$  For  $n = 1$ ,  $\text{RHS} = (1+x)^1 = 1+x$  and  $\text{LHS} = (1+nx) = (1+x)$ .  
Hence  $S(1)$  is true.



(4) Prove that  $(1+x)^n \geq (1+nx)$ , for all natural number  $n$ , where  $x > -1$ .

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : (1+x)^n \geq (1+nx)$$

where  $x > -1$  For  $n = 1$ ,  $\text{RHS} = (1+x)^1 = 1+x$  and  $\text{LHS} = (1+nx) = (1+x)$ .  
Hence  $S(1)$  is true.

Assume  $S(k)$  is true

$$(1+x)^k \geq (1+kx)$$

where  $x > -1$

(4) Prove that  $(1+x)^n \geq (1+nx)$ , for all natural number  $n$ , where  $x > -1$ .

**Solution :** Let  $S(n)$  be the given statement

$$S(n) : (1+x)^n \geq (1+nx)$$

where  $x > -1$  For  $n = 1$ ,  $\text{RHS} = (1+x)^1 = 1+x$  and  $\text{LHS} = (1+nx) = (1+x)$ .  
Hence  $S(1)$  is true.

Assume  $S(k)$  is true

$$(1+x)^k \geq (1+kx)$$

where  $x > -1$  Now we show that  $S(k+1)$  is true  $(1+x)^{k+1} \geq (1+(k+1)x)$  where  $x > -1$ .

Now

$$\begin{aligned}(1+x)^{k+1} &= (1+x)^k(1+x) \\ &\geq (1+kx)(1+x) \\ &= 1+kx+x+kx^2 \\ &\geq 1+kx+x \\ &= 1+(k+1)x\end{aligned}$$

Hence  $S(k+1)$  is true.

So  $S(k+1)$  is true whenever  $S(k)$  is true. By Principal of Mathematical Induction  $S(n)$  is true for all natural number  $n$ .