



Unit 0. Course Overview, Homework

Course > 0, Project 0 (1 week)

> Homework 0 >

8. (Optional Ungraded Warmup) 1D
Optimization via Calculus

Audit Access Expires May 11, 2020

You lose all access to this course, including your progress, on May 11, 2020.

Upgrade by Mar 25, 2020 to get unlimited access to the course as long as it exists on the site. **Upgrade now**

8. (Optional Ungraded Warmup) 1D Optimization via Calculus

(Optional) Review: 1D Optimization via Calculus

0 points possible (ungraded)

(For this problem, you are welcome to use any computational tools that would be helpful.)

Let $f(x) = \frac{1}{3}x^3 - x^2 - 3x + 10$ defined on the interval $[-4, 4]$.

Let x_1 and x_2 be the critical points of f , and let's impose that $x_1 < x_2$. Fill in the next two boxes with the values of x_1 and x_2 , respectively: (Recall that the **critical points** of f are those $x \in \mathbb{R}$ such that $f'(x) = 0$.)

 $x_1 =$

✓ Answer: -1

 $x_2 =$

✓ Answer: 3

Fill in the next two boxes with the values of $f''(x_1)$ and $f''(x_2)$, respectively:

 $f''(x_1) =$

✓ Answer: -4

 $f''(x_2) =$

✓ Answer: 4

Solution:

Observe that

$$f'(x) = x^2 - 2x - 3 = (x - 3)(x + 1).$$

Hence the **critical points** are $x_1 = -1$ and $x_2 = 3$. The **second derivative** is

$$f''(x) = 2x - 2$$

so that

$$f''(x_1) = -4, \quad f''(x_2) = 4.$$

You have used 1 of 3 attempts

i Answers are displayed within the problem

(Optional) Review: 1D Optimization via Calculus (Continued)

0 points possible (ungraded)

(For this problem, you are welcome to use any computational tools that would be helpful.)

Recall that x_1 and x_2 are the critical points of the function

$$f(x) = \frac{1}{3}x^3 - x^2 - 3x + 10.$$

According to the second derivative test, x_1 is a ...

☒ Local Maximum☐ Local Minimum☐ None of the above

and x_2 is a

☐ Local Maximum☒ Local Minimum☐ None of the above



At what value of x is the (global) minimum value of $f(x)$ attained on the interval $[-4, 4]$?

✓ Answer: -4

At what value of x the (global) maximum value of $f(x)$ attained on the interval $[-4, 4]$?

✓ Answer: -1

Solution:

The previous problem implies that f is concave at x_1 and convex at x_2 , so x_1 is a **local maximum** and x_2 is a **local minimum**. To figure out the *global* extrema, we need to test the critical points as well as the endpoints: -4 and 4 . We compute that

$$f(x_1) = \frac{35}{3} \approx 11.6666, \quad f(x_2) = 1$$


$$f(-4) = -\frac{46}{3} \approx -15.33333, \quad f(4) = 10/3 \approx 3.3333$$

Hence the **maximum value** of f on $[-4, 4]$ is $\frac{35}{3} \approx 11.6666$ and the **minimum value** is $-\frac{46}{3} \approx -15.33333$.

Remark: It is very important to remember to test the endpoints when doing optimization.

Submit

You have used 1 of 2 attempts

 Answers are displayed within the problem

(Optional)Strict Concavity

0 points possible (ungraded)

Which of the following functions are strictly concave? (Choose all that apply.) (Recall that a twice-differentiable function $f : I \rightarrow \mathbb{R}$, where I is a subset of \mathbb{R} , is **strictly concave** if $f''(x) < 0$ for all $x \in I$.)

☐ $f_1(x) = x$ on \mathbb{R}
☒ $f_2(x) = -e^{-x}$ on \mathbb{R}
☒ $f_3(x) = x^{0.99}$ on the interval $(0, \infty)$
☐ $f_4(x) = x^2$ on \mathbb{R}
**Solution:**

- $f_1(x) = x$ is **not** strictly concave because $f_1''(x) = 0$.
- $f_2(x) = -e^{-x}$ is strictly concave because $f_2''(x) = -e^{-x} < 0$ for all $x \in \mathbb{R}$.
- $f_3(x) = x^{0.99}$ is strictly concave because $f_3''(x) = (0.99)(-0.01)x^{-1.01} < 0$ for all $x \in (0, \infty)$.
- $f_4(x) = x^2$ is **not** strictly concave because $f_4''(x) = 2 > 0$. In fact, this function is strictly *convex*.

Submit

You have used 1 of 2 attempts

i Answers are displayed within the problem

Discussion



Add a Post

Hide Discussion

Topic: Unit 0. Course Overview, Homework 0, Project 0 (1 week):Homework 0 / 8. (Optional Ungraded Warmup) 1D Optimization via Calculus

Show all posts

by recent activity

 Tool for Strict Concavity	8
 Buggy Grader algo? On strict concavity, the first has zero second derivative, the third and fourth have a positive c...	2

Learn About Verified Certificates

© All Rights Reserved