



[Unit 4 Unsupervised Learning \(2](#)

[Course](#) > [weeks](#))

> [Lecture 15. Generative Models](#) >

9. Gaussian Generative models

Audit Access Expires May 11, 2020

You lose all access to this course, including your progress, on May 11, 2020.

9. Gaussian Generative models

Gaussian Generative Models



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Multivariate Gaussian Random Vector

A random vector $\mathbf{X} = (X^{(1)}, \dots, X^{(d)})^T$ is a **Gaussian vector**, or **multivariate Gaussian or normal variable**, if any linear combination of its components is a (univariate) Gaussian variable or a constant (a "Gaussian" variable with zero variance), i.e., if $\alpha^T \mathbf{X}$ is (univariate) Gaussian or constant for any constant non-zero vector $\alpha \in \mathbb{R}^d$.

The distribution of \mathbf{X} , the **d -dimensional Gaussian or normal distribution**, is completely specified by the vector mean

$\mu = \mathbf{E}[\mathbf{X}] = (\mathbf{E}[X^{(1)}], \dots, \mathbf{E}[X^{(d)}])^T$ and the $d \times d$ covariance matrix Σ . If Σ is invertible, then the pdf of \mathbf{X} is

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{\sqrt{(2\pi)^d \det(\Sigma)}} e^{-\frac{1}{2}(\mathbf{x}-\mu)^T \Sigma^{-1}(\mathbf{x}-\mu)}, \quad \mathbf{x} \in \mathbb{R}^d$$

where $\det(\Sigma)$ is the determinant of the Σ , which is positive when Σ is invertible.

If $\mu = \mathbf{0}$ and Σ is the identity matrix, then \mathbf{X} is called a **standard normal random vector**.

Note that when the covariant matrix Σ is diagonal, the pdf factors into pdfs of univariate Gaussians, and hence the components are independent.

Gaussian Distribution

1/1 point (graded)

Recall that the likelihood of x being generated from a multi-dimensional Gaussian with mean μ and all the components being uncorrelated and having the same standard deviation σ is:

$$P(x|\mu, \sigma^2) = \frac{1}{(2\pi\sigma^2)^{d/2}} \exp\left(-\frac{1}{2\sigma^2} \|x - \mu\|^2\right)$$

Let $x = \begin{bmatrix} \frac{1}{\sqrt{\pi}} \\ 2 \end{bmatrix}$ be a vector in the two-dimensional space.

Let G be a two-dimensional Gaussian distribution with mean μ and standard deviation σ taking values as follows

$$\mu = \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \sigma = \sqrt{\frac{1}{2\pi}}$$

Calculate the likelihood $p(x|\mu, \sigma^2)$ of x being sampled from the Gaussian distribution G with mean μ and variance σ^2 taking values as given above.

Enter the value of $\log p(x|\mu, \sigma^2)$ below. (Note that we use \log for the natural logarithm, i.e. \log_e .)

✓ Answer: -1

Solution:

Note that the likelihood of vector x being sampled from a Gaussian distribution G with mean μ and variance σ^2 is given as follows

$$p(x|\mu, \sigma^2) = \frac{1}{2\pi\sigma^2} \exp\left(-\frac{1}{2\sigma^2} \|x - \mu\|^2\right)$$

Substituting the value of $\sigma = \sqrt{\frac{1}{2\pi}}$ from above, we have

$$p(x|\mu, \sigma^2) = \frac{1}{2\pi \frac{1}{2\pi}} \exp\left(-\frac{1}{2 \frac{1}{2\pi}} \|x - \mu\|^2\right)$$

$$p(x|\mu, \sigma^2) = \exp\left(-\pi\|x - \mu\|^2\right)$$

Substituting the value of $x = \begin{bmatrix} \frac{1}{\sqrt{\pi}} \\ 2 \end{bmatrix}$ and $\mu = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$, we have

$$p(x|\mu, \sigma^2) = \exp\left(-\pi\left[\left(\frac{1}{\sqrt{\pi}} - 0\right)^2 + (2 - 2)^2\right]\right)$$

$$p(x|\mu, \sigma^2) = \exp\left(-\pi\frac{1}{\pi}\right)$$

$$p(x|\mu, \sigma^2) = \exp(-1)$$

$$\log(p(x|\mu, \sigma^2)) = \log(\exp(-1)) = -1$$

You have used 2 of 2 attempts

i Answers are displayed within the problem

Discussion

Topic: Unit 4 Unsupervised Learning (2 weeks) :Lecture 15.
Generative Models / 9. Gaussian Generative models

Show all posts

by recent activity

- | | |
|---|----------|
| ? <u>Isn't $x - \mu ^2$ = to SQRT of differences squared?</u> | 3 |
| ? <u>$P(X \mu, \sigma)$</u>
<u>Is not $P(X \mu, \sigma)$ always equal to 0? NVM I can see below it refers to $p(X \mu, \sigma)$ - but...</u> | 1 |

© All Rights Reserved