



[Unit 1 Linear Classifiers and Course](#) > [Generalizations \(2 weeks\)](#)
3. Linear Classifiers Mathematically Revisited

[Lecture 2. Linear Classifier and Perceptron](#) >

Audit Access Expires May 11, 2020

You lose all access to this course, including your progress, on May 11, 2020. Upgrade by Mar 25, 2020 to get unlimited access to the course as long as it exists on the site. **[Upgrade now](#)**

3. Linear Classifiers Mathematically Revisited

Linear Classifiers Mathematically Revisited



Video[Download video file](#)**Transcripts**[Download SubRip \(.srt\) file](#)[Download Text \(.txt\) file](#)

Inner product and Orthogonal vectors

1/1 point (graded)

What is the inner product of $[0, 1, 1]$ and $[1, 1, 1]$?

✓ Answer: 2

Solution:

$$0 \cdot 1 + 1 \cdot 1 + 1 \cdot 1 = 2$$

You have used 1 of 3 attempts

i Answers are displayed within the problem

Linear Classifier Practice

1/1 point (graded)

We saw in the lecture above that for a linear classifier h , $h(x; \theta) = \text{sign}(\theta \cdot x)$, i.e. the sign of the dot product of θ and x . Now consider θ which is given by

$$\theta = (1, -1) \tag{3.1}$$

Which of the following points would be classified as positive by θ ? Please choose all correct answers.

☒ $(1, -1)$ ☒ $(1, 0)$ ☐ $(0, 1)$ ☐ $(0, 0)$ **Solution:**

X is positively classified by the classifier if and only if $x \cdot \theta > 0$. The dot product of $(1, -1)$ with θ is positive. Also, $(1, 0) \cdot \theta$ is positive. On the other hand, $(0, 1) \cdot \theta$ and $(0, 0) \cdot \theta$ are nonpositive. Thus the first and second points are positively classified by θ .

You have used 1 of 3 attempts

i Answers are displayed within the problem

Offset Added

1/1 point (graded)

Again, we have a linear classifier with θ given by

$$\theta = (1, -1) \tag{3.2}$$

and the offset, θ_0 given by $\theta_0 = -1$ Now which of the following points would be classified as positive by θ ? Please choose all correct answers.

☒ $(1, -1)$ ☐ $(1, 0)$ ☐ $(0, 1)$ ☐ $(0, 0)$ **Solution:**

X is positively classified by the classifier if and only if $x \cdot \theta + \theta_0 > 0$. The dot product of $(1, -1)$ with θ is 2, and adding -1 makes it still positive. However, $x \cdot \theta + \theta_0 \leq 0$ for other data points.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 1 Linear Classifiers and Generalizations (2 weeks):Lecture 2. Linear Classifier and Perceptron / 3. Linear Classifiers Mathematically Revisited

Add a Post

Show all posts

by recent activity

- ? why theta is orthogonal to the decision boundary? 4

Sorry if you are dissappointed, but I do not understand why theta is orthogonal to the decision...
- 💬 Suggestion 3

It would be more didactic if drawings of both cenarios (Linear Classifier Practice and Offset A...

✓	<u>Linear classifier</u>	5
	<u>X is positively classified, if $x \cdot \theta > 0$ and negatively classified, if $x \cdot \theta < 0$. What about, if $x \cdot t$...</u>	
?	<u>Degree of freedom</u>	2
	<u>What does the author mean by degree of freedom here?</u>	
💬	<u>Confusion at 12.40 regarding θ_0</u>	2
💬	<u>Can Anyone explains this with a numerical example?</u>	4
	<u>> There are multiple parameter vectors θ that defined exactly the same classifier. Each lin...</u>	
✓	<u>Exchange label values</u>	3
	<u>In the lecture, we assume that all points that (after the seperation) above the line have a labe...</u>	
?	<u>about the norm of the θ vector</u>	2

Learn About Verified Certificates

© All Rights Reserved