



BRAC University

Department of Mathematics and Natural Sciences

LECTURE ON

Real Analysis II (MAT321)

Theorems on Uniform Convergence

Continuity, Integration, Differentiation

FEBRUARY 11, 2026

CONDUCTED BY

Partho Sutra Dhor

Lecturer, BRAC University, Dhaka-1212

✉ partho.dhor@bracu.ac.bd | ✉ parthosutradhor@gmail.com

For updates subscribe on  [@ParthoSutraDhor](#)

Convergence of Sequence in Real Numbers

Convergence of a Sequence

A sequence $\{a_n\}$ is said to **converge** to a real number a if for every real number $\varepsilon > 0$, there exists a natural number $N \in \mathbb{N}$ such that for all $n \geq N$, the terms of the sequence satisfy the inequality

$$|a_n - a| < \varepsilon.$$

If such a number a exists, it is called the **limit** of the sequence, and we write

$$\lim a_n = a \quad \text{or} \quad a_n \rightarrow a \text{ as } n \rightarrow \infty.$$

Cauchy Criterion for Convergence

A sequence $\{a_n\}$ is said to satisfy the **Cauchy criterion** if for every real number $\varepsilon > 0$, there exists a natural number $N \in \mathbb{N}$ such that for all $m, n \geq N$, the terms of the sequence satisfy the inequality

$$|a_n - a_m| < \varepsilon.$$

Note: A sequence that satisfies the Cauchy criterion is called a **Cauchy sequence**. In the real numbers, every Cauchy sequence converges to a limit, and every convergent sequence is a Cauchy sequence.

Limit and Continuity of a function

Limit of a Function

Let $f : A \rightarrow \mathbb{R}$ be a function defined on a subset A of the real numbers, and let c be a limit point of A . We say that the **limit** of $f(x)$ as x approaches c is equal to L , and we write

$$\lim_{x \rightarrow c} f(x) = L,$$

if for every $\varepsilon > 0$, there exists a $\delta > 0$ such that for all $x \in A$ with $0 < |x - c| < \delta$, we have

$$|f(x) - L| < \varepsilon.$$

Continuity of a Function

A function $f : A \rightarrow \mathbb{R}$ is said to be **continuous** at a point $c \in A$ if

$$\lim_{x \rightarrow c} f(x) = f(c).$$

In other words, f is continuous at c if for every $\varepsilon > 0$, there exists a $\delta > 0$ such that for all $x \in A$ with $|x - c| < \delta$, we have

$$|f(x) - f(c)| < \varepsilon.$$

Poinwise Convergence and Uniform Convergence

Pointwise Convergence

A sequence of functions $\{f_n\}$ defined on a domain D is said to **converge pointwise** to a function f on D if for **every** $x \in D$ and for every $\varepsilon > 0$, there exists a natural number $N \in \mathbb{N}$ (which may depend on x and ε) such that for all $n \geq N$, the following inequality holds:

$$|f_n(x) - f(x)| < \varepsilon.$$

In this case, we write

$$\lim_{n \rightarrow \infty} f_n(x) = f(x) \quad \text{for all } x \in D.$$

Uniform Convergence

A sequence of functions $\{f_n\}$ defined on a domain D is said to **converge uniformly** to a function f on D if for every $\varepsilon > 0$, there exists a natural number $N \in \mathbb{N}$ such that for all $n \geq N$ and for all $x \in D$, the following inequality holds:

$$|f_n(x) - f(x)| < \varepsilon.$$

In this case, we write

$$f_n \rightarrow f \text{ uniformly on } D.$$

Note: In uniform convergence, the choice of N depends only on ε and not on x , which is a stronger condition than pointwise convergence.

Uniform Convergence implies Pointwise Convergence

Theorem

If a sequence of functions $\{f_n\}$ converges uniformly to a function f on a domain D , then it also converges pointwise to f on D . 

Uniform Limit of Continuous Functions is Continuous

Theorem

Let $\{f_n\}$ be a sequence of functions defined on a domain D , and suppose that each f_n is continuous at a point $c \in D$. If $\{f_n\}$ converges uniformly to a function f on D , then f is also continuous at the point c .

❓ Problem

Show that $f_n(x) = x^n$ is not uniformly convergent on $[0, 1]$.



Interchange of Limits under Uniform Convergence

💡 Interchange of Limits for Sequence

Let $\{f_n\}$ be a sequence of functions that converges uniformly to a function f on a set $D \subset \mathbb{R}$, and let x_0 be a limit point of D . Suppose that for each $n = 1, 2, 3, \dots$,

$$\lim_{x \rightarrow x_0} f_n(x) = a_n$$

exists. Then:

1. the sequence $\{a_n\}$ converges, and
- 2.

$$\lim_{x \rightarrow x_0} f(x) = \lim_{n \rightarrow \infty} a_n.$$

💡 Interchange of Limits for Series

Let $D \subset \mathbb{R}$ and let $\sum_{n=1}^{\infty} f_n$ be a series of functions defined on D whose partial sums

$$S_N(x) = \sum_{n=1}^N f_n(x)$$

converge uniformly on D to a function $f : D \rightarrow \mathbb{R}$. Let x_0 be a limit point of D . Assume that for each $n = 1, 2, 3, \dots$ the limit

$$\lim_{\substack{x \rightarrow x_0 \\ x \in D}} f_n(x) = a_n$$

exists. Then:

1. the numerical series $\sum_{n=1}^{\infty} a_n$ converges, and
- 2.

$$\lim_{\substack{x \rightarrow x_0 \\ x \in D}} f(x) = \sum_{n=1}^{\infty} a_n.$$

When Pointwise Convergence is Uniform Convergence

Dini's Theorem

Let K be compact, and $f_n : K \rightarrow \mathbb{R}$ continuous. Suppose $f_n \rightarrow f$ pointwise on K , f is continuous, and the convergence is monotone (either $f_n \uparrow f$ or $f_n \downarrow f$ pointwise). Then $f_n \rightarrow f$ uniformly on K .

Problem

Show that $f_n(x) = \tan^{-1}(nx)$ is uniformly convergent on $[a, b]$ for $a > 0$, but not uniformly convergent on $[0, b]$.

❓ Problem

Let $x \geq 0$ and define $f_1(x) = \sqrt{x}$, $f_{n+1}(x) = \sqrt{x + f_n(x)}$. Discuss uniform convergence on (i) $[a, b]$ with $0 < a < b$ and (ii) $[0, 1]$.

Uniform Convergence and Integration

💡 Uniform Convergence and Integration

Let $\{f_n\}$ be a sequence of functions that converges uniformly to f on $[a, b]$. If each function f_n is integrable on $[a, b]$, then f is integrable on $[a, b]$, and the sequence

$$\left\{ \int_a^x f_n(t) dt \right\}$$

converges uniformly to

$$\int_a^x f(t) dt$$

on $[a, b]$, that is,

$$\int_a^x f(t) dt = \lim_{n \rightarrow \infty} \int_a^x f_n(t) dt, \quad \forall x \in [a, b].$$

🔗 Uniform Convergence of a Series and Integration

If a series $\sum_{n=1}^{\infty} f_n$ converges uniformly to f on $[a, b]$, and each term $f_n(x)$ is integrable on $[a, b]$, then f is integrable on $[a, b]$, and the series

$$\sum_{n=1}^{\infty} \left(\int_a^x f_n(t) dt \right)$$

converges uniformly to

$$\int_a^x f(t) dt$$

on $[a, b]$, that is,

$$\int_a^x f(t) dt = \sum_{n=1}^{\infty} \left(\int_a^x f_n(t) dt \right), \quad \forall x \in [a, b].$$

Problem

Define on $[0, 1]$:

$$f_n(x) = \begin{cases} n(1 - nx), & 0 < x < \frac{1}{n}, \\ 0, & \text{otherwise.} \end{cases}$$

Compute

$$\int_0^1 f_n(x) dx$$

and compare it with the pointwise limit of $\{f_n\}$.

🔍 Problem

Let

$$f_n(x) = nxe^{-nx^2} \quad \text{on } [0, 1].$$

Compute

$$\int_0^1 f_n(x) dx$$

and compare it with the pointwise limit of $\{f_n\}$.

Uniform Convergence and Differentiation

💡 Uniform Convergence of Derivatives

Let $\{f_n\}$ be a sequence of functions defined on the closed interval $[a, b]$ such that $f_n \rightarrow f$ pointwise on $[a, b]$, and assume that each f_n is differentiable. If $\{f'_n\}$ converges uniformly on $[a, b]$ to a function G , then f is differentiable on $[a, b]$ and

$$f'(x) = G(x).$$

💡 Uniform Convergence of Derivatives with One-Point

Let $\{f_n\}$ be a sequence of differentiable functions on $[a, b]$ such that it converges at least at one point $x_0 \in [a, b]$. If the sequence of derivatives $\{f'_n\}$ converges uniformly to a function G on $[a, b]$, then the sequence $\{f_n\}$ converges uniformly on $[a, b]$ to a function f and

$$f'(x) = G(x).$$

💡 Term-by-Term Differentiation of a Series

Let $\sum_{n=1}^{\infty} f_n$ be a series of differentiable functions on $[a, b]$ such that it converges at least at one point $x_0 \in [a, b]$. If the series of derivatives

$$\sum_{n=1}^{\infty} f'_n$$

converges uniformly to a function G on $[a, b]$, then the given series

$$\sum_{n=1}^{\infty} f_n$$

converges uniformly on $[a, b]$ to a function f and

$$f'(x) = G(x).$$

🔍 Problem

Let

$$f_n(x) = \frac{\cos(nx)}{n} \quad (x \in \mathbb{R}).$$

1. Show that $f_n \rightarrow 0$ uniformly on $\mathbb{R}$.
2. Find where f'_n converges.
3. Construct a modified example where $f_n \rightarrow 0$ uniformly but $\{f'_n\}$ is unbounded.

② Uniform Convergence and Differentiation

State the theorem that relates uniform convergence and differentiation for a sequence of functions. Hence verify the result in detail for the sequence of functions $\{f_n\}$ where

$$f_n(x) = x - \frac{x^n}{n}, \quad x \in [0, 1].$$

Thank You!

We'd love your questions and feedback.


Partho Sutra Dhor

Lecturer, BRAC University, Dhaka-1212

✉ partho.dhor@bracu.ac.bd | ✉ parthosutradhor@gmail.com

 **@ParthoSutraDhor**

(Lectures, walkthroughs, and course updates)



Scan for the channel

References

- [1] Stephen Abbott, *Understanding Analysis*, 2nd Edition, Springer, 2015.
- [2] Walter Rudin, *Principles of Mathematical Analysis*, 3rd Edition, McGraw-Hill, 1976.