



BRAC University

Department of Mathematics and Natural Sciences

LECTURE ON

Real Analysis II (MAT321)

Continuity in Metric Space

Continuity, Uniform continuity, Homeomorphisms

MARCH 13, 2026

CONDUCTED BY

Partho Sutra Dhor

Lecturer, BRAC University, Dhaka-1212

✉ partho.dhor@bracu.ac.bd | ✉ parthosutradhor@gmail.com

For updates subscribe on  [@ParthoSutraDhor](#)

Continuity of Functions in Metric Spaces

Continuity at a Point in $\mathbb{R}$

Let $f : A \rightarrow \mathbb{R}$ be a function defined on a subset A of $\mathbb{R}$, and let $c \in A$. We say that f is **continuous at the point** a if for every $\varepsilon > 0$, there exists a $\delta > 0$ such that

$$\forall x \in A \text{ and } |x - a| < \delta \implies |f(x) - f(a)| < \varepsilon.$$

Continuity at a Point in Metric Space

Let (X, d_X) and (Y, d_Y) be metric spaces, and let $f : X \rightarrow Y$ be a function. We say that f is **continuous at the point** $a \in X$ if for every $\varepsilon > 0$, there exists a $\delta > 0$ such that

$$\forall x \in X \text{ and } d_X(x, a) < \delta \implies d_Y(f(x), f(a)) < \varepsilon.$$

Equivalent Definition using Open Spheres

Let (X, d_X) and (Y, d_Y) be metric spaces, and let $f : X \rightarrow Y$ be a function. We say that f is **continuous at the point** $a \in X$ if for every open sphere $S_\varepsilon(f(a))$ in Y , there exists an open sphere $S_\delta(a)$ in X such that

$$f(S_\delta(a)) \subseteq S_\varepsilon(f(a)).$$

Equivalent Definition using Open Set

Let (X, d_X) and (Y, d_Y) be metric spaces, and let $f : X \rightarrow Y$ be a function. We say that f is **continuous at the point** $a \in X$ if for every open set U in Y containing $f(a)$, there exists an open set V in X containing a such that

$$f(V) \subseteq U.$$

Continuous Functions

Continuity using Open Sets

Let (X, d_X) and (Y, d_Y) be metric spaces, and let $f : X \rightarrow Y$ be a function. We say that f is continuous at all points of X i.e. **continuous** if for any open set U in Y , the preimage $f^{-1}(U)$ is an open set in X .

Continuity using Closed Sets

Let (X, d_X) and (Y, d_Y) be metric spaces, and let $f : X \rightarrow Y$ be a function. We say that f is continuous at all points of X i.e. **continuous** if for any closed set C in Y , the preimage $f^{-1}(C)$ is a closed set in X .

Examples of Continuous Functions

💡 Continuity in Discrete Metric Space

Let (X, d_X) be a discrete metric space, and let (Y, d_Y) be any metric space. Then every function $f : X \rightarrow Y$ is continuous at every point of X .

💡 Continuity of a constant function

Let (X, d_X) and (Y, d_Y) be metric spaces, and let $f : X \rightarrow Y$ be a constant function defined by $f(x) = c$ for all $x \in X$, where $c \in Y$ is a fixed point. Then f is continuous at every point of X .

Continuity using Sequences

💡 Continuity at a Point using Sequences

Let (X, d_X) and (Y, d_Y) be metric spaces, and let $f : X \rightarrow Y$ be a function. We say that f is **continuous at the point** $a \in X$ if and only if for every sequence (x_n) in X that converges to a , the sequence $(f(x_n))$ in Y converges to $f(a)$.

Uniform Continuity

Uniform Continuity

Let (X, d_X) and (Y, d_Y) be metric spaces, and let $f : X \rightarrow Y$ be a function. We say that f is **uniformly continuous** if for every $\varepsilon > 0$, there exists a $\delta > 0$, that depends only on ε , such that

$$\forall x, y \in X \text{ and } d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \varepsilon.$$

Cauchy sequence with uniform continuity

Let (X, d_X) and (Y, d_Y) be metric spaces, and let $f : X \rightarrow Y$ be a uniformly continuous function. Then for every Cauchy sequence (x_n) in X , the sequence $(f(x_n))$ in Y is also a Cauchy sequence.

② Not uniform continuity with Cauchy sequence

Give an example of a function $f : X \rightarrow Y$ that is continuous but not uniformly continuous, and a Cauchy sequence (x_n) in X such that the sequence $(f(x_n))$ is not a Cauchy sequence in Y .

Homeomorphisms

Homeomorphism

Let (X, d_X) and (Y, d_Y) be metric spaces. A function $f : X \rightarrow Y$ is called a **homeomorphism** if it is a bijection and both f and its inverse $f^{-1} : Y \rightarrow X$ are continuous.

Homeomorphic Spaces

Let (X, d_X) and (Y, d_Y) be metric spaces. We say that X and Y are **homeomorphic** if there exists a homeomorphism $f : X \rightarrow Y$. In this case, we write $X \cong Y$.

Thank You!

We'd love your questions and feedback.


Partho Sutra Dhor

Lecturer, BRAC University, Dhaka-1212

✉ partho.dhor@bracu.ac.bd | ✉ parthosutradhor@gmail.com

 **@ParthoSutraDhor**

(Lectures, walkthroughs, and course updates)



Scan for the channel

References

- [1] Erwin Kreyszig, *Introductory Functional Analysis with Applications*, John Wiley & Sons, 1978.
- [2] Walter Rudin, *Principles of Mathematical Analysis*, 3rd Edition, McGraw-Hill, 1976.