



BRAC University

Department of Mathematics and Natural Sciences

LECTURE ON

Real Analysis (MAT221)

Riemann Integrability

Cauchy Criterion, Integration of Continuous Functions

AUGUST 17, 2026

CONDUCTED BY

Partho Sutra Dhor

Lecturer, BRAC University, Dhaka-1212

✉ partho.dhor@bracu.ac.bd | ✉ parthosutradhor@gmail.com

For updates subscribe on  [@ParthoSutraDhor](#)

Lower Sums

Lower Sum

Let f be bounded on $[a, b]$, and let $P = \{x_0, x_1, \dots, x_n\}$ be a partition of $[a, b]$.

The **lower sum** of f with respect to P is

$$L(f, P) = \sum_{i=1}^n m_i(x_i - x_{i-1}).$$

Upper Sums

Upper Sum

Let f be bounded on $[a, b]$, and let

$$P = \{x_0, x_1, \dots, x_n\}$$

be a partition of $[a, b]$.

The **upper sum** of f with respect to P is

$$U(f, P) = \sum_{i=1}^n M_i(x_i - x_{i-1}).$$

Lower Integral

Lower Integral

Let f be bounded on $[a, b]$.

The **lower integral** of f over $[a, b]$ is

$$\int_a^b f = \inf \{L(f, P) : P \text{ is a partition of } [a, b]\}.$$

Upper Integral

Upper Integral

Let f be bounded on $[a, b]$.

The **upper integral** of f over $[a, b]$ is

$$\overline{\int_a^b} f = \sup \{U(f, P) : P \text{ is a partition of } [a, b]\} .$$

Riemann Integrability

Riemann Integrable Function

A bounded function

$$f : [a, b] \rightarrow \mathbb{R}$$

is called **Riemann integrable** on $[a, b]$ if

$$\underline{\int_a^b} f = \overline{\int_a^b} f.$$

The common value is called the **Riemann integral** of f and is denoted by

$$\int_a^b f(x) dx.$$

🔍 Problem

Show directly that the constant function

$$f(x) = k$$

is integrable over any closed interval $[a, b]$. What is $\int_a^b f$?

Cauchy Criterion for Riemann Integrability

💡 Cauchy Criterion

A bounded function f on $[a, b]$ is Riemann integrable if and only if for every $\epsilon > 0$, there exists a partition P of $[a, b]$ such that

$$U(f, P) - L(f, P) < \epsilon.$$

❓ Problem

Consider $f(x) = 2x + 1$ over the interval $[1, 3]$. Let P be the partition consisting of the points $\{1, \frac{3}{2}, 2, 3\}$.

- (a) Compute $L(f, P)$, $U(f, P)$, and $U(f, P) - L(f, P)$.
- (b) What happens to the value of $U(f, P) - L(f, P)$ when we add the point $\frac{5}{2}$ to the partition?
- (c) Find a partition P' of $[1, 3]$ for which

$$U(f, P') - L(f, P') < 2.$$

Sequential Criterion

Prove that a bounded function f is integrable on $[a, b]$ if and only if there exists a sequence of partitions $(P_n)_{n=1}^{\infty}$ satisfying

$$\lim_{n \rightarrow \infty} [U(f, P_n) - L(f, P_n)] = 0.$$

❓ Problem

Using the previous theorem show that the function

$$f(x) = 2x^2 + 3$$

is integrable on $[0, 1]$.



Problem

Show that the Dirichlet function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = \begin{cases} 1, & \text{if } x \in \mathbb{Q}, \\ 0, & \text{if } x \notin \mathbb{Q}, \end{cases}$$

is not Riemann integrable on $[0, 1]$.



Continuity and Uniform Continuity

Continuity

A function f is continuous on an interval I if for every $c \in I$ and for every $\epsilon > 0$, there exists a $\delta > 0$ such that for all $x \in I$, if $|x - c| < \delta$, then $|f(x) - f(c)| < \epsilon$. Here δ may depend on both ϵ and c .

Uniform Continuity

A function f is uniformly continuous on an interval I if for every $\epsilon > 0$, there exists a $\delta > 0$ such that for all x, y in the domain of f , if $|x - y| < \delta$, then $|f(x) - f(y)| < \epsilon$.

Continuous Functions are Riemann Integrable

💡 **Continuity $\implies$ Riemann Integrable**

If a function f is continuous on $[a, b]$, then f is Riemann integrable on $[a, b]$.

Thank You!

We'd love your questions and feedback.


Partho Sutra Dhor

Lecturer, BRAC University, Dhaka-1212

✉ partho.dhor@bracu.ac.bd | ✉ parthosutradhor@gmail.com

 **@ParthoSutraDhor**

(Lectures, walkthroughs, and course updates)



Scan for the channel

References

- [1] Stephen Abbott, *Understanding Analysis*, 2nd Edition, Springer, 2015.
- [2] Terence Tao, *Analysis I*, 3rd Edition, Texts and Readings in Mathematics, Hindustan Book Agency, 2016.