

ACETAMINOPHEN 325 MG- acetaminophen tablet
Reliable 1 Laboratories LLC

Acetaminophen USP 325 mg Tablets

Drug Facts

Active ingredient (in each tablet)

Acetaminophen USP, 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual or menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg in 24 hours, which is the maximum daily amount
- child takes more than 5 tablets in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	take 2 tablets every 4-6 hours while symptoms last, not more than 12 tablets in 24 hours
children 6 to 11 years	take 1 tablet every 4-6 hours while symptoms last, not more than 5 tablets in 24 hours
children under 6 years	do not use

Other information

- store at 15° to 30°C (59° to 86°F)

Inactive ingredients

povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid

Questions or comments?

call **516-341-0666**, 8:30 am - 4:30 pm ET, Monday - Friday

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*Reliable 1 Laboratories LLC is not affiliated with the owner of the trademark Tylenol®.

Distributed by: **Reliable 1 Laboratories LLC, Valley Stream, NY 11580**

www.reliable1labs.com

NDC 69618-010-01

Regular Strength


Acetaminophen USP 325 mg

PAIN RELIEVER

FEVER REDUCER

100 TABLETS

*Compare to Active Ingredient in Regular Strength TYLENOL®



Drug Facts		Purpose
Active ingredient (in each tablet)		
Acetaminophen USP, 325 mg Pain reliever/Fever reducer		
Uses		
■ temporarily relieves minor aches and pains due to: ■ muscular aches ■ headache ■ minor pain of arthritis ■ backache ■ toothache ■ the common cold ■ menstrual cramps		
Warnings		
Liver warning: This product contains acetaminophen. Severe liver damage may occur if adult takes more than 4,000 mg in 24 hours, which is the maximum daily amount. Do not drink alcohol or take other products containing acetaminophen while taking this product.		
Child warning: Do not give children under 16 years of age aspirin or products containing aspirin without a doctor's advice. Children and teenagers should not take aspirin for colds or chickenpox or flu unless directed by a doctor.		
Allergy alert: Acetaminophen may cause severe skin reactions. Some may include: ■ rash ■ hives ■ skin redness or sores ■ swelling of the eyes, face, mouth, tongue, or throat. If a skin reaction occurs, stop use and seek medical help right away.		
Directions ■ do not take more than directed (see overdose warning)		
adults & children 12 years and over take 2 tablets every 4 - 6 hours while symptoms last; not more than 12 tablets in 24 hours		
children 6 to 11 years take 1 tablet every 4 - 6 hours while symptoms last; not more than 5 tablets in 24 hours		
children under 6 years do not use		
Other information		
■ store at 15° to 30°C (59° to 86°F).		
Inactive ingredients povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid		
Questions or comments? call 516-341-0666, 8:30 am - 4:30 pm ET, Monday - Friday		

Drug Facts continued on back of label

TAMPER EVIDENCE: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*Reliable 1 Laboratories LLC is not affiliated with the owner of the trademark TYLENOL®.

Distributed by: Reliable 1 Laboratories LLC
Valley Stream, NY 11580
www.reliable1labs.com

3 69618 01001 7

Lot No.:
PEEL HERE
FOR MORE
DRUG FACTS
Exp. Date:

Drug Facts (continued)

Do not use with any other drug containing

acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if the user has liver disease or is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
 - pain gets worse or lasts more than 5 days in children
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition. These could be signs of a serious condition.
- If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Oversdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning)

take 2 tablets every 4 - 6 hours

while symptoms last; not more

than 12 tablets in 24 hours

take 1 tablet every 4 - 6 hours

while symptoms last; not more

than 5 tablets in 24 hours

children under 6 years do not use

ACETAMINOPHEN 325 MG

acetaminophen tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:69618-010

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	AP;012
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69618-010-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2015	
2	NDC:69618-010-10	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2015	



ACETAMINOPHEN

325mg Tablet

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/01/2015	

Labeler - Reliable 1 Laboratories LLC (079718111)

Registrant - Reliable 1 Laboratories LLC (079718111)