

# CSCI E-82a

## Probabilistic Programming and AI

### Lecture 2

# Markov Graphical Models

Steve Elston



HARVARD  
Extension School


Copyright 2019, Stephen F Elston. All rights reserved.

# Outline

- Why Markov Graphical Models?
- Properties of Markov Graphical Models
- Cliques of Graphical Models
- Potentials for Markov Graphical Models
- Independencies and the Hammerly Clifford Theorem
- Potentials and the Hammerly Clifford Theorem
- Independencies and Separation in MRFs
- Markov Blanket
- Pairwise Markov Property
- Transforming DAGs to MRFs and Moralization
- Independencies and Separation in MRFs
- DAGs vs MRFs
- Summary

# Why Markov Graphical Models?

The previous lesson addressed **directed acyclic graphical models (DAGs)**



Attribution, Wojtek Przytula, et. al. 2002

# Why Markov Graphical Models?


The previous lesson addressed **directed acyclic graphical models (DAGs)**

- DAGs **cannot represent certain independency structures**:
  - Non-directional dependency on neighbors
    - Spatial relationships
    - Social networks
    - Image data
    - Molecular structure
    - Many more.....
  - Cyclical structures
  - Etc...
- How can these independencies be represented?
- **Markov graphical model** or **Markov random field (MRF)** model


# Why Markov Graphical Models?

Markov are related to DAGs

- DAGs have directed edges
- Markov graphical models have undirected edges
- DAGs can be transformed into equivalent Markov graph



**DAG**



**Undirected Markov graph**

# Why Markov Graphical Models?

- Markov random field models arise in statistical physics
  - Solid state physics
  - e.g. Ising model of magnetism
- Markov random field models can represent complex **independency structure**
- But, the we pay a price in **computational complexity**
- If a DAG represents the independency structure, use it!
- Otherwise, we use a MRF, if computationally feasible

# Why Markov Graphical Models?

Solve protein folding problem

<http://www.yaroslavvb.com/papers/yanover-linear.pdf>





Figure 2: **(a)** Cow actin binding protein (PDB code 1pne). **(b)** A closer view of its 6 C-terminal residues. Given the protein backbone (black) and the amino acid sequence, side-chain prediction is the problem of predicting the native side-chain conformation (gray). **(c)** Problem representation as a graphical model for those C-terminal residues shown in (b) (nodes located at  $C^\alpha$  atom positions, edges drawn in black).

Attribution; Yanover, Meltzer, Weiss 2006

# Focus on representation with graphical models



Schematic of intelligent agent using directed graphical model



# Properties of Markov Graphical Models

- Markov random field models have **undirected edges**
- Model distributions in MRFs using **potentials**
  - Recall that DAGs model distributions using **CPDs**
- Potentials are not distributions and must be normalized by a **partition function**
- There is a **potential** for **each clique of the graph**
- Potentials define the **strength of the interaction** between the nodes in a clique
  - For example, people who interact directly in a social network are more likely to influence each other (like a social clique)


# Cliques of Graphical Models

**Definition:** A **clique** is a subset of vertices of an undirected graph such that **every two distinct vertices in the clique are adjacent**

- The **subgraphs** of a clique **must be complete**
- **Independency structure** is determined by the cliques of the MRF
- A node can be in multiple cliques
- A clique can be as small as one node

# Cliques of Graphical Models


**Definition:** A **maximal clique** is a clique which cannot be enlarged without violating the clique property




- Maximal Cliques required for inference algorithms
- Example: consider the undirected graph with cliques, (A,B), (A,C), (B,D), (C,D)
- Enlarging any of these cliques **violates the clique property**, since the vertices would not be adjacent
- Therefore, the cliques (A,B), (A,C), (B,D), (C,D) and **maximal cliques**

# Cliques of Graphical Models - Example


**Example:** undirected graph with 4 cliques defined




**Clique 1,  
(A,B)**



**Clique 2,  
(A,C)**




**Clique 3,  
(B,D)**



**Clique 4,  
(C,D)**

# Potentials for Markov Graphical Models

Distribution for Markov random fields is modeled by **potentials**



- Each clique has a potential
- The **product of the potentials is also a potential**:  
 $\tilde{p}(A, B, C, D) = \phi(A, B)\phi(A, C)\phi(B, D)\phi(C, D)$
- The product of the potentials can be **transformed to a distribution by a normalization**:

$$p(A, B, C, D) = \frac{1}{Z} \tilde{p}(A, B, C, D)$$

- Where the normalization is the **partition function**:

$$Z = \sum_{A, B, C, D} \tilde{p}(A, B, C, D)$$

# Potentials for Markov Graphical Models

How to formulate a **distribution given the potentials** for complex graphs?

- The general formulation for a multivariate distribution is

$$p(x_1, \dots, x_n) = \frac{1}{Z} \prod_{c \in C} \phi_c(x_c)$$

- Where,  $c$  is a clique in the set of cliques
- The partition function is given by:

$$Z = \sum_{x_1, \dots, x_n} \prod_{c \in C} \phi_c(x_c)$$

# Potentials for Markov Graphical Models

Computing the partition function presents a significant problem

$$Z = \sum_{x_1, \dots, x_n} \prod_{c \in C} \phi_c(x_c)$$

- Partition function has high computational complexity
  - Compute a product for each data sample
  - Sum the products over the data samples
- Computation complexity makes exact solution of many MRF problems impractically difficult

# Potentials for Markov Graphical Models

How does all this relate to Bayesian networks?

- Recall, we can express the distribution for a Bayesian network using global semantics:

$$P(X) = \prod_{i=1:d} P(X_i | \{parents(X_i)\})$$

- The potentials are the conditional probability distributions (CPDs)
- But, what happened to the partition function?
- For Bayesian networks  $Z = 1.0$



# Independencies and the Hammerly Clifford Theorem

How can we **model independencies** in undirected graphical models?

- The **Hammerly-Clifford theorem** provides a tool to map the **conditional independence properties** of an undirected graph  $G$
- Further, the Hammerly-Clifford theorem gives a practical way to formulate potentials

# Independencies and the Hammerly Clifford Theorem

**Hammerly-Clifford theorem:** Let  $p(x)$  be a strictly positive distribution and let  $G$  be an undirected graph, the **conditional independence properties** of  $p(x)$  are satisfied if and only if the distribution can be represented as a product of factors, one factor representing each maximal clique,  $c$ , of  $G$ :

$$p(x \mid \Theta) = \frac{1}{Z(\theta)} \prod_{c \in G} \psi_c(x_c \mid \theta_c)$$

Where,

$$Z(\theta) = \sum_x \prod_{c \in G} \psi_c(x_c \mid \theta_c)$$

And,  $Z(\theta)$  is the partition function that ensures  $p(x \mid \Theta)$  is in the range  $\{0, 1\}$

# Potentials and the Hammerly Clifford Theorem

The Hammerly Clifford Theorem provides a practical way to formulate potentials

- Use a **Gibbs Distribution**:

$$p(x \mid \theta) = \frac{1}{Z(\theta)} \exp\left(-\sum_c E(x_c \mid \theta_c)\right)$$

- Where,  $E(x_c) = \text{Energy of clique } c$
- The potential for a clique is then:

$$\phi(x_c \mid \theta_c) = \exp\left(-E(x_c \mid \theta_c)\right)$$

- Given the minus sign, **the lower the energy of the state of a clique, c, the higher the probability**

# Independencies and Separation in MRFs

How can we **model independencies or separation** in undirected graphical models?

- **Definition:** For a graph  $G$ , **disjoint** subsets  $A$  and  $B$  are separated by subset  $S$  if every path from  $A$  to  $B$  passes through  $S$  then  $S$  **separates**  $A$  and  $B$ . Or, if  $S = \emptyset$ , then no path exists from  $A$  to  $B$ , and  $A$  and  $B$  are **separated**
- **Definition:** For a graph  $G$  with disjoint sets  $A$ ,  $B$  and  $S$ , where  $S$  separates  $A$  and  $B$ , then  $A \perp B \mid S$ , known as the **global Markov property**

# Independencies and Separation in MRFs

How can we **model independencies or separation** in undirected graphical models?

- **Definition:** Given subsets  $X$ ,  $Y$  and  $Z$ ,  $X$  and  $Y$  are conditionally independent or **D-separated** conditioned on the subset  $Z$  if they are separated on the moralized graph
- **Definition:** A graph  $G$  is a **dependency map** or **D-map** of a distribution  $P$  if the graph contains every conditional independence in  $P$ . We can represent this relationship as:

$$(X \perp Y \mid Z_G) \Leftarrow (X \perp Y \mid Z_P)$$


# Independencies and Separation in MRFs

There are two significant properties of independence maps in undirected graphs that are **soundness** and **completeness**

- **Theorem:** For any graph  $G$  that factorizes a distribution  $P$  then  $I(G) \subseteq I(P)$ . This relationship is known as the **soundness** property
- **Claim:** For any graph  $G$ , with subsets  $X$ ,  $Y$  and  $Z$ , that factorizes a distribution  $P$ , if  $(X \perp Y \mid Z) \subseteq I(P)$  then  $d\text{-sep}_G(X; Y \mid Z)$ . This relationship is known as the **completeness property**

# Independencies and Separation in MRFs - Example

Example of **modeling independencies or separation** in undirected graphical models



There are **5 maximal cliques**

$\{A, B, C\}$

$\{C, R\}$

$\{Q, R, S\}$

$\{S, X\}$


$\{X, Y, Z\}$

Cliques  $\{A, B, C\}$  and  $\{X, Y, Z\}$  are separated by clique  $\{Q, R, S\}$

Or,  $\{A, B, C\} \perp \{X, Y, Z\} \mid \{Q, R, S\}$

# Markov Blanket

For a MRF, any node is conditionally independent of all others given its **Markov Blanket**



**Definition:** A subset  $U$  is a **Markov blanket** of  $X$  in the set of nodes  $\chi$  of the graph  $G$  if  $X \notin U$  and if  $U$  is a minimal set of nodes such that:

$$(X \perp \chi - \{X\} - U \mid U) \in I(P)$$

- Where  $\chi - \{X\} - U$  is the set of nodes not in  $X$  or  $U$
- This definition is a result of the D-separation property for MRFs



## Pairwise Markov Property

The **pairwise Markov property** connects the local and global Markov properties of MRFs

**Definition:** Two nodes are conditionally independent given the other nodes in the graph if there is no direct edge between them. This property is the **pairwise Markov property**

- The pairwise Markov property relates to the **global Markov property** and the **local Markov property** in a somewhat circular fashion:

$$Global \rightarrow Local \rightarrow Pairwise \rightarrow Global$$


# Transforming DAGs to MRFs and Moralization

How is the structure of a DAG related to the structure of MRF

- The process of **moralization** is the key step in transforming a DAG to an MRF
- **Definition:** An **immorality** in a directed graph  $G$  occurs where either;  
a) there is a directed edge between  $X$  and  $Y$ , or b)  $X$  and  $Y$  are both parents of the same node  $Z$
- **Definition:** A **moral graph**,  $M(G)$ , of a BN structure,  $G$  is the **undirected graph** over  $X$  that contains an undirected edge between  $X$  and  $Y$  if; a) there is a directed edge between  $X$  and  $Y$ , or b)  $X$  and  $Y$  are both parents of the same node  $Z$

# Transforming DAGs to MRFs and Moralization - Example

## Example of an immorality



- Consider a DAG with a **v-structure or collider**
- With Z not observed the independency can be expressed:


$$P(X, U \mid Z) = P(X \mid Z) P(U \mid Z)$$

- The path from X to U is blocked by Z
- Therefore, this relationship is an **immorality** since X and U are parents of Z
- We **moralize** the graph by **marrying** X and U with an undirected edge


# Transforming DAGs to MRFs and Moralization - Example

## Example of transforming a DAG to a MRF


- Start with the original DAG
- Not observing Z blocks a path to W and U and X are independent
- Therefore, the **ancestral graph** does not contain W
- Moralized undirected graph



Original DAG



Ancestral graph



Moralized undirected graph

# Transforming DAGs to MRFs and Moralization


How is the structure of a DAG related to the structure of MRF

- We can relate the I-map between a DAG and a MRF through a corollary
- **Corollary:** Given a distribution  $P_B$  such that  $B$  is a parameterization on a graph  $G$ , then  $M(G)$  is an I-map for  $P_B$
- However, all of this **does not mean** that the independency structure of the DAG and resulting MRF will be the same


# Independencies and Separation in MRFs - Example

Example of **modeling independencies or separation** in undirected graphical models

- Start with DAG
- Transform to undirected moralized graph



**DAG**




**Moralized MN**

# Independencies and Separation in MRFs - Example

Example of **modeling independencies or separation** in undirected graphical models

- Define maximal cliques on moralized undirected graph
- Each clique has a potential



# Independencies and Separation in MRFs - Example

Example of **modeling independencies or separation** in undirected graphical models

- Now, factorize the unconditional distribution into potentials

$$\begin{aligned} P(I, D, G, S, L) &= \frac{1}{Z} \phi(D, I, G) \phi(G, L) \phi(I, S) \\ &= \frac{1}{Z} \exp\{-\mathbb{E}(D, I, G) - \mathbb{E}(G, L) - \mathbb{E}(I, S)\} \end{aligned}$$

And the partition function is:

$$Z = \sum_{I, D, G, S, L} \phi(D, I, G) \phi(G, L) \phi(I, S)$$


The distribution is modeled at the product of potentials on the undirected graph



## DAGs vs MRFs - Example



Example of **modeling independencies or separation** in undirected graphical models

- Wait! Have we **lost the independence map** of the DAG?
- Yes, **the potential of the maximal cliques have a different map!**



## DAGs vs MRFs


Directed acyclic graphs and Markov networks **cannot always represent the same independency structures**



- Why is this the case?
- Consider a DAG with a **v-structure or collider**
- **X and Y are unconditionally independent:**  $X \perp Y$ .
- Now consider the moralized Markov network
- There is **no independency!**

## DAGs vs MRFs

Directed acyclic graphs and Markov networks **cannot always represent the same independency structures**



- Consider the Markov network with 4 maximal cliques, and potentials:

$$\phi(W, X) \phi(X, Y) \phi(Y, Z) \phi(Z, W)$$

Independencies are:

$$\phi(X, Y) \perp \phi(Z, W) \mid \{\phi(W, X), \phi(Y, Z)\}$$

$$\phi(W, X) \perp \phi(Y, Z) \mid \{\phi(X, Y), \phi(Z, W)\}$$

- Multiple DAGs are possible on this skeleton, eg.
- But, **no DAG can represent the independencies!**

## DAGs vs MRFs

Directed acyclic graphs and Markov networks **cannot always represent the same independency structures**

There are actually **4 possible cases of independency maps**:

- Representable by a DAG, but not a MRF
- Representable by a MRF, but not a DAG
- Representable by both a DAG and MRF
- Not representable by either a DAG or MRF – fortunately, rare in practice

# Vocabulary Summary

A **clique** is a subset of vertices of an undirected graph such that **every two distinct vertices in the clique are adjacent**

A **maximal clique** is a clique which cannot be enlarged without violating the clique property

For a graph  $G$ , **disjoint** subsets  $A$  and  $B$  are separated by subset  $S$  if every path from  $A$  to  $B$  passes through  $S$  then  $S$  **separates**  $A$  and  $B$ . Or, if  $S = \emptyset$ , then no path exists from  $A$  to  $B$ , and  $A$  and  $B$  are **separated**

For a graph  $G$  with disjoint sets  $A$ ,  $B$  and  $S$ , where  $S$  separates  $A$  and  $B$ , then  $A \perp B \mid S$  known as the **global Markov property**

Given subsets  $X$ ,  $Y$  and  $Z$ ,  $X$  and  $Y$  are conditionally independent or **D-separated** conditioned on the subset  $Z$  if they are separated on the moralized graph

A graph  $G$  is a **dependency map** or **D-map** of a distribution  $P$  if the graph contains every conditional independence in  $P$ .

A subset  $U$  is a **Markov blanket** of  $X$  in the set of nodes  $\chi$  of the graph  $G$  if  $X \notin U$  and if  $U$  is a minimal set of nodes such that:  $(X \perp \chi - \{X\} - U \mid U) \in I(P)$

Two nodes are conditionally independent given the other nodes in the graph if there is no direct edge between them. This property is the **pairwise Markov property**

**Moralization** is a key step in turning DAG to an MRF

An **immorality** in a directed graph  $G$  occurs where either; a) there is a directed edge between  $X$  and  $Y$ , or b)  $X$  and  $Y$  are both parents of the same node  $Z$

A **moral graph**,  $M(G)$ , of a BN structure,  $G$  is the **undirected graph** over  $X$  that contains an undirected edge between  $X$  and  $Y$  if; a) there is a directed edge between  $X$  and  $Y$ , or b)  $X$  and  $Y$  are both parents of the same node  $Z$