

Lecture 1: Fundamentals of Magnetic Resonance Imaging

Administrative: Exercise registration poll

LOGBEITRAG: "Semesterstart mit StudOn / Start of semester with StudOn" unter "Support > StudOn-News"

[Repository](#) > [5. Faculty of Engineering](#) > [5.6 AIBE](#) > [Computational Imaging Lab](#) > [Campo Courses: Winter 2024/25](#) > [Computational Magnetic Resonance Imaging Vorlesung](#) > [Lab exercise](#) > [Sign up for an HPC account](#)



Sign up for an HPC account

If you want to participate in the exercise, you will need to have an HPC account.

[Booking Objects](#) | [Info](#) | [Reservations](#) | [Settings](#) | [Participants](#)

[Add Object](#)



Booking Objects

(1 - 1 of 1)

Title	Description	Available	Actions
Sign up for an HPC account		121/150	Actions ▼

(1 - 1 of 1)

Outline




B_0



B_1




G



Nuclear spin

$$\vec{L} = \vec{r} \times (m\vec{v})$$





Angular momentum of nucleus

Magnetic dipole moment



$$\vec{\tau} = \vec{\mu} \times \vec{B}$$





$$\vec{\mu} = \gamma \vec{I}$$

Gyromagnetic ratio



Element	(MHz/T)
^1H	42.58
^3He	-32.43
^{23}Na	11.26
^{31}P	17.24

Thermal motion: Random orientation






Interaction with B_0 : Magnetization



$$M = \frac{\rho \gamma^2 \hbar B_0}{4kT}$$

$$\vec{M} = \sum \mu_i$$





Interaction with B_0 : Precession

$$\vec{\tau} = \vec{\mu} \times \vec{B}$$


$$\omega_0 = \gamma B_0 \text{ Larmor frequency (} ^1\text{H)}$$

$$\frac{\gamma}{2\pi} = 42.6(\text{MHz}/T)$$

Angular momentum of
top in gravitation field




Nolan 2010







Interaction with B_0 : Precession




Interaction with radiofrequency field B1

$$f_0 = \frac{\gamma}{2\pi} B_0$$




Interaction with radiofrequency field B1

$$f_0 = \frac{\gamma}{2\pi} B_0$$





$$\frac{d\vec{M}}{dt} = \gamma \left(\vec{M} \times \vec{B}_{1+} \right)$$




Interaction with radiofrequency field B1

$$f_0 = \frac{\gamma}{2\pi} B_0$$




Signal reception

$$U_{ind} = - \frac{d\Phi}{dt}$$




Hahn 1950




Contrast


PD_w




T1_w





T2_w





Longitudinal Relaxation





Longitudinal Relaxation



Longitudinal Relaxation





Longitudinal Relaxation





Longitudinal Relaxation



$$M_z = M_0 \left(1 - e^{-t/T_1}\right)$$





Transversal Relaxation



Transversal Relaxation





Transversal Relaxation




Transversal Relaxation

$$M_{xy} = M_0 e^{-t/T_2}$$



Transversal Relaxation

$$M_{xy} = M_0 e^{-t/T_2^*}$$




$$\frac{1}{T_2^*} = \frac{1}{T_2'} + \frac{1}{T_2}$$

T_2 Tissue property, irreversible



T_2' Field inhomogeneity
(magnet, susceptibility),
reversible

$$T_2^* < T_2$$


PD contrast





$$M_z = M_0 \left(1 - e^{-t/T_1} \right)$$






PD contrast




$$M_{xy} = M_0 e^{-t/T_2}$$





T₂ contrast




PD contrast





$$M_z = M_0 \left(1 - e^{-t/T_1} \right)$$




T₁ contrast




$$M_z = M_0 \left(1 - e^{-t/T_1}\right)$$







How do we get an image?




Hahn 1950







Interaction with gradient fields G





$$\omega_0 = \gamma B_0$$

$$\oplus B_0$$








Interaction with gradient fields G







$$\omega(x) = \omega_0 + \gamma G_x x \oplus B_0$$




$$B(x) = B_0 + G_x x$$







Interaction with gradient fields G




$$B(y) = B_0 + G_y y$$




Interaction with gradient fields G







$$\omega(z) = \omega_0 + \gamma G_z z \oplus B_0$$




$$B(z) = B_0 + G_z z$$



Interaction with gradient fields G





$$\omega(z) = \omega_0 + \gamma G_z z \oplus B_0$$





$$B(z) = B_0 + G_z z$$



Pulse sequences





Pulse sequences





Pulse sequences





Pulse sequences





Pulse sequences




Pulse sequences



Pulse sequences



Pulse sequences





$$\phi(x, t) = \gamma x \int_0^t G_x d\tau \equiv k_x x$$

$$\phi(y, G_y) = \gamma y \int_0^{T_y} G_y d\tau \equiv k_y y$$

MR pulse sequence and k-space

Slice selection





$$s(k_x, k_y) = \int_x \int_y m(x, y) e^{-ik_x x} e^{-ik_y y} dx dy$$

$$\phi(x, t) = \gamma x \int_0^t G_x d\tau \equiv k_x x$$

$$\phi(y, G_y) = \gamma y \int_0^{T_y} G_y d\tau \equiv k_y y$$

Summary




B_0




B_1



G




Next week




Outlook lab exercise


label 1, CSF



label 2, GM




label 3, WM




SDw



T1w



T2w



FLAIR



MPRAGE

