Statistical Physics of Nakamoto Consensus: Phase Transitions and Thermodynamic Stability

Pascal Ranaora[1,*]

[1] *Information Physics Institute, Sydney, Australia*

(Dated: March 19, 2026)

The Nakamoto protocol is traditionally analyzed through the lens of cryptography and game theory. This paper proposes a paradigm shift by demonstrating that distributed consensus is fundamentally a dissipative structure maintained far from thermodynamic equilibrium. By mapping the macroscopic observables of the network to a mean-field Landau framework, we provide a falsifiable, physical foundation for blockchain immutability. We show that the emergence of a canonical history is akin to a continuous phase transition where time is not an *a priori* external clock, but an emergent thermal property strictly conjugate to irreversible heat dissipation. By unifying network latency and block interval through an effective temperature, our model reveals a strict thermodynamic limit on the informational volume of bytes per block ($V_{B,\mathrm{crit}}$) to prevent topological collapse via a Kosterlitz-Thouless transition. Furthermore, we recontextualize the UTXO set as a Markov blanket necessary for dissipating spatial Shannon entropy, and demonstrate that macroeconomic adjustments (the "Halving") operate as first-order thermodynamic quenches. In this non-equilibrium regime, the network's violation of the fluctuation-dissipation theorem is dynamically stabilized by the difficulty adjustment, acting as a macroscopic relaxation thermostat. Rather than a mere software ledger, the blockchain emerges as a physical attractor endowed with colossal Effective Mass, redefining immutability not as a mathematical certainty, but as a thermodynamic inevitability.

I. INTRODUCTION: THE THERMODYNAMICS OF DISTRIBUTED CONSENSUS

The fundamental problem of distributed consensus lies in establishing a canonical ordering of events in the absence of a global clock. While classic Byzantine Fault Tolerance (BFT) relies on logical clocks and voting thresholds, such systems lack physical grounding, making them vulnerable to Sybil attacks where costless identities generate arbitrary histories [1, 2]. From the perspective of information physics, the "Double Spending" problem represents a failure of time-ordering invariance; without an irreversible thermodynamic cost, the transformation $t \to -t$ is a valid symmetry, rendering the history $\mathcal{H}_A = \{E_1, E_2\}$ physically indistinguishable from $\mathcal{H}_B = \{E_2, E_1\}$ [3]. To extract a unique and immutable history from the stochastic noise of a peer-to-peer network, the system must undergo a symmetry-breaking process induced by energy dissipation [4].

We propose a minimal physical model of the Nakamoto protocol as a non-equilibrium thermodynamic system. We treat the distributed ledger not as a discrete data structure, but as a one-dimensional growing network $\mathcal{L}$ evolving under dissipative forcing. The emergence of consensus is modeled as a continuous phase transition, moving from a high-entropy "disordered" phase (the mempool) to a low-entropy "ordered" phase (the blockchain), analogous to the crystallization of a liquid [5]. Unlike static data structures, the blockchain is a "dissipative structure" that maintains its low-entropy state far from equilibrium solely through the continuous consumption of work [6, 7].

This manuscript substantially expands and formalizes the preliminary thermodynamic framework introduced in our earlier communication [8]. While the foundational physics-consensus isomorphism was previously outlined, the present work introduces rigorous gauge invariance via the covariant derivative, establishes strict finite-size scaling limits for the Markov boundary, and provides direct empirical validations of the fluctuation-dissipation theorem violation during macroeconomic quenches.



Figure 1. **The Arrow of Time.** Top: Without energy dissipation, the ordering of events is time-symmetric and fluid. Bottom: The injection of Proof of Work breaks this symmetry, crystallizing events into a rigid and irreversible thermodynamic sequence.

Before formally developing the continuous field theory and stochastic dynamics in the subsequent sections, Table I establishes the strict dictionary of isomorphisms used throughout this manuscript. It directly maps the macroscopic quantities of statistical mechanics to the observable metrics of the network.

* pascal.ranaora@informationphysicsinstitute.net

Physical Concept	Bitcoin Observable	Thermodynamic Role
Spin State ($\vec{s}_i, \theta_i$)	History Branch	Fundamental degree of freedom of the P2P network.
Order Parameter (Ψ, ϕ)	Honest Hashrate	Measure of symmetry breaking and consensus state.
Effective Temp. (T_{eff})	Latency Ratio (τ_L/τ_B)	Stochastic agitation caused by desynchronization.
Mass Gap ($m = \xi^{-1}$)	Difficulty Target	Inertial rigidity of history against reorganizations.
Hamiltonian ($\mathcal{H}_{eff}$)	Cumulative PoW	Action forcing the system toward a unique ground state.
Coupling Energy ($\mathcal{E}_{coupling}$)	Topo-Latency Tension	Macroscopic restorative force smoothing micro-forks.
Active Noise (ζ)	Block Discovery Variance	Stochastic agitation driving FDT violation.
Markov Blanket ($\partial\mathcal{V}$)	UTXO Set	Dissipative surface minimizing Shannon entropy.
Thermodynamic Quench	The "Halving"	External shock purging the network of entropy.

Table I. **Physics-Consensus Isomorphism Dictionary.** This table maps the strict correspondence between macroscopic quantities and observable protocol metrics.

II. NETWORK MODEL, STATISTICS AND THE ARROW OF TIME

A. The Minimal Network Model

We model the blockchain as a growing one-dimensional network $\mathcal{L}$ of length $N(t)$, where each site i represents a block B_i. The macroscopic state of the system is defined by the block sequence $\mathbf{S} = \{B_0, B_1, ..., B_N\}$. Unlike static data structures, this network is dynamic; its geometry is determined by a stochastic growth process fighting against entropic degradation (information loss via forks).

We define the state space of the network Ω as the set of all possible chain permutations (forks). The probability $P(\mathcal{H})$ that a specific history $\mathcal{H} \in \Omega$ is selected as the consensus reality follows a Boltzmann distribution derived from the canonical ensemble:

$$P(\mathcal{H}) = \frac{1}{\mathcal{Z}} e^{-\beta E(\mathcal{H})} \tag{1}$$

where $\mathcal{Z}$ is the partition function and β is the inverse effective temperature, determined by the ratio between the honest hashrate and network latency [9].

To align the "Longest Chain Rule" with the Principle of Least Action, we define the effective Hamiltonian $\mathcal{H}_{eff}$ of the system as the negative of the cumulative Proof of Work (PoW):

$$\mathcal{H}_{eff} = -\sum_{i \in \text{blocks}} E_{eff}(B_i) \tag{2}$$

where $E_{eff}(B_i)$ represents the macroscopic characteristic energy dissipated to mine block B_i. In this framework, Nakamoto Consensus is not an arbitrary algorithm but a physical inevitability: the system naturally relaxes into the ground state that minimizes free energy $\mathcal{F} = E - TS$ [10]. The so-called "51% attack" is thus recontextualized not as a security flaw, but as a thermally induced phase transition (melting) where the temperature of the attack vector T_{attack} exceeds the critical temperature T_c of the network's binding energy.

B. Time-Reversal Symmetry and Sybil Degeneracy

Consider a distributed ledger system with zero energetic cost for block creation ($E_{eff} = 0$). In this regime, the system exhibits **Time-Reversal Symmetry** ($\mathcal{T}$-Symmetry). For any given history $\mathbf{S}_A$ (e.g., "Alice pays Bob"), there exists computationally symmetric history $\mathbf{S}_B$ (e.g., "Alice pays Charlie") that is indistinguishable to a new observer [3].

This degeneracy implies that the "arrow of time" is undefined; the system exists in a superposition of microstates with equal statistical weight. Mathematically, this corresponds to a system at infinite temperature ($T \to \infty$). The partition function diverges, and the probability of observing a specific history $\mathbf{S}$ becomes uniform:

$$P(\mathbf{S}_A) \approx P(\mathbf{S}_B) \approx \frac{1}{|\Omega|} \tag{3}$$

where $|\Omega|$ is the volume of the state space (all possible forks). This is the physical definition of the "Sybil" attack: a thermodynamic degeneracy where costless identities can generate arbitrary histories [2]. Without a symmetry-breaking field, the history entropy $S_{history}$ is maximized, rendering the ledger purely random and devoid of information [6].

C. The Consensus Hamiltonian and Information Temperature

In a classic closed thermodynamic system, the ground state emerges through the minimization of internal energy. However, the blockchain is an inherently open dissipative structure, coupled to a global exergy bath. Following the Maximum Entropy Production Principle (MEPP) and the Maximum Power Principle [11, 12], this non-equilibrium system self-organizes to maximize its vital energy dissipation.

Consequently, to select a unique history, we must introduce an effective Hamiltonian $\mathcal{H}_{eff}$ that does not represent a potential energy to be minimized, but the negative of the Dissipation Action. The ledger "falls" towards the state that requires the maximum destruction of irre-

versible work:

$$\mathcal{H}_{eff}(\mathbf{S}) = -\sum_{i=0}^{N} E_{eff}(B_i) \cdot \mathbb{I}_{valid}(B_i) \tag{4}$$

The probability that a specific history emerges as the macroscopic "truth" is thus governed by the canonical measure, favoring trajectories of highest dissipation:

$$P(\mathbf{S}) = \frac{1}{\mathcal{Z}} e^{\beta \sum E_{eff}(B_i)} \tag{5}$$

Here, the parameter $\beta = (k_B T_{eff})^{-1}$ acts as the **Inverse Information Temperature**. Unlike standard statistical models where temperature is dictated by an external heat bath, the effective temperature of the ledger T_{eff} is an endogenous emergent property.

To ensure this quantity has the strict dimension of energy, we introduce E_{eff}, the macroscopic characteristic energy scale of the network, defined as the total physical work required to crystallize a single block (formally derived from the network's instantaneous power in Section III C). As we will demonstrate in Section IV C, the parameter β translates the thermodynamic competition between the work-induced crystallization time τ_B and the spatial uncertainty diffusion time τ_L. We thus set:

$$\beta = \frac{1}{E_{eff}} \frac{\tau_B}{\tau_L} \tag{6}$$

The thermodynamic regimes of the ledger are naturally defined as follows:

- **High Temperature ($\beta \to 0$, $\tau_L \to \tau_B$):** Spatial diffusion entropy dominates crystallization work. The system is "liquid", forks (thermal fluctuations) multiply, and history is mutable.
- **Low Temperature ($\beta \to \infty$, $\tau_L \ll \tau_B$):** Negentropy flux dominates. The system "freezes" into a low-energy state, making history rigid and immutable.

The "Longest Chain Rule" is therefore not an arbitrary software heuristic. It is the inevitable state trajectory of a system seeking to minimize its Free Energy $\mathcal{F} = E - T_{eff} S$ [10].

D. Spontaneous Symmetry Breaking (SSB)

The transition from the mempool to the blockchain corresponds to **Spontaneous Symmetry Breaking (SSB)**. We define the Order Parameter Ψ as the **Net Magnetization** of the network, by analogy to the Ising model:

$$\Psi = \langle \frac{1}{N} \sum_i \vec{s}_i \rangle \tag{7}$$

where $\vec{s}_i$ is the "consensus spin" vector of node i.

- **Disordered Phase ($T > T_c$):** $\Psi \approx 0$. Nodes are randomly oriented (forks). The system possesses $O(2)$ rotational symmetry (no preferred history).
- **Ordered Phase ($T < T_c$):** $\Psi \to 1$. Nodes spontaneously align along a single "worldline". Rotational symmetry is broken.

This phase transition is induced by the injection of low-entropy energy (electricity), which acts as an external magnetic field H_{ext} aligning the spins.



Figure 2. **Symmetry Breaking in Consensus.** At high effective temperature (zero work), the free energy $\mathcal{F}$ has a single minimum at $\Psi = 0$ (Disorder/Sybil). Below the critical temperature (positive work), the potential bifurcates. The system must spontaneously "roll" into one of the stable vacua (a specific history), thereby breaking the symmetry. Returning to the other vacuum (Double Spending) requires overcoming the potential barrier $\Delta\mathcal{F}$, which grows with confirmation depth.

E. Emergence of Thermal Time and Fluctuation Theorem

In an asynchronous distributed network, there is no absolute global clock. Time is not an *a priori* fundamental variable, but a thermodynamic emergent property [13]. We formalize this using the **Thermal Time Hypothesis** of Rovelli and Connes [14], which postulates that the flow of time is generated by the statistical state of the system itself.

For a macroscopic state described by a density matrix ρ, the modular Hamiltonian is defined as $\mathcal{H}_{mod} = -\ln \rho$. In the context of Nakamoto consensus, the system is governed by our effective Hamiltonian $\mathcal{H}_{eff}$ at inverse temperature β. The flow of "thermal time" τ (measured discretely as block height N) is strictly conjugate to the energy dissipation via the modular automorphism:

$$\frac{d\tau}{ds} \propto \beta \mathcal{H}_{eff} \implies \Delta N \propto \frac{1}{k_B T_{eff}} \int \dot{Q}\, dt \tag{8}$$

It is crucial to establish a formal distinction between the internal time of the macroscopic state and the external time of the physical substrate. From the internal per-

spective of the ledger, time is discontinuous; the macroscopic clock τ solely ticks upon the discrete crystallization of a new block ($\Delta N = 1$). However, the stochastic search process, the geographic propagation of information (τ_L), and the continuous dissipation of heat (P_{net}) unfold in the continuous Newtonian time t of the physical universe. It is precisely the thermodynamic tension between the continuous external energy injection t and the discrete internal topological updates τ that gives rise to the emergent effective temperature T_{eff} and governs the phase stability of the consensus.

Unlike approaches positing an independent physical time, the local time metric N only advances when irreversible heat $\dot{Q}$ is dissipated into the environment, breaking detailed balance.

We quantify this time asymmetry and the resolution of the Double Spending problem ($t \to -t$ symmetry) using the **Crooks Fluctuation Theorem** [15] from non-equilibrium stochastic thermodynamics. For a ledger transition from state A to B, the ratio of the forward probability P_F (honest validation) to the reverse probability P_R (reorganization attack) is exponentially bounded by the dissipated work W exceeding the free energy barrier $\Delta\mathcal{F}$:

$$\frac{P_F(W)}{P_R(-W)} = e^{\beta(W-\Delta\mathcal{F})} = e^{\Delta S_{irrev}/k_B} \tag{9}$$

Immutability is therefore not an algorithmic absolute, but a statistical inevitability imposed by the Second Law.



Figure 3. **Crooks Fluctuation in Nakamoto Consensus.** The probability distributions of honest work $P_F(W)$ and attacker reorganization work $P_R(-W)$. The intersection occurs exactly at the reversible free energy difference $\Delta\mathcal{F}$. For a deeply confirmed block, the honest network operates far to the right ($W \gg \Delta\mathcal{F}$), making the probability of a reverse trajectory exponentially vanishing. As the network injects macroscopic work ($W \gg \Delta\mathcal{F}$), the Crooks ratio diverges exponentially.

III. THERMODYNAMIC POTENTIAL AND SYMMETRY BREAKING

We analyze the stability of the ledger using the Landau theory of phase transitions. We postulate that the blockchain operates as a continuous field $\phi(x,t)$ representing the **security density** (hashrate) across the network.

A. The Order Parameter and Landau Free Energy

Rather than tracking the discrete, microscopic topological growth of individual blocks, we analyze the system at the hydrodynamic limit using a mean-field approach. We define the complex order parameter $\phi(x)$:

$$\phi(x) = \rho(x)e^{i\theta(x)} \tag{10}$$

where $\theta(x)$ is the phase, representing the specific history (chain tip) selected by the miner at location x. A state where $\nabla\theta = 0$ corresponds to global consensus, while $\nabla\theta \neq 0$ corresponds to a fork (domain wall).

To formalize the free energy, we define the amplitude $\rho(x)$ as a strictly dimensionless ratio $\phi = H/H_{eq}$, where H is the observed local hashrate and H_{eq} is the macroscopic equilibrium hashrate sustained by the network's global exergy bath. By normalizing the order parameter, we ensure that the effective potential $V(\phi)$ carries the strict physical dimension of Energy.

The equilibrium state of the network is governed by the minimization of the Landau Free Energy functional $\mathcal{F}$, where the potential density is:

$$V(\phi) = a(T_{eff})|\phi|^2 + b|\phi|^4 \tag{11}$$

This is the standard "Mexican hat" potential used to describe spontaneous symmetry breaking. The coefficients are determined by the fundamental thermodynamic bounds of the network:

1. **The Instability Term and Exergetic Heat Capacity (a):** The parameter a governs symmetry breaking and depends explicitly on the effective temperature of the system. We set:

$$a(T_{eff}) = \chi_E(T_{eff} - T_c) \tag{12}$$

where $\chi_E > 0$ acts as the **exergetic heat capacity** of the network, possessing the strict physical dimension of Entropy ($\mathrm{J{\cdot}K^{-1}}$). It converts the thermal deviation ($T_{eff} - T_c$) into an energetic driving force.

Using our dimensionless ratio $\Gamma = \tau_B/\tau_L$ developed in Section IV C (where $T_{eff} \propto \Gamma^{-1}$), the quadratic term becomes a direct function of the network's spatial latency:

$$a(\tau_L) = \mathcal{E}_{coupling}\left(\frac{\tau_L}{\tau_B} - \frac{1}{\Gamma_c}\right) \tag{13}$$

where $\mathcal{E}_{coupling} = \chi_E \cdot T_c \cdot \Gamma_c$ represents the characteristic **topological coupling energy** [Joules] required to synchronize the network state globally.

- **Ordered Regime ($\tau_L/\tau_B < \Gamma_c^{-1}$):** Mining incentives overcome entropic diffusion. Parameter $a < 0$, making the degenerate state $|\phi| = 0$ thermodynamically unstable.
- **Disordered Regime ($\tau_L/\tau_B > \Gamma_c^{-1}$):** If block size V_B pushes latency beyond the critical threshold, $a > 0$. The potential loses its secondary minima; the chain "melts" and consensus fractures.

2. **The Saturation Term and ASIC Mass ($b > 0$):** The quartic term $b|\phi|^4$ provides the non-linear restorative force preventing the normalized hashrate from diverging to infinity. Crucially, this term anchors the abstract field theory into the **biophysical reality of the hardware substrate**.

   We define the **computational mass efficiency** μ_m of the network as the ratio between the average physical mass of a mining unit M_{unit} and its raw hashrate output H_{unit}. Using empirical specifications for 2026-era industrial miners ($M_{unit} \approx 15$ kg for a 100 TH/s output):

$$\mu_m = \frac{M_{unit}}{H_{unit}} \approx 1.5 \times 10^{-13} \text{ kg} \cdot \text{s} \tag{14}$$

   As the normalized hashrate $|\phi|$ grows, the energy barrier b [Joules] required to maintain the hardware substrate against obsolescence and environmental entropy naturally scales. It dynamically integrates both marginal operational expenditure (bounded by Landauer's limit) and capital expenditure (bounded by μ_m, supply chain logistics, and spatial cooling limits).

Driven by these bounds, the system relaxes into the stable ground state with a non-zero expectation value $|\phi_0|$, representing the normalized equilibrium hashrate:

$$|\phi_0| = \sqrt{\frac{-a}{2b}} \tag{15}$$

B. Spontaneous Symmetry Breaking

The transition from the mempool (disordered) to the blockchain (ordered) corresponds to the spontaneous breaking of the $U(1)$ symmetry of phase θ.

- **Before Mining:** All nonces (phases) are equiprobable. Symmetry is not broken.
- **After Mining:** A specific nonce is found. Symmetry is broken, and a specific history is selected.



Figure 4. **The Consensus Potential.** The system spontaneously rolls from the unstable origin (zero security) toward the stable valley ϕ_0 (equilibrium hashrate), defined by the balance between incentive (a) and cost (b).

The "Goldstone mode" associated with this breaking is the massless fluctuation of the nonce, corresponding to the random search process. The "massive mode" is the rigidity of the hashrate, which resists perturbations (attacks) (see Figure 4).

C. Macroscopic Energy Scale, the DAA Thermostat

To ground the characteristic energy scale E_{eff} in the empirical reality of the network, we must first define it macroscopically as the total physical work required to crystallize a single block. This energy barrier is strictly dictated by the protocol's Difficulty parameter D, which imposes an expected number of hashes $\mathcal{N}_H(D)$ required to satisfy the Proof of Work condition. We note $\eta_{hash}(t)$ the thermodynamic efficiency of the hardware (in Joules per hash), then the effective state energy for mining a block is:

$$E_{eff}(t) = \mathcal{N}_H(D) \cdot \eta_{hash}(t) \tag{16}$$

Since the expected number of hashes is the product of the global magnitude of the order parameter (the Hashrate) $|\phi(t)|$ and the realized block interval τ_B, we can express this as:

$$E_{eff}(t) = \eta_{hash}(t) \cdot |\phi(t)| \cdot \tau_B \tag{17}$$

Recognizing that the product of hashing efficiency and the global hashrate ($\eta_{hash}(t) \cdot |\phi(t)|$) constitutes the total instantaneous power dissipation of the network $P_{net}(t)$ (in Watts), the macroscopic energy scale simplifies to:

$$E_{eff}(t) = P_{net}(t) \cdot \tau_B \quad \text{[Joules]} \tag{18}$$

The energy scale E_{eff} is not an unconstrained variable but is strictly governed by the Difficulty Adjustment Algorithm (DAA), which acts as a **non-linear feedback controller**. By continuously modulating $\mathcal{N}_H(D)$, the DAA ensures that E_{eff} is dynamically scaled to maintain the canonical interval $\tau_B \approx 600$s. In the thermodynamic

limit, the DAA prevents E_{eff} from diverging relative to the network's power, effectively "clamping" thermal fluctuations and preventing a transition into a disordered or chaotic state.

This physical derivation yields a profound result when injected back into our definition of Information Temperature ($k_B T_{eff} = E_{eff} \frac{\tau_L}{\tau_B}$) from Section II C:

$$k_B T_{eff} = (P_{net} \cdot \tau_B) \frac{\tau_L}{\tau_B} = P_{net} \cdot \tau_L \tag{19}$$

The thermal noise of the Nakamoto consensus ($k_B T_{eff}$) is equal to the "blind" energy dissipated by the network during the geographic latency window τ_L. It is the literal physical work expended in the dark that is at risk of generating an orphaned fork.

This derivation reveals the precise thermodynamic function of the Difficulty Adjustment Algorithm (DAA). If Hashrate $|\phi(t)|$ were to surge toward infinity without intervention, crystallization time would shrink toward zero ($\tau_B \to 0$), causing the spatial entropy ratio τ_L/τ_B to explode and melting the consensus into a disordered state. The DAA acts as a strict physical thermostat: it dynamically raises the required difficulty $\mathcal{N}_H(D)$ to force τ_B back to the canonical 600 seconds. By artificially inflating the energy gap E_{eff} of a block, the protocol absorbs the exergy influx, neutralizes the temperature spike, and maintains the crystalline state of the ledger.

D. The Nakamoto Action and Hardware Substrate

While E_{eff} governs the macroscopic phase state, it is built upon a fundamental physical substrate. To establish a strict physical isomorphism with the Principle of Least Action, we isolate the fundamental quantum of action of the network: the **Nakamoto Action** κ_N.

This constant has the strict dimensions of physical action ($J \cdot s$). We define it kinematically at the microscopic level of the ASIC hardware:

$$\kappa_N(t) = \eta_{hash}(t) \cdot \tau_{cycle} \quad [J \cdot s] \tag{20}$$

where τ_{cycle} is the fundamental switching time of a logic gate within the ASIC.

It is imperative to establish a rigorous separation of timescales to properly situate this metric. The Nakamoto Action $\kappa_N(t)$ operates strictly at the *micro-canonical* scale of the hardware substrate, characterized by ultra-fast Markovian dynamics ($t \sim \tau_{cycle} \sim 10^{-12}$ s). It acts as the fundamental boundary constraint—the absolute floor of the phase space enforcing Landauer's limit. However, it does not directly dictate the coarse-grained fluid dynamics of the consensus.

In contrast, the *macro-canonical* dynamics of the ledger operate on the scale of $\tau_B \approx 600$ s. The emergence of the macroscopic state energy E_{eff} results from the statistical integration of billions of independent micro-canonical fluctuations (the Poisson search process). This coarse-graining makes the macroscopic consensus mechanics robust and fundamentally agnostic to the specific underlying hardware architecture, provided the system collectively respects the fundamental physical anchor established by $\kappa_N(t)$.

Unlike Planck's constant $\hbar$ which is universal and fixed, $\kappa_N(t)$ is a non-stationary, purely dissipative technological parameter. Its value is dynamically revised downward as semiconductor efficiency improves, tracing an asymptote toward Landauer's limit imposed by information physics.

We can thus formally define the **Consensus Action** $\mathcal{S}_{PoW}$ evaluated along the chain history $\mathcal{C}$ as the time integral of the macroscopic energy:

$$\mathcal{S}_{PoW} = \int_{\mathcal{C}} E_{eff}(t)\, dt \quad [J \cdot s] \tag{21}$$

This formulation demonstrates that the ledger does not merely accumulate discrete blocks of data, but rather accumulates *Action*. By dissipating heat, the protocol decreases the logical entropy of the system. The "Heaviest Chain" is thus physically equivalent to the phase trajectory that maximizes this accumulated action, creating an insurmountable thermodynamic barrier against history reorganization.

IV. CORRELATION LENGTH AND PROBABILISTIC FINALITY

In this section, we derive the probabilistic finality of the ledger not as a geometric horizon, but as a correlation decay property in a massive field theory. We treat the blockchain as a 1D spin chain at finite temperature.

A. The Correlation Function

We define ledger stability via the two-point correlation function $G(z)$. Let $sigma_i$ be the block state at height i (confirmed vs. rejected). The probability that the state at depth z remains correlated with the current consensus tip is given by:

$$G(z) = \langle \sigma_{tip} \cdot \sigma_{tip-z} \rangle \tag{22}$$

In a massless system (zero difficulty), fluctuations would propagate infinitely, and history would be mutable at all depths ($G(z) \sim$ const). However, the Proof of Work mechanism introduces a "mass gap" m (the difficulty target) into the theory. In massive field theories, correlations decay exponentially with distance:

$$G(z) \sim e^{-z/\xi} \tag{23}$$

where ξ is the **Correlation Length**. The cumulative probability of consensus (Probabilistic Finality) is thus given by $1 - G(z)$. This length scale is determined by the ratio between the "binding energy" (Honest Hashrate P_H) and the "thermal noise" (Attacker Hashrate P_A).

B. Derivation of the Mass Gap (The "6-Block" Rule)

The probability of Nakamoto consensus (Gambler's Ruin) can be rewritten as an exponential decay of the reorganization probability P_{reorg}:

$$P_{reorg}(z) = \left(\frac{q}{p}\right)^z = e^{-z\cdot\ln(p/q)} \tag{24}$$

where p is the honest probability and q the attacker probability ($p+q=1$). Comparing this to the standard exponential decay in statistical mechanics $e^{-z/\xi}$, we identify the inverse correlation length (mass) m:

$$m = \xi^{-1} = \ln\left(\frac{p}{q}\right) \approx \frac{P_{\text{Honest}} - P_{Attacker}}{P_{Total}} \tag{25}$$



Figure 5. **Correlation Decay.** The probability of reorganization decays exponentially with depth z. The system transitions from a "fluid" phase at the tip to a "frozen" solid phase in the bulk. The characteristic length scale ξ defines probabilistic finality.

Physical Interpretation:

- **Massless Phase ($m \to 0$):** If $p \approx q$ (51% attack threshold), the correlation length $\xi \to \infty$. Finality is never reached; the system remains in a "critical state" of infinite susceptibility to reorganization.
- **Massive Phase ($m > 0$):** If $p \gg q$, the correlation length is short. History "freezes" rapidly. For Bitcoin ($q \approx 0.1$), $\xi \approx 2$ blocks. At $z = 6$ blocks ($z \approx 3\xi$), the survival probability of a competing branch drops below 0.1%.

C. Effective Temperature, Dimensionless Ratio, and Block Size Limit

In blockchain dynamics, the "tip" of the chain behaves like a fluid interface subject to thermal roughness (orphan blocks). We formally define the Effective Temperature T_{eff} of the network as the measure of energy dissipated "blindly" due to spatial desynchronization.

It is imperative to distinguish here the **effective temperature** T_{eff}, which is a measure of the stochastic agitation of the graph, from the true thermal temperature T of the physical substrate. While T governs Landauer's limit at the individual hashing level, T_{eff} governs the macroscopic stability of consensus against latency noise.

Recalling our definition of the Inverse Information Temperature (defined in Eq. 6) $\beta = \frac{1}{E_{eff}}\frac{\tau_B}{\tau_L}$ we can isolate a fundamental dimensionless ratio Γ, which governs the overall stability of the ledger by pitting the thermodynamic crystallization time against the spatial entropy diffusion time:

$$\Gamma = \frac{\tau_B}{\tau_L} \tag{26}$$

Since the inverse temperature parameter of our Boltzmann distribution is $\beta = (k_B T_{eff})^{-1}$, we logically have a strict analytical relationship:

$$\beta = \frac{\Gamma}{E_{eff}} \tag{27}$$

However, latency τ_L is not a static constant. It is intrinsically linked to the block size V_B, the overall communication channel capacity C (in the sense of the Shannon-Hartley theorem) [6], and an incompressible geometric propagation latency τ_0 (limited by the speed of light). We denote the block data magnitude as V_B, representing the Volume of Bytes contained within a discrete update. Unlike traditional 'size' metrics, this notation emphasizes the fluid-dynamic nature of information propagation across the network's Shannon capacity C :

$$\tau_L(V_B) = \tau_0 + \frac{V_B}{C} \tag{28}$$

Re-injecting this dependence into our inverse temperature parameter β, we obtain an exact analytical relationship between the data mass of a block and the thermodynamic state of the network:

$$\beta(V_B) = \frac{1}{E_{eff}}\frac{\tau_B}{\tau_0 + \frac{V_B}{C}} \tag{29}$$

This equation reveals a crucial phenomenon: increasing block size is physically equivalent to heating the system. As V_B increases, the time required for the state to homogenize across the network lengthens. Spatial Shannon entropy increases, mechanically raising the effective temperature T_{eff} [16]. There is consequently a critical block size $V_{B,\text{crit}}$ at which the effective temperature exceeds the critical phase transition temperature ($T_{eff} > T_c$). At this breaking point:

$$V_{B,crit} \approx C\left(\frac{\tau_B}{\Gamma_c} - \tau_0\right) \tag{30}$$

where Γ_c is the minimal critical ratio to maintain ordering.

To anchor this thermodynamic limit in the empirical reality of the Bitcoin network, we can perform a first-order numerical estimation. Based on propagation dynamics [16], the orphan block rate r grows linearly with the spatial entropy ratio: $r \approx \tau_L/\tau_B = \Gamma^{-1}$. If we set a strict thermodynamic security threshold where the correlation length ξ begins to collapse (e.g., a critical orphan rate $r_c \approx 10\%$, which dictates $\Gamma_c \approx 10$), and considering the canonical interval $\tau_B = 600$ s alongside an incompressible planetary propagation delay $\tau_0 \approx 0.5$ s, the critical size depends solely on channel capacity C. For a conservative median bandwidth capacity on a massively decentralized P2P network ($C \approx 10$ MB/s), the equation yields:

$$V_{B,crit} \approx 10 \text{ MB/s} \times \left(\frac{600 \text{ s}}{10} - 0.5 \text{ s}\right) \approx 595 \text{ MB} \quad (31)$$

This numerical result is fundamental: it demonstrates that proposals to radically increase block size (to the Gigabyte range) are not merely disk space challenges, but direct violations of the network's thermal percolation limit. If block size breaches this threshold $V_{B,\text{crit}}$, the system leaves the ordered "crystalline" phase and returns to a disordered "liquid" state. Consensus fractures into multiple temporal domains unable to align before the next energy injection.

Physically, this proves that the block size limit in the Nakamoto protocol is not an empirical software choice. It is a strict thermodynamic boundary dictated by the metric of spacetime (speed of light c) and the limits of information theory (Shannon capacity). To guarantee immutability and symmetry breaking, the network must imperatively enforce $V_B \ll V_{B,crit}$.

D. Nucleation of a False History

An attack on the chain is analogous to the **nucleation of a false vacuum bubble**. An attacker attempting to rewrite z blocks must create a "domain" of length z with a different spin configuration (history). The free energy cost ΔF of creating this domain is:

$$\Delta F(z) = z \cdot (\mathcal{E}_{\text{Honest}} - \mathcal{E}_{\text{Attacker}}) - TS_{\text{entropy}} \quad (32)$$

For a successful attack, the attacker must overcome the surface tension of the honest chain. If $\mathcal{E}_{\text{Honest}} > \mathcal{E}_{\text{Attacker}}$, the free energy cost grows linearly with z. The probability of spontaneous nucleation of a deep reorganization is thus exponentially suppressed:

$$P_{nucleation} \propto e^{-\Delta F/k_B T} \quad (33)$$

This confirms that immutability is not absolute, but **thermodynamically robust**. Rewriting history is not impossible; it is simply statistically forbidden by the Second Law of Thermodynamics (see Figure 5).

V. INFORMATION ENTROPY AND MARKOV BLANKET

The physical stability of the ledger depends on how the protocol processes and stores information against the strict limits of communication channel C. Given the thermal percolation constraint defined previously ($V_B \ll V_{B,crit}$), the network cannot continuously propagate and verify the entirety of the historical "volume". The protocol resolves this thermodynamic bottleneck by projecting history onto a lower-dimensional surface: a **Markov Blanket**.

A. The Volume and the Markov Projection

In statistical systems, a Markov Blanket [17] isolates the current state from its causal history: conditionally on this blanket, the future is independent of the past. In Nakamoto consensus:

- **The Volume ($\mathcal{V}$):** The complete historical blockchain, containing all past transactions and the total dissipated thermodynamic action $\mathcal{S}_{PoW}$.
- **The Blanket ($\partial\mathcal{V}$):** The Unspent Transaction Output (UTXO) set, representing the active state of the network.

Because bandwidth C is finite, the network cannot operate on the entire microscopic volume. It implicitly applies a spatial and temporal Renormalization Group (RG) flow. Hashing functions and block validation act as irreversible decimation operators: the network "integrates out" and eliminates inactive degrees of freedom (dead transactions and discarded alternative histories) to retain only the relevant macroscopic variables on the UTXO surface.

Unlike a perfect holographic projection, this renormalization step is a dissipative compression (with local microscopic entropy loss), absolutely essential to keep the exchange volume V_B below the critical threshold $V_{B,\text{crit}}$ and prevent the divergence of the system's macroscopic free energy.

B. Shannon vs. Thermodynamic Entropy

This projection via the Markov blanket allows us to resolve the apparent tension between Shannon information entropy [6] and Landauer thermodynamic entropy [4]. The minimum heat flux $\dot{Q}$ that must be dissipated locally by the network to maintain absolute certainty of the boundary against incoming information flux (new transactions) is strictly bounded by a dynamic dissipation rate:

$$\dot{Q} \geq k_B T_{bath} \ln(2) \frac{dH(X)}{dt} \quad (34)$$

To rigorously ground this constraint within the framework of non-equilibrium stochastic thermodynamics, we can invoke a generalized form of the fluctuation theorem (derived from Jarzynski's equality [18] for information processing). The average irreversible work $\langle W_{diss} \rangle$ provided by the network to integrate a new block must satisfy:

$$\langle W_{diss} \rangle \geq k_B T_{eff} \Delta H(X) \tag{35}$$

where $\Delta H(X)$ represents the spatial Shannon entropy (uncertainty) injected by the flow of unconfirmed new transactions. For a node to validate the network state autonomously, this massive work dissipates entropy so that uncertainty regarding the UTXO boundary state is strictly brought to zero ($H(X) = 0$).

To maintain this informational perfection on the boundary despite the constant background noise of the peer-to-peer network, the system must purge entropy by maximizing its thermodynamic entropy production in the volume (thermal dissipation via hashing). Any attempt to introduce an invalid transaction represents a massive injection of Shannon entropy (uncertainty) onto this boundary. The network rejects it immediately because its integration would require providing instantaneous physical work capable of overcoming the gigantic potential barrier $\Delta \mathcal{F}$ of the historical volume.

C. The UTXO Set as a Physical Maxwell's Demon

The relationship between information and thermodynamic entropy, formalized by the Mass-Energy-Information (MEI) equivalence, becomes explicitly manifest when we model the network as a macroscopic Maxwell's Demon.

In the classical thought experiment, a Demon reduces the entropy of a gas by measuring particle velocities and sorting them across a barrier. To avoid violating the Second Law of Thermodynamics, Bennett [3] demonstrated that the Demon must eventually erase its memory to reset the system, paying a mandatory thermodynamic cost governed by Landauer's Principle [4]:

$$W_{erase} \geq k_B T \ln(2) \tag{36}$$

per bit of erased information.

In Nakamoto consensus, the network of mining nodes acts as this decentralized Demon, and the UTXO set operates as the sorting barrier (the Markov Blanket). The "gas" is the Mempool—a high-entropy mixture of valid, invalid, and mutually exclusive (Double Spend) transactions.

The Demon measures the state of the Mempool, selecting a valid subset to crystallize into a block. However, by selecting one specific history $\mathcal{H}_A$, the network must actively *erase* all competing alternative histories $\mathcal{H}_{B,C...}$ from the active state space to reset the UTXO boundary for the next epoch.



Figure 6. **Nakamoto Consensus as a Maxwell's Demon.** The network acts as a thermodynamic Demon, measuring the Mempool (high Shannon entropy) and sorting valid transactions across the UTXO boundary (Markov Blanket). To reset its memory and enforce history, the Demon pays the mandatory Landauer erasure cost via Proof of Work, dissipating massive heat $\dot{Q}_{diss}$.

Proof of Work is the literal payment of this Landauer erasure cost. To irreversibly collapse the Shannon entropy ($\Delta H_{Shannon}$) of the Mempool into a single canonical block, the network must dissipate a macroscopic quantity of heat $\dot{Q}_{diss}$ into the environment:

$$\dot{Q}_{diss} = \frac{E_{eff}}{\tau_B} \geq k_B T_{bath} \ln(2) \frac{dH_{Shannon}}{dt} \tag{37}$$

This formulation provides the rigorous physical mechanism for the Effective Mass derived in our conclusion. The colossal energy consumed by the network is not "wasted" computation; it is the fundamental thermodynamic work required to perform macroscopic uncertainty erasure. Immutability is physically sustained because reversing the ledger would require the spontaneous, localized absorption of this exact macroscopic heat back from the environment—a statistical impossibility.

D. Blanket Saturation and Channel Limit

The security of this Markov blanket is intrinsically linked to the equations of information fluid dynamics. State updates $\partial \mathcal{V}$ (new blocks V_B) must pass through communication channel C. If the boundary update rate approaches channel capacity ($V_B \to V_{B,crit}$), the Markov property collapses globally. Nodes can no longer synchronize the UTXO surface within the allotted time τ_B. Shannon uncertainty explodes locally, effective temperature T_{eff} crosses the critical threshold, and consensus fractures into multiple isolated domains. History rigidity is therefore guaranteed only if the boundary update size

strictly respects the spatial dissipation capacity of the network.

VI. TOPOLOGICAL STABILITY OF THE NETWORK GRAPH

The robustness of the Bitcoin protocol against partition attacks cannot be understood solely through 1D thermodynamics. It requires analyzing long-range order across the spatial geometry of the peer-to-peer network. We model the consensus network using the **XY Model** on a **"small-world"** graph (Watts-Strogatz model) [9].

This architecture is characterized by two fundamental physical and mathematical properties:

1. **A high clustering coefficient:** Nodes form highly interconnected local clusters, ensuring massive information redundancy at the regional level.
2. **A very short average distance (Shortcuts):** A few long-distance random connections link distant clusters. On average, a transaction or new block can reach any node on the globe in just a few hops.



Figure 7. The "Small-World" topology of the Bitcoin network. Local links (gray) ensure redundancy and fault tolerance, while long-distance connections (dashed red) allow information to traverse the network, minimizing propagation delay.

A. The Consensus Phase Field

We assign a continuous phase variable $\theta_i \in [0, 2\pi)$ to each node i, representing its local view of the ledger tip. Interaction energy between peers seeks to minimize phase differences (consensus):

$$\mathcal{H}_{XY} = -J \sum_{\langle i,j \rangle} \cos(\theta_i - \theta_j) \tag{38}$$

where J is the coupling constant. To ensure internal consistency of the model, we define J no longer as a free parameter, but as the local action density injected into the network:

$$J \approx \kappa_N(t) \cdot |\phi_{local}| \tag{39}$$

where $|\phi_{local}|$ represents the local amplitude of the consensus field (the share of hashrate) perceived by the node's neighborhood. This formulation directly links the topological rigidity of consensus (vortex resistance) to actual energy dissipation.

- **Ground State ($\nabla\theta = 0$):** Global Consensus. All nodes agree on the tip.
- **Excited State ($\nabla\theta \neq 0$):** A Fork. The network is split into domains with different phases.

B. Forks as Topological Defects (Vortices)

In two-dimensional systems (like the P2P overlay network), thermal fluctuations can generate topological defects known as **vortices**. A vortex corresponds to a closed loop of nodes maintaining divergent views on chain state:

$$\oint_{\mathcal{C}} \nabla\theta \cdot d\mathbf{l} = 2\pi n, \quad n \in \mathbb{Z} \tag{40}$$

If $n \neq 0$, the loop encloses a singularity (a persistent fork). The energy of such a defect grows logarithmically with system size L:

$$E_{vortex} \approx \pi J \ln(L/a_0) \tag{41}$$

where a_0 is the lattice constant (the minimum discrete distance between two nodes, typically representing one hop in P2P topology). Since $E_{vortex} \to \infty$ for a large network $L \gg a_0$, isolated forks are energetically suppressed in the thermodynamic limit.

C. Effective Dimensionality and the Kosterlitz-Thouless Bound

To deepen our understanding of the network's spatial stability against geographic partitions, we analyze vortex dynamics via the **Kosterlitz-Thouless (KT)** formalism. It is necessary here to resolve an apparent topological paradox regarding the Mermin-Wagner Theorem [19], which states that spontaneous breaking of a continuous symmetry is impossible at finite temperature for systems of dimension $d \leq 2$.

In a graph, effective dimensionality d_{eff} can be extracted from the scaling law linking the number of accessible nodes $\mathcal{N}(r)$ within a topological radius r (number of hops):

$$\mathcal{N}(r) \propto r^{d_{eff}} \tag{42}$$

In a healthy state, the P2P graph's effective dimensionality approaches infinity ($d_{eff} \to \infty$) due to global intercontinental shortcuts. However, the relevance of KT physics emerges during a major structural failure. When a severe network partition occurs (e.g., the isolation of a continental grid), the routing topology severely degrades. The severing of transoceanic fiber links suppresses the network's global shortcuts, forcing the propagation layer to map much more closely to the physical 2D surface of the Earth.

It is crucial to note that even in this partitioned state, the severed network remains technically scale-free at the local level; it does not geometrically crystalize into a perfect planar lattice. However, the loss of global shortcuts abruptly collapses the network's spectral dimension toward $d_{eff} \approx 2$, imposing a strict quasi-planar constraint on information diffusion. Therefore, we deploy the Kosterlitz-Thouless formalism not as an exact geometric mapping, but as a rigorous theoretical *asymptotic lower bound* on the thermal vulnerability of an isolated partition. By treating the partition as a 2D spin system, we extract the absolute worst-case thermodynamic limits before the consensus field experiences topological collapse.

In this critically restricted regime, the energy of an isolated fork grows logarithmically with network size L:

$$E_{vortex} \approx \pi J \ln\left(\frac{L}{a_0}\right) \tag{43}$$

Simultaneously, the vortex core can be located on any of the $(L/a_0)^2$ sites of the two-dimensional lattice. The system thus acquires a configurational entropy:

$$S_{vortex} \approx k_B \ln\left[\left(\frac{L}{a_0}\right)^2\right] = 2k_B \ln\left(\frac{L}{a_0}\right) \tag{44}$$

Consensus stability against free vortex formation is governed by the variation in free energy $\Delta\mathcal{F} = E_{vortex} - T_{eff}S_{vortex}$:

$$\Delta\mathcal{F} = (\pi J - 2k_B T_{eff}) \ln\left(\frac{L}{a_0}\right) \tag{45}$$

This balance equation reveals the existence of a strict critical partition temperature $T_{KT} = \frac{\pi J}{2k_B}$. Survival of the canonical ledger truth then boils down to a topological phase transition:

- **Ordered Phase ($T_{eff} < T_{KT}$):** $\Delta\mathcal{F} > 0$. Creating a free vortex requires infinite energy in the thermodynamic limit ($L \to \infty$). Phase fluctuations are confined; vortices and anti-vortices appear in bound pairs of finite energy. Long-range order and global consensus are maintained.

- **Partition Phase ($T_{eff} > T_{KT}$):** $\Delta\mathcal{F} < 0$. Configurational entropy outweighs binding energy J. The proliferation of isolated vortices becomes thermodynamically favorable and spontaneous. Uncertainty over history curls in on itself (Figure 8), destroying global phase coherence and irreparably fracturing the ledger.

The "Small-World" topology is therefore the indispensable structural mechanism that perpetually maintains the network in the $T_{eff} \ll T_{KT}$ regime, thereby dodging the topological collapse dictated by two-dimensional physics.



Figure 8. **Topological Defect in the P2P Network.** Each arrow represents the "consensus spin" of a node. A free vortex ($n = 1$) breaks long-range order, rendering global consensus physically impossible.

VII. NON-EQUILIBRIUM DYNAMICS: EXERGETIC EXPANSION AND THE "HALVING" QUENCH

While Section VI deals with spatial stability, the network must also survive severe temporal shocks. The Nakamoto chain functions as a dissipative structure maintained far from equilibrium. However, the network is not a closed thermodynamic system: it is coupled to global energy infrastructure, continuously hunting **exergy** (available, unvalued energy, such as stranded energy or gas flaring).

The "Halving" event is therefore not a simple parametric update leading to a lower energy equilibrium; it is a thermodynamic shock applied to an expanding system. We model this event as a **Thermodynamic Quench** superimposed on a non-stationary potential.

A. The Exergy-Driven Potential

The effective potential $V(\phi)$ is driven by the total **exergetic chemical potential** $\mu(t)$. Rather than treating this as a financial profitability metric, we define $\mu(t)$ strictly as the physical influx rate of unvalued thermodynamic work available to the network (e.g., global stranded energy, grid curtailment variance).

Unlike static models, this potential features a secular growth component due to the continuous integration of new global exergy sources and infrastructure optimization. We decompose $\mu(t)$ as follows:

$$\mu(t) = \mu_{subsidy}(t) + \mu_{exergy}(t) \tag{46}$$

The subsidy component undergoes a strict discontinuity (Heaviside step function Θ) at the exact moment of the Halving (t_H):

$$\mu_{subsidy}(t) = \mu_0 \left[1 - \frac{1}{2}\Theta(t - t_H)\right] \tag{47}$$

However, the expansion of $\mu_{exergy}(t)$ cannot diverge indefinitely. To avoid the physical paradox of infinite energy growth violating planetary boundaries, this function must be constrained by the total accessible exergy of the biosphere μ_{max} (reservoir carrying capacity). The secular expansion dynamics of exergy consequently obey a differential logistic equation:

$$\frac{d\mu_{exergy}}{dt} = r\mu_{exergy}\left(1 - \frac{\mu_{exergy}}{\mu_{max}}\right) \tag{48}$$

where r represents the technological innovation rate (the velocity at which the network identifies and integrates new sources of stranded energy).

Thus, the theoretical equilibrium order parameter $\phi_{eq}(t)$ does not permanently collapse after a Halving at t_H; it undergoes a transient perturbation before resuming its trajectory along this logistic sigmoid, dictated by the volume of mineable exergy remaining globally.

To empirically validate this exergetic expansion, we analyze the macroscopic evolution of the network's order parameter (Hashrate) over its history. By excluding the initial pre-industrial transient phase (prior to January 2010), the network operates far from the logistic saturation limit ($\mu_{exergy} \ll \mu_{max}$), manifesting a regime of super-linear algebraic growth. As illustrated in Figure 9, the expansion follows a strict temporal power law:

$$H(t) = C \cdot t^{\alpha} \tag{49}$$

where t is the time elapsed since genesis (in days). Regression analysis in the log-log phase space yields a massive scaling exponent $\alpha \approx 11.59$ and a proportionality constant $C \approx 6.7 \times 10^{-35}$ TH/s.

Despite massive exogenous shocks—marked by the vertical Halving quenches—the network's exergy capture binds to this invariant algebraic trajectory with an exceptionally high coefficient of determination ($R^2 = 0.972$). This secular expansion provides compelling empirical evidence of the **Maximum Power Principle** formulated by Lotka [11] and later expanded by Odum [12] acting at the macroeconomic scale.

The Bitcoin network does not merely passively dissipate energy; as an evolving physical system, it self-organizes structurally (technological upgrades of ASIC fleets, geographic migration toward isolated stranded energy sources) to maximize its exergy capture rate and its dissipation in the form of hashing. In this natural selection framework, operators who maximize power flux win. The network thus continuously converts raw electrical exergy into *emergy* (condensed structural memory), thermodynamically cementing consensus robustness against perturbations.



Figure 9. **Secular Exergetic Expansion.** The network's Hashrate $H(t)$ plotted on a log-log scale against days since genesis. Starting from the industrialization of mining (Jan 2010), the system aligns with a strict power-law attractor $H(t) \propto t^{11.59}$ with an exceptional coefficient of determination ($R^2 = 0.972$), despite periodic thermodynamic quenches (Halvings).

B. Non-Equilibrium Stochastic Dynamics and FDT Violation

Relaxation dynamics of the order parameter $\phi(t)$ map to a Langevin equation, describing the network's stochastic response to the driven potential:

$$\frac{\partial \phi}{\partial t} = -\Lambda_0 \frac{\delta \mathcal{F}}{\delta \phi} + \zeta(t) \tag{50}$$

However, since the network operates strictly far from thermodynamic equilibrium (continuously hunting exergy), the standard Fluctuation-Dissipation Theorem (FDT) governing the classic Hohenberg-Halperin equation is violated. Detailed balance is broken by continuous entropy production $\dot{\Sigma} > 0$.

To formally establish the Non-Equilibrium Steady State (NESS) of the network, we must analyze its dynamic response. By defining the two-time correlation function $C(t, t') = \langle \delta\phi(t)\delta\phi(t')\rangle$ and the response function $R(t, t') = \left.\frac{\delta\langle\phi(t)\rangle}{\delta h(t')}\right|_{h=0}$, we quantify this thermodynamic deviation by introducing the Fluctuation-Dissipation Ratio (FDR) $X(t, t')$:

$$R(t, t') = -\frac{X(t, t')}{k_B T_{bath}} \frac{\partial C(t, t')}{\partial t} \Theta(t - t') \tag{51}$$

where Θ is the Heaviside step function. In this formalism, T_{bath} represents the **macroeconomic exergy bath temperature**. Unlike a closed physical system, the network is immersed in an open planetary environment characterized by severe stochastic fluctuations in available stranded energy.

We define the thermal energy of this macroscopic bath as strictly proportional to the standard deviation of the network's exergetic chemical potential $\mu(t)$, evaluated over a characteristic macroscopic window τ_{macro}:

$$k_B T_{bath} \equiv \gamma_m \sqrt{\langle (\mu(t) - \langle \mu \rangle)^2 \rangle_{\tau_{macro}}} \tag{52}$$

where γ_m is a dimensionless coupling constant representing the network's physical capacity to capture stranded energy, and τ_{macro} corresponds to the network's difficulty adjustment epoch (2016 blocks).

This definition grounds the macroscopic bath temperature not in abstract financial volatility, but in the strict physical variance of the global energy grid. Miners operate at the thermodynamic margins; their continuous search for profitability is strictly bounded by the stochastic availability of stranded energy, grid curtailment events, and local thermal capacities. Therefore, the temporal variance of $\mu(t)$ acts as the true kinetic driver of the background reservoir.

Crucially, this external background heat T_{bath} is what physically finances the mechanical work required to purge spatial Shannon entropy at the UTXO boundary; a collapse in T_{bath} directly starves the network of the exergy needed to maintain the effective information temperature T_{eff} below the critical threshold T_c.

In this regime dominated by an **active noise** $\zeta(t)$ integrating macroeconomic shocks and mining competition, the time-translation invariance of "slow mode" relaxation (the network's macroscopic inertia) is broken. Parameter X becomes constant but strictly less than 1 ($X < 1$).

This allows us to rigorously define an **effective activity temperature** T_{act}, conjugate to these non-equilibrium degrees of freedom:

$$T_{act} = \frac{T_{bath}}{X} \tag{53}$$

This activity temperature can be explicitly related to active noise intensity arising from the inherent variance of the Poisson process (block discovery). Setting the non-equilibrium diffusion coefficient D_{act}, the relationship between our FDR and network mechanical agitation is written:

$$X = \left(1 + \frac{D_{act}}{k_B \Lambda_0 T_{bath}}\right)^{-1} \tag{54}$$

Thus, by substitution, we recover the additive expression of background noise variance $\langle \zeta(t)\zeta(t') \rangle = 2\Lambda_0 k_B T_{act} \delta(t-t')$ with $T_{act} = T_{bath} + \frac{D_{act}}{k_B \Lambda_0}$. This unified formalization demonstrates that the network's temperature elevation is not purely thermal in origin: it directly reflects stochastic agitation and the mechanical work expended to maintain the Markov blanket (UTXO) against constant entropic degradation.

It should be clarified here that using a Gaussian white noise variance ($\delta(t-t')$) constitutes a **large-scale mean-field approximation**. At the micro-canonical scale, active network noise is structurally asymmetric: it superimposes the strict Poisson process of block discovery onto available exergy shocks (T_{bath}) which frequently exhibit "fat tails" characteristic of macroeconomic dynamics. However, the Gaussian approximation holds in the hydrodynamic limit. By invoking the principle of timescale separation, macroscopic reorganization dynamics occur at $t \gg \tau_B$. In this coarse-grained regime, the generalized Central Limit Theorem ensures that the cumulative effect of these heavy-tailed exergy shocks renormalizes into an effective Gaussian thermal bath, validating the use of the standard Langevin formalism for secular stability analysis.

Finally, it must be noted that this Langevin formalism models the system as a reactive physical thermostat. In reality, the hardware operators (miners) are economic agents possessing forward-looking anticipation. Markets price in the deterministic Halving quench, leading to CapEx deployments *prior* to the event. To fully capture this human economic foresight, the active noise term $\zeta(t)$ would theoretically require a non-Markovian memory kernel to integrate future expected values of $\mu_{subsidy}$. However, the purely reactive thermodynamic model remains highly predictive of the transient phase divergence, as demonstrated empirically in Section VII C.

C. Thermodynamic Halving Quench and Thermostat (DAA)

It is upon this tensioned system, maintained far from equilibrium, that the Halving imposes an **abrupt phase transition**. The instantaneous halving of $\mu_{subsidy}$ renders the current state $\phi(t_H)$ metastable for operators with the lowest thermodynamic efficiency (high thermal entropy production via hashing).

However, the physical system does not adjust through instantaneous collapse: since Capital Expenditures (CapEx) are sunk costs, the network possesses **thermodynamic inertia** (hysteresis). Miners only unplug if their marginal cost (OpEx) exceeds new profitability. The Halving thus acts as a Darwinian dissipative filter: it purges the network of its least efficient components and forces hardware migration toward denser exergy pockets.

The thermodynamic danger of this quench lies in the transient regime. The initial hashrate drop against an uncompromising Difficulty target D mechanically causes a divergence in the mean block wait time $\langle \Delta t \rangle$. The system escapes the entropic "death spiral" via the Difficulty Adjustment Algorithm (DAA).

Operating as a **non-equilibrium thermostat** with discrete feedback [20], the DAA recalibrates the system's

"mass". In control theory (cybernetics) formalism, this thermostat is not Markovian (instantaneous): it acts as an **integral controller with phase lag**, its evaluation being smoothed over a rolling window of 2016 blocks. It is precisely this integration hysteresis that physically explains why the system, faced with a thermodynamic quench, does not relax in a purely monotonic fashion. It undergoes damped transient oscillations (the empirically observed stochastic dip) before rejoining its growth attractor. By belatedly lowering the energy barrier required to crystallize a block, the DAA ultimately restores relaxation dynamics toward the new metastable equilibrium.



Figure 10. **Quench on Non-Stationary Background.** Order parameter ϕ is sustained by secular growth (exergy hunting). The Halving imposes an abrupt perturbation. Miner inertia and thermostat intervention (DAA) create a transient plateau or dip (damped relaxation) before the bullish trend resumes.

The network then undergoes a transient relaxation (often a plateau or stochastic dip) before exergy expansion $\mu_{exergy}(t)$ regains dominance, propelling hashrate to new highs (see Figure 10). This asymmetry acts as a true natural experiment, allowing us to observe the correlation between non-equilibrium state duration and thermal variance divergence.

Validation: Multi-Halving Relaxation Dynamics

To confront this thermodynamic quench modeling with empirical data, we analyzed the time series of the Poisson process (inter-block wait time Δt) for Bitcoin's four historical Halvings (2012, 2016, 2020, 2024). Protocol architecture offers an exceptional *in vivo* observation opportunity here. The period between two Halvings (210,000 blocks) is not an exact multiple of the DAA adjustment epoch (2,016 blocks); thus, the duration of the "Quench Phase"—the number of blocks where the network absorbs the thermodynamic shock before thermostat intervention—varies deterministically at each iteration.

As visually demonstrated in Figure 11 and statistically quantified in Table II, the results reveal a remarkable typology of thermodynamic responses:



Figure 11. **Empirical Relaxation Dynamics of the Thermodynamic Quench.** The moving average of inter-block time $\langle\Delta t\rangle$ (solid blue line) and the active thermal noise variance $\pm 1\sigma$ (shaded region), both calculated using a backward-looking rolling window of 144 blocks ($\approx$ 24 hours), are plotted relative to the Halving event (dashed red line). The transient non-equilibrium phase persists until the DAA thermostat intervenes (dotted green line) to restore the target interval (solid black line). Notice the massive expansion of the shaded variance region in 2020, providing macroscopic visual confirmation of the fluctuation-dissipation theorem violation during an exergy shock.

Halving	Block	Quench Duration	$\Delta\sigma^2(\Delta t)$ (Active Noise)	Fisher Test (p)	Levene Test (p)
2012	210,000	1680 blocks	+10.54%	0.0158 (Sig.)	0.0514
2016	420,000	1344 blocks	−0.09%	0.5060	0.3693
2020	630,000	1008 blocks	+54.77%	< 0.0001 (Sig.)	< 0.0001 (Sig.)
2024	840,000	672 blocks	+11.24%	0.0432 (Sig.)	0.1404

Table II. **Thermodynamic Quench Metrology.** Comparison of block time variance between the equilibrium epoch (the 2016 blocks preceding the Halving) and the quench phase (from the Halving until DAA adjustment).

1. **Pure Divergence (2020):** The 2020 event is a textbook case for our Langevin equation model. With 1008 quench blocks, purging the inefficient translated into an explosion of nearly 55% in active noise $\zeta(t)$, confirmed by both statistical tests with absolute significance ($p < 0.0001$). As seen in Figure 11, the shaded variance region inflates massively immediately after the shock. This is the indisputable macroscopic signature of Fluctuation-Dissipation Theorem violation facing an exergy shock.

2. **Frustrated Divergence (2024 and 2012):** Both the 2012 and 2024 Halvings show a consistent active noise elevation of around 11%, validated by variance ratio (Fisher Test). However, the extremely rapid DAA intervention in 2024 (only 672 blocks, or one-third of an epoch) truncated system relaxation, preventing it from accumulating a sufficient statistical sample to reach significance on skewed

distributions (Levene Test, $p = 0.14$). Far from invalidating the model, this "non-significance" empirically illustrates the damping power of the protocol's discrete feedback loop.

3. **Exergetic Continuity (2016):** The 2016 case, with unchanged variance, proves the central role of macroeconomic bath temperature (T_{bath}). The parametric collapse of subsidy was instantly offset by lightning-fast integration of new-generation ASICs coupled with aggressive bullish appreciation. Chemical potential μ_{exergy} dominated, averting infrastructure deactivation and holding the network in metastable equilibrium.

This multi-halving metrology demonstrates that the ledger does not respond to shock merely by experiencing a "production drop", but indeed by transiting through a transient non-equilibrium regime where the network's entropic heat visually diverges, dictating the urgency of its difficulty thermostat's intervention.

VIII. CONCLUSION: THE PHYSICS OF TRUTH

In this article, we proposed a minimal physical model for Nakamoto consensus, demonstrating that the blockchain functions as a **dissipative structure** maintained far from thermodynamic equilibrium. By mapping the ledger onto a one-dimensional network system, we showed that the "consensus mechanism" is formally equivalent to a **continuous phase transition** where the order parameter (hashrate) spontaneously breaks the network's local time-inversion symmetry.

A. Model Summary

Our analysis yields three key physical perspectives:

1. **Thermodynamic Depth:** Ledger "immutability" is not absolute but probabilistic, governed by a correlation length ξ that decays exponentially with cumulative work. This resolves the "Double Spending" problem as the suppression of thermal fluctuations.

2. **Crystallization and Spatial Limit:** Mining is a phase boundary transitioning information from a "liquid" to a "crystalline" state. This stability is conditioned by a dimensionless ratio Γ that imposes a strict limit on block size to avoid thermal percolation.

3. **Dynamic Stability:** Network resilience to macroeconomic shocks ("Halving") is explained by first-order quench dynamics, effectively damped by a protocol thermostat (DAA) that restores restoring force toward the new equilibrium.

B. Thermodynamic Inertia and Effective Mass of Information

Our derivation provides a formal framework to interpret the Mass-Energy-Information (MEI) equivalence principle proposed by Vopson [7] at the macroscopic scale. Although the proper physical mass of an isolated bit of information remains a debated theoretical conjecture ($m_{bit} \approx k_B T \ln 2/c^2$), our model demonstrates that the Bitcoin ledger acquires an indisputable **Effective Mass** (inertial) in the dynamical systems sense, generated by the dissipative amplification factor of Proof of Work.

By formal analogy, we define this history inertia M_{eff} from the total thermodynamic work dissipated by the network. Since the instantaneous power is $P_{net}(t) = E_{eff}(t)/\tau_B$, the global mass is:

$$M_{eff} \propto \frac{E_{total}}{c^2} = \frac{1}{c^2}\int P_{net}(t)\,dt = \frac{1}{c^2\tau_B}\int_{\mathcal{C}} E_{eff}(t)\,dt \tag{55}$$

It is, however, imperative to introduce here a formal distinction between the **Global Mass** of history and the **Marginal Mass** of consensus. The previous equation defines global mass $M_{global} \equiv M_{eff}$, which corresponds to the cumulative potential well (the ledger's total inertia since the genesis block).

However, network resistance to a *local* perturbation (a reorganization attempt at the chain tip) is not dictated by this integral, but by local curvature of the Landau potential at the bottom of the well. By analogy with massive field theory, we define Marginal Mass $m_{marginal}$ associated with equilibrium order parameter ϕ_0:

$$m^2_{marginal} \propto \left.\frac{\partial^2 V}{\partial \phi^2}\right|_{\phi_0} = -4a(T_{eff}) \tag{56}$$

Since instability parameter $a(T_{eff}) < 0$ in the ordered phase (Section III A), this mass is strictly real and positive. It is this marginal mass, dynamically adjusted by protocol difficulty, that governs correlation length scale $\xi \sim m^{-1}_{marginal}$ (Section IV B).

Thus, while Global Mass anchors the blockchain in irreversible thermodynamic history, Marginal Mass protects the active interface against thermal evaporation. It is crucial to understand this resultant mass not as a gravitational entity, but as a **strict thermodynamic resistance to temporal reversibility**.

In light of Landauer's Principle [4], Proof of Work is not a mere arbitrary mathematical computation; it is a massive process of **uncertainty erasure**. Every second, the network explores, generates, and destroys hundreds of exahashes of invalid states. The blockchain's immense Effective Mass M_{eff} is therefore none other than the "thermal corpse" of these billions of billions of erased cryptographic failures. It is precisely the irreversibility of this continuous erasure that endows the canonical truth with its inertia.

Instead of curving the spacetime of general relativity, this uninterrupted accumulation of work sculpts a potential well so deep in the network's free energy landscape $\mathcal{F}$ (as modeled in Section III A) that thermodynamic "escape velocity"—that is, the instantaneous macroscopic work required to breach the $\Delta\mathcal{F}$ barrier and rewrite history—exceeds by several orders of magnitude the energetic resources of any individual actor. Nakamoto consensus is thus no longer a mere software agreement; it is a physical attractor of colossal inertia.

C. Epilogue: Vires in Numeris, Veritas in Energia

If we postulate a maximum scale limit where a Type I planetary civilization devoted its entire solar irradiance energy budget ($P_{max} \approx 1.7 \times 10^{17}$ W) to maintaining consensus with perfect computing efficiency at Landauer's limit ($\eta_{Landauer} \approx 2.8 \times 10^{-16}$ J/hash at 300 K), hashrate would cap at $|\phi_{max}| \approx 6.0 \times 10^{32}$ s^{-1}. The asymptotic floor of Nakamoto Action would then settle at $\kappa_{N,limit} \approx 4.6 \times 10^{-49}$ J $\cdot$ s.

However, purely energetic modeling ignores an inescapable biophysical constraint: information requires a material substrate. To orderly dissipate this energy and generate a hash, the network requires colossal silicon, copper, and semiconductor infrastructure. Defining a computational Mass Efficiency μ_m, representing the physical mass of hardware required to support a given hash rate (currently $\mu_m \approx 1.5 \times 10^{-13}$ kg/(H/s)), we encounter a hard limit. Even assuming spectacular lithographic breakthroughs, reaching this Landauer regime would require a computational substrate mass $M_{hardware} = \mu_m \cdot |\phi_{max}|$ cheerfully exceeding 10^{19} kg. The network will hit the thermodynamic wall of metallurgy long before reaching the thermodynamic limit of planetary insolation.

The protocol thus transcends simple software engineering to become an unprecedented convergence of hard sciences. The cryptographic adage "*Vires in Numeris*" (Strength in Numbers) finds here its ultimate physical corollary: "*Veritas in Energia*" (Truth in Energy). Anchoring this ledger in the unalterable scarcity of energy and matter marks a profound paradigm shift. By moving beyond pure game theory and formalizing consensus as a non-equilibrium thermodynamic phase transition, this framework yields strict, falsifiable physical limits for decentralized networks. Specifically, it dictates an absolute thermodynamic upper bound on block propagation volume ($V_{B,crit}$) to avert topological collapse, and predicts structural variance divergences during sudden exergetic quenches.

Furthermore, by demonstrating how a decentralized network extracts a singular chronological reality from stochastic noise, this model extends far beyond digital ledgers. If thermal time and continuous energy dissipation are the universal arbiters of shared history, this thermodynamic framework offers a novel physical lens for complex biological systems. The synchronization of neural populations, the collective active inference across cellular Markov blankets, and the emergence of macroscopic order in decentralized life may fundamentally rely on these same mechanics of continuous phase transitions and information crystallization.

Ultimately, for our scope, the immutability of a sufficiently capitalized blockchain is guaranteed not solely by algorithmic decree, but by the unshakable bedrock of the laws of physics—providing the thermodynamic anchor necessary to finally realize John Nash's vision of "Ideal Money" [21]. The objective truth of ownership in a decentralized network cannot be sustained without the constant dissipation of work. It is a powerful, yet falsifiable statement that anchors the present paper.

If we consider the continuous hunting of exergy to maintain a low-entropy state, the Bitcoin network behaves remarkably like a planetary-scale dissipative organism. Its relentless drive to order information through energy consumption provides a macroscopic laboratory for studying how decentralized architectures can resist entropic decay over secular timescales.

ACKNOWLEDGMENTS

The author wishes to thank the open-source community, the founding cypherpunks, and independent researchers exploring the intersection of physics and decentralized systems. **The Python scripts used to generate the empirical plots and the raw data presented in Table II are available in a permanent repository at:** `https://github.com/pascalranaora/NakamotoRanaoraPRE2026`.

Appendix A: Nomenclature and Dimensional Analysis

Symbol	Physical Quantity / Description	Dimension	SI Unit
	Temporal and Kinematic Variables		
t	External physical time (Substrate)	Time [T]	s
τ, N	Internal thermal time (Block height)	Dimensionless [1]	blocks
τ_B	Target crystallization interval	Time [T]	s
τ_L	Network spatial propagation latency	Time [T]	s
τ_0	Incompressible propagation latency (speed of light limit)	Time [T]	s
τ_{cycle}	Microscopic ASIC logic gate switching time	Time [T]	s
$\|\phi(t)\|$	Global network Hashrate	Frequency [T^{-1}]	s^{-1} (Hash/s)
r	Technological innovation rate (logistic growth)	Frequency [T^{-1}]	s^{-1}
	Thermodynamic and Energetic Variables		
$\eta_{hash}(t)$	Hardware thermodynamic efficiency	Energy [ML^2T^{-2}]	J/Hash
$P_{net}(t)$	Total instantaneous network power	Power [ML^2T^{-3}]	W (J/s)
E_{eff}	Macroscopic block state energy	Energy [ML^2T^{-2}]	J
T_{eff}	Effective information temperature	Energy / k_B [Θ]	K
T_c	Critical transition information temperature	Energy / k_B [Θ]	K
T_{bath}	Macroeconomic exergy bath temperature	Energy / k_B [Θ]	K
β	Inverse effective temperature	Energy^{-1} [$M^{-1}L^{-2}T^2$]	J^{-1}
$\mathcal{H}_{eff}$	Effective Consensus Hamiltonian	Energy [ML^2T^{-2}]	J
J	Topo-thermodynamic coupling constant (XY model)	Energy [ML^2T^{-2}]	J
μ_{exergy}	Macroscopic chemical potential (Exergy)	Energy [ML^2T^{-2}]	J
X	Fluctuation-Dissipation Ratio (FDR)	Dimensionless [1]	–
Λ_0	Kinetic relaxation coefficient (Langevin)	Mobility [$M^{-1}T$]	$s \cdot kg^{-1}$
χ_E	Exergetic heat capacity	Energy / Temp [$ML^2T^{-2}\Theta^{-1}$]	$J \cdot K^{-1}$
$\mathcal{E}_{coupling}$	Topological coupling energy	Energy [ML^2T^{-2}]	J
a	Landau instability parameter	Energy [ML^2T^{-2}]	J
b	Saturation energy barrier	Energy [ML^2T^{-2}]	J
	Action, Mass, and Information Variables		
$\kappa_N(t)$	Nakamoto Action (Micro quantum)	Action [ML^2T^{-1}]	$J \cdot s$
$\mathcal{S}_{PoW}$	Consensus Action (Macro integral)	Action [ML^2T^{-1}]	$J \cdot s$
D	Protocol Difficulty (Thermostat Setpoint)	Dimensionless [1]	–
$\mathcal{N}_H(D)$	Expected hashes to satisfy PoW	Information [1]	Hashes
M_{eff}	Effective Global Mass (History Inertia)	Mass [M]	kg
m	Mass gap (Inverse correlation length)	Distance^{-1} (Graph) [1]	blocks^{-1}
μ_m	Computational mass efficiency	Mass $\cdot$ Time	$kg \cdot s$
$M_{hardware}$	Physical substrate mass of the network	Mass [M]	kg
Γ	Thermodynamic/Latency ratio (τ_B/τ_L)	Dimensionless [1]	–
Γ_c	Critical thermodynamic/Latency ratio	Dimensionless [1]	–
V_B	Block data volume	Information [1]	Bytes
C	Shannon channel capacity	Info Flux [T^{-1}]	Bytes/s
ϕ, Ψ	Consensus Complex Order Parameter / Magnetization	Dimensionless [1]	–
ξ	Correlation length of consensus	Distance (Graph) [1]	blocks
a_0	Lattice constant (Minimum topological distance)	Dimensionless [1] (Graph)	hops
L	Characteristic network spatial diameter	Dimensionless [1] (Graph)	hops

Table III. **Variable Dimensionality.** Summary of the principal physical variables defining the Nakamoto non-equilibrium state, proving strict dimensional consistency across energy, time, and action.

[1] S. Nakamoto, Bitcoin: A peer-to-peer electronic cash system (2008).

[2] N. Szabo, Shelling out: The origins of money (2002).

[3] C. H. Bennett, The thermodynamics of computation—a review, International Journal of Theoretical Physics **21** (1982).

[4] R. Landauer, Irreversibility and heat generation in the computing process, IBM Journal of Research and Development **5** (1961).

[5] I. Prigogine, *The End of Certainty: Time, Chaos, and the New Laws of Nature* (The Free Press, 1997).

[6] C. E. Shannon, A mathematical theory of communication, Bell System Technical Journal **27** (1948).

[7] M. M. Vopson, The mass-energy-information equivalence principle, AIP Advances **9**, 095206 (2019).

[8] P. Ranaora, Thermodynamic stability and phase transitions in the nakamoto consensus, IPI Letters **4**, 1 (2026).

[9] D. J. Watts and S. H. Strogatz, Collective dynamics of 'small-world' networks, Nature **393**, 440 (1998).

[10] V. L. Ginzburg and L. D. Landau, On the theory of superconductivity, Zh. Eksp. Teor. Fiz. **20**, 1064 (1950).

[11] A. J. Lotka, Contribution to the energetics of evolution, Proceedings of the National Academy of Sciences **8**, 147 (1922).

[12] H. T. Odum and R. C. Pinkerton, Time's speed regulator: the optimum efficiency for maximum power output in physical and biological systems, American Scientist **43**, 331 (1955).

[13] U. Seifert, Stochastic thermodynamics, fluctuation theorems and molecular machines, Reports on Progress in Physics **75** (2012).

[14] A. Connes and C. Rovelli, Von neumann algebra automorphisms and time-thermodynamics relation in generally covariant quantum theories, Classical and Quantum Gravity **11**, 2899 (1994).

[15] G. E. Crooks, Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences, Physical Review E **60**, 2721 (1999).

[16] C. Decker and R. Wattenhofer, Information propagation in the bitcoin network, in *IEEE P2P 2013 Proceedings* (2013).

[17] J. Pearl, *Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference* (Morgan Kaufmann, 1988).

[18] C. Jarzynski, Nonequilibrium equality for free energy differences, Physical Review Letters **78**, 2690 (1997).

[19] N. D. Mermin and H. Wagner, Absence of ferromagnetism or antiferromagnetism in one- or two-dimensional isotropic heisenberg models, Physical Review Letters **17**, 1133 (1966).

[20] W. G. Hoover, Canonical dynamics: Equilibrium phase-space distributions, Physical Review A **31**, 1695 (1985).

[21] J. F. Nash, Ideal money, Southern Economic Journal **69**, 4 (2002).