Octonion Quantum Chromodynamics

B. C. Chanyal
(1), P. S. Bisht⁽¹⁾, Tianjun Li⁽²⁾ and O. P. S. Negi
(1.2)* April 3, 2012

(1) Department of Physics Kumaun University S. S. J. Campus Almora - 263601 (U.K.), India.

 (2) Institute of Theoretical Physics Chinese Academy of Sciences
 Zhong Guan Cun East Street 55
 P. O. Box 2735
 Beijing 100190, P. R. China.

Email- bcchanyal@gmail.com ps_bisht 123@rediffmail.com tli@itp.ac.cn ops_negi@yahoo.co.in

Abstract

Starting with the usual definitions of octonions, an attempt has been made to establish the relations between octonion basis elements and Gell-Mann λ matrices of SU(3) symmetry on comparing the multiplication tables for Gell-Mann λ matrices of SU(3) symmetry and octonion basis elements. Consequently, the quantum chromo dynamics (QCD) has been reformulated and it is shown that the theory of strong interactions could be explained better in terms of non-associative octonion algebra. Further, the octonion automorphism group SU(3) has been suitably handled with split basis of octonion algebra showing that the $SU(3)_C$ gauge theory of colored quarks carries two real gauge fields which are responsible for the existence of two gauge potentials respectively associated with electric charge and magnetic monopole and supports well the idea that the colored quarks are dyons.

Key Words: Octonions, Quantum Chromodynamics (QCD), SU(3) symmetry, λ matrices.

PACS No.: 12.10 Dm, 12.60.-i, 14.80 Hv.

^{*}Address from Feb. 21- April 20, 2012:- Institute of Theoretical Physics, Chinese Academy of Sciences, Zhong Guan Cun East Street 55, P. O. Box 2735, Beijing 100190, P. R. China.

1 Introduction

In spite of the symmetry, conservation laws and gauge fields describe elementary particle in terms of their field quanta and interactions. Nevertheless, the role of number system (hyper complex numbers) has been an important factor in understanding the various theories of physics from macroscopic to microscopic level. In fact, there has been a revival in the formulation of natural laws in terms of numbers. So, according to celebrated Hurwitz theorem there exists [1] four division algebra consisting of \mathbb{R} (real numbers), \mathbb{C} (complex numbers), \mathbb{H} (quaternions) [2, 3] and \mathbb{O} (octonions) [4, 5, 6]. All these four algebra's are alternative with totally anti symmetric associators. Real and complex numbers are limited only up to two dimensions, quaternions are extended to four dimensions (one real and three imaginaries) while octonions represent eight dimensions (one real and seven imaginaries). Real and complex numbers are commutative and associative. Quaternion are associative but not commutative while its next generalization to octonions is neither commutative nor associative. Rather, the laws of alternatively and distributivity are obeyed by octonions. Quaternions and octonions are extensively used in the various branches of physics and mathematics. The octonion analysis has also played an important role in the context of various physical problems [7, 8, 9, 10, 11, 12, 13, 14] of higher dimensional supersymmetry, super gravity and super strings etc while the quaternions have an important role to unify [15, 16, 17] electromagnetism and weak forces to represent the electroweak $SU(2) \times U(1)$ sector of standard model. Likewise, the octonions are extensively studied [18, 19] for the description of color quarks and played an important role for unification programme of fundamental interactions in terms of successful gauge theories. Furthermore, the quaternionic formulation of Yang-Mill's field equations and octonion reformulation of quantum chromo dynamics (QCD) has also been developed [20] by taking magnetic monopoles [21, 22, 23] and dyons (particles carrying electric and magnetic charges) [24, 25, 26, 27] into account. It is shown that the three quaternion units explain the structure of Yang-Mill's field while the seven octonion units provide the consistent structure of $SU(3)_C$ gauge symmetry of quantum chromo dynamics. Keeping in view the potential importance of monopoles and dyons and their possible role in quark confinement, in this paper, we have made an attempt to construct SU(3) gauge theory suitably handled with octonions for colored quarks. So, starting from the usual definitions of octonions, we have established the suitable connection between octonion basis elements and SU(3) symmetry after comparing the multiplication tables for Gell-Mann λ matrices of SU(3) symmetry and octonion basis elements. Consequently, the quantum chromo dynamics (QCD) has been reformulated and it is shown that the theory of strong interactions could be explained better in terms of non-associative octonion algebra. Further, the octonion automorphism group SU(3) has been reconnected to the split basis of octonion algebra and it is shown that the $SU(3)_C$ gauge theory of colored quarks describes two real gauge fields identified as the gauge strengths of two types of chromo charges showing the presence of electric charge and and magnetic monopoles. So, it is concluded that the present formalism of colored quarks suitably describes the existence of dyons:particles carry the simultaneous existence of electric charge and magnetic monopoles.

2 Octonion Definition

An octonion x is expressed [28, 29, 30] as

$$x = (x_0, x_1, ..., x_7) = x_0 e_0 + \sum_{A=1}^{7} x_A e_A \qquad (A = 1, 2, ..., 7)$$
 (1)

where $e_A(A = 1, 2,, 7)$ are imaginary octonion units and e_0 is the multiplicative unit element. The octet $(e_0, e_1, e_2, e_3, e_4, e_5, e_6, e_7)$ is known as the octonion basis and its elements satisfy the following multiplication rules

$$e_0 = 1$$
, $e_0 e_A = e_A e_0 = e_A$, $e_A e_B = -\delta_{AB} e_0 + f^{ABC} e_C$. $(\forall A, B, C = 1, 2, ..., 7)$ (2)

The structure constants f^{ABC} are completely antisymmetric and take the value 1. i.e. $f^{ABC} = +1\{\forall (ABC) = (123), (471), (257), (165), (624), (543), (736)\}$. Here the octonion algebra \mathcal{O} is described over the algebra of real numbers having the vector space of dimension 8. Octonion algebra [14] is non associative and multiplication rules for its basis elements given by equations (2) are then generalized in the following table as [28, 29, 30];

•	e_1	e_2	e_3	e_4	e_5	e_6	e_7
e_1	-1	e_3	$-e_2$	e_7	$-e_6$	e_5	$-e_4$
e_2	$-e_3$	-1	e_1	e_6	e_7	$-e_4$	$-e_5$
e_3	e_2	$-e_1$	-1	$-e_5$	e_4	e_7	$-e_6$
e_4	$-e_7$	$-e_6$	e_5	-1	$-e_3$	e_2	e_1
e_5	e_6	$-e_7$	$-e_4$	e_3	-1	$-e_1$	e_2
e_6	$-e_5$	e_4	$-e_7$	$-e_2$	e_1	-1	e_3
e_7	e_4	e_5	e_6	$-e_1$	$-e_2$	$-e_3$	-1

Table 1- Octonion Multiplication table.

Hence, we get the following relations among octonion basis elements i.e.

$$[e_A, e_B] = 2f^{ABC}e_C; \quad \{e_A, e_B\} = -\delta_{AB}e_0; \quad e_A(e_Be_C) \neq (e_Ae_B)e_C;$$
 (3)

where brackets [] and {} are respectively used for commutation and the anti commutation relations while δ_{AB} is the usual Kronecker delta-Dirac symbol. Octonion conjugate is thus defined as,

$$\bar{x} = x_0 e_0 - \sum_{A=1}^{7} x_A e_A \qquad (A = 1, 2,, 7).$$
 (4)

An Octonion can be decomposed in terms of its scalar (Sc(x)) and vector (Vec(x)) parts as

$$Sc(x) = \frac{1}{2}(x+\bar{x}) = x_0; \quad Vec(x) = \frac{1}{2}(x-\bar{x}) = \sum_{A=1}^{7} x_A e_A.$$
 (5)

Conjugates of product of two octonions as well as its own conjugate are defined as

$$(\overline{xy}) = \overline{y}\overline{x} ; \overline{(\overline{x})} = x;$$
 (6)

while the scalar product of two octonions is defined as

$$\langle x, y \rangle = \sum_{\alpha=0}^{7} x_{\alpha} y_{\alpha} = \frac{1}{2} (x \bar{y} + y \bar{x}) = \frac{1}{2} (\bar{x} y + \bar{y} x);$$
 (7)

which can be written in terms of octonion units as

$$\langle e_A, e_B \rangle = \frac{1}{2} (e_A \overline{e_B} + e_B \overline{e_A}) = \frac{1}{2} (\overline{e_A} e_B + \overline{e_B} e_A) = \delta_{AB}.$$
 (8)

The norm of the octonion N(x) is defined as

$$N(x) = \overline{x}x = x\,\overline{x} = \sum_{\alpha=0}^{7} x_{\alpha}^{2} e_{0}; \tag{9}$$

which is zero if x = 0, and is always positive otherwise. It also satisfies the following property of normed algebra

$$N(xy) = N(x)N(y) = N(y)N(x).$$
(10)

As such, for a nonzero octonion x, we define its inverse as

$$x^{-1} = \frac{\bar{x}}{N(x)} \tag{11}$$

which shows that

$$x^{-1}x = xx^{-1} = 1.e_0; (xy)^{-1} = y^{-1}x^{-1}.$$
 (12)

3 SU(3) Generators (Gell-Mann Matrices)

The Gell-Mann λ matrices are used for the representations of the infinitesimal generators of the special unitary group called SU(3). This group consists eight linearly independent generators $G_A \forall (A = 1, 2, 3, \dots, 8)$ which satisfy the following commutation relation as

$$[G_A, G_B] = iF^{ABC}G_C \tag{13}$$

where F^{ABC} is the structure constants. It is completely antisymmetric (i.e. $F^{123}=+1$; $F^{147}=F^{165}=F^{246}=F^{257}=F^{354}=F^{367}=\frac{1}{2}$ and $F^{458}=F^{678}=\frac{\sqrt{3}}{2}$). Independent generators $G_A(\forall A=1,2,3,......8)$ of SU(3) symmetry group are related with the 3×3 Gell-Mann λ matrices as

$$G_A = \frac{\lambda_A}{2} \tag{14}$$

where $\lambda_A(\forall A = 1, 2, 3, \dots 8)$ are defined as

$$\lambda_{1} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \qquad \lambda_{2} = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix};
\lambda_{3} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \qquad \lambda_{4} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix};
\lambda_{5} = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}; \qquad \lambda_{6} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix};
\lambda_{7} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}; \qquad \lambda_{8} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & -2 \end{pmatrix} \tag{15}$$

and satisfy the following properties

$$(\lambda_A)^{\dagger} = \lambda_A;$$

$$Tr(\lambda_A) = 0 \quad Tr(\lambda_A \lambda_B) = 2\delta_{AB};$$

$$[\lambda_A, \lambda_B] = 2iF^{ABC}\lambda_C \ (\forall A, B, C = 1, 2, 3.....8).$$
(16)

As such, we may summarize the multiplication rules for the generators (in terms of λ matrices) of SU(3) symmetry in the following table as;

•	λ_1	λ_2	λ_3	λ_4	λ_5	λ_6	λ_7	λ_8
λ_1	Λ_1	$i\lambda_3$	$-i\lambda_2$	$\frac{i}{2}\lambda_7$	$-\frac{i}{2}\lambda_6$	$\frac{i}{2}\lambda_5$	$-\frac{i}{2}\lambda_4$	$\frac{1}{\sqrt{3}}\lambda_1$
λ_2	$-i\lambda_3$	Λ_2	$i\lambda_1$	$\frac{i}{2}\lambda_6$	$\frac{i}{2}\lambda_7$	$-\frac{i}{2}\lambda_4$	$-\frac{i}{2}\lambda_5$	$\frac{1}{\sqrt{3}}\lambda_2$
λ_3	$i\lambda_2$	$-i\lambda_1$	Λ_3	$-\frac{i}{2}\lambda_5$	$\frac{i}{2}\lambda_4$	$\frac{i}{2}\lambda_7$	$-\frac{i}{2}\lambda_6$	$\frac{1}{\sqrt{3}}\lambda_3$
λ_4	$-\frac{i}{2}\lambda_7$	$-\frac{i}{2}\lambda_6$	$\frac{i}{2}\lambda_5$	Λ_4	$-\frac{i}{2}\lambda_3$	$\frac{i}{2}\lambda_2$	$\frac{i}{2}\lambda_1$	$-\frac{\sqrt{3}}{2}i\lambda_5$
λ_5	$\frac{i}{2}\lambda_6$	$-\frac{i}{2}\lambda_7$	$-\frac{i}{2}\lambda_4$	$\frac{i}{2}\lambda_3$	Λ_5	$-\frac{i}{2}\lambda_1$	$rac{i}{2}\lambda_2$	$\frac{\sqrt{3}}{2}i\lambda_4$
λ_6	$-\frac{i}{2}\lambda_5$	$\frac{i}{2}\lambda_4$	$-\frac{i}{2}\lambda_7$	$-\frac{i}{2}\lambda_2$	$\frac{i}{2}\lambda_1$	Λ_6	$\frac{i}{2}\lambda_3$	$-\frac{\sqrt{3}}{2}i\lambda_7$
λ_7	$\frac{i}{2}\lambda_4$	$\frac{i}{2}\lambda_5$	$\frac{i}{2}\lambda_6$	$-\frac{i}{2}\lambda_1$	$-\frac{i}{2}\lambda_2$	$-\frac{i}{2}\lambda_3$	Λ_7	$\frac{\sqrt{3}}{2}i\lambda_6$
λ_8	$-\frac{1}{\sqrt{3}}\lambda_1$	$-\frac{1}{\sqrt{3}}\lambda_2$	$-\frac{1}{\sqrt{3}}\lambda_3$	$\frac{\sqrt{3}}{2}i\lambda_5$	$-\frac{\sqrt{3}}{2}i\lambda_4$	$\frac{\sqrt{3}}{2}i\lambda_7$	$-\frac{\sqrt{3}}{2}i\lambda_6$	Λ_8

Table 2- Multiplication table for Gell-Mann λ matrices of SU(3) symmetry.

where the $\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4, \Lambda_5, \Lambda_6, \Lambda_7, \Lambda_8$ are described in terms of 3×3 matrices as

$$\Lambda_{1} = \Lambda_{2} = \Lambda_{3} = \Lambda_{123} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \Longrightarrow (\lambda_{1})^{2} = (\lambda_{2})^{2} = (\lambda_{3})^{2} = \Lambda_{123};$$

$$\Lambda_{4} = \Lambda_{5} = \Lambda_{45} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \Longrightarrow (\lambda_{4})^{2} = (\lambda_{5})^{2};$$

$$\Lambda_{6} = \Lambda_{7} = \Lambda_{67} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \Longrightarrow (\lambda_{6})^{2} = (\lambda_{7})^{2};$$

$$\Lambda_{8} = \frac{4}{3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \Longrightarrow (\lambda_{8})^{2} = \frac{4}{3} \hat{1}.$$
(17)

where $\hat{1}$ is 3×3 unit matrix.

4 Relation between Octonions and SU(3)Generators

Comparing Table-1 and Table-2, we may observe a resemblance between the octonions and the Gell-Mann λ matrices of SU(3) symmetry on using simultaneously the relations (2) and (17) in the following table as

Octonions basis	SU(3) generators
$e_1 \longmapsto$	$i\lambda_1$
$e_2 \longmapsto$	$i\lambda_2$
$e_3 \longmapsto$	$i\lambda_3$
$e_4 \longmapsto$	$rac{i}{2}\lambda_4$
$e_5 \longmapsto$	$rac{i}{2}\lambda_5$
$e_6 \longmapsto$	$-\frac{i}{2}\lambda_6$
$e_7 \longmapsto$	$-\frac{i}{2}\lambda_7$
$e_0 \longmapsto$	$\frac{\sqrt{3}}{2}\lambda_8$

Table 3- Relation between Octonion basis and SU(3) generators.

As such, we have the freedom to establish [20] a connection between the octonion basis elements e_A and 3×3 Gell-Mann λ matrices of SU(3) in the following manner i.e.

$$e_{1} \Rightarrow i\lambda_{1}, \ e_{2} \Rightarrow i\lambda_{2}, \ e_{3} \Rightarrow i\lambda_{3} \longmapsto e_{A} \iff i\lambda_{A}; \quad (\forall A = 1, 2, 3);$$

$$e_{4} \Rightarrow \frac{i}{2}\lambda_{4}, \ e_{5} \Rightarrow \frac{i}{2}\lambda_{5}, \iff e_{A} = \frac{i}{2}\lambda_{A}; \quad (\forall A = 4, 5,);$$

$$e_{6} \Rightarrow \frac{i}{2}\lambda_{6}, \ e_{7} \Rightarrow -\frac{i}{2}\lambda_{7}, \iff e_{A} = -\frac{i}{2}\lambda_{A}; \quad (\forall A = 6, 7,);$$

$$e_{0} \iff \frac{\sqrt{3}}{2}\lambda_{8}. \tag{18}$$

These results are similar to the those derived earlier by Günaydin Gürsey [18] for octonion units and λ matrices of SU(3) symmetry. Here, equation (18) satisfies the Cayley algebra followed by the octonion

multiplication rule $e_A \cdot e_B = -\delta_{AB} + f_{ABC}e_C$. So, we have the freedom to establish the following relations among structure constants of octonions and SU(3) symmetry as

$$F^{ABC} \Rightarrow f^{ABC} \ (\forall ABC = 123);$$

$$F^{ABC} \Rightarrow \frac{1}{2} f^{ABC} \ (\forall ABC = 147, 246, 257, 435, 516, 673);$$

$$F^{ABC} \Rightarrow \frac{\sqrt{3}}{2} f^{ABC} \ (\forall ABC = 458, 678). \tag{19}$$

Hence, we get

$$[e_{A}, e_{B}] \Rightarrow i [\lambda_{A}, \lambda_{B}] \qquad (\forall ABC = 123);$$

$$[e_{A}, e_{B}] \Rightarrow \frac{i}{2} [\lambda_{A}, \lambda_{B}] \qquad (\forall ABC = 147, 246, 257, 435, 516, 673);$$

$$[e_{A}, e_{B}] \Rightarrow \frac{\sqrt{3}}{2} i [\lambda_{A}, \lambda_{B}] \qquad (\forall ABC = 458, 678);$$

$$(20)$$

which are the commutation relations among octonions basis elements and Gell-Mann λ matrices of SU(3) symmetry the so called Eight fold way. The benefit to write the octonions in terms of Gell-Mann λ matrices of SU(3) symmetry may be described as

- \diamond Non-associativity of octonions does not effect the invariance of the symmetry group SU(2) spin (or isospin) multiplets for the given values of structure constants f^{ABC} .
- \diamond It is better to describe the SU(3) symmetry in terms of compact notations of octonions. Accordingly, the theory of strong interactions could be described better in terms of non associative Cayley algebra.
- \diamond The eighth Gell-Mann λ matrix could be designated in terms of hyper charge which may have the direct link with the scalar octonion unit e_0 .
- \diamond It may be concluded that the algebra of strong interactions corresponds to the SU(3) automorphisms of the octonion algebra which is in support of the results obtained earlier by Günaydin [19].

5 Octonions and QCD

The color group SU(3) corresponds to the local symmetry whose gauging gives rise to Quantum Chromodynamics (QCD). There are two different types of SU(3) symmetry. The first one is the symmetry that acts on the different colors of quarks. This symmetry is an exact gauge symmetry mediated by the gluons. Other SU(3) symmetry is a flavor symmetry which rotates different flavors of quarks to each other, or flavor SU(3). Flavor SU(3) is an approximate symmetry of the vacuum of QCD, and is not a fundamental symmetry at all. It is an accidental consequence of the small mass of the three lightest quarks. Here, we are interested in exact SU(3) symmetry of colors in terms of octonion algebra. For this, let us substitute the values of octonion units e_A in terms of λ_A from equation (18) in to equation (1) so that we may express an octonion x as

$$x = x_0 \left(\frac{\sqrt{3}}{2}\lambda_8\right) + x_1(i\lambda_1) + x_2(i\lambda_2) + x_3(i\lambda_3) + x_4\left(\frac{i}{2}\lambda_4\right) + x_5\left(\frac{i}{2}\lambda_5\right) + x_6\left(-\frac{i}{2}\lambda_6\right) + x_7\left(-\frac{i}{2}\lambda_7\right)$$
(21)

which can further be reduced as

$$x = x_0 O_0 + x_1 O_1 + x_2 O_2 + x_3 O_3 + x_4 O_4 + x_5 O_5 + x_6 O_6 + x_7 O_7$$
(22)

where $\mathcal{O}_0 \to \frac{\sqrt{3}}{2}\lambda_8$, $\mathcal{O}_1 \to i\lambda_1$, $\mathcal{O}_2 \to i\lambda_2$, $\mathcal{O}_3 \to i\lambda_3$, $\mathcal{O}_4 \to \frac{i}{2}\lambda_4$, $\mathcal{O}_5 \to \frac{i}{2}\lambda_5$, $\mathcal{O}_6 \to -\frac{i}{2}\lambda_6$, $\mathcal{O}_7 \to -\frac{i}{2}\lambda_7$. Thus the octonion conjugate be written as as

$$\bar{x} = x_0 \mathcal{O}_0 - x_1 \mathcal{O}_1 - x_2 \mathcal{O}_2 - x_3 \mathcal{O}_3 - x_4 \mathcal{O}_4 - x_5 \mathcal{O}_5 - x_6 \mathcal{O}_6 - x_7 \mathcal{O}_7. \tag{23}$$

Here the new octonion units \mathcal{O}_A associated with the SU(3) symmetry satisfy the octonion algebra

$$\mathcal{O}_A \cdot \mathcal{O}_B = -\delta_{AB} + f_{ABC} \mathcal{O}_C. \tag{24}$$

As such, we may reformulate the theory of strong interactions, the quantum Chromodynamics (QCD) based on colour $SU(3)_c$ whose generators satisfy the non-associative algebra of octonions. Let us consider the triplet (u, d, s) (i.e. the up, down, and strange) flavors of quarks as the three objects of the group namely the SU(3) group of flavor symmetry. Each quark has been described in terms of three colors namely the red, blue and green. The dynamics of the quarks and gluons are controlled by the quantum Chromodynamics Lagrangian. The gauge invariant QCD Lagrangian is described as

$$\mathcal{L} = \bar{\psi}_{j} (i\gamma^{\mu} (D_{\mu})_{jk} - m \, \delta_{jk}) \psi_{k} - \frac{1}{4} \, G^{a}_{\mu\nu} G^{\mu\nu}_{a}$$

$$= \bar{\psi}_{j} (i\gamma^{\mu} \partial_{\mu} - m) \psi_{j} - g \, G^{a}_{\mu} \bar{\psi}_{j} \gamma^{\mu} T^{a}_{jk} \psi_{k} - \frac{1}{4} \, G^{a}_{\mu\nu} G^{\mu\nu}_{a}$$
(25)

where (j, k = 1, 2, 3) are labeled for three quark fields associated with three colors (namely the red, blue and green) so that we have

$$\psi_j = \begin{pmatrix} \psi_R \\ \psi_B \\ \psi_G \end{pmatrix}; \qquad \bar{\psi}_j = (\bar{\psi}_R, \bar{\psi}_B, \bar{\psi}_G)$$
 (26)

which is a dynamical function of space-time, in the fundamental representation of the SU(3) gauge group, indexed by (j, k = 1, 2, 3). In equation (25) G^a_μ is the octet of gluon fields which is also a dynamical function of space-time in the adjoint representation of the SU(3) gauge group, indexed by a, b, ... = 1, 2,, 8; the γ^μ are the Dirac matrices connecting the spinor representation to the vector representation of the Lorentz group; and T^a_{jk} are the generators connecting the fundamental, anti-fundamental and adjoint representations of the SU(3) gauge group. In our case, the octonion units connecting to Gell-Mann λ matrices provide one such representation for the generators of SU(3) gauge group. In equation (25), the symbol $G^a_{\mu\nu}$ represents the gauge invariant gluon field strength tensor, analogous to the electromagnetic field strength tensor $F_{\mu\nu}$ in Electrodynamics. It is described by

$$G_{\mu\nu}^{a} = \partial_{\mu}G_{\nu}^{a} - \partial_{\nu}G_{\mu}^{a} - g f^{abc} G_{\mu}^{b} G_{\mu}^{c} (\forall a, b, c = 1, 2, 3, ..., 8)$$
(27)

where f^{abc} are the structure constants of SU(3) groups as described above in terms of octonions. In equation (25), the constants m and g control the quark mass and coupling constants of the theory, subject to renormalization in the full quantum theory. Here we may introduce a local phase transformation in color space. Under SU(3) symmetry the spinor ψ transforms as

$$\psi \longrightarrow \psi' = U\psi = \exp\{i\lambda \cdot \alpha(x)\}\psi; \qquad (\lambda = 1, 2, \dots, 8)$$
 (28)

where

$$\lambda.\alpha(x) = \lambda_1\alpha_1 + \lambda_2\alpha_2 + \lambda_3\alpha_3 + \lambda_4\alpha_4 + \lambda_5\alpha_5 + \lambda_6\alpha_6 + \lambda_7\alpha_7 + \lambda_8\alpha_8 \tag{29}$$

which on using Table-3, may directly be written in the following form in terms of octonions i.e.

$$\lambda.\alpha(x) = -ie_1\alpha_1 - ie_2\alpha_2 - ie_3\alpha_3 - 2ie_4\alpha_4 - 2ie_5\alpha_5 - 2ie_6\alpha_6 - 2ie_7\alpha_7 + \frac{2}{\sqrt{3}}e_0\alpha_8. \tag{30}$$

As such, the the quantum Chromodynamics (QCD) may be reformulated in terms of octonions and non-associative algebra in order to explain its interesting consequences like

- ♦ Quarks confinement
- \diamond Color blindness of nature
- \diamond Asymptotic freedom
- ♦ Calculation for the masses of mesons and baryons etc.

6 Split octonions

The split octonions [18, 19, 20, 29] are a non associative extension of split quaternions. They differ from the octonion in the signature of quadratic form. The split octonion have a signature (4,4) whereas the octonions have positive signature (8,0). The Cayley algebra of octonions over the field of complex number is visualized as the algebra of split octonions with its following basis element,

$$u_{0} = \frac{1}{2} (e_{0} + ie_{7}), \qquad u_{0}^{\star} = \frac{1}{2} (e_{0} - ie_{7}),$$

$$u_{j} = \frac{1}{2} (e_{j} + ie_{j+3}), \qquad u_{j}^{\star} = \frac{1}{2} (e_{j} - ie_{j+3}) (\forall j = 1, 2, 3)$$
(31)

where (\star) denotes the complex conjugation and $(i = \sqrt{-1})$, the imaginary unit, commutes with all e_A ($\forall A = 1, 2, 3, ..., 7$). In equation (31) $u_0, u_0^{\star}, u_j, u_j^{\star}$ are defined as the bi-valued representations of quaternion units e_0 , e_1 , e_2 , e_3 which satisfy the following [31] multiplication rule

$$e_i e_k = -\delta_{ik} + \epsilon_{ikl} e_l \ (\forall j, k, l = 1, 2, 3) \tag{32}$$

where ϵ_{jkl} are the three index Levi-Civita symbols. The split octonion basis elements $u_0, u_0^{\star}, u_j, u_j^{\star}$ satisfy the following multiplication rule

$$u_{i}u_{j} = \epsilon_{ijk}u_{k}^{\star}; \qquad u_{i}^{\star}u_{j}^{\star} = -\epsilon_{ijk}u_{k}^{\star} \qquad (\forall i, j, k = 1, 2, 3)$$

$$u_{i}u_{j}^{\star} = -\delta_{ij}u_{0}; \qquad u_{i}u_{0} = 0; \qquad u_{i}^{\star}u_{0} = u_{i}^{\star}$$

$$u_{i}^{\star}u_{j} = -\delta_{ij}u_{0}; \qquad u_{i}u_{0}^{\star} = u_{0}; \qquad u_{i}^{\star}u_{0}^{\star} = 0$$

$$u_{0}u_{i} = u_{i}; \qquad u_{0}^{\star}u_{i} = 0; \qquad u_{0}u_{i}^{\star} = 0$$

$$u_{0}^{\star}u_{i}^{\star} = u_{i}; \qquad u_{0}^{2} = u_{0}; \qquad u_{0}^{\star2} = u_{0}^{\star}; \qquad u_{0}u_{0}^{\star} = u_{0}^{\star}u_{0} = 0.$$

$$(33)$$

So, we may introduce a convenient realization for the basis elements $(u_0, u_0^{\star}, u_j, u_j^{\star})$ in term of Pauli's spin matrices as

$$u_{0} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}; \qquad u_{0}^{\star} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix};$$

$$u_{j} = \begin{bmatrix} 0 & 0 \\ e_{j} & 0 \end{bmatrix}; \qquad u_{j}^{\star} = \begin{bmatrix} 0 & -e_{j} \\ 0 & 0 \end{bmatrix}. \qquad (\forall j = 1, 2, 3)$$
(34)

The Cayley's split octonion algebra may also be expressed via2 × 2 Zorn's vector matrix realizations as

$$A = au_0^{\star} + bu_0 + x_i u_i^{\star} + y_i u_i = \begin{pmatrix} a & -\overrightarrow{x} \\ \overrightarrow{y} & b \end{pmatrix}$$
 (35)

where a and b are scalars and \overrightarrow{x} and \overrightarrow{y} are 3-vectors with the product defined [18, 32] as

$$\begin{pmatrix} a & \overrightarrow{x} \\ \overrightarrow{y} & b \end{pmatrix} \begin{pmatrix} c & \overrightarrow{u} \\ \overrightarrow{v} & d \end{pmatrix} = \begin{pmatrix} ac + \overrightarrow{x} \cdot \overrightarrow{v} & a\overrightarrow{u} + d\overrightarrow{x} - \overrightarrow{y} \times \overrightarrow{v} \\ c\overrightarrow{y} + b\overrightarrow{v} - \overrightarrow{x} \times \overrightarrow{u} & bd + \overrightarrow{y} \cdot \overrightarrow{u} \end{pmatrix}. \tag{36}$$

Hence, the split octonion conjugate equation may be written via 2×2 Zorn's vector matrix realizations as

$$\overline{A} = au_0 + bu_0^{\star} - x_i u_i^{\star} - y_i u_i = \begin{pmatrix} b & \overrightarrow{x} \\ -\overrightarrow{y} & a \end{pmatrix}. \tag{37}$$

So, the norm of A is defined as

$$A\overline{A} = \overline{A}A = (ab + \overrightarrow{x}.\overrightarrow{y})\hat{1}$$
(38)

where $\hat{1}$ as the unit matrix of order 2×2 .

7 Split-Octonion SU(3) Gauge Theory

The automorphism group of the Octonion algebra is the 14-dimensional exceptional G_2 group that admits a SU(3) subgroup leaving invariant the idempotents u_0 and u_0^* described by equation (31). This $SU(3)_C$ was identified as the color group acting on the quark and anti-quark triplets [19, 32]. As such, the automorphism group SU(3) of the quantum mechanical Hilbert space should be considered as an exact

symmetry and can not be identified as the symmetry of broken unitary spin gauge group. It is like the $SU(3)_C$ color gauge group of quantum chromo-dynamics (QCD). Therefore, in order to describe the SU(3) gauge theory suitably handled with split octonions, let us start with the split octonion equivalent of any four vector A_{μ} and its conjugate in terms of the following 2 × 2 Zorn matrix realization as

$$Z(A) = \begin{bmatrix} x_4 & -\overrightarrow{x} \\ \overrightarrow{y} & y_4 \end{bmatrix}; \qquad Z(\overline{A}) = \begin{bmatrix} x_4 & \overrightarrow{x} \\ -\overrightarrow{y} & y_4 \end{bmatrix}. \tag{39}$$

Rather, the Octonion covariant derivative or \mathcal{O} - derivative of an octonion K is defined [32, 33, 34] as

$$K_{\parallel\mu} = K_{,\mu} + [\Im_{\mu}, K]$$
 (40)

where \Im_{μ} is the Octonion affinity. It is the object that makes $K_{\parallel\mu}$ transform like an octonion under \mathcal{O} transformations i.e.

$$K' = U K U^{-1}
K'_{\parallel \mu} = U K_{\parallel \mu} U^{-1}
\mathfrak{S}'_{\mu} = U \mathfrak{S}_{\mu} U^{-1} - \frac{\partial U}{\partial x^{\mu}} U^{-1}$$
(41)

where the U(x) are octonions which define local (Octonion) unitary transformations and are isomorphic to the rotation group O(3). Thus, equation (41) describes SU(2) nature of octonion \mathcal{O} transformations resulting to the octonion affinity (gauge potential) \mathfrak{F}_{μ} of Yang-Mill's type field and is expressed as.

$$\Im_{\mu} = -L_{\mu j} u_{j}^{\star} - K_{\mu j} u_{j} = \begin{bmatrix} 0_{2} & L_{\mu i} e_{j} \\ -K_{\mu j} e_{j} & 0_{2} \end{bmatrix} (\forall j = 1, 2, 3)$$

$$(42)$$

where the quaternion units $e_j = -i\sigma_j$ are suitably handled [31] with Pauli spin matrices σ_j . Now, we have the freedom to extend SU(2) gauge theory to the case of SU(3) Yang Mills gauge theory of colored quarks by replacing the Pauli spin matrices to Gellmann λ matrices. So, from equation (42), it is clear that octonion covariant derivative (40) is subjected by two real (or one complex) gauge potential transformations. Hence, G^a_μ the octet of gluon fields describing Lagrangian (25) is either a complex gauge field or comprises the order pair of two real gauge fields. So, we may write the covariant derivative D_μ for SU(3) Lagrangian (25) as

$$D_{\mu} = \partial_{\mu} + \mathbb{V}_{\mu} \tag{43}$$

where \mathbb{V}_{μ} is the octonion form of generalized four potential described as

$$V_{\mu} = e_0 \left(A_{\mu}^{\alpha} e_{\alpha} \right) + i e_7 \left(B_{\mu}^{\alpha} g_{\alpha} \right) \qquad (\forall \mu = 0, 1, 2, 3; \alpha = 1, 2, \dots, 8.)$$
(44)

The beauty of the equation (44) reinforces the SU(3) symmetry of colored quarks with two gauge potentials as the consequence of automorphism group of split octonions in terms of 2×2 Zorn vector matrix realization. Here, the two gauge potentials A^{α}_{μ} and $B^{\alpha}_{\mu}(\forall \mu = 0, 1, 2, 3; \alpha = 1, 2,, 8.)$ may be identified as the gauge potentials for two chromo charges supposed to be responsible for the existence of electric

and magnetic chromo-charges. It may, therefore, be concluded that octonion colored quarks are dyons: the particles which carry the simultaneous existence of electric and magnetic charges [24, 25, 26, 27]. Substituting the value of $SU(3)_C$ octonion gauge potential V_{μ} (44) in to the equation (43), we may write the covariant derivative D_{μ} as

$$D_{\mu} = \partial_{\mu} + e_0 \left(A^{\alpha}_{\mu} e_{\alpha} \right) + i e_7 \left(B^{\alpha}_{\mu} g_{\alpha} \right)$$

$$= u_0^* \left(\partial_{\mu} + A^{\alpha}_{\mu} e_{\alpha} + B^{\alpha}_{\mu} g_{\alpha} \right) + u_0 \left(\partial_{\mu} + A^{\alpha}_{\mu} e_{\alpha} - B^{\alpha}_{\mu} g_{\alpha} \right)$$
(45)

which may is equivalently [34] be written as

$$D_{\mu} = \begin{pmatrix} \partial_{\mu} + \left(e_{\alpha} A_{\mu}^{\alpha} + g_{\alpha} B_{\mu}^{\alpha} \right) & 0 \\ 0 & \partial_{\mu} + \left(e_{\alpha} A_{\mu}^{\alpha} - g_{\alpha} B_{\mu}^{\alpha} \right) \end{pmatrix}. \tag{46}$$

It yields to

$$[D_{\mu}, D_{\nu}] = \begin{pmatrix} G^{\alpha}_{\mu\nu} e_{\alpha} + \mathsf{G}^{\alpha}_{\mu\nu} g_{\alpha} & 0\\ 0 & G^{\alpha}_{\mu\nu} e_{\alpha} - \mathsf{G}^{\alpha}_{\mu\nu} g_{\alpha} \end{pmatrix} \longmapsto \mathbb{G}^{\alpha}_{\mu\nu}$$
(47)

where

$$G^{\alpha}_{\mu\nu} = \partial_{\mu}A^{\alpha}_{\nu} - \partial_{\nu}A^{\alpha}_{\mu} + e_{\alpha} \left[A^{\alpha}_{\mu}, A^{\alpha}_{\nu} \right] \longmapsto E^{\alpha}_{\mu\nu};$$

$$G^{\alpha}_{\mu\nu} = \partial_{\mu}B^{\alpha}_{\nu} - \partial_{\nu}B^{\alpha}_{\mu} + g_{\alpha} \left[B^{\alpha}_{\mu}, B^{\alpha}_{\nu} \right] \longmapsto H^{\alpha}_{\mu\nu};$$

$$(48)$$

are two SU(3) non-Abelian gauge field strengths in term of electric $(E^{\alpha}_{\mu\nu})$ and magnetic field $(H^{\alpha}_{\mu\nu})$ field strengths obtained respectively from electric A^{α}_{μ} and magnetic B^{α}_{μ} gauge potentials. Operating the covariant derivative D_{μ} (46) to the generalized field strength $\mathbb{G}^{\alpha}_{\mu\nu}$ of dyons (47), we get

$$D_{\mu}\mathbb{G}^{\alpha}_{\ \mu\nu} = \begin{pmatrix} \partial_{\mu}G^{\alpha}_{\mu\nu}e_{\alpha} + \partial_{\mu}\mathsf{G}^{\alpha}_{\mu\nu}g_{\alpha} & 0\\ 0 & \partial_{\mu}G^{\alpha}_{\mu\nu}e_{\alpha} - \partial_{\mu}\mathsf{G}^{\alpha}_{\mu\nu}g_{\alpha} \end{pmatrix} = \mathbb{J}^{\alpha}_{\nu}$$
(49)

where $\mathbb{J}^{\alpha}_{\nu}$ describes the generalized octonion gauge current of dyons in term of 2×2 Zorn matrix realization of split octonion SU(3) gauge theory. It also comprises the electric and magnetic four currents of dyons as

$$\mathbb{J}_{\nu}^{\alpha} = \begin{pmatrix} J_{\nu}^{\alpha} e_{\alpha} + K_{\nu}^{\alpha} g_{\alpha} & 0\\ 0 & J_{\nu}^{\alpha} e_{\alpha} - K_{\nu}^{\alpha} g_{\alpha} \end{pmatrix}.$$
(50)

Here $J^{\alpha}_{\nu} = \partial_{\mu} G^{\alpha}_{\mu\nu}$ and $K^{\alpha}_{\nu} = \partial_{\mu} G^{\alpha}_{\mu\nu}$ are the four currents respectively associated with the presence of electric and magnetic charges. So, it is concluded concluded that split octonion SU(3) gauge theory of colored quarks describes dyons which are the particles carrying the simultaneous existence of electric and magnetic monopoles.

ACKNOWLEDGMENT: One of us (OPSN) acknowledges the financial support from Third World Academy of Sciences, Trieste (Italy) and Chinese Academy of Sciences, Beijing under UNESCO-TWAS Associateship Scheme. He is also thankful to Prof. Yue-Liang Wu for his hospitality and research facilities at ITP and KITPC, Beijing (China).

References

- [1] L. E. Dickson, "On Quaternions and Their Generalization and the History of the Eight Square Theorem", Ann. Math., 20, (1919), 155.
- [2] W. R. Hamilton, "Elements of Quaternions", Chelsea Publications Co., New York, (1969).
- [3] P.G.Tait, "An elementary Treatise on Quaternions", Oxford Univ. Press (1875).
- [4] A. Cayley, "An Jacobi's elliptic functions, in reply to the Rev. B. Bornwin; and on quaternion", Phil. Mag., 26 (1845), 208.
- [5] R. P. Graves, "Life of Sir William Rowan Hamilton", 3 volumes, Arno Press, New York, (1975).
- [6] J. C. Baez, "The Octonions", Bull. Amer. Math. Soc., 39, (2001), 145.
- [7] S. Catto, "Exceptional Projective Geometries and Internal Symmetries", eprint: arXiv:hep-th/0302079.
- [8] S. Catto, "Symmetries in Science: from the rotation group to quantum algebras", 6 (1993), 129, Ed. B. Gruber, Plenum Press.
- [9] R. Foot and G. C. Joshi, "Space-time symmetries of superstring and Jordan Algebras", Int. J. Theor. Phys., 28, (1989) 1449.
- [10] R. Foot and G. C. Joshi, "String theories and the Jordan algebras", Phys. Lett., B199 (1987), 203.
- [11] T. Kugo and P. Townsend, "Supersymmetry and the Division Algebras", Nucl Phys., B221 (1983), 357.
- [12] C. A. Manogue and A. Sudbery, Phys. Rev., "General solutions of covariant superstring equations of motion", <u>D40</u> (1989), 4073.
- [13] J. Schray, "Octonions and Supersymmetry", Ph. D. thesis, Department of Physics, Oregon State University, (1994) (un published).
- [14] K. S. Abdel-Khalek Mostafa, "Ring Division Algebra, Self Duality and Supersymmetry", eprint:arXiv: hep-th/0002155.
- [15] K. Morita, "Quaternionic Variational Formalism for Poincaré Gauge Theory and Supergravity", Prog. Theor. Phys., 73 (1985), 999.
- [16] K. Morita, "Gauge Theories over Quaternions and Weinberg-Salam Theory", Prog. Theor. Phys., 65 (1981), 2071.
- [17] K. Morita, "Quaternionic Weinberg-Salam Theory", Prog. Theor. Phys., <u>67</u> (1982), 1860.

- [18] M. Günaydin and Feza Gürsey; "Quark structure and octonions", J. Math. Phys., <u>14</u> (1973), 1651.
- [19] M. Günaydin, "Octonionic Hilbert spaces, the Poincaré group and SU(3)", J. Math. Phys., **17** (1976), 1875.
- [20] Pushpa, P. S. Bisht, Tianjun Li and O. P. S. Negi, "Quaternion Octonion Reformulation of Quantum Chromodynamics", Int. J. Theor. Phys., **50** (2011), 594.
- [21] P. A. M. Dirac, "Quantised Singularities in the Electromagnetic Field", Proc. Royal Society, A133 (1931), 60.
- [22] G. 't Hooft, "Magnetic monopoles in unified gauge theories", Nucl. Phys., <u>B79</u> (1974), 276.
- [23] A. M. Polyakov, "Particle spectrum in quantum field theory", JETP Lett., <u>20</u> (1974), 194.
- [24] J. Schwinger, "Dyons Versus Quarks", Science, 166 (1969), 690.
- [25] D. Zwanzinger, "Quantum Field Theory of Particles with Both Electric and Magnetic Charges", Phys. Rev., 176 (1968), 1489.
- [26] B. Julia and A. Zee, "Poles with both magnetic and electric charges in non-Abelian gauge theory", Phys. Rev., <u>D11</u> (1975), 2227.
- [27] O. P. S. Negi and H. Dehnen, "Gauge Formulation for Two potential Theory of Dyons", Int. J. Theor. Phys., 50 (2011), 2446.
- [28] B. C. Chanyal, P. S. Bisht, O. P. S. Negi, "Generalized Octonion Electrodynamics", Int. J. Theor. Phys., 49 (2010), 1333.
- [29] B. C. Chanyal, P. S. Bisht, O. P. S. Negi, "Generalized Split-Octonion Electrodynamics", Int. J. Theor. Phys., 50 (2011), 1919.
- [30] P. S. Bisht, B. Pandey and O. P. S. Negi, "Interpretations of octonion wave equations", FIZIKA (Zagreb), **B17** (2008), 405.
- [31] Pushpa, P. S. Bisht, and O. P. S. Negi, "Spontaneous symmetry breaking in presence of electric and magnetic charges", Int. J. Theor. Phys., **50** (2011), 1927.
- [32] Carlos Castro, "On the noncommutative and nonassociative geometry of octonionic space time, modified dispersion relations and grand unification", J. Math. Phys., 48 (2007), 073517.
- [33] S. Marques and C. G. Oliveira, "An extension of quaternionic metrics to octonions", J. Math. Phys., 26, (1983), 3131.
- [34] Shalini Dangwal, P S Bisht, O P S Negi, "Octonionic Gauge Formulation for Dyonic Fields", eprint:arXiv: hep-th/0608061.