

Deformations of instantons

M. F. ATIYAH[†], N. J. HITCHIN[†], AND I. M. SINGER^{†‡}

[†] Mathematical Institute, Oxford, OX1 3LB, England; and [‡] Department of Mathematics, University of California, Berkeley, California 94720; and Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Contributed by I. M. Singer, April 20, 1977

ABSTRACT A study is made of the self-dual Yang-Mills fields in Euclidean 4-space. For $SU(2)$ gauge theory it is rigorously shown that the solutions depend on $8k - 3$ parameters, where k is the Pontrjagin index.

There has been considerable interest recently in the instanton or pseudo-particle solutions of the classical Yang-Mills equations in Euclidean 4-space (refs. 1, 2, and 3). In geometrical terms, these equations are the variational equations for the norm-square $\|F\|^2$ of the curvature F of a fiber-bundle with group G and connection A over R^4 . In physics terminology, $\|F\|^2$ is the action, F the gauge field, A the gauge potential, and G the gauge group. The cases studied in most detail are for $G = SU(n)$ and, particularly, $G = SU(2)$.

The connection A is assumed to be asymptotically flat in an appropriate sense so that $F \rightarrow 0$ at ∞ and $\|F\|^2 < \infty$. Since the variational equations are conformally invariant with respect to change in the metric on R^4 , the most natural geometrical restriction to impose on A at ∞ is that it extends to a connection for a bundle over the 4-sphere S^4 . The topological type of such a bundle is then determined by a homotopy class of maps $S^3 \rightarrow G$, which is given by an integer k when $G = SU(n)$ (and more generally for any simple compact nonabelian Lie group): this is referred to by physicists as the Pontrjagin index (differing from the topologist's terminology by a factor of 2).

Using the duality $*$ -operator on R^4 or S^4 , we can decompose F into $F^+ \oplus F^-$, where $*F^+ = F^+$ and $*F^- = -F^-$. Clearly, $\|F\|^2 = \|F^+\|^2 + \|F^-\|^2$, while the Pontrjagin index k is given by[§]

$$k = \frac{1}{8\pi^2} (\|F^+\|^2 - \|F^-\|^2).$$

Hence, $\|F^+\|^2 \geq 8\pi^2 k$, and the minimum is attained only if $F^- = 0$ or $F^+ = 0$. Solutions with $F^- = 0$ are called self-dual solutions and have been constructed for all $k \geq 0$. For $k = 0$ we have the trivial solution $F = 0$, for $k = 1$ we have the "instanton," and for $k > 1$ we have "multi-instantons." The most general explicit solutions constructed so far are those of Jackiw *et al.* (ref. 3), which depend on $5k + 4$ parameters.[¶] Our main result is that the complete set of solutions depends on $8k - 3$ parameters. This confirms some preliminary results of Jackiw and Rebbi (ref. 4) and Schwartz (ref. 5).

RESULTS

If a connection A yields a self-dual Yang-Mills field F , then so does any connection $g(A)$ where g is a bundle automorphism (or gauge transformation). The space of all solutions A modulo the action of this gauge group will, as usual in such geometric problems, be called the space of moduli. Our main result can now be formulated as a precise theorem:

THEOREM. *The space of moduli of self-dual $SU(2)$ -Yang-Mills fields over S^4 , with Pontrjagin index $k \geq 1$, is a manifold of dimension $8k - 3$.*

The standard deformation theory approach to such problems is to consider the linearized equations modulo the infinitesimal gauge transformations. Here this leads to a three-step elliptic complex

$$0 \rightarrow \mathcal{G} \xrightarrow{D_0} \mathcal{G} \otimes \Omega^1 \xrightarrow{D_1} \mathcal{G} \otimes \Omega^2 \rightarrow 0$$

where Ω^1 denotes 1-forms, Ω^2 denotes anti-self-dual 2-forms on S^4 , and \mathcal{G} is the Lie algebra of G . The operator D_0 is the covariant derivative and D_1 is the anti-self-dual part of the covariant derivative ($D_1 D_0 = 0$ because we are using a self-dual connection). The index theorem of Atiyah-Singer (ref. 6) yields the alternating sum formula $h^0 - h^1 + h^2 = 3 - 8k$. Here h^0 is just the dimension of the null space of D_0 , and this is zero unless the $SU(2)$ -bundle is trivial (which is excluded for $k \geq 1$). h^1 is the potential number of moduli and h^2 is the dimension of the null space of D_1^* . Fortunately, in our case, a Bochner type vanishing theorem works very well and we find $h^2 = 0$. This gives $h^1 = 8k - 3$, showing that this is the dimension of the solutions of the linearized problem.

Now we appeal to the general theorem of Kuranishi (ref. 7), which, when $h^2 = 0$, guarantees that the infinitesimal variations really integrate to give genuine local variations. Moreover, the Kuranishi theorem asserts that the family of solutions thus obtained is (locally) complete and effective (non-redundant). This leads to the theorem as stated above.

Note. The theorem does not assert that, for each k , the space of moduli is connected: In principle it may have several components. For $k = 1$ it is in fact connected and is the hyperbolic 5-space. See Yang (ref. 8) for a different discussion of the case $k = 1$.

FURTHER REMARKS

The above arguments apply equally to $SU(n)$, provided we have an irreducible connection (so that $h^0 = 0$), i.e., one that does not come trivially from $SU(n - 1)$. We then find $h^1 = 4nk - n^2 + 1$. Moreover, the existence of irreducible $SU(n)$ solutions can be deduced from this formula provided $k \geq (n - 1)/2$. In the opposite direction one can deduce nonexistence for $k < n/4$.

Similar methods, i.e., index theorem plus vanishing theorem, yield a formula for the dimension d of the space of zero-eigenvalue fermions (harmonic spinors): one finds $d = k$.

The problem of explicitly constructing the $(8k - 3)$ -parameter families of solutions, whose existence is asserted by our theorem, can be treated by converting it into a problem in algebraic geometry (M. F. Atiyah and R. Ward, unpublished).

The 4-sphere can be replaced by other 4-manifolds M (compact, oriented, Riemannian). If M is also a spin-manifold, then we have the two spin $SU(2)$ -bundles, P^+ , P^- . One can show that P^+ (with the Riemannian connection) is self-dual if

[§] Physicists use a different norm and get a factor $1/(16\pi^2)$.

[¶] The formula is a little different for $k = 1, 2$.

and only if the metric on M is an Einstein metric ($R_{ij} = \lambda g_{ij}$) and that P^- is then anti-self-dual. The deformation theory above can then be applied to P^+ to obtain the number of moduli. The vanishing theorem for a self-dual field ($h^2 = 0$) still applies provided that the conformal Weyl tensor of M is self-dual and that the scalar curvature is positive.

I.M.S. was partially supported by the National Science Foundation.

The costs of publication of this article were defrayed in part by the payment of page charges from funds made available to support the research which is the subject of the article. This article must therefore

be hereby marked "advertisement" in accordance with 18 U. S. C. §1734 solely to indicate this fact.

1. Belavin, A., Polyakov, A., Schwartz, A. & Tyupkin, Y. (1975) *Phys. Lett. B* **59**, 85.
2. Witten, E. (1977) *Phys. Rev. Lett.* **38**, 121.
3. Jackiw, R., Nohl, C. & Rebbi, C. (1977) *Phys. Rev. D* **15**, 1642.
4. Jackiw, R. & Rebbi, C. (1977) *Phys. Lett. B* **67**, 189.
5. Schwarz, A. S. (1977) *Phys. Lett. B* **67**, 172.
6. Atiyah, M. F. & Singer, I. M. (1968) *Ann. Math.* **87**, 484-530.
7. Kuranishi, M. (1965) *Proceedings of the Conference on Complex Analysis* (Springer-Verlag, Inc., New York), p. 142.
8. Yang, C. N. (1977) *Phys. Rev. D*, in press.