

Axial-Anomaly-Induced Fermion Fractionization and Effective Gauge-Theory Actions in Odd-Dimensional Space-Times

A. J. Niemi

The Institute for Advanced Study, Princeton, New Jersey 08540,^(a) and Research Institute for Theoretical Physics, University of Helsinki, SF-00170 Helsinki 17, Finland

and

G. W. Semenoff

Department of Physics, University of British Columbia, Vancouver, British Columbia V6T 2A6, Canada
(Received 2 September 1983)

A new quantum field-theoretical technique is developed and used to explore the relationship between even-space-time-dimensional axial anomalies and background-field-induced fermion numbers and Euler-Heisenberg effective actions in odd-dimensional space-times.

PACS numbers: 11.15.Tk, 11.30.Ly

It is by now well known that topological objects possess surprising quantum properties such as fractional fermion number.¹⁻⁸ Examples exist in 1+1 dimensions where kinks (solitons) provide the topologically nontrivial setting and in 3+1 dimensions where magnetic monopoles are used as the background field. An important experimental realization of this phenomenon has already appeared in the physics of linearly conjugated polymers.^{2,3}

In general, fermion fractionization is amenable to an elegant mathematical analysis: In a system where fermions interact with classical background fields the induced fermion number is the spectral asymmetry of the pertinent Dirac Hamiltonian; schematically,

$$N = -\frac{1}{2} \left\{ \sum_{E_n > 0} 1 - \sum_{E_n < 0} 1 \right\} \quad (1)$$

(Ref. 7), where E_n are the energy eigenvalues. Various techniques have been developed to compute (1)⁴⁻⁸ and quite recently a direct topological analysis has been applied to several (1+1)-⁶ and (3+1)-dimensional⁷ field-theoretical models. This approach is based on the derivation of certain trace identities which relate N to an asymptotic surface integral, and the result involves only the topological properties of the background fields.

In an independent line of study it has been found that odd-space-time-dimensional gauge field theories can exhibit unexpected structures such as the appearance of topological mass terms in 2+1 dimensions.^{9,10} These extra terms, related to Chern-Simons secondary characteristic classes, were originally introduced by requiring gauge symmetry but have recently been observed to arise through radiative corrections.¹¹

Here we shall explore the relationship between fermion-number fractionization and the topological properties generic to odd-space-time-dimensional gauge theories. As an illustration of the general structure we shall first analyze the (2+1)-dimensional QED. We then indicate how this analysis can be generalized to higher odd-dimensional space-times with non-Abelian gauge fields. Applications envisaged involve the Abrikosov-Nielsen-Olesen vortex which appears naturally in the physics of type-II superconductors, and also the string theory of hadrons and certain cosmological scenarios where stringlike structures appear. Furthermore, three-dimensional Euclidean bosonic field theories may represent the high-temperature behavior of (3+1)-dimensional quantum field theories.

We shall compute the fermion number induced by static classical background fields. In the course of this calculation we show that the trace-identity analysis presented in Refs. 6 and 7 is not always applicable. The operators involved are singular and in general exhibit anomalies. In the (2+1)-dimensional case these anomalies are governed by the two-dimensional Abelian Euclidean axial-anomaly equation¹²:

$$\begin{aligned} (i/2) \partial^k \text{tr} \{ \gamma^k \gamma^5 D(x, x) \} \\ = M \text{tr} \{ \gamma^5 D(x, x) \} + (e/4\pi) \epsilon^{ij} F_{ij}. \end{aligned} \quad (2)$$

(The effective dimensional reduction from 2+1 to 2 dimensions is a consequence of time translation invariance of the background field.) Here $D(x, y)$ is the (regulated) Euclidean Dirac propagator,

$$(i \not{\partial} + e \not{A} - M) D(x, y) = \delta(x - y), \quad (3)$$

and γ^i ($i = 1, 2$) and γ^5 are the two-dimensional γ

matrices. In the models studied in Refs. 6 and 7 the anomaly term is absent since there are no odd-dimensional anomalies. However, we find that here the entire fractional fermion number arises from this term. It is the two-dimensional Pontryagin density; thus the relationship between (1) and the topology of background fields. We shall use a simple covariance argument to deduce a low-momentum approximation of the induced current, and therefore the Euler-Heisenberg effective action. This effective action contains the topological mass term for (2+1)-dimensional gauge fields.^{9,10}

The (2+1)-dimensional QED Lagrangian is

$$\mathcal{L} = -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} + \bar{\psi}(i\not{\partial} + e\not{A} - m)\psi. \quad (4)$$

we find

$$\delta\Gamma/\delta A_\mu = e\langle j^\mu(x) \rangle = -e(i/2)\text{tr}\{S_>(x,x)\gamma^\mu + S_<(x,x)\gamma^\mu\}. \quad (8)$$

Here Γ is the effective action for the gauge fields obtained by eliminating the Fermi fields by functional integration, and the trace is over Dirac indices. The propagator (7) satisfies

$$(i\not{\partial} + e\not{A} - m)S(x,y) = \delta(x-y). \quad (9)$$

By defining

$$S(x,y) = \int_C (dE/2\pi) \exp[-iE(x^0 - y^0)] S_E(x,y), \quad (10)$$

where C is a causal contour in the complex E plane, and by performing a Wick rotation $E \rightarrow iE$, we find

$$\langle j^\mu(x) \rangle = -\frac{1}{2} \lim_{\epsilon \rightarrow 0^+} \int (dE/2\pi) e^{-iE\epsilon} \text{tr}\{S_E(x,x)\gamma^\mu + S_{-E}(x,x)\gamma^\mu\}. \quad (11)$$

We define

$$H\psi_n = E_n\psi_n, \quad H = \alpha^k(i\nabla^k + eA^k) + \beta m, \quad (12)$$

where $\beta = \gamma^0$, $\alpha^k = \gamma^0\gamma^k$, and H is the Dirac Hamiltonian of the system (4). We get

$$\langle j^0(x) \rangle = -\frac{1}{2} \lim_{\epsilon \rightarrow 0^+} \sum_{E_n} \exp(-|E_n|\epsilon) \psi_n^\dagger(x) \psi_n(x) \text{sgn}\{E_n\} \quad (13)$$

and

$$N = \int d^2x \langle j^0(x) \rangle = -\frac{1}{2} \lim_{\epsilon \rightarrow 0^+} \sum_{E_n} \exp(-|E_n|\epsilon) \text{sgn}\{E_n\}. \quad (14)$$

Equation (14) is a heat-kernel regularization of (1). We shall now evaluate (13) and (14). For this we observe that

$$\text{tr}\{S_E(xx)\gamma^0 + S_{-E}(xx)\gamma^0\} = 2(m/\sigma) \text{tr}\langle x|\gamma^0/(i\not{\partial} + e\not{A} - \sigma)|x\rangle, \quad (15)$$

where $\sigma^2 = m^2 + E^2$. We then identify $M = \sigma$ and $\gamma^5 = \gamma^0$ in (2) and get

$$\langle j^0(x) \rangle = \frac{1}{2} \int \frac{dE}{2\pi} \left\{ i \frac{m}{\sigma^2} \partial^k \text{tr}[\gamma^k \gamma^5 D(x,x)] - \frac{m}{\sigma^2} \frac{e}{2\pi} \epsilon^{ij} F_{ij} \right\}. \quad (16)$$

By evaluating the integral over two-space in (14) we find that the contribution from the first term in

We assume that $A_\mu(x)$ is a static classical background field and $A_0=0$. The magnetic flux is

$$\Phi = (1/4\pi) \int d^2x \epsilon^{ij} F_{ij} = (1/2\pi) \oint dx^i A_i. \quad (5)$$

(In the case of vortices in the Abelian-Higgs model we would have $\Phi = n/q$, where n is the topological quantum number of the vortex and q is the charge of the scalar field.) We first verify (1) and then evaluate the induced fermion number density and the fermion number.

Consider the current

$$j^\mu(x) = \frac{1}{2} [\bar{\psi}, \gamma^\mu \psi]. \quad (6)$$

By introducing the causal propagator

$$iS(x,y) = \theta(x^0 - y^0) \langle \psi(x) \bar{\psi}(y) \rangle - \theta(y^0 - x^0) \langle \bar{\psi}(y) \psi(x) \rangle = i\theta(x^0 - y^0) S_>(x,y) - i\theta(y^0 - x^0) S_<(x,y), \quad (7)$$

(16) vanishes and we have

$$N = \frac{e}{4\pi} \int \frac{dE}{2\pi} \frac{m}{m^2 + E^2} \int d^2x \epsilon^{ij} F_{ij} = \frac{1}{2} e \Phi, \quad (17)$$

which is the relation between the fractional fermion number and the magnetic flux. It should be noted that N is directly proportional to the parameters: In odd space-time dimensions there is no chirality. Hence the extra degree of freedom necessary for more complicated relations^{4,7,8} is missing.

By covariance we conclude that the part of (16) responsible for fractional fermion number is

$$\langle j^\mu(x) \rangle = (e/8\pi) \epsilon^{\mu\nu\sigma} F_{\nu\sigma}, \quad (18)$$

which is gauge invariant and conserved. If we evaluate (16) with a space-time constant F_{ij} we find that the first term vanishes. Consequently in the low-momentum limit (11) is exactly given by (18). Substituting (18) into (8) we find the functional antiderivative with respect to A_μ . The re-

sult is the Euler-Heisenberg effective action

$$\Gamma = \int d^3x \left\{ -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + (e^2/16\pi) \epsilon^{\mu\nu\sigma} F_{\mu\nu} A_\sigma \right\}. \quad (19)$$

Notice that the last, radiatively induced term is the topological mass term (Chern-Simons invariant) studied in Refs. 9 and 10. We conclude that, if absent at the classical level, this term is generated through quantum corrections.¹¹ Consequently the theory (4) exhibits a mass gap necessary for confinement.^{13,14} Furthermore, the existence of this mass gap, even as $m \rightarrow 0$, indicates a dynamical breaking of parity.⁹⁻¹¹ Finally, we also observe that if this effective bosonic field theory supports topological excitations they will carry a fermion number given by (17).

We can readily extend the previous results to arbitrary odd-dimensional space-times with non-Abelian background fields. As an example we consider a $(4+1)$ -dimensional $SU(N)$ invariant gauge theory with N_f Dirac fermions. As before we assume a static magnetic background gauge field with $A_0=0$. The four-dimensional non-Abelian Euclidean axial-anomaly equation is¹⁵

$$(i/2) \partial^k \text{tr} \{ \gamma^k \gamma^5 \lambda^a D(x, x) \} = \frac{1}{2} \text{tr} \{ \gamma^5 \lambda^a, -e \not{A} + M \} D(x, x) + (N_f g^2/16\pi^2) \text{tr} \{ \lambda^a * F^{ij} F_{ij} \}. \quad (20)$$

In order to apply this equation we must restrict the background gauge field so that

$$[\lambda^a, A_i(x)] = 0. \quad (21)$$

We then deduce the general result by requiring both Lorentz and gauge covariance. With (21), Eq. (15) remains valid and repeating the previous steps we find

$$\langle j^a_0(x) \rangle = - (N_f g^2/64\pi^2) \text{tr} \{ \lambda^a * F^{ij} F_{ij} \}. \quad (22)$$

The induced non-Abelian charge is

$$Q^a = - (N_f g^2/64\pi^2) \int d^4x \text{tr} \{ \lambda^a * F^{ij} F_{ij} \}. \quad (23)$$

Notice that we have arrived at a finite result in spite of the nonrenormalizability of the original field theory. The radiatively induced term also has a topological interpretation: It is the Chern-Simons invariant in five dimensions.

In Ref. 11 it has been observed that Witten's anomaly¹⁷ is also present in $(2+1)$ -dimensional $SU(2)$ gauge theory. Consistency with the results of Refs. 9 and 10 requires that the number of Dirac fermions, N_f , must be even. It is straightforward to check from (26) that this is also true in $4+1$ dimensions.

In the case of an Abelian anomaly equation we simply set $\lambda^a = 1$ in (20) and (23) reduces to

$$N = -\frac{1}{4} N_f q, \quad (24)$$

where q is the Pontryagin index of the background field.

By implementing both Lorentz and gauge covariance in (22) we deduce the low-momentum approximation of the current,

$$\langle j^a_\mu(x) \rangle = - (N_f g^2/128\pi^2) \epsilon_{\mu\nu\rho\sigma\delta} \text{tr} \{ \lambda^a F^{\nu\rho} F^{\sigma\delta} \}. \quad (25)$$

This current then yields the Euler-Heisenberg effective action¹⁶

$$\Gamma = \int d^5x \left\{ -\frac{1}{4} \text{tr} [F^{\mu\nu} F_{\mu\nu}] - (N_f g^2/96\pi^2) \epsilon^{\alpha\beta\gamma\delta\eta} A_\alpha \partial_\beta A_\gamma \partial_\delta A_\eta + \frac{3}{2} A_\alpha A_\beta A_\gamma \partial_\delta A_\eta + \frac{3}{5} A_\alpha A_\beta A_\gamma A_\delta A_\eta \right\}. \quad (26)$$

The analysis presented here together with the results in Ref. 11 then suggest the following relations between axial anomalies in general space-time dimensions: In $2n+2$ dimensions the Abelian anomaly term is given by the Pontryagin density which is a total divergence of a current. Take that component of this current that does not belong to a given $(2n+1)$ -dimensional subspace. This gives the $(2n+1)$ -dimensional Chern-Simons invariant.^{9,10} The $2n$ -dimensional non-Abelian anomaly term is then a functional derivative of

the $(2n+1)$ -dimensional Chern-Simons invariant with respect to that component of the gauge field that does not belong to the $2n$ -dimensional subspace. Furthermore, the radiatively induced $(2n+1)$ -dimensional Euler-Heisenberg effective action is the pertinent Chern-Simons invariant which is entirely responsible for the fermion number in that dimensionality. We have shown that this conjecture is true for $n=1,2$ and expect that our result is valid for all integer values of n .¹⁸

The authors thank Roman Jackiw for helpful communications. One of us (G.W.S.) thanks A. N. Redlich for helpful discussions and communicating his unpublished work. He further acknowledges the kind hospitality of the Research Institute for Theoretical Physics of the University of Helsinki where part of this work was done. This work was supported in part by the Natural Sciences and Engineering Research Council of Canada and the Academy of Finland. This work was also supported in part by the U. S. Department of Energy through Grant No. DE-AC02-76ER02220.

After completing this manuscript we became aware of an investigation by B. Zumino, Wu Yang-Shi, and A. Zee (to be published) who study left-right anomalies and arrive at relations between axial anomalies in different dimensions that are very similar to the ones found here for $V-A$ anomalies.

^(a)Present address.

¹R. Jackiw and C. Rebbi, Phys. Rev. D **13**, 3398 (1976).

²W. P. Su, J. R. Schrieffer, and A. J. Heeger, Phys. Rev. Lett. **42**, 1698 (1979), and Phys. Rev. B **22**, 2099 (1980).

³R. Jackiw and J. R. Schrieffer, Nucl. Phys. **B190** [FS3], 253 (1981).

⁴J. Goldstone and F. Wilczek, Phys. Rev. Lett. **47**, 986 (1981).

⁵W. A. Bardeen, S. Elitzur, Y. Frishman, and E. Rabinovici, Nucl. Phys. **B218**, 445 (1983).

⁶A. J. Niemi and G. W. Semenoff, MIT Report No. 1086, 1983 (to be published).

⁷M. B. Paranjape and G. W. Semenoff, MIT Report No. 1091, 1983 (to be published).

⁸R. Jackiw, in *Quantum Structure of Space and Time*, edited by M. J. Duff and C. Isham (Cambridge Univ. Press, Cambridge, England, 1982).

⁹S. Deser, R. Jackiw, and S. Templeton, Phys. Rev. Lett. **48**, 975 (1982), and Ann. Phys. (N.Y.) **140**, 372 (1982).

¹⁰R. Jackiw, in *Gauge Theories of the Eighties*, edited by R. Raitio and J. Lindfors (Springer-Verlag, Berlin, 1983).

¹¹A. N. Redlich, MIT Report No. 1107, 1983 (to be published).

¹²S. L. Adler, in *Lectures on Elementary Particles and Quantum Field Theory*, edited by S. Deser, M. Grisaru, and H. Pendleton (MIT Press, Cambridge, Mass., 1970); S. Treiman, R. Jackiw, and D. Gross, *Lectures on Current Algebra and Its Applications* (Princeton Univ. Press, Princeton, N.J., 1972).

¹³A. M. Polyakov, Nucl. Phys. **B120**, 429 (1977).

¹⁴R. P. Feynman, Nucl. Phys. **B188**, 479 (1981).

¹⁵W. A. Bardeen, Phys. Rev. **184**, 1848 (1969); we have set the axial background gauge field to zero.

¹⁶P. H. Frampton and T. W. Kephart, Phys. Rev. Lett. **50**, 1347 (1983); our result does not agree with the one deduced from this reference.

¹⁷E. Witten, Phys. Lett. **117B**, 324 (1982).

¹⁸For $n=1$, see R. Jackiw, Rev. Mod. Phys. **52**, 661 (1980).