

**Low-Energy Effective Action
of the Supersymmetric $CP(N - 1)$ Model
in the Large- N Limit**

Pavel A. Bolokhov^a, Mikhail Shifman^b and Alexei Yung^{b,c}

*^aTheoretical Physics Department, St.Petersburg State University, Ulyanovskaya 1,
Peterhof, St.Petersburg, 198504, Russia*

*^bWilliam I. Fine Theoretical Physics Institute, University of Minnesota,
Minneapolis, MN 55455, USA*

^cPetersburg Nuclear Physics Institute, Gatchina, St. Petersburg 188300, Russia

Abstract

We perform a supersymmetric completion ...

1 Introduction

2 Supersymmetrization

Our potential is,

$$V_{\text{eff}} = -\frac{N}{4\pi} \left\{ (|\sqrt{2}\sigma|^2 + iD) \log (|\sqrt{2}\sigma|^2 + iD) - iD - |\sqrt{2}\sigma|^2 \log |\sqrt{2}\sigma|^2 \right\}. \quad (2.1)$$

It can be cast in integral forms which will prove useful below,

$$\begin{aligned} \frac{4\pi}{N} V_{\text{eff}} &= - \int_0^{iD} dt \ln (|\sqrt{2}\sigma|^2 + t) = \\ &= - |\sqrt{2}\sigma|^2 \left(\ln |\sqrt{2}\sigma|^2 \cdot x + \int_0^x \ln (1+x) dx \right). \end{aligned} \quad (2.2)$$

It can also be represented as a series,

$$\frac{4\pi}{N} V_{\text{eff}} = -iD \ln |\sqrt{2}\sigma|^2 + \sum_{k \geq 1} \frac{(-1)^k}{k(k+1)} iD \left(\frac{iD}{|\sqrt{2}\sigma|^2} \right)^k. \quad (2.3)$$

The first term comes from Witten's superpotential. The first term of the series (with $k = 1$) is taken into account by the kinetic term. Thus we can write our effective action as

$$\begin{aligned} \frac{4\pi}{N} \mathcal{L} &= - \int d^4\theta \frac{1}{2} \left| \ln \sqrt{2}\Sigma \right|^2 - i \int d^2\tilde{\theta} \left(\sqrt{2}\Sigma \ln \sqrt{2}\Sigma - \sqrt{2}\Sigma \right) + \\ &+ \frac{i}{2} \int d^2\tilde{\theta} S \sum_{k \geq 1} \frac{(-1)^k}{k(k+1)(k+2)} \left(\frac{S}{\sqrt{2}\Sigma} \right)^k + \text{h.c.}, \end{aligned} \quad (2.4)$$

where

$$S = \frac{i}{2} \overline{D}_R D_L \ln \sqrt{2}\Sigma, \quad \overline{S} = \frac{i}{2} \overline{D}_L D_R \ln \sqrt{2}\Sigma. \quad (2.5)$$

The series in the second line can be written as a D -term. We get

$$\begin{aligned} \frac{4\pi}{N} \mathcal{L} &= - \int d^4\theta \frac{1}{2} \left| \ln \sqrt{2}\Sigma \right|^2 - i \int d^2\tilde{\theta} \left(\sqrt{2}\Sigma \ln \sqrt{2}\Sigma - \sqrt{2}\Sigma \right) + \\ &+ \frac{1}{2} \int d^4\theta \ln \sqrt{2}\Sigma \sum_{k \geq 1} \frac{(-1)^k}{k(k+1)(k+2)} \left(\frac{S}{\sqrt{2}\Sigma} \right)^k + \text{h.c.} \end{aligned} \quad (2.6)$$

Here we can see explicitly that the only F -term is the Witten's superpotential, while all other terms are D -terms.

The last term can also be re-written as an analytic expression,

$$\begin{aligned} \frac{4\pi}{N} \mathcal{L} = & - \int d^4\theta \frac{1}{2} \left| \ln \sqrt{2}\Sigma \right|^2 - i \int d^2\tilde{\theta} \left(\sqrt{2}\Sigma \ln \sqrt{2}\Sigma - \sqrt{2}\Sigma \right) - \\ & - \frac{i}{4} \int d^2\tilde{\theta} \left(\frac{(S + \sqrt{2}\Sigma)^2}{S} \ln \left(1 + \frac{S}{\sqrt{2}\Sigma} \right) - \sqrt{2}\Sigma \right) + \text{h.c.} \quad (2.7) \end{aligned}$$

We wrote this expression containing the logarithm as a (twisted) superpotential, but it can be easily put into a D -term form should a need arise.

Before proceeding with finding the component version of the above expressions, let us first write out the component expansion of the “known” part of the action — that is, of the Witten's superpotential and the “superkinetic” term,

$$\begin{aligned} \frac{4\pi}{N} \mathcal{L}_{(0)} = & - \int d^4\theta \left(\frac{1}{2} \left| \ln \sqrt{2}\Sigma \right|^2 + \right. \\ & \left. + i d^2\tilde{\theta} \left(\sqrt{2}\Sigma \ln \sqrt{2}\Sigma - \sqrt{2}\Sigma \right) + i d^2\bar{\tilde{\theta}} \left(\sqrt{2}\bar{\Sigma} \ln \sqrt{2}\bar{\Sigma} - \sqrt{2}\bar{\Sigma} \right) \right) = \\ = & - iD \log |\sqrt{2}\sigma|^2 - \frac{1}{2} \frac{(iD)^2}{|\sqrt{2}\sigma|^2} - F_{03} \log \frac{\sqrt{2}\sigma}{\sqrt{2}\bar{\sigma}} + \\ & + \frac{|\partial_\mu \sigma|^2 + \frac{1}{2} F_{03}^2 + \frac{1}{2} \left(\bar{\lambda}_R i \overleftrightarrow{\mathcal{D}}_L \lambda_R + \bar{\lambda}_L i \overleftrightarrow{\mathcal{D}}_R \lambda_L \right)}{|\sqrt{2}\sigma|^2} - \quad (2.8) \\ & - 2 \frac{i\sqrt{2}\sigma \bar{\lambda}_R \lambda_L + i\sqrt{2}\bar{\sigma} \bar{\lambda}_L \lambda_R}{|\sqrt{2}\sigma|^2} + 2 \frac{\bar{\lambda}_R \lambda_L \bar{\lambda}_L \lambda_R}{|\sqrt{2}\sigma|^4} + \\ & + \frac{(iD + F_{03}) i\sqrt{2}\sigma \bar{\lambda}_R \lambda_L + (iD - F_{03}) i\sqrt{2}\bar{\sigma} \bar{\lambda}_L \lambda_R}{|\sqrt{2}\sigma|^4}. \end{aligned}$$

The long derivatives here are (*not sure why they have different signs, shouldn't*)

$$\begin{aligned} \mathcal{D}_L &= \partial_L - \partial_L \ln \sqrt{2}\sigma \\ \mathcal{D}_R &= \partial_R + \partial_R \ln \sqrt{2}\bar{\sigma}, \quad (2.9) \end{aligned}$$

with conjugate expressions for the left-ward derivatives $\overleftarrow{\mathcal{D}}_L$ and $\overleftarrow{\mathcal{D}}_R$. These derivatives include the Christoffel symbol $1/\sigma$ which arises from the fact that we have a Kähler potential $|\log \Sigma|^2$. In coordinates $\Sigma, \overline{\Sigma}$ the latter potential generates a metric $1/|\sigma|^2$ which is seen in the denominators in the expressions above. However, the Riemann tensor is zero for this metric, reflecting the observation that the target space is in fact flat — if one adopts $\log \Sigma$ and $\log \overline{\Sigma}$ as the coordinates, the metric disappears completely. Still, we prefer to work with the coordinates Σ and $\overline{\Sigma}$.

The first two terms in the component expansion in (2.8) form the leading part of the effective potential (2.3). The third term is the anomaly, *may need more discussion*. The terms on the fourth line in (2.8) are the kinetic terms, while the terms on the rest of the lines are Yukawa-like and quartic couplings.

We now calculate the component expansion of the series term in (2.4) in the two-derivative approximation. That is, we only retain terms with no more than two spacetime derivatives for bosons, and terms with no more than one spacetime derivative for fermions. We do so by performing the superspace integration on each of the terms in the series and then assembling the terms into logarithms. Rather than listing the resulting expression for the series by itself, we present the component

expression for the whole Lagrangian (2.4), which appears more compact:

$$\begin{aligned}
\mathcal{L}_{\text{two deriv}} &= \frac{|\partial_\mu \sigma|^2}{|\sqrt{2}\sigma|^2} - F_{03} \log \frac{\sqrt{2}\sigma}{\sqrt{2}\bar{\sigma}} + V_{\text{eff}} - \\
&- \frac{F_{03}^2}{|\sqrt{2}\sigma|^2} \left(2 \frac{\ln(1+x) - x}{x^2} + \frac{1}{2} \frac{1}{1+x} \right) - \\
&- \frac{\bar{\lambda}_R i \overleftrightarrow{\mathcal{D}}_L \lambda_R + \bar{\lambda}_L i \overleftrightarrow{\mathcal{D}}_R \lambda_L}{|\sqrt{2}\sigma|^2} \frac{\ln(1+x) - x}{x^2} - \\
&- 2 \frac{i\sqrt{2}\sigma \bar{\lambda}_R \lambda_L + i\sqrt{2}\bar{\sigma} \lambda_L \lambda_R}{|\sqrt{2}\sigma|^2} \frac{\ln(1+x)}{x} + \\
&+ 4 \frac{\bar{\lambda}_R \lambda_L \bar{\lambda}_L \lambda_R}{|\sqrt{2}\sigma|^4} \left(\frac{\ln(1+x) - x}{x^2} + \frac{1}{1+x} \right) + \\
&+ \frac{1}{4} \square \log |\sqrt{2}\sigma|^2 \cdot \frac{(1-x^2) \ln(1+x) - x}{x^2} - \\
&- 2 \frac{F_{03}(i\sqrt{2}\sigma \bar{\lambda}_R \lambda_L - i\sqrt{2}\bar{\sigma} \lambda_L \lambda_R)}{|\sqrt{2}\sigma|^4} \frac{\ln(1+x) - x}{x^2}.
\end{aligned} \tag{2.10}$$

Here we have introduced a shorthand for the series expansion quantity

$$x = \frac{iD}{|\sqrt{2}\sigma|^2}. \tag{2.11}$$

We notice that the effect of adding the series to Eq. (2.8) is the multiplication of all terms by certain functions of x . In fact, the whole expression (2.8) is the leading order approximation (or subleading for some of the terms) in x of the two-derivative Lagrangian (2.10).

- Confirm the form of the long derivatives

3 Conclusion

Acknowledgments

This work is supported in part by DOE grant DE-FG02-94ER40823. The work of A.Y. was supported by FTPI, University of Minnesota, by RFBR Grant No. 13-02-00042a and by Russian State Grant for Scientific Schools RSGSS-657512010.2.

References