

Two-dimensional — four-dimensional correspondence: Kinks and Monopoles in $\mathcal{N} = 2$ theories

Pavel A. Bolokhov

University of Minnesota Duluth

April 8, 2015

Confinement is customary thought of as medium where (chromo)electric charges are bound by flux tubes via condensation of monopoles.

Confinement is customary thought of as medium where (chromo)electric charges are bound by flux tubes via condensation of monopoles.

A number of theories have been proposed, such as the Seiberg-Witten theory, which are shown to have confinement

Confinement is customary thought of as medium where (chromo)electric charges are bound by flux tubes via condensation of monopoles.

A number of theories have been proposed, such as the Seiberg-Witten theory, which are shown to have confinement

However, in these theories, confinement is essentially abelian.

Confinement is customary thought of as medium where (chromo)electric charges are bound by flux tubes via condensation of monopoles.

A number of theories have been proposed, such as the Seiberg-Witten theory, which are shown to have confinement

However, in these theories, confinement is essentially abelian.

This has problems with our experience with QCD. The low-energy group is $U(1)^{N-1}$ which leads to the existence of $N - 1$ Abelian strings, and therefore $N - 1$ infinite towers of mesons.

Confinement is customary thought of as medium where (chromo)electric charges are bound by flux tubes via condensation of monopoles.


A number of theories have been proposed, such as the Seiberg-Witten theory, which are shown to have confinement

However, in these theories, confinement is essentially abelian.


This has problems with our experience with QCD. The low-energy group is $U(1)^{N-1}$ which leads to the existence of $N - 1$ Abelian strings, and therefore $N - 1$ infinite towers of mesons.

In real-world QCD confinement is *non-abelian*.

Certain supersymmetry SQCD scenarios allow condensation of quarks to cause confinement of the monopoles




Certain supersymmetry SQCD scenarios allow condensation of quarks to cause confinement of the monopoles



Vortex strings are allowed to end on a monopole which they therefore confine. The low-energy dynamics of these strings is described by an effective two-dimensional worldsheet theory


Certain supersymmetry SQCD scenarios allow condensation of quarks to cause confinement of the monopoles



Vortex strings are allowed to end on a monopole which they therefore confine. The low-energy dynamics of these strings is described by an effective two-dimensional worldsheet theory

The strings represent the vacua of this theory, while the monopoles play the role of *kinks* of the worldsheet theory

Certain supersymmetry SQCD scenarios allow condensation of quarks to cause confinement of the monopoles



Vortex strings are allowed to end on a monopole which they therefore confine. The low-energy dynamics of these strings is described by an effective two-dimensional worldsheet theory

The strings represent the vacua of this theory, while the monopoles play the role of *kinks* of the worldsheet theory

For the study of confinement of monopoles, therefore, it is instructive to investigate the dynamics of two-dim (typically CP^{N-1}) effective worldsheet theory

Two-dimensional models have interesting similarities to 4-d gauge theories

CP^{N-1} theory has been shown to have chiral symmetry breaking, mass gap, asymptotic freedom, *etc* and all these properties are much easier to show in two dimensions

In general, two-dimensional theories are simpler and two-dimensional methods are more powerful

There is also a two-dimensional — four-dimensional duality of their BPS spectra

$\mathcal{N} = 2$ $N_c = N_f$ SYM in four dimensions at the root of the first baryonic Higgs branch



$\mathcal{N} = (2, 2)$ CP^{N-1} theory in two dimensions



supports non-Abelian vortex strings



kinks interpolate between worldsheet vacua

monopoles



kinks

strings



vacua

At weak coupling the Seiberg-Witten theory contains Quarks,
W-bosons, Monopoles, Dyons and bound states

At strong coupling — we consider CP^{N-1} theory
Strong coupling spectrum is accessible via the mirror theory

CP^{N-1} theory with *twisted* masses:

$$r \left(|\mathcal{D}_\mu n^l|^2 + |\sigma - m^l|^2 |n^l|^2 + i D (|n^l|^2 - 1) + \dots \right) \\ + \frac{1}{4e^2} F_{\mu\nu}^2 + \frac{1}{e^2} |\partial_\mu \sigma|^2 + \frac{1}{2e^2} D^2 + \dots,$$

in the $e^2 \rightarrow \infty$ limit

The theory has N vacua — both classically and exactly

CP^{N-1} theory with *twisted* masses:

$$r \left(|\mathcal{D}_\mu n^l|^2 + |\sigma - m^l|^2 |n^l|^2 + i D (|n^l|^2 - 1) + \dots \right) \\ + \frac{1}{4e^2} F_{\mu\nu}^2 + \frac{1}{e^2} |\partial_\mu \sigma|^2 + \frac{1}{2e^2} D^2 + \dots,$$

in the $e^2 \rightarrow \infty$ limit

CP^{N-1} model with \mathcal{Z}_N twisted masses

$$m_l = m_0 \cdot e^{2\pi i l / N}$$

in this case $\mathcal{Z}_N \subset U_R(1)$ remains unbroken

Exact superpotential

The theory possesses an “exact” superpotential of Veneziano-Yankielowicz type

$$\mathcal{W}_{\text{eff}} = -i\tau\hat{\sigma} + \frac{1}{2\pi} \sum_l (\hat{\sigma} - m_l) \left(\ln \frac{\hat{\sigma} - m_l}{\mu} - 1 \right)$$

μ = UV cut-off scale

Exact superpotential

Vacuum values:

$$\mathcal{W}_{\text{eff}} = -\frac{1}{2\pi} \left(N \sigma_p + \sum_l m_l \ln (\sigma_p - m_l) \right)$$

where

$$\sigma_p = \sigma_0 \cdot e^{2\pi i p/N}$$

$$\sigma_0 = \sqrt[N]{1 + m_0^N}$$

vacuum equation

$$\prod_l (\sigma - m_l) = 1.$$

Mirror dual of \mathbb{CP}^{N-1}

The mirror dual is the affine Toda theory

$$\mathcal{W}_{\text{mirror}}^{\mathbb{CP}^{N-1}} = -\frac{1}{2\pi} \left(x_1 + x_2 + \dots + x_n + \sum m_l \ln x_l \right),$$

$$x_1 x_2 \dots x_n = 1$$

Only the superpotential is known in that theory

The vacuum values coincide with $\mathcal{W}_{\text{eff}}(\sigma_p)$ upon identification

$$x_l^{(p)} = \sigma_p - m_l$$

Based on work of K.Hori and C.Vafa, N mirror kinks can be found at strong coupling

$$|m_k| \leq 1$$

Limiting to the sector interpolating between 0^{th} and 1^{st} vacua

$$\mathcal{Z} = \mathcal{W}(\sigma_1) - \mathcal{W}(\sigma_0) + i m_k, \quad k = 0, \dots, N-1,$$



The spectrum

$$\mathcal{Z} = \mathcal{W}(\sigma_1) - \mathcal{W}(\sigma_0) + i m_k, \quad k = 0, \dots, N-1,$$

describes N dyonic kinks (of either \mathbb{CP}^{N-1} or the mirror theory) in the fundamental of $\text{SU}(N)$

The spectrum

$$\mathcal{Z} = \mathcal{W}(\sigma_1) - \mathcal{W}(\sigma_0) + i m_k, \quad k = 0, \dots, N-1,$$

describes N dyonic kinks (of either \mathbb{CP}^{N-1} or the mirror theory) in the fundamental of $\text{SU}(N)$

Question:

What happens to them at weak coupling?

The spectrum

$$\mathcal{Z} = \mathcal{W}(\sigma_1) - \mathcal{W}(\sigma_0) + i m_k, \quad k = 0, \dots, N-1,$$

describes N dyonic kinks (of either \mathbb{CP}^{N-1} or the mirror theory) in the fundamental of $\text{SU}(N)$

Question:

What happens to them at weak coupling?

What states do they correspond to?

Previously known picture

Weak coupling:

$$\mathcal{Q}_{ik}, \quad \mathcal{M}_{ik}, \quad \mathcal{D}_{ik}^n$$

$$\mathcal{D}_{ik}^n + Q \quad \text{—} \quad \text{bound states}$$

$$\mathcal{Q}_{ik} = i(m_i - m_k)$$

$$\mathcal{M}_{ik} \text{ — purely topological kink interpolating from } (k) \rightarrow (i)$$

$$\mathcal{D}_{ik}^n = \mathcal{M}_{ik} + i n (m_i - m_k) \text{ — tower of dyonic kinks upon quasiclassical quantization}$$

CP^2 theory

We focus on CP^2 model — the first non-trivial theory.

We find the curves of marginal stability (*c.m.s.*) — analogues of wall crossing — for this model, where the spectrum can change due to decays

CP^2 theory


We focus on CP^2 model — the first non-trivial theory.

We find the curves of marginal stability (*c.m.s.*) — analogues of wall crossing — for this model, where the spectrum can change due to decays

c.m.s. are a supersymmetric version of a “phase transition”

there are practically no phase transitions in supersymmetric theories except, perhaps, when supersymmetry is broken, or for theories with $N_c \rightarrow \infty$

Moduli space of \mathbb{CP}^2 — plane of m_0




there are three \mathbb{Z}_3 -equivalent sectors

So we choose only one topological sector:

$$\text{kinks:} \quad (0) \longrightarrow (1)$$

and one sector in m_0 -plane:



the other two sectors are completely equivalent

other kinks have the same masses, just central charges shifted by a phase

Weak coupling spectrum

$$\mathcal{Q}_{ik}$$

$$\mathcal{M}_{10}$$

$$\mathcal{D}_{10}^{(n)}$$

$$\mathcal{D}_{10}^{(n)} \mathcal{Q}_{02}$$

Weak coupling spectrum

decays



\mathcal{Q}_{ik}

\mathcal{M}_{10}

$\mathcal{D}_{10}^{(n)}$

$\mathcal{D}_{10}^{(n)} \mathcal{Q}_{02}$

Weak coupling spectrum

decays



\mathcal{Q}_{ik}

\mathcal{M}_{10}

decay all
but one



$\mathcal{D}_{10}^{(n)}$

$\mathcal{D}_{10}^{(n)} \mathcal{Q}_{02}$

Weak coupling spectrum

decays



\mathcal{Q}_{ik}

\mathcal{M}_{10}

decay all
but one



$\mathcal{D}_{10}^{(n)}$

all decay



$\mathcal{D}_{10}^{(n)} \mathcal{Q}_{02}$

Weak coupling spectrum

decays

 \mathcal{Q}_{ik}

stays

 \mathcal{M}_{10} decay all
but one $\mathcal{D}_{10}^{(n)}$

all decay

 $\mathcal{D}_{10}^{(n)} \mathcal{Q}_{02}$

Strong coupling spectrum

$$\mathcal{W}_1 = \mathcal{W}_0 + i m_0$$

$$\mathcal{W}_1 = \mathcal{W}_0 + i m_1$$

$$\mathcal{W}_1 = \mathcal{W}_0 + i m_2$$

Strong coupling spectrum

$$\mathcal{W}_1 = \mathcal{W}_0 + i m_0$$

$$\mathcal{W}_1 = \mathcal{W}_0 + i m_1$$

$$\mathcal{W}_1 = \mathcal{W}_0 + i m_2 \quad \text{--- the lightest}$$

Strong coupling spectrum

$$\mathcal{W}_1 = \mathcal{W}_0 + i m_0$$

$$\mathcal{W}_1 = \mathcal{W}_0 + i m_1$$

decays \Leftarrow $\mathcal{W}_1 = \mathcal{W}_0 + i m_2$ — the lightest

Strong coupling spectrum

$$\mathcal{W}_1 = \mathcal{W}_0 + i m_0 \quad \text{--- next lightest}$$

$$\mathcal{W}_1 = \mathcal{W}_0 + i m_1$$

$$\text{decays} \iff \mathcal{W}_1 = \mathcal{W}_0 + i m_2 \quad \text{--- the lightest}$$

Strong coupling spectrum

$$\mathcal{M}_{10} \Leftarrow \mathcal{W}_1 - \mathcal{W}_0 + i m_0 \quad \text{--- next lightest}$$

$$\mathcal{W}_1 - \mathcal{W}_0 + i m_1$$

$$\text{decays} \Leftarrow \mathcal{W}_1 - \mathcal{W}_0 + i m_2 \quad \text{--- the lightest}$$

Strong coupling spectrum

$$\mathcal{M}_{10} \iff \mathcal{W}_1 - \mathcal{W}_0 + i m_0 \quad \text{--- next lightest}$$

$$\mathcal{W}_1 - \mathcal{W}_0 + i m_1 \quad \text{--- the heavier}$$

$$\text{decays} \iff \mathcal{W}_1 - \mathcal{W}_0 + i m_2 \quad \text{--- the lightest}$$

Strong coupling spectrum


$$\mathcal{M}_{10} \iff \mathcal{W}_1 - \mathcal{W}_0 + i m_0 \quad \text{--- next lightest}$$

$$\mathcal{D}_{10} \iff \mathcal{W}_1 - \mathcal{W}_0 + i m_1 \quad \text{--- the heavier}$$

$$\text{decays} \iff \mathcal{W}_1 - \mathcal{W}_0 + i m_2 \quad \text{--- the lightest}$$

Strong coupling spectrum

$$\begin{array}{llllll}
 \mathcal{M}_{10} & \Leftarrow & \mathcal{W}_1 - \mathcal{W}_0 + i m_0 & - & \text{next lightest} & \leftarrow \text{build their own tower} \\
 \mathcal{D}_{10} & \Leftarrow & \mathcal{W}_1 - \mathcal{W}_0 + i m_1 & - & \text{the heavier} & \leftarrow \\
 \text{decays} & \Leftarrow & \mathcal{W}_1 - \mathcal{W}_0 + i m_2 & - & \text{the lightest} &
 \end{array}$$

$c.m.s.$ 

Decays

Simple decays:

$$\mathcal{D}^{(n)} Q \longrightarrow \mathcal{D}^{(n)} + Q \quad n < 1$$

$$\mathcal{D}^{(n)} Q \longrightarrow \mathcal{D}^{(n-1)} + \tilde{Q} \quad n > 1$$

Decays

Simple decays:

$$\mathcal{D}^{(n)} Q \longrightarrow \mathcal{D}^{(n)} + Q \quad n < 1$$

$$\mathcal{D}^{(n)} Q \longrightarrow \mathcal{D}^{(n-1)} + \tilde{Q} \quad n > 1$$

Topological decays:

$$\mathcal{D}_{10}^{(1)} Q \longrightarrow \overline{\mathcal{M}}_{02} + \overline{\mathcal{D}}_{21}^{(1)}$$

$$\mathcal{V}_{10}^2 \longrightarrow \overline{\mathcal{D}}_{02} + \overline{\mathcal{M}}_{21}$$

Generalizing to higher N

Two states become part of the “monopole” tower

Generalizing to higher N

Two states become part of the “monopole” tower

All extra states decay

Generalizing to higher N

Two states become part of the “monopole” tower

All extra states decay

(Indeed for even N there are *no bound states*)

Generalizing to higher N

Two states become part of the “monopole” tower

All extra states decay

(Indeed for even N there are *no bound states*)

We can infer that most likely they decay for all N

SQCD in four dimensions

there are also N states at strong coupling

SQCD in four dimensions

there are also N states at strong coupling

our results indicate that two of them are part of one tower of dyons

SQCD in four dimensions

there are also N states at strong coupling

our results indicate that two of them are part of one tower of dyons

other states decay before reaching the weak coupling

Thank you