

Congruence Classes of Supporting the Erdős-Straus Conjecture I: Tame Solutions ¹

Xiaoping Xu

HLM, Institute of Mathematics, Academy of Mathematics & System Sciences
Chinese Academy of Sciences, Beijing 100190, P.R. China
& School of Mathematics, University of Chinese Academy of Sciences,
Beijing 100049, P.R. China

Abstract

In 1948, Erdős and Straus formulated a conjecture : for any positive integer $n > 2$, there exist positive integers n_1, n_2 and n_3 such that

$$\frac{4}{n} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3},$$

which is still open. It is known that one only needs to prove the conjecture for any prime number n such that $n \equiv 1 \pmod{24}$. If $n = 24m + 1$ and $n_1 \leq n_2, n_3$, then $n_1 = 6m + k$ with $1 \leq k \leq 12m$. A solution (n_1, n_2, n_3) of the above equation is called a *tame solution* if n_2 and n_3 are factors of $(6m + k)(24m + 1)$. We call $n = 24m + 1$ *wild* if it does not have any tame solution. In this paper, we derive twenty-eight families of tame solutions of the above equation. Numeric test shows that they cover all the tame primes.

Keywords: Erdős-Straus conjecture; Egyptian fraction; congruence class; tame solution; wild solution; wild prime.

1 Introduction

Ancient Egyptians used sums of unit fractions (whose numerators are 1) to express fractions due to their ways of distributing food. For instance, $5/8$ was interpreted by ancient Egyptians as distributing five pancakes fairly among eight people. They cut first four pancakes into halves and then cut the last one into eight equal pieces. So each person got the same share: $1/2 + 1/8$ pancakes. This amazingly interpreted the mathematical equation

$$\frac{5}{8} = \frac{1}{2} + \frac{1}{8}. \quad (1.1)$$

So an Egyptian fraction is a sum of distinct unit fractions. By repeatedly applying the simple fact

$$\frac{1}{k} = \frac{1}{k+1} + \frac{1}{k(k+1)}, \quad (1.2)$$

one can easily prove that any fraction is a Egyptian fraction. However, it is very difficult to determine if a fraction can be expressed as a sum of fixed number of unit fractions (cf. [3] for an excellent exposition and extensive references). For example, it is difficult to know if a fraction can be written as a sum of two unit fractions (e.g., cf. [5, 6, 7, 10, 16]).

¹2010 Mathematical Subject Classification. Primary 11D68: Secondary 11D85, 11A67, 11B75

In 1948, Erdős and Straus formulated a conjecture : for any positive integer $n > 2$, there exist positive integers n_1, n_2 and n_3 such that

$$\frac{4}{n} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3}, \quad (1.3)$$

which is still open up to now. For each n , the number of solution can go to infinity as n does. Elscholtz and Tao [3] found excellent bounds for it.

Mordell [15] proved that the conjecture holds for positive integers in 834 congruence classes modulo 840. Terzi [23] used computer to verify that the conjecture holds for positive integers in the congruence classes modulo 120120 except 198 classes. Kotsireas [12] verified the conjecture for every $n < 10^{10}$. There are other interesting partial results on the conjecture or related works (e.g., cf. [1, 2, 4, 11, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25]).

It is known that one only needs to prove the conjecture for any prime number n such that $n \equiv 1 \pmod{24}$, which will be reviewed in next section. Suppose that $n = 24m + 1$ and $n_1 \leq n_2, n_3$. Then $n_1 = 6m + k$ with $1 \leq k \leq 12m$ and

$$\frac{4}{24m + 1} = \frac{1}{6m + k} + \frac{4k - 1}{(6m + k)(24m + 1)}. \quad (1.4)$$

If

$$\frac{1}{n_2} = \frac{\mathfrak{S}_1}{(6m + k)(24m + 1)} \quad \text{and} \quad \frac{1}{n_2} = \frac{\mathfrak{S}_2}{(6m + k)(24m + 1)} \quad (1.5)$$

for some positive integers $\mathfrak{S}_1$ and $\mathfrak{S}_2$ such that $\mathfrak{S}_1 + \mathfrak{S}_2 = 4k - 1$, we call the triple (n_1, n_2, n_3) a *tame solution* of the Erdős-Straus equation (1.3) for the positive integer $n = 24m + 1$. Moreover, we call $n = 24m + 1$ *wild* if it does not have any tame solution. Based on our definition in the earlier version of this work [26], Howerton [8] showed that there are only fourteen wild primes of the form $24m + 1 \leq 2.4 \times 10^{11}$. In this paper, we derive twenty-eight families of tame solutions of Erdős-Straus equation (1.3). Numeric test shows that they cover all the tame primes. Furthermore, we call $\mathfrak{S}_1$ and $\mathfrak{S}_2$ the *numerator summands* for the tame solution (n_1, n_2, n_3) . The paper is organized as follows.

In Section 2, we present some basic known facts about the Erdős-Straus equation and prove that a tame solution (n_1, n_2, n_3) of the equation for a prime $n = 24m + 1$ must satisfy (1.4), (1.5) and $\mathfrak{S}_1, \mathfrak{S}_2$ are factors of $6m + k$. In Section 3, we solve the equation for a prime of the form $24m + 1$ with $\mathfrak{S}_1 = 2(2j + 1)$ and $\mathfrak{S}_2 = 4\ell + 1$. In Section 4, we solve the equation for a prime of the form $24m + 1$ with $\mathfrak{S}_1 = 4(2j + 1)$ and $\mathfrak{S}_2 = 4\ell + 3$. In Section 5, we solve the equation for a prime of the form $24m + 1$ with $\mathfrak{S}_1 = 2^{r+3}(2j + 1)$ and $\mathfrak{S}_2 = 4\ell + 3$.

2 Basics of the Erdős-Straus Equation

In this section, we list some basic known facts about the Erdős-Straus equation (1.3). Then we prove that the numerator summands of the tame solution of the equation for a prime $n = 24m + 1$ must be factors of $6m + k$.

As shown in [21], a simple calculation

$$\frac{4}{2k} = \frac{1}{k} + \frac{1}{k + 1} + \frac{1}{k(k + 1)}, \quad (2.1)$$

$$\frac{4}{3k} = \frac{1}{3k} + \frac{1}{k + 1} + \frac{1}{k(k + 1)}, \quad (2.2)$$

and

$$\frac{4}{3k-1} = \frac{1}{3k-1} + \frac{1}{k} + \frac{1}{k(3k-1)} \quad (2.3)$$

shows that Erdős-Straus equation holds for all n except those $n \equiv 1 \pmod{6}$. Throughout this paper, we denote by $\mathbb{N}$ the set of nonnegative integers. So we only need to solve (1.3) for $n \in 1 + 6\mathbb{N}$. Further calculations

$$\frac{4}{4k-1} = \frac{1}{k} + \frac{1}{k(4k-1)+1} + \frac{1}{k(4k-1)(k(4k-1)+1)} \quad (2.4)$$

and

$$\frac{4}{24k-11} = \frac{1}{6k-2} + \frac{1}{(3k-1)(24k-11)} + \frac{1}{(6k-2)(24k-11)} \quad (2.5)$$

show that Erdős-Straus equation holds for all n except those $n \equiv 1 \pmod{24}$. This fact is known to some people (e.g., cf. [14]). If (n_1, n_2, n_3) is a solution of the Erdős-Straus equation (1.3) for n and k is another positive integer, then

$$\frac{4}{nk} = \frac{1}{n_1k} + \frac{1}{n_2k} + \frac{1}{n_3k} \quad (2.6)$$

naturally holds. So it is enough to solve the equation for any prime number n . From now on, we always assume that

$$n = 24m + 1 \quad \text{is a prime and } m \in \mathbb{N}. \quad (2.7)$$

Moreover, we can assume a tame solution (n_1, n_2, n_3) of (1.3) for n satisfying $n_1 \leq n_2, n_3$. Then we have

$$\frac{4}{3(24m+1)} \leq \frac{1}{n_1} \leq \frac{4}{24m+1}. \quad (2.8)$$

Equivalently

$$6m+1 \leq n_1 \leq 18m. \quad (2.9)$$

Thus

$$n_1 = 6m + k \quad \text{with } 1 \leq k \leq 12m. \quad (2.10)$$

In particular,

$$\frac{4}{24m+1} = \frac{1}{6m+k} + \frac{4k-1}{(6m+k)(24m+1)} \quad (2.11)$$

and

$$\frac{1}{n_2} = \frac{\mathfrak{S}_1}{(6m+k)(24m+1)} \quad \text{and} \quad \frac{1}{n_2} = \frac{\mathfrak{S}_2}{(6m+k)(24m+1)} \quad (2.12)$$

for some positive integers $\mathfrak{S}_1$ and $\mathfrak{S}_2$ such that

$$\mathfrak{S}_1 + \mathfrak{S}_2 = 4k - 1. \quad (2.13)$$

Denote

$$\mathfrak{S} = \text{l.c.m}(\mathfrak{S}_1, \mathfrak{S}_2), \quad (2.14)$$

the least common multiple of $\mathfrak{S}_1$ and $\mathfrak{S}_2$. Let $c \in \mathbb{Z}$ such that $m + c\mathfrak{S} > 0$. Observe

$$(6(m+c\mathfrak{S})+k)(24(m+c\mathfrak{S})+1) = (6m+k)(24m+1) + 6c\mathfrak{S}(48m+24c\mathfrak{S}+4k+1). \quad (2.15)$$

By Assumption (2.12),

$$(6m+k)(24m+1) = n_2\mathfrak{S}_1 = n_3\mathfrak{S}_2. \quad (2.16)$$

The above two expressions imply that

$$\frac{\mathfrak{S}_1}{(6(m+c\mathfrak{S})+k)(24(m+c\mathfrak{S})+1)} \quad \text{and} \quad \frac{\mathfrak{S}_2}{(6(m+c\mathfrak{S})+k)(24(m+c\mathfrak{S})+1)} \quad (2.17)$$

are unit fractions. Moreover,

$$4(6(m+c\mathfrak{S})+k) - (24(m+c\mathfrak{S})+1) = 4(6m+1) - (2m+1) = 4k-1 = \mathfrak{S}_1 + \mathfrak{S}_2. \quad (2.18)$$

Therefore,

$$\begin{aligned} \frac{4}{24(m+c\mathfrak{S})+1} &= \frac{1}{6(m+c\mathfrak{S})+k} + \frac{\mathfrak{S}_1}{(6(m+c\mathfrak{S})+k)(24(m+c\mathfrak{S})+1)} \\ &\quad + \frac{\mathfrak{S}_1}{(6(m+c\mathfrak{S})+k)(24(m+c\mathfrak{S})+1)} \end{aligned} \quad (2.19)$$

is a solution of the Erdős-Straus equation (1.3); that is,

$$\{24(m+c\mathfrak{S})+1 \mid c \in \mathbb{Z} \text{ such that } m+c\mathfrak{S} > 0\} \quad \text{are tame numbers.} \quad (2.20)$$

This partially explains why Mordell [15] and Terzi [23] had their modulo conditions.

Note that

$$4k-1 \leq 48m-1 \quad (2.21)$$

by (2.10). Suppose $\mathfrak{S}_i \nmid (6m+k)$. Since $24m+1$ is a prime, we must have $(24m+1) \mid \mathfrak{S}_i$. So the above equation yields

$$\mathfrak{S}_i = 24m+1. \quad (2.22)$$

Without loss of generality, we may assume $i = 2$. Then

$$\mathfrak{S}_1 = 4k-1 - \mathfrak{S}_2 = 4k-24m-2 \leq 24m-2. \quad (2.23)$$

Thus

$$\mathfrak{S}_1 \mid (6m+k). \quad (2.24)$$

According to (2.23), we can write

$$\mathfrak{S}_1 = 4j+2 \quad \text{with } j \in \mathbb{N}. \quad (2.25)$$

Based on (2.11) and (2.13), we have

$$\mathfrak{S}_1 + 24m+1 = \mathfrak{S}_1 + \mathfrak{S}_2 = 4k-1 = 4(6m+k) - (24m+1). \quad (2.26)$$

By (2.24),

$$\mathfrak{S}_1 \mid [2(24m+1)] \implies (2j+1) \mid (24m+1). \quad (2.27)$$

If $j = 0$, $24m+3 = 4k-1 \implies k = 6m+1$ is odd, which contradicts (2.24) and (2.25). So $j > 0$. Then (2.23) and (2.25) show that $2j+1 \geq 3$ is a proper factor of $24m+1$. This contradicts the assumption that $24m+1$ is a prime.

Theorem 2.1 *For a prime n of the form $24m+1$, its tame solution (n_1, n_2, n_3) must satisfies (2.10)-(2.13) and*

$$\mathfrak{S}_1 \mid (6m+k), \quad \mathfrak{S}_2 \mid (6m+k). \quad (2.28)$$

Since

$$\mathfrak{S}_1 + \mathfrak{S}_2 = 4k - 1 = 4(6m + k) - (24m + 1) \quad (2.29)$$

and $n = 24m + 1$ is a prime, we must have

$$\text{g.c.d}(\mathfrak{S}_1, \mathfrak{S}_2) = 1. \quad (2.30)$$

In the rest of this paper, we always derive a necessary condition of m on the tameness of a prime $24m + 1$, but the condition is also sufficient for any m satisfying the condition such that $24m + 1$ is tame.

3 Cases with the $\mathfrak{S}_1 = 2(2j + 1)$ and $\mathfrak{S}_2 = 4\ell + 1$

First we have

$$k = j + \ell + 1 \quad (3.1)$$

by (2.13). Moreover,

$$6m + k = 6m + j + \ell + 1. \quad (3.2)$$

According to the first expression in (2.28),

$$2 | (6m + j + \ell + 1). \quad (3.3)$$

So

$$j + \ell \equiv 1 \pmod{2}. \quad (3.4)$$

3.1 Case $j = 2s$ and $\ell = 6t + 1$

In this case,

$$\mathfrak{S}_1 = 2(4s + 1), \quad \mathfrak{S}_2 = 24t + 5 \quad (3.5)$$

and

$$6m + j + \ell + 1 = 2(3(m + t) + s + 1) \quad (3.6)$$

According to the first expression in (2.28),

$$(4s + 1) | (3(m + t) + s + 1); \quad (3.7)$$

that is,

$$(3(m + t) + s + 1) \equiv 0 \pmod{4s + 1}. \quad (3.8)$$

So

$$3(m + t) + s + 1 \equiv 4s + 1 \pmod{12s + 1}. \quad (3.9)$$

Equivalently,

$$3(m + t) \equiv 3s \pmod{4s + 1}. \quad (3.10)$$

Subcase 1. $s \not\equiv 2 \pmod{3}$.

In this subcase, $4s + 1 \not\equiv 2 \pmod{3}$ and the above equation implies

$$m + t \equiv s \pmod{4s + 1}; \quad (3.11)$$

that is,

$$m + t = s + c(4s + 1) \quad \text{for some } c \in \mathbb{N}. \quad (3.12)$$

So

$$6m + j + \ell + 1 = 2(3c + 1)(4s + 1). \quad (3.13)$$

According to the second expression in (2.28) and (2.30),

$$(24t + 5)|(3c + 1) \implies c = 16t + 3 + d(24t + 5) \quad \text{for } d \in \mathbb{N}. \quad (3.14)$$

Thus

$$m = s - t + (4s + 1)[16t + 3 + d(24t + 5)] \quad \text{for some } c \in \mathbb{N}. \quad (3.15)$$

and

$$6m + j + \ell + 1 = 2(3d + 2)(4s + 1)(24t + 5). \quad (3.16)$$

Expression (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{2(3d + 2)(4s + 1)(24t + 5)} \\ &+ \frac{2(4s + 1) + (24t + 5)}{2(3d + 2)(4s + 1)(24t + 5)(24m + 1)} \\ &= \frac{1}{2(3d + 2)(4s + 1)(24t + 5)} + \frac{1}{(3d + 2)(24t + 5)(24m + 1)} \\ &+ \frac{1}{2(3d + 2)(4s + 1)(24m + 1)}. \end{aligned} \quad (3.17)$$

Subcase 2. $s = 3s_1 + 2$ with $s_1 \in \mathbb{N}$.

In this subcase,

$$4s + 1 = 12s_1 + 9 = 3(4s_1 + 3). \quad (3.18)$$

So (3.10) is equivalent to

$$3(m + t) \equiv 9s_1 + 6 \pmod{3(4s_1 + 3)}, \quad (3.19)$$

which yields

$$m + t \equiv 3s_1 + 2 \pmod{4s_1 + 3}; \quad (3.20)$$

that is,

$$m + t = 3s_1 + 2 + c(4s_1 + 3) \quad \text{for some } c \in \mathbb{N}. \quad (3.21)$$

So

$$6m + j + \ell + 1 = 6(c + 1)(4s + 1). \quad (3.22)$$

According to the second expression in (2.28) and (2.30),

$$(24t + 5)|(c + 1) \implies c = -1 + (d + 1)(24t + 5) \quad \text{for } d \in \mathbb{N}. \quad (3.23)$$

Thus

$$m = -s_1 - t - 1 + (d + 1)(4s_1 + 3)(24t + 5) \quad (3.24)$$

$$\mathfrak{S}_1 = 6(4s_1 + 3) \quad (3.25)$$

and

$$6m + j + \ell + 1 = 6(d + 1)(4s_1 + 3)(24t + 5). \quad (3.26)$$

Expression (2.11) becomes

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6(d+1)(4s_1+3)(24t+5)} + \frac{6(4s_1+3) + (24t+5)}{6(d+1)(4s_1+3)(24t+5)(24m+1)} \\
&= \frac{1}{6(d+1)(4s_1+3)(24t+5)} + \frac{1}{(d+1)(24t+5)(24m+1)} \\
&\quad + \frac{1}{6(d+1)(4s_1+3)(24m+1)}. \tag{3.27}
\end{aligned}$$

Theorem 3.1 *For any positive integer m in (3.15), we have the tame solution (3.17) of the Erdős-Straus equation. When m is in (3.24), we have the tame solution (3.27) of the Erdős-Straus equation.*

3.2 Case $j = 6s$ and $\ell = 6t + 3$

In this case,

$$\mathfrak{S}_1 = 2(12s + 1), \quad \mathfrak{S}_2 = 24t + 13 \tag{3.28}$$

by (3.1), and

$$6m + j + \ell + 1 = 2(3(m + s + t) + 2) \tag{3.29}$$

According to the first expression in (2.28),

$$(12s + 1) | (3(m + s + t) + 2); \tag{3.30}$$

that is,

$$3(m + s + t) + 2 \equiv 0 \pmod{12s + 1}. \tag{3.31}$$

So

$$3(m + s + t) + 2 \equiv 2(12s + 1) \pmod{12s + 1}. \tag{3.32}$$

Equivalently,

$$3(m + t) \equiv 21s \pmod{12s + 1}. \tag{3.33}$$

Hence

$$m + t \equiv 7s \pmod{12s + 1}. \tag{3.34}$$

Under this condition,

$$m + t = 7s + c(12s + 1) \quad \text{for some } c \in \mathbb{N} \tag{3.35}$$

and

$$6m + j + \ell + 1 = 2(3c + 2)(12s + 1). \tag{3.36}$$

According to the second expression in (2.28) and (2.30),

$$(24t + 13) | (3c + 2) \implies c = 16t + 8 + d(24t + 13) \quad \text{for } d \in \mathbb{N}. \tag{3.37}$$

Now

$$m = 7s - t + (16t + 8 + d(24t + 13))(12s + 1) \tag{3.38}$$

and

$$6m + j + \ell + 1 = 2(3d + 2)(12s + 1)(24t + 13). \tag{3.39}$$

Expression (2.11) becomes

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{2(3d+2)(12s+1)(24t+13)} + \frac{2(12s+1) + (24t+13)}{2(3d+2)(12s+1)(24t+13)(24m+1)} \\
&= \frac{1}{2(3d+2)(12s+1)(24t+13)} + \frac{1}{(3d+2)(24t+13)(24m+1)} \\
&\quad + \frac{1}{2(3d+2)(12s+1)(24m+1)}. \tag{3.40}
\end{aligned}$$

Theorem 3.2 *For any positive integer m in (3.38), we have the tame solution (3.40) of the Erdős-Straus equation.*

3.3 Case $j = 6s$ and $\ell = 6t + 5$

In this case,

$$\mathfrak{S}_1 = 2(12s+1), \quad \mathfrak{S}_2 = 24t+21 = 3(8t+7) \tag{3.41}$$

by (4.1), and

$$6m + j + \ell + 1 = 6(m + s + t + 1) \tag{3.42}$$

According to the first expression in (2.28),

$$(12s+1) | (m + s + t + 1); \tag{3.43}$$

that is,

$$m = -s - t - 1 + c(12s+1) \quad \text{for some } c \in \mathbb{N}. \tag{3.44}$$

Moreover,

$$6m + j + \ell + 1 = 6c(12s+1). \tag{3.45}$$

According to the second expression in (2.28) and (2.30),

$$(24t+21) | [6c(12s+1)] \implies (8t+7) | c \implies c = (d+1)(8t+7) \tag{3.46}$$

for some $d \in \mathbb{N}$. Thus

$$m = -s - t - 1 + (d+1)(12s+1)(8t+7) \tag{3.47}$$

and

$$6m + j + \ell + 1 = 6(d+1)(12s+1)(8t+7). \tag{3.48}$$

Expression (2.11) becomes

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6(d+1)(12s+1)(8t+7)} \\
&\quad + \frac{2(12s+1) + 3(8t+7)}{6(d+1)(12s+1)(8t+7)(24m+1)} \\
&= \frac{1}{6(d+1)(12s+1)(8t+7)} + \frac{1}{3(d+1)(8t+7)(24m+1)} \\
&\quad + \frac{1}{2(d+1)(12s+1)(24m+1)}. \tag{3.49}
\end{aligned}$$

Theorem 3.3 *For any positive integer m in (3.47), we have the tame solution (3.49) of the Erdős-Straus equation.*

3.4 Case $j = 6s + 1$ and $\ell = 6t + 4$

In this case,

$$\mathfrak{S}_1 = 6(4s + 1), \quad \mathfrak{S}_2 = 24t + 17 \quad (3.50)$$

by (3.1), and

$$6m + j + \ell + 1 = 6(m + s + t + 1) \quad (3.51)$$

According to the first expression in (2.28),

$$(4s + 1)|(m + s + t + 1); \quad (3.52)$$

that is,

$$m = -s - t - 1 + c(4s + 1) \quad \text{for some } c \in \mathbb{N}. \quad (3.53)$$

Moreover,

$$6m + j + \ell + 1 = 6c(4s + 1). \quad (3.54)$$

According to the second expression in (2.28) and (2.30),

$$(24t + 17)|[6c(4s + 1)] \implies (24t + 17)|c \implies c = (d + 1)(24t + 17) \quad (3.55)$$

for some $d \in \mathbb{N}$. Thus

$$m = -s - t - 1 + (d + 1)(4s + 1)(24t + 17) \quad (3.56)$$

and

$$6m + j + \ell + 1 = 6(d + 1)(4s + 1)(24t + 17). \quad (3.57)$$

Expression (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6(d + 1)(4s + 1)(24t + 17)} + \frac{6(4s + 1) + (24t + 17)}{6(d + 1)(4s + 1)(24t + 17)(24m + 1)} \\ &= \frac{1}{6(d + 1)(4s + 1)(24t + 17)} + \frac{1}{(d + 1)(24t + 17)(24m + 1)} \\ &\quad + \frac{1}{6(d + 1)(4s + 1)(24m + 1)}. \end{aligned} \quad (3.58)$$

Theorem 3.4 *For any positive integer m in (3.56), we have the tame solution (3.58) of the Erdős-Straus equation.*

3.5 Case $j = 6s + 2$ and $\ell = 6t + 3$

In this case,

$$\mathfrak{S}_1 = 2(12s + 5), \quad \mathfrak{S}_2 = 24t + 13 \quad (3.59)$$

by (3.1), and

$$6m + j + \ell + 1 = 6(m + s + t + 1) \quad (3.60)$$

According to the first expression in (2.28),

$$(12s + 5)|(m + s + t + 1); \quad (3.61)$$

that is,

$$m = -s - t - 1 + c(12s + 5) \quad \text{for some } c \in \mathbb{N}. \quad (3.62)$$

Moreover,

$$6m + j + \ell + 1 = 6c(12s + 5). \quad (3.63)$$

According to the second expression in (2.28) and (2.30),

$$(24t + 13)|c \implies c = (d + 1)(24t + 13) \quad (3.64)$$

for some $d \in \mathbb{N}$. Thus

$$m = -s - t - 1 + (d + 1)(12s + 5)(24t + 13) \quad (3.65)$$

and

$$6m + j + \ell + 1 = 6(d + 1)(12s + 5)(24t + 13). \quad (3.66)$$

Expression (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6(d + 1)(12s + 5)(24t + 13)} \\ &+ \frac{2(12s + 5) + (24t + 13)}{6(d + 1)(12s + 5)(24t + 13)(24m + 1)} \\ &= \frac{1}{6(d + 1)(12s + 5)(24t + 13)} + \frac{1}{3(d + 1)(24t + 13)(24m + 1)} \\ &+ \frac{1}{6(d + 1)(12s + 5)(24m + 1)}. \end{aligned} \quad (3.67)$$

Theorem 3.5 *For any positive integer m in (3.65), we have the tame solution (3.67) of the Erdős-Straus equation.*

3.6 Case $j = 6s + 3$ and $\ell = 6t$

In this case,

$$\mathfrak{S}_1 = 2(12s + 7), \quad \mathfrak{S}_2 = 24t + 1 \quad (3.68)$$

by (3.1), and

$$6m + j + \ell + 1 = 2(3(m + s + t) + 2). \quad (3.69)$$

According to the first expression in (2.28),

$$(12s + 7)|(3(m + s + t) + 2); \quad (3.70)$$

that is,

$$3(m + s + t) + 2 \equiv 0 \pmod{12s + 7}. \quad (3.71)$$

So

$$3(m + s + t) + 2 \equiv 2(12s + 7) \pmod{12s + 7}. \quad (3.72)$$

Equivalently,

$$3(m + t) \equiv 21s + 12 \pmod{12s + 7}. \quad (3.73)$$

So

$$m + t \equiv 7s + 4 \pmod{12s + 7}. \quad (3.74)$$

Under this condition,

$$m + t = 7s + 4 + c(12s + 7) \quad \text{for some } c \in \mathbb{N} \quad (3.75)$$

and

$$6m + j + \ell + 1 = 2(3c + 2)(12s + 7). \quad (3.76)$$

The second expression in (2.28) and (2.30) imply

$$(24t + 1)|(3c + 2) \implies c = 16t + d(24t + 1) \quad \text{for } d \in \mathbb{N}. \quad (3.77)$$

Now

$$m = 7s - t + 4 + (16t + d(24t + 1))(12s + 7). \quad (3.78)$$

and

$$6m + j + \ell + 1 = 2(3d + 2)(12s + 7)(24t + 1). \quad (3.79)$$

Expression (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{2(3d + 2)(12s + 7)(24t + 1)} \\ &+ \frac{2(12s + 7) + (24t + 1)}{2(3d + 2)(12s + 7)(24t + 1)(24m + 1)} \\ &= \frac{1}{2(3d + 2)(12s + 7)(24t + 1)} + \frac{1}{(3d + 2)(24t + 1)(24m + 1)} \\ &+ \frac{1}{2(3d + 2)(12s + 7)(24m + 1)}. \end{aligned} \quad (3.80)$$

Theorem 3.6 *For any positive integer m in (3.78), we have the tame solution (3.80) of the Erdős-Straus equation.*

3.7 Case $j = 6s + 3$ and $\ell = 6t + 2$

In this case,

$$\mathfrak{S}_1 = 2(12s + 7), \quad \mathfrak{S}_2 = 24t + 9 = 3(8t + 3) \quad (3.81)$$

by (3.1), and

$$6m + j + \ell + 1 = 6(m + s + t + 1) \quad (3.82)$$

According to the first expression in (2.28),

$$(12s + 7)|(m + s + t + 1); \quad (3.83)$$

that is,

$$m + s + t + 1 \equiv 0 \pmod{12s + 7}. \quad (3.84)$$

So

$$m + t \equiv 11s + 6 \pmod{12s + 7}. \quad (3.85)$$

Under this condition,

$$m + t = 11s + 6 + c(12s + 7) \quad \text{for some } c \in \mathbb{N} \quad (3.86)$$

and

$$6m + j + \ell + 1 = 6(c + 1)(12s + 7). \quad (3.87)$$

The second expression in (2.28) and (2.30) imply

$$(8t + 3)|(c + 1). \quad (3.88)$$

Thus

$$c = (d+1)(8t+3) - 1 \quad \text{for } 0 < d \in \mathbb{N}. \quad (3.89)$$

Now

$$m = -t - s - 1 + (d+1)(12s+7)(8t+3) \quad (3.90)$$

and

$$6m + j + \ell + 1 = 6(d+1)(12s+7)(8t+3). \quad (3.91)$$

Expression (2.11) becomes

$$\begin{aligned} \frac{4}{24m+1} &= \frac{1}{6(d+1)(12s+7)(8t+3)} \\ &\quad + \frac{2(12s+7)+3(8t+3)}{6(d+1)(12s+7)(8t+3)(24m+1)} \\ &= \frac{1}{6(d+1)(12s+7)(8t+3)} + \frac{1}{3(d+1)(8t+3)(24m+1)} \\ &\quad + \frac{1}{2(d+1)(12s+7)(24m+1)}. \end{aligned} \quad (3.92)$$

Theorem 3.7 *For any positive integer m in (3.90), we have the tame solution (3.92) of the Erdős-Straus equation.*

3.8 Case $j = 6s + 3$ and $\ell = 6t + 4$

In this case,

$$\mathfrak{S}_1 = 2(12s+7), \quad \mathfrak{S}_2 = 24t+17 \quad (3.93)$$

by (3.1), and

$$6m + j + \ell + 1 = 2(3(m+s+t) + 4) \quad (3.94)$$

According to the first expression in (2.28),

$$(12s+7)|(3(m+s+t) + 4); \quad (3.95)$$

that is,

$$3(m+s+t) + 4 \equiv 0 \pmod{12s+7}. \quad (3.96)$$

So

$$3(m+s+t) + 4 \equiv 12s+7 \pmod{12s+7}. \quad (3.97)$$

Equivalently,

$$3(m+t) \equiv 9s+3 \pmod{12s+7}. \quad (3.98)$$

So

$$m+t \equiv 3s+1 \pmod{12s+7}. \quad (3.99)$$

Under this condition,

$$m+t = 3s+1 + c(12s+7) \quad \text{for some } c \in \mathbb{N} \quad (3.100)$$

and

$$6m + j + \ell + 1 = 2(3c+1)(12s+7). \quad (3.101)$$

According to the second expression in (2.28) and (2.30),

$$(24t + 17)|(3c + 1) \implies c = 16t + 11 + d(24t + 17) \quad \text{for } d \in \mathbb{N}. \quad (3.102)$$

Now

$$m = 3s + 1 - t + (16t + 11 + d(24t + 17))(12s + 7) \quad (3.103)$$

and

$$6m + j + \ell + 1 = 2(3d + 2)(12s + 7)(24t + 17). \quad (3.104)$$

Expression (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{2(3d + 2)(12s + 7)(24t + 17)} \\ &+ \frac{2(12s + 7) + (24t + 17)}{2(3d + 2)(12s + 7)(24t + 17)(24m + 1)} \\ &= \frac{1}{2(3d + 2)(12s + 7)(24t + 17)} + \frac{1}{(3d + 2)(24t + 17)(24m + 1)} \\ &+ \frac{1}{2(3d + 2)(12s + 7)(24m + 1)}. \end{aligned} \quad (3.105)$$

Theorem 3.8 *For any positive integer m in (3.103), we have the tame solution (3.105) of the Erdős-Straus equation.*

3.9 Case $j = 6s + 5$ and $\ell = 6t$

In this case,

$$\mathfrak{S}_1 = 2(12s + 11), \quad \mathfrak{S}_2 = 24t + 1 \quad (3.106)$$

by (3.1), and

$$6m + j + \ell + 1 = 6(m + s + t + 1) \quad (3.107)$$

According to the first expression in (2.28),

$$(12s + 11)|(m + s + t + 1); \quad (3.108)$$

that is,

$$m + s + t + 1 \equiv 0 \pmod{12s + 11}. \quad (3.109)$$

So

$$m \equiv -s - t - 1 \pmod{12s + 11}. \quad (3.110)$$

Under this condition,

$$m = -s - t - 1 + c(12s + 11) \quad \text{for some } c \in \mathbb{N} \quad (3.111)$$

and

$$6m + j + \ell + 1 = 6c(12s + 11). \quad (3.112)$$

The second expression in (2.28) and (2.30) imply

$$(24t + 1)|c \implies c = (d + 1)(24t + 1) \quad \text{for } 0 < d \in \mathbb{N}. \quad (3.113)$$

Now

$$m = -s - t - 1 + (d + 1)(24t + 1)(12s + 11). \quad (3.114)$$

and

$$6m + j + \ell + 1 = 6(d+1)(12s+11)(24t+1). \quad (3.115)$$

Expression (2.11) becomes

$$\begin{aligned} \frac{4}{24m+1} &= \frac{1}{6(d+1)(12s+11)(24t+1)} \\ &+ + \frac{2(12s+11) + (24t+1)}{6(d+1)(12s+11)(24t+1)(24m+1)} \\ &= \frac{1}{(d+1)(12s+11)(24t+1)} + \frac{1}{3(d+1)(24t+1)(24m+1)} \\ &+ \frac{1}{6(d+1)(12s+11)(24m+1)}. \end{aligned} \quad (3.116)$$

Theorem 3.9 *For any positive integer m in (3.145), we have the tame solution (3.147) of the Erdős-Straus equation.*

4 Cases $\mathfrak{S}_1 = 4j$, j is odd and $\mathfrak{S}_2 = 4\ell + 3$

This is the case when

$$\mathfrak{S}_1 = 4j, \quad k = j + \ell + 1 \quad \text{with } j, \ell \in \mathbb{N} \quad (4.1)$$

and $\ell \geq 1$ by (2.13). Moreover,

$$6m + k = 6m + j + \ell + 1. \quad (4.2)$$

According to the first expression in (2.28),

$$4 | (6m + j + \ell + 1). \quad (4.3)$$

So

$$j + \ell \equiv 1 \pmod{2}. \quad (4.4)$$

4.1 Case $m = 2m_1$, $j = 4s + 1$ and $\ell = 12t + 3$

Now

$$\mathfrak{S}_1 = 4(4s+1), \quad \mathfrak{S}_2 = 48t+11 \quad (4.5)$$

and

$$6m' + j' + \ell' = 4(3(m_1 + t) + s + 1). \quad (4.6)$$

According to the first expression in (2.28),

$$(4s+1) | (3(m_1 + t) + s + 1); \quad (4.7)$$

that is

$$3(m_1 + t) + s + 1 \equiv 0 \pmod{4s+1}. \quad (4.8)$$

Note

$$3(m_1 + t) + s + 1 \equiv 4s + 1 \pmod{4s+1}. \quad (4.9)$$

Equivalently,

$$3(m_1 + t) \equiv 3s \pmod{4s + 1}. \quad (4.10)$$

Subcase 1. $s \not\equiv 2 \pmod{3}$.

In this subcase,

$$m_1 + t \equiv s \pmod{4s + 1}. \quad (4.11)$$

Hence

$$m_1 + t = s + c(4s + 1) \quad \text{for some } c \in \mathbb{N}. \quad (4.12)$$

Observe that

$$6m + j + \ell; = 4(3c + 1)(4s + 1). \quad (4.13)$$

According to the second expression in (2.28) and (2.30),

$$(48t + 11)|(3c + 1). \quad (4.14)$$

Therefore,

$$c = 32t + 7 + d(48t + 11) \quad \text{for some } d \in \mathbb{N}. \quad (4.15)$$

This implies

$$m = 2m_1 = 2[s - t + (32t + 7 + d(48t + 11))(4s + 1)] \quad (4.16)$$

and

$$6m + j + \ell + 1 = 4(3d + 2)(4s + 1)(48t + 11). \quad (4.17)$$

Expression (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{4(3d + 2)(4s + 1)(48t + 11)} \\ &+ \frac{4(4s + 1) + (48t + 11)}{4(3d + 2)(4s + 1)(48t + 11)(24m + 1)} \\ &= \frac{1}{4(3d + 2)(4s + 1)(48t + 11)} + \frac{1}{(3d + 2)(48t + 11)(24m + 1)} \\ &+ \frac{1}{4(3d + 2)(4s + 1)(24m + 1)}. \end{aligned} \quad (4.18)$$

Subcase 2. $s \equiv 2 \pmod{3}$.

In this subcase, $s = 3s_1 + 2$ for some $s_1 \in \mathbb{N}$. Then (4.10) is equivalent to

$$m_1 + t \equiv 3s_1 + 2 \pmod{4s_1 + 3}. \quad (4.19)$$

Hence

$$m_1 + t = 3s_1 + 2 + c(4s_1 + 3) \quad \text{for some } c \in \mathbb{N}. \quad (4.20)$$

Moreover,

$$3(m_1 + t) + s + 1 = 3(m_1 + t + s_1 + 1) = 3(c + 1)(4s_1 + 3). \quad (4.21)$$

According to the second expression in (2.28) and (2.30),

$$(48t + 11)|(c + 1). \quad (4.22)$$

Thus

$$c = (d + 1)(48t + 11) - 1 \quad \text{for some } d \in \mathbb{N}. \quad (4.23)$$

Now

$$m = -2s_1 - 2t - 2 + 2(d + 1)(4s_1 + 3)(48t + 11) \quad (4.24)$$

and

$$6m + j + \ell + 1 = 12(d + 1)(4s_1 + 3)(48t + 11). \quad (4.25)$$

Expression (2.11) becomes

$$\begin{aligned} & \frac{4}{24m + 1} = \frac{1}{12(d + 1)(4s_1 + 3)(48t + 11)} \\ & + \frac{12(4s_1 + 3) + (48t + 11)}{12(d + 1)(4s_1 + 3)(48t + 11)(24m + 1)} \\ = & \frac{1}{12(d + 1)(4s_1 + 3)(48t + 11)} + \frac{1}{(d + 1)(48t + 11)(24m + 1)} \\ & + \frac{1}{12(d + 1)(4s_1 + 3)(24m + 1)}. \end{aligned} \quad (4.26)$$

Theorem 4.1 *For any positive integer m in (4.16), we have the tame solution (4.18) of the Erdős-Straus equation. When m is of the form (4.24), we have the tame solution (4.26) of the Erdős-Straus equation.*

4.2 Case $m = 2m_1 + 1, j = 12s + 5$ and $\ell = 4t$

In this case,

$$\mathfrak{S}_1 = 4(12s + 5), \quad \mathfrak{S}_2 = 16t + 3 \quad (4.27)$$

by (4.1), and

$$6m + j + \ell + 1 = 12m_1 + 6 + 12s + 5 + 4t + 1 = 4(3(m_1 + s + 1) + t) \quad (4.28)$$

According to the second expression in (2.28),

$$(16t + 3) | (3(m_1 + s + 1) + t); \quad (4.29)$$

that is

$$3(m_1 + s + 1) + t \equiv 0 \pmod{16t + 3}. \quad (4.30)$$

Note

$$3(m_1 + s + 1) + t \equiv 16t + 3 \pmod{16t + 11}. \quad (4.31)$$

Equivalently,

$$3(m_1 + s) \equiv 15t \pmod{16t + 3}. \quad (4.32)$$

Subcase 1. $t \not\equiv 0 \pmod{3}$.

In this subcase, the above equation is equivalent to

$$m_1 + s \equiv 5t \pmod{16t + 3}. \quad (4.33)$$

Hence

$$m_1 = 5t - s + c(16t + 3) \quad \text{for some } c \in \mathbb{N}. \quad (4.34)$$

Observe that

$$6m + j + \ell + 1 = 4(3c + 1)(16t + 3). \quad (4.35)$$

According to the first expression in (2.28) and (2.30),

$$(12s + 5)|(3c + 1). \quad (4.36)$$

Therefore,

$$c = 8s + 3 + d(12s + 5) \quad \text{for some } d \in \mathbb{N}. \quad (4.37)$$

This implies

$$m_1 = 5t - s + (8s + 3 + d(12s + 5))(16t + 3) \quad (4.38)$$

$$\implies m = 2[5t - s + (8s + 3 + d(12s + 5))(16t + 3)] + 1 \quad (4.39)$$

and

$$6m + j + \ell + 1 = 4(3d + 2)(12s + 5)(16t + 3). \quad (4.40)$$

Expression (2.11) becomes

$$\begin{aligned} & \frac{4}{24m + 1} = \frac{1}{4(3d + 2)(12s + 3)(16t + 3)} \\ & + \frac{4(12s + 5) + (16t + 3)}{4(3d + 2)(12s + 5)(16t + 3)(24m + 1)} \\ & = \frac{1}{4(3d + 2)(12s + 5)(16t + 3)} + \frac{1}{(3d + 2)(16t + 3)(24m + 1)} \\ & + \frac{1}{4(3d + 2)(12s + 5)(24m + 1)}. \end{aligned} \quad (4.41)$$

Subcase 2. $t \equiv 0 \pmod{3}$.

In this subcase, $t = 3t_1$ for some $t_1 \in \mathbb{N}$. Equation (3.42) becomes

$$3(m_1 + s) \equiv 45t_1 \pmod{3(16t_1 + 1)}. \quad (4.42)$$

So

$$m_1 + s \equiv 15t_1 \pmod{16t_1 + 1}; \quad (4.43)$$

that is,

$$m_1 + s = 15t_1 + c(16t_1 + 1) \quad \text{for some } c \in \mathbb{N}. \quad (4.44)$$

Observe that

$$6m + j + \ell + 1 = 12(c + 1)(16t_1 + 1). \quad (4.45)$$

According to the first expression in (2.28) and (2.30),

$$(12s + 5)|(c + 1). \quad (4.46)$$

Therefore,

$$c = (d + 1)(12s + 5) - 1 \quad \text{for some } d \in \mathbb{N}. \quad (4.47)$$

This implies

$$m_1 = -s - t_1 - 1 + (d + 1)(12s + 5)(16t_1 + 1) \quad (4.48)$$

$$\implies m = 2[-s - t_1 + (d+1)(12s+5)(16t_1+1)] - 1 \quad (4.49)$$

and

$$6m + j + \ell + 1 = 12(d+1)(12s+5)(16t_1+1). \quad (4.50)$$

Expression (2.11) becomes

$$\begin{aligned} \frac{4}{24m+1} &= \frac{1}{12(d+1)(12s+5)(16t_1+1)} \\ &+ \frac{4(12s+5) + 3(16t_1+1)}{12(d+1)(12s+5)(16t_1+1)(24m+1)} \\ &= \frac{1}{12(d+1)(12s+5)(16t_1+9)} + \frac{1}{3(d+1)(16t_1+9)(24m+1)} \\ &+ \frac{1}{4(d+1)(12s+5)(24m+1)}. \end{aligned} \quad (4.51)$$

Theorem 4.2 *For any positive integer m in (4.39), we have the tame solution (4.41) of the Erdős-Straus equation. When m is in (4.49), we have the tame solution (4.51) of the Erdős-Straus equation.*

4.3 Case $m = 2m_1, j = 12s + 7$ and $\ell = 12t + 4$

Now

$$\mathfrak{S}_1 = 4(12s+7), \quad \mathfrak{S}_2 = 48t+19 \quad (4.52)$$

by (4.1), and

$$6m + j + \ell + 1 = 12(m_1 + s + t + 1). \quad (4.53)$$

According to the first expression in (2.28),

$$(12s+7)|(m_1 + s + t + 1); \quad (4.54)$$

that is

$$m_1 + s + t + 1 \equiv 0 \pmod{12s+7}. \quad (4.55)$$

Hence

$$m_1 = -s - t - 1 + c(12s+7) \quad \text{for some } c \in \mathbb{N}. \quad (4.56)$$

Observe that

$$6m + j + \ell + 1 = 12c(12s+7). \quad (4.57)$$

According to the second expression in (2.28) and (2.30),

$$(48t+19)|c. \quad (4.58)$$

Therefore,

$$c = (d+1)(48t+19) \quad \text{for some } d \in \mathbb{N}. \quad (4.59)$$

This implies

$$m = 2m_1 = 2[-s - t - 1 + (d+1)(12s+7)(48t+19)] \quad (4.60)$$

and

$$6m + j + \ell + 1 = 12(d+1)(12s+7)(48t+19). \quad (4.61)$$

Expression (2.11) becomes

$$\begin{aligned}
& \frac{4}{24m+1} = \frac{1}{12(d+1)(12s+7)(48t+19)} \\
& + \frac{4(12s+7) + (48t+19)}{12(d+1)(12s+7)(48t+19)(24m+1)} \\
= & \frac{1}{12(d+1)(12s+7)(48t+19)} + \frac{1}{3(d+1)(48t+19)(24m+1)} \\
& + \frac{1}{12(d+1)(12s+7)(24m+1)}. \tag{4.62}
\end{aligned}$$

Theorem 4.3 *For any positive integer m in (4.60), we have the tame solution (4.62) of the Erdős-Straus equation.*

4.4 Case $m = 2m_1 + 1, j = 12s + 1$ and $\ell = 12t + 4$

Now

$$\mathfrak{S}_1 = 4(12s + 1), \quad \mathfrak{S}_2 = 48t + 19 \tag{4.63}$$

by (4.1), and

$$6m + j + \ell + 1 = 12(m_1 + s + t + 1). \tag{4.64}$$

According to the first expression in (2.28),

$$(12s + 1) | (m_1 + s + t + 1); \tag{4.65}$$

that is

$$m_1 + s + t + 1 \equiv 0 \pmod{12s + 1}. \tag{4.66}$$

Hence

$$m_1 = -s - t - 1 + c(12s + 1) \quad \text{for some } c \in \mathbb{N}. \tag{4.67}$$

Observe that

$$6m + j + \ell + 1 = 12c(12s + 1). \tag{4.68}$$

According to the second expression in (2.28) and (2.30),

$$(48t + 19) | c. \tag{4.69}$$

Therefore,

$$c = (d + 1)(48t + 19) \quad \text{for some } d \in \mathbb{N}. \tag{4.70}$$

This implies

$$m = 2m_1 + 1 = 2[-s - t + (d + 1)(12s + 1)(48t + 19)] - 1 \tag{4.71}$$

and

$$6m + j + \ell + 1 = 12(d + 1)(12s + 1)(48t + 19). \tag{4.72}$$

Expression (2.11) becomes

$$\begin{aligned}
& \frac{4}{24m+1} = \frac{1}{12(d+1)(12s+1)(48t+19)} \\
& + \frac{4(12s+1) + (48t+19)}{12(d+1)(12s+1)(48t+19)(24m+1)} \\
= & \frac{1}{12(d+1)(12s+1)(48t+19)} + \frac{1}{3(d+1)(48t+19)(24m+1)} \\
& + \frac{1}{12(d+1)(12s+1)(24m+1)}. \tag{4.73}
\end{aligned}$$

Theorem 4.4 *For any positive integer m in (4.70), we have the tame solution (4.72) of the Erdős-Straus equation.*

4.5 Case $m = 2m_1 + 1, j = 4s + 3$ and $\ell = 12t + 2$

In this case,

$$\mathfrak{S}_1 = 4(4s + 3), \quad \mathfrak{S}_2 = 48t + 11 \tag{4.74}$$

by (4.1), and

$$6m + j + \ell + 1 = 4(3(m_1 + t + 1) + s). \tag{4.75}$$

According to the first expression in (2.28),

$$(4s + 3) | (3(m_1 + t + 1) + s); \tag{4.76}$$

that is

$$3(m_1 + t + 1) + s \equiv 0 \pmod{4s + 3}. \tag{4.77}$$

Note

$$3(m_1 + t + 1) + s \equiv 4s + 3 \pmod{4s + 3}. \tag{4.78}$$

Equivalently,

$$3(m_1 + t) \equiv 3s \pmod{4s + 3}. \tag{4.79}$$

Subcase 1. $s \equiv 0 \pmod{3}$.

In this subcase, $s = 3s_1$ for some $s_1 \in \mathbb{N}$. Moreover, the above equation becomes

$$3(m_1 + t) \equiv 9s_1 \pmod{12s_1 + 3}. \tag{4.80}$$

So

$$m_1 + t \equiv 3s_1 \pmod{4s_1 + 1}. \tag{4.81}$$

Thus

$$m_1 = 3s_1 - t + c(4s_1 + 1) \quad \text{for some } c \in \mathbb{N}. \tag{4.82}$$

In particular,

$$6m + j + \ell + 1 = 12(m_1 + t + s_1 + 1) = 12(c + 1)(4s_1 + 1). \tag{4.83}$$

The second expression in (2.28) and (2.30) yield

$$(48t + 11)|(c + 1) \implies c = (d + 1)(48t + 11) - 1 \quad \text{for some } d \in \mathbb{N}. \quad (4.84)$$

Equation (4.71) shows

$$m_1 = -s_1 - t - 1 + (d + 1)(4s_1 + 1)(48t + 11). \quad (4.85)$$

$$\implies m = 2[-s_1 - t + (d + 1)(4s_1 + 1)(48t + 11)] - 1. \quad (4.86)$$

Furthermore,

$$6m + j + \ell + 1 = 12(d + 1)(4s_1 + 1)(48t + 11). \quad (4.87)$$

Therefore, (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{12(d + 1)(4s_1 + 1)(48t + 11)} \\ &+ \frac{12(4s_1 + 1) + (48t + 11)}{12(d + 1)(4s_1 + 1)(48t + 11)(24m + 1)} \\ &= \frac{1}{12(d + 1)(4s_1 + 1)(48t + 11)} + \frac{1}{(d + 1)(48t + 11)(24m + 1)} \\ &+ \frac{1}{12(d + 1)(4s_1 + 1)(24m + 1)}. \end{aligned} \quad (4.88)$$

Subcase 2. $s \not\equiv 0 \pmod{3}$.

By (4.68),

$$m_1 + t \equiv s \pmod{4s + 3}. \quad (4.89)$$

Thus

$$m_1 \equiv s - t \pmod{4s + 3}. \quad (4.90)$$

Hence

$$m_1 = s - t + c(4s + 3) \quad \text{for some } c \in \mathbb{N}. \quad (4.91)$$

Observe that

$$6m + j + \ell + 1 = 4(3c + 1)(4s + 3). \quad (4.92)$$

According to the second expression in (2.28) and (2.30),

$$(48t + 11)|(3c + 1). \quad (4.93)$$

Therefore,

$$c = 32t + 7 + d(48t + 11) \quad \text{for some } d \in \mathbb{N}. \quad (4.94)$$

This implies

$$m_1 = s - t + (32t + 7 + d(48t + 11))(4s + 3) \quad (4.95)$$

$$\implies m = 2[s - t + (32t + 7 + d(48t + 11))(4s + 3)] + 1 \quad (4.96)$$

and

$$6m + j + \ell + 1 = 4(3d + 2)(4s + 3)(48t + 11). \quad (4.97)$$

Expression (2.11) becomes

$$\begin{aligned}
& \frac{4}{24m+1} = \frac{1}{4(3d+2)(4s+3)(48t+11)} \\
& + \frac{4(4s+3) + (48t+11)}{4(3d+2)(4s+3)(48t+11)(24m+1)} \\
& = \frac{1}{4(3d+2)(4s+3)(48t+11)} + \frac{1}{(3d+2)(48t+11)(24m+1)} \\
& + \frac{1}{4(3d+2)(4s+3)(24m+1)}. \tag{4.98}
\end{aligned}$$

In summary, we have:

Theorem 4.5 *For any positive integer m in (4.86), we have the tame solution (4.88) of the Erdős-Straus equation. When m is in (4.96), we have the tame solution (4.98) of the Erdős-Straus equation.*

4.6 Case $m = 2m_1, j = 12s + 11$ and $\ell = 4t$

In this case,

$$\mathfrak{S}_1 = 4(12s + 11), \quad \mathfrak{S}_2 = 16t + 3 \tag{4.99}$$

by (4.1), and

$$6m + j + \ell + 1 = 4(3(m_1 + s + 1) + t). \tag{4.100}$$

According to the second expression in (2.28),

$$(16t + 3) | (3(m_1 + s + 1) + t); \tag{4.101}$$

that is

$$3(m_1 + s + 1) + t \equiv 0 \pmod{16t + 3}. \tag{4.102}$$

Note

$$3(m_1 + s + 1) + t \equiv 16t + 3 \pmod{16t + 3}. \tag{4.103}$$

Equivalently,

$$3(m_1 + s) \equiv 15t \pmod{16t + 3}. \tag{4.104}$$

Subcase 1. $t \not\equiv 0 \pmod{3}$.

In this subcase, the above equation is equivalent to

$$m_1 + s \equiv 5t \pmod{16t + 3}. \tag{4.105}$$

Thus

$$m_1 = 5t - s + c(16t + 3) \quad \text{for some } c \in \mathbb{N}. \tag{4.106}$$

In particular,

$$6m + j + \ell + 1 = 4(3c + 1)(16t + 3). \tag{4.107}$$

The first expression in (2.28) and (2.30) yield

$$(12s + 11) | (3c + 1) \implies c = 8s + 7 + d(12s + 11) \quad \text{for some } d \in \mathbb{N}. \tag{4.108}$$

Equation (5.91) shows

$$m_1 = 5t - s + (8s + 7 + d(12s + 11))(16t + 3) \quad (4.109)$$

$$\implies m = 2[5t - s + (8s + 7 + d(12s + 11))(16t + 3)]. \quad (4.110)$$

Furthermore,

$$6m + j + \ell + 1 = 4(3d + 2)(12s + 11)(16t + 3). \quad (4.111)$$

Therefore, (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{4(3d + 2)(12s + 11)(16t + 3)} \\ &+ \frac{4(12s + 11) + (16t + 3)}{4(3d + 2)(12s + 11)(16t + 3)(24m + 1)} \\ &= \frac{1}{4(3d + 2)(12s + 11)(16t + 3)} + \frac{1}{4(3d + 2)(16t + 3)(24m + 1)} \\ &+ \frac{1}{4(3d + 2)(12s + 11)(24m + 1)}. \end{aligned} \quad (4.112)$$

Subcase 2. $t \equiv 0 \pmod{3}$.

In this subcase, $t = st_1$ for some $t_1 \in \mathbb{N}$. Now (4.104) becomes

$$3(m_1 + s + 1) \equiv 45t_1 + 3 \pmod{3(16t_1 + 1)}; \quad (4.113)$$

equivalently,

$$m_1 + s \equiv 15t_1 \pmod{16t_1 + 1}; \quad (4.114)$$

Thus

$$m_1 = 15t_1 - s + c(16t_1 + 1) \quad \text{for some } c \in \mathbb{N}. \quad (4.115)$$

In particular,

$$6m + j + \ell + 1 = 12(c + 1)(16t_1 + 1). \quad (4.116)$$

The first expression in (2.28) and (2.30) yield

$$(12s + 11)|(c + 1) \implies c = (d + 1)(12s + 11) - 1 \quad \text{for some } d \in \mathbb{N}. \quad (4.117)$$

Equation (5.91) shows

$$m_1 = -s - t - 1 + (d + 1)(12s + 11)(16t_1 + 1) \quad (4.118)$$

$$\implies m = 2[-s - t_1 - 1 + (d + 1)(12s + 11)(16t_1 + 1)]. \quad (4.119)$$

Furthermore,

$$6m + j + \ell + 1 = 12(d + 1)(12s + 11)(16t_1 + 1). \quad (4.120)$$

Therefore, (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{12(d + 1)(12s + 11)(16t_1 + 1)} \\ &+ \frac{4(12s + 11) + 3(16t_1 + 1)}{12(d + 1)(12s + 11)(16t_1 + 1)(24m + 1)} \\ &= \frac{1}{12(d + 1)(12s + 11)(16t_1 + 1)} + \frac{1}{3(d + 1)(16t_1 + 1)(24m + 1)} \\ &+ \frac{1}{4(d + 1)(12s + 11)(24m + 1)}. \end{aligned} \quad (4.121)$$

In summary, we have:

Theorem 4.6 *When m is in (4.110), the Erdős-Straus equation (4.112) holds. If m is in (4.119), the Erdős-Straus equation (4.121) holds.*

4.7 Section Summary

In the above sections, $m = g(x, d, y)$, $\mathfrak{S}_1 = \eta(x)$ and $\mathfrak{S}_2 = h(y)$ are integral polynomials, where $x = s$ or s_1 and $y = 2t$ or $y = 2t_1$. If we change y to odd positive integer, they still give a solution of the Erdős-Straus equation. In summary, we have the following table of solutions:

Table 4.6

m	$\mathfrak{S}_1$	$\mathfrak{S}_2$
$2x - y + 2(16y + 7 + d(24y + 11))(4x + 1)$	$4(4x + 1)$	$24y + 11$
$-2x - y - 2 + 2(d + 1)(4x + 3)(24y + 11)$	$12(4x + 3)$	$24y + 11$
$5y - 2x + 2(8x + 3 + d(12x + 5))(8y + 3) + 1$	$4(12x + 5)$	$8y + 3$
$-2x - y - 1 + 2(d + 1)(12x + 5)(8y + 1)$	$4(12x + 5)$	$3(8y + 1)$
$-2x - y - 2 + 2(d + 1)(12x + 7)(24y + 19)$	$4(12x + 7)$	$24y + 19$
$-2x - y - 1 + 2(d + 1)(12x + 1)(24y + 19)$	$4(12x + 1)$	$24y + 19$
$-2x - y + 2(d + 1)(4x + 1)(24y + 11) - 1$	$12(4x + 1)$	$24y + 11$
$2x - y + 2(16y + 7 + d(24y + 11))(4x + 3) + 1$	$4(4x + 3)$	$24y + 11$
$5y + 2[-x + (8x + 7 + d(12s + 11))(8y + 3)]$	$4(12x + 11)$	$8y + 3$
$-2x - y - 2 + 2(d + 1)(12x + 11)(8y + 1)$	$4(12x + 11)$	$3(8y + 1)$

5 Cases $\mathfrak{S}_1 = 2^{r+3}j'$, j' is odd and $\mathfrak{S}_2 = 4\ell + 3$

In this situation, $j = 2^{r+1}j'$.

$$\mathfrak{S}_1 = 2^{r+3}j', \quad \mathfrak{S}_2 = 4\ell + 3. \quad (5.1)$$

Moreover,

$$6m + j + \ell + 1 = 6m + 2^{r+1}j' + \ell + 1. \quad (5.2)$$

According to the first expression in (2.28),

$$6m + 2^{r+1}j' + \ell + 1 \equiv 0 \pmod{2^{r+3}j'}. \quad (5.3)$$

So

$$\ell \text{ is odd.} \quad (5.4)$$

5.1 $j = 3 \cdot 2^{r+1}(2s+1)$ and $\ell = 3 \cdot 2^{r+3}t + 5$

In this case,

$$6m + j + \ell + 1 = 6m + 3 \cdot 2^{r+1}(2s+1) + 3 \cdot 2^{r+3}t + 6 = 6[m + 2^r(2s+4t+1) + 1]. \quad (5.5)$$

Now (5.3) becomes,

$$6[m + 2^r(2s+4t+1) + 1] \equiv 0 \pmod{2^{r+3}3(2s+1)}; \quad (5.6)$$

equivalently,

$$m + 2^r(2s+4t+1) + 1 \equiv 0 \pmod{2^{r+2}(2s+1)}; \quad (5.7)$$

Thus

$$m = 2^{r+2}(2s+1)c - 2^r(2s+4t+1) - 1 \quad \text{for some } 0 < c \in \mathbb{N}. \quad (5.8)$$

Note

$$\mathfrak{S}_2 = 4\ell + 3 = 3 \cdot 2^{r+5}t + 23, \quad 6m + j + \ell + 1 = 3 \cdot 2^{r+3}(2s+1)c. \quad (5.9)$$

According to the second expression in (2.28) and (2.30),

$$(3 \cdot 2^{r+5}t + 23) | (3 \cdot 2^{r+3}(2s+1)c) \implies (3 \cdot 2^{r+5}t + 23) | c \quad (5.10)$$

$$\implies c = (d+1)(3 \cdot 2^{r+5}t + 23) \quad \text{for some } d \in \mathbb{N}. \quad (5.11)$$

Hence

$$m = (d+1)2^{r+3}(2s+1)(3 \cdot 2^{r+5}t + 23) - 2^r(2s+4t+1) - 1 \quad (5.12)$$

and

$$6m + j + \ell + 1 = 3(d+1)2^{r+3}(2s+1)(3 \cdot 2^{r+5}t + 23). \quad (5.13)$$

Therefore, (2.11) becomes

$$\begin{aligned} \frac{4}{24m+1} &= \frac{1}{3(d+1)2^{r+3}(2s+1)(3 \cdot 2^{r+5}t + 23)} \\ &+ \frac{(3 \cdot 2^{r+3}(2s+1) + (3 \cdot 2^{r+5}t + 23))}{3(d+1)2^{r+3}(2s+1)(3 \cdot 2^{r+5}t + 23)d(24m+1)} \\ &= \frac{1}{3(d+1)2^{r+3}(2s+1)(3 \cdot 2^{r+5}t + 23)} + \frac{1}{(d+1)(3 \cdot 2^{r+5}t + 23)(24m+1)} \\ &+ \frac{1}{3(d+1)2^{r+3}(2s+1)d(24m+1)}. \end{aligned} \quad (5.14)$$

Theorem 5.1 When m is in (5.12), the Erdős-Straus equation (5.14) holds.

5.2 $j = 2^{r+1}(2s+1)$, $3 \nmid (2s+1)$ and $\ell = 3 \cdot 2^{r+3}t + 5$

In this case,

$$6m + j + \ell + 1 = 6m + 2^{r+1}(2s+1) + 3 \cdot 2^{r+3}t + 6 = 6(m + 2^{r+2}t + 1) + 2^{r+1}(2s+1). \quad (5.15)$$

Now (5.3) becomes,

$$6(m + 2^{r+2}t + 1) + 2^{r+1}(2s+1) \equiv 0 \pmod{2^{r+3}(2s+1)}; \quad (5.16)$$

equivalently,

$$3(m + 2^{r+2}t + 1) + 2^r(2s + 1) \equiv 0 \pmod{2^{r+2}(2s + 1)}. \quad (5.17)$$

Note

$$3(m + 2^{r+2}t + 1) + 2^r(2s + 1) \equiv 2^{r+2}(2s + 1) \pmod{2^{r+2}(2s + 1)}; \quad (5.18)$$

equivalently,

$$3(m + 2^{r+2}t + 1) \equiv 2^r(2s + 1)3 \pmod{2^{r+2}(2s + 1)}. \quad (5.19)$$

Hence

$$m + 2^{r+1}t + 1 \equiv 2^r(2s + 1) \pmod{2^{r+1}(2s + 1)}, \quad (5.20)$$

that is,

$$m = 2^r(2s + 1) + 2^{r+2}(2s + 1)c - 2^{r+2}t - 1 \quad \text{for some } c \in \mathbb{N}. \quad (5.21)$$

Moreover,

$$\mathfrak{S}_2 = 4\ell + 3 = 3 \cdot 2^{r+5}t + 23, \quad 6m + j + \ell + 1 = 2^{r+3}(2s + 1)(3c + 1). \quad (5.22)$$

According to the second expression in (2.28) and (2.30),

$$(3 \cdot 2^{r+5}t + 23) | (2^{r+3}(2s + 1)(3c + 1)) \implies (3 \cdot 2^{r+5}t + 23) | (3c + 1). \quad (5.23)$$

Thus

$$c = 2^{r+6}t + 15 + d(3 \cdot 2^{r+5}t + 23) \quad \text{for some } d \in \mathbb{N}. \quad (5.24)$$

Now

$$m = 2^r(2s + 1)[4(2^{r+6}t + 15 + d(3 \cdot 2^{r+5}t + 23)) + 1] - 2^{r+2}t - 1. \quad (5.25)$$

and

$$6m + j + \ell + 1 = (2d + 2)2^{r+3}(2s + 1)(3 \cdot 2^{r+5}t + 23). \quad (5.26)$$

Therefore, (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{(3d + 2)2^{r+3}(2s + 1)(3 \cdot 2^{r+5}t + 23)} \\ &\quad + \frac{2^{r+3}(2s + 1) + (3 \cdot 2^{r+5}t + 23)}{(3d + 2)2^{r+3}(2s + 1)(3 \cdot 2^{r+5}t + 23)(24m + 1)} \\ &= \frac{1}{(3d + 2)2^{r+3}(2s + 1)(3 \cdot 2^{r+5}t + 23)} \\ &\quad + \frac{1}{(3d + 2)(3 \cdot 2^{r+5}t + 23)(24m + 1)} \\ &\quad + \frac{1}{(3d + 2)2^{r+3}(2s + 1)(24m + 1)}. \end{aligned} \quad (5.27)$$

Theorem 5.2 *When m is in (5.25), the Erdős-Straus equation (5.27) holds.*

5.3 $j = 2^{2r+2}(6s+1)$ and $\ell = 3 \cdot 2^{2r+4}t + 1$

In this case,

$$6m+j+\ell+1 = 6m+2^{2r+2}(6s+1)+3 \cdot 2^{2r+4}t+2 = 6[m+2^{2r+2}(s+2t)+(2^{2r+1}+1)/3]. \quad (5.28)$$

Now (5.3) becomes,

$$6[m+2^{2r+2}(s+2t)+(2^{2r+1}+1)/3] \equiv 0 \pmod{2^{2r+4}(6s+1)}; \quad (5.29)$$

equivalently,

$$3[m+2^{2r+2}(s+2t)+(2^{2r+1}+1)/3] \equiv 0 \pmod{2^{2r+3}(6s+1)}. \quad (5.30)$$

So

$$m+2^{2r+2}(s+2t)+(2^{2r+1}+1)/3 \equiv 0 \pmod{2^{2r+3}(6s+1)}. \quad (5.31)$$

Thus

$$m = 2^{2r+3}(6s+1)c - 2^{2r+2}(s+2t) - (2^{2r+1}+1)/3 \quad \text{for some } 0 < c \in \mathbb{N}. \quad (5.32)$$

Note

$$\mathfrak{S}_2 = 4\ell + 3 = 3 \cdot 2^{2r+6}t + 7, \quad 6m + j + \ell + 1 = 2^{2r+4}3(6s+1)c. \quad (5.33)$$

According to the second expression in (2.28) and (2.30),,

$$(3 \cdot 2^{2r+6}t + 7) | (3 \cdot 2^{2r+4}(6s+1)c) \quad (5.34)$$

$$\implies c = (d+1)(3 \cdot 2^{r+6}t + 7) \quad \text{for some } d \in \mathbb{N}. \quad (5.35)$$

Hence

$$m = (d+1)2^{2r+3}(6s+1)(3 \cdot 2^{2r+6}t + 7) - 2^{2r+2}(s+2t) - (2^{2r+1}+1)/3 \quad (5.36)$$

and

$$6m + j + \ell + 1 = 3(d+1)2^{2r+4}(6s+1)(3 \cdot 2^{2r+6}t + 7)d. \quad (5.37)$$

Therefore, (2.11) becomes

$$\begin{aligned} \frac{4}{24m+1} &= \frac{1}{3(d+1)2^{2r+4}(6s+1)(3 \cdot 2^{2r+6}t + 7)} \\ &+ \frac{2^{2r+4}(6s+1) + (3 \cdot 2^{2r+6}t + 7)}{(d+1)2^{2r+4}(6s+1)(3 \cdot 2^{2r+6}t + 7)(24m+1)} \\ &= \frac{1}{3(d+1)2^{2r+4}(6s+1)(3 \cdot 2^{2r+6}t + 7)} \\ &+ \frac{1}{3(d+1)(3 \cdot 2^{2r+6}t + 7)(24m+1)} \\ &+ \frac{1}{3(d+1)2^{2r+4}(6s+1)(24m+1)}. \end{aligned} \quad (5.38)$$

Theorem 5.3 When m is in (5.36), the Erdős-Straus equation (5.38) holds.

5.4 $j = 2^{2r+2}(6s+5)$ and $\ell = 3 \cdot 2^{2r+4}t + 1$

In this case,

$$6m+j+\ell+1 = 6m+2^{2r+2}(6s+5)+3 \cdot 2^{2r+4}t+2 = 6(m+2^{2r+3}t)+2^{2r+2}(6s+5)+2. \quad (5.39)$$

Now (5.3) becomes,

$$6(m+2^{2r+3}t)+2^{2r+2}(6s+5)+2 \equiv 0 \pmod{2^{2r+4}(6s+5)}; \quad (5.40)$$

equivalently,

$$3(m+2^{2r+3}t)+2^{2r+1}(6s+5)+1 \equiv 0 \pmod{2^{2r+3}(6s+5)}. \quad (5.41)$$

Note

$$3(m+2^{2r+3}t)+2^{2r+1}(6s+5)+1 \equiv 2^{2r+4}(6s+5) \pmod{2^{2r+3}(6s+5)}. \quad (5.42)$$

So

$$3(m+2^{2r+3}t) \equiv 2^{2r+1}7(6s+5)-1 \pmod{2^{2r+3}(6s+5)}. \quad (5.43)$$

$$3(m+2^{2r+3}t) \equiv 3[2^{2r+2}(7s+6)-(2^{2r+1}+1)/3] \pmod{2^{2r+3}(6s+5)}. \quad (5.44)$$

Hence

$$m+2^{2r+3}t \equiv 2^{2r+2}(7s+6)-(2^{2r+1}+1)/3 \pmod{2^{2r+3}(6s+5)}. \quad (5.45)$$

Thus

$$m = 2^{2r+3}(6s+5)c + 2^{2r+2}(7s-2t+6) - (2^{2r+1}+1)/3 \quad \text{for some } c \in \mathbb{N}. \quad (5.46)$$

Observe

$$\mathfrak{S}_2 = 4\ell + 3 = 3 \cdot 2^{2r+6}t + 7, \quad 6m+j+\ell+1 = (3c+2)2^{2r+4}(6s+5). \quad (5.47)$$

According to the second expression in (2.28) and (2.30),

$$(3 \cdot 2^{2r+6}t + 7) | ((3c+2)2^{2r+4}(6s+5)) \implies (3 \cdot 2^{2r+6}t + 7) | (3c+2) \quad (5.48)$$

$$\implies c = 2^{2r+7}t + 4 + d(3 \cdot 2^{2r+6}t + 7) \quad \text{for some } d \in \mathbb{N}. \quad (5.49)$$

Hence

$$m = 2^{2r+3}(6s+5)(2^{2r+7}t + 4 + d(3 \cdot 2^{2r+6}t + 7)) + 2^{2r+2}(7s-2t+6) - (2^{2r+1}+1)/3 \quad (5.50)$$

and

$$6m+j+\ell+1 = (3d+2)2^{2r+4}(6s+5)(3 \cdot 2^{2r+6}t + 7). \quad (5.51)$$

Therefore, (2.11) becomes

$$\begin{aligned} \frac{4}{24m+1} &= \frac{1}{(3d+2)2^{2r+4}(6s+5)(3 \cdot 2^{2r+6}t + 7)} \\ &+ \frac{2^{2r+4}(6s+5) + (3 \cdot 2^{2r+6}t + 7)}{(3d+2)2^{2r+4}(6s+5)(3 \cdot 2^{2r+6}t + 7)(24m+1)} \\ &= \frac{1}{(3d+2)2^{2r+4}(6s+5)(3 \cdot 2^{2r+6}t + 7)} \\ &+ \frac{1}{(3d+2)(3 \cdot 2^{2r+6}t + 7)(24m+1)} \\ &+ \frac{1}{(3d+2)2^{2r+4}(6s+5)(24m+1)}. \end{aligned} \quad (5.52)$$

Theorem 5.4 When m is in (5.50), the Erdős-Straus equation (5.52) holds.

5.5 $j = 2^{2r+2}(6s + 5)$ and $\ell = 3 \cdot 2^{2r+4}t + 3$

In this case,

$$6m + j + \ell + 1 = 6m + 2^{2r+2}(6s + 5) + 3 \cdot 2^{2r+4}t + 4 = 6[m + 2^{2r+2}(s + 2t) + (5 \cdot 2^{2r+1} + 2)/3]. \quad (5.53)$$

Now (5.3) becomes,

$$6[m + 2^{2r+2}(s + 2t) + (5 \cdot 2^{2r+1} + 2)/3] \equiv 0 \pmod{2^{2r+4}(6s + 5)}; \quad (5.54)$$

equivalently,

$$3[m + 2^{2r+2}(s + 2t) + (5 \cdot 2^{2r+1} + 2)/3] \equiv 0 \pmod{2^{2r+3}(6s + 5)}. \quad (5.55)$$

So

$$m + 2^{2r+2}(s + 2t) + (5 \cdot 2^{2r+1} + 2)/3 \equiv 0 \pmod{2^{2r+3}(6s + 5)}. \quad (5.56)$$

Thus

$$m = 2^{2r+3}(6s + 5)c - 2^{2r+2}(s + 2t) - (5 \cdot 2^{2r+1} + 2)/3 \quad \text{for some } 0 < c \in \mathbb{N}. \quad (5.57)$$

Note

$$\mathfrak{S}_2 = 4\ell + 3 = 3 \cdot 2^{2r+6}t + 15, \quad 6m + j + \ell + 1 = 3 \cdot 2^{2r+4}(6s + 5)c. \quad (5.58)$$

According to the second expression in (2.28) and (2.30),

$$(2^{2r+6}3t + 15)|(3 \cdot 2^{2r+4}(6s + 1)c) \quad (5.59)$$

$$\implies c = (d + 1)(2^{r+6}t + 5) \quad \text{for some } d \in \mathbb{N}. \quad (5.60)$$

Hence

$$m = (d + 1)2^{2r+3}(6s + 5)(2^{2r+6}t + 5)d - 2^{2r+2}(s + 2t) - (5 \cdot 2^{2r+1} + 2)/3 \quad (5.61)$$

and

$$6m + j + \ell + 1 = (d + 1)2^{2r+4}(6s + 5)(3 \cdot 2^{2r+4}t + 15)d. \quad (5.62)$$

Therefore, (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{(d + 1)2^{2r+4}(6s + 5)(3 \cdot 2^{2r+6}t + 15)} \\ &+ \frac{2^{2r+4}(6s + 5) + (3 \cdot 2^{2r+6}t + 15)}{(d + 1)2^{2r+4}(6s + 5)(3 \cdot 2^{2r+6}t + 15)(24m + 1)} \\ &= \frac{1}{(d + 1)2^{2r+4}(6s + 1)(3 \cdot 2^{2r+6}t + 15)} \\ &+ \frac{1}{(d + 1)(3 \cdot 2^{2r+6}t + 15)(24m + 1)} \\ &+ \frac{1}{(d + 1)2^{2r+4}(6s + 5)d(24m + 1)}. \end{aligned} \quad (5.63)$$

Theorem 5.5 *When m is in (5.61), the Erdős-Straus equation (5.63) holds.*

5.6 $j = 2^{2r+1}(6s + 1)$ and $\ell = 3 \cdot 2^{2r+3}t + 1$

In this case,

$$6m + j + \ell + 1 = 6m + 2^{2r+1}(6s + 1) + 3 \cdot 2^{2r+3}t + 2 = 6(m + 2^{2r+2}t) + 2^{2r+1}(6s + 1) + 2. \quad (5.64)$$

Now (5.3) becomes,

$$6(m + 2^{2r+2}t) + 2^{2r+1}(6s + 1) + 2 \equiv 0 \pmod{2^{2r+3}(6s + 1)}; \quad (5.65)$$

equivalently,

$$3(m + 2^{2r+2}t) + 2^{2r}(6s + 1) + 1 \equiv 0 \pmod{2^{2r+2}(6s + 1)}. \quad (5.66)$$

Note

$$3(m + 2^{2r+2}t) + 2^{2r}(6s + 1) + 1 \equiv 2^{2r+3}(6s + 1) \pmod{2^{2r+2}(6s + 1)} \quad (5.67)$$

$$\implies 3(m + 2^{2r+2}t) \equiv 2^{2r}7(6s + 1) - 1 \pmod{2^{2r+2}(6s + 1)} \quad (5.68)$$

$$\implies 3(m + 2^{2r+2}t) \equiv 3[2^{2r+1}(7s + 1) + (2^{2r} - 1)/3] \pmod{2^{2r+2}(6s + 1)} \quad (5.69)$$

$$\implies m + 2^{2r+2}t \equiv 2^{2r+1}(7s + 1) + (2^{2r} - 1)/3. \pmod{2^{2r+2}(6s + 1)} \quad (5.70)$$

So

$$m = c(2^{2r+2}(6s + 1)) + 2^{2r+1}(7s - 2t + 1) + (2^{2r} - 1)/3 \quad \text{for some } c \in \mathbb{N}. \quad (5.71)$$

Now

$$\mathfrak{S}_2 = 4\ell + 3 = 3 \cdot 2^{2r+5}t + 7, \quad 6m + j + \ell + 1 = 3(3c + 2)2^{2r+3}(6s + 1). \quad (5.72)$$

According to the second expression in (2.28) and (2.30),

$$(3 \cdot 2^{2r+5}t + 7)|(3(3c + 2)2^{2r+3}(6s + 1)) \implies (3 \cdot 2^{2r+5}t + 7)|(3c + 2) \quad (5.73)$$

$$\implies c = 2^{2r+6}t + 4 + d(3 \cdot 2^{2r+5}t + 7) \quad \text{for some } d \in \mathbb{N}. \quad (5.74)$$

Hence

$$m = 2^{2r+2}(6s + 1)(2^{2r+6}t + 4 + d(3 \cdot 2^{2r+5}t + 7)) + 2^{2r+1}(7s - 2t + 1) + (2^{2r} - 1)/3 \quad (5.75)$$

and

$$6m + j + \ell + 1 = 3(3d + 2)2^{2r+3}(6s + 1)(3 \cdot 2^{2r+5}t + 7). \quad (5.76)$$

Therefore, (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{3(3d + 2)2^{2r+3}(6s + 1)(3 \cdot 2^{2r+5}t + 7)} \\ &+ \frac{2^{2r+3}(6s + 1) + (3 \cdot 2^{2r+5}t + 7)}{3(3d + 2)2^{2r+3}(6s + 1)(3 \cdot 2^{2r+5}t + 7)(24m + 1)} \\ &= \frac{1}{3(3d + 2)2^{2r+3}(6s + 1)(3 \cdot 2^{2r+5}t + 7)} \\ &+ \frac{1}{3(3d + 2)(3 \cdot 2^{2r+5}t + 7)(24m + 1)} \\ &+ \frac{1}{3(3d + 2)2^{2r+3}(6s + 1)(24m + 1)}. \end{aligned} \quad (5.77)$$

Theorem 5.6 When m is in (5.75), the Erdős-Straus equation (5.77) holds.

5.7 $j = 2^{2r+1}(6s+1)$ and $\ell = 3 \cdot 2^{2r+3}t + 3$

In this case,

$$6m+j+\ell+1 = 6m+2^{2r+1}(6s+1)+3 \cdot 2^{2r+3}t+4 = 6[m+2^{2r+1}(s+2t)+(2^{2r}+2)/3]. \quad (5.78)$$

Now (5.3) becomes,

$$6[m+2^{2r+1}(s+2t)+(2^{2r}+2)/3] \equiv 0 \pmod{2^{2r+3}(6s+1)}; \quad (5.79)$$

equivalently,

$$3[m+2^{2r+1}(s+2t)+(2^{2r}+2)/3] \equiv 0 \pmod{2^{2r+2}(6s+1)}. \quad (5.80)$$

So

$$m+2^{2r+1}(s+2t)+(2^{2r}+2)/3 \equiv 0 \pmod{2^{2r+2}(6s+1)}. \quad (5.81)$$

Thus

$$m = 2^{2r+2}(6s+1)c - 2^{2r+1}(s+2t) - (2^{2r}+2)/3. \quad \text{for some } c \in \mathbb{N}. \quad (5.82)$$

Note

$$\mathfrak{S}_2 = 4\ell + 3 = 3 \cdot 2^{2r+5}t + 15, \quad 6m+j+\ell+1 = 3 \cdot 2^{2r+3}(6s+1)c. \quad (5.83)$$

According to the second expression in (2.28) and (2.30),

$$(3 \cdot 2^{2r+5}t + 15) | (3 \cdot 2^{2r+3}(6s+1)c) \quad (5.84)$$

$$\implies c = (d+1)(2^{r+5}t + 5) \quad \text{for some } d \in \mathbb{N}. \quad (5.85)$$

Hence

$$m = (d+1)2^{2r+2}(6s+1)(2^{2r+5}t + 5) - 2^{2r+1}(s+2t) - (2^{2r}+2)/3 \quad (5.86)$$

and

$$6m+j+\ell+1 = (d+1)2^{2r+3}(6s+1)(3 \cdot 2^{2r+5}t + 15). \quad (5.87)$$

Therefore, (2.11) becomes

$$\begin{aligned} \frac{4}{24m+1} &= \frac{1}{(d+1)2^{2r+3}(6s+1)(3 \cdot 2^{2r+5}t + 15)} \\ &+ \frac{2^{2r+3}(6s+1) + (3 \cdot 2^{r+5}t + 15)}{(d+1)2^{2r+3}(6s+1)(3 \cdot 2^{2r+5}t + 15)(24m+1)} \\ &= \frac{1}{(d+1)2^{2r+3}(6s+1)(3 \cdot 2^{r+5}t + 15)} \\ &+ \frac{1}{(d+1)(3 \cdot 2^{r+5}t + 15)(24m+1)} \\ &+ \frac{1}{(d+1)2^{2r+3}(6s+1)d(24m+1)}. \end{aligned} \quad (5.88)$$

Theorem 5.7 *When m is in (5.86), the Erdős-Straus equation (5.88) holds.*

5.8 $j = 2^{2r+1}(6s + 5)$ and $\ell = 3 \cdot 2^{2r+3}t + 1$

In this case,

$$6m + j + \ell + 1 = 6m + 2^{2r+1}(6s + 5) + 3 \cdot 2^{2r+3}t + 2 = 6[m + 2^{2r+1}(s + 2t) + (5 \cdot 2^{2r} + 1)/3]. \quad (5.89)$$

Now (5.3) becomes,

$$6[m + 2^{2r+1}(s + 2t) + (5 \cdot 2^{2r} + 1)/3] \equiv 0 \pmod{2^{2r+3}(6s + 5)}; \quad (5.90)$$

equivalently,

$$3[m + 2^{2r+1}(s + 2t) + (5 \cdot 2^{2r} + 1)/3] \equiv 0 \pmod{2^{2r+2}(6s + 5)}. \quad (5.91)$$

So

$$m + 2^{2r}(s + 2t) + (5 \cdot 2^{2r} + 1)/3 \equiv 0 \pmod{2^{2r+2}(6s + 5)}. \quad (5.92)$$

Thus

$$m = 2^{2r+2}(6s + 5)c - 2^{2r+1}(s + 2t) - (5 \cdot 2^{2r} + 1)/3 \quad \text{for some } c \in \mathbb{N}. \quad (5.93)$$

Note

$$\mathfrak{S}_2 = 4\ell + 3 = 3 \cdot 2^{2r+5}t + 7, \quad 6m + j + \ell + 1 = 3 \cdot 2^{2r+3}(6s + 5)c. \quad (5.94)$$

According to the second expression in (2.28) and (2.30),

$$(3 \cdot 2^{2r+5}t + 7) | (3 \cdot 2^{2r+3}(6s + 5)c) \quad (5.95)$$

$$\implies c = (d + 1)(3 \cdot 2^{2r+5}t + 7) \quad \text{for some } d \in \mathbb{N}. \quad (5.96)$$

Hence

$$m = (d + 1)2^{2r+2}(6s + 5)(3 \cdot 2^{2r+5}t + 7) - 2^{2r+1}(s + 2t) - (5 \cdot 2^{2r} + 1)/3 \quad (5.97)$$

and

$$6m + j + \ell + 1 = 3(d + 1)2^{2r+3}(6s + 5)(3 \cdot 2^{2r+5}t + 7). \quad (5.98)$$

Therefore, (2.11) becomes

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{3(d + 1)2^{2r+3}(6s + 5)(3 \cdot 2^{2r+5}t + 7)} \\ &+ \frac{2^{2r+3}(6s + 5) + (3 \cdot 2^{2r+5}t + 7)}{3(d + 1)2^{2r+3}(6s + 5)(3 \cdot 2^{2r+5}t + 7)(24m + 1)} \\ &= \frac{1}{3(d + 1)2^{2r+3}(6s + 5)(3 \cdot 2^{2r+5}t + 7)} + \frac{1}{3(d + 1)(3 \cdot 2^{2r+5}t + 7)(24m + 1)} \\ &+ \frac{1}{3(d + 1)2^{2r+3}(6s + 5)(24m + 1)}. \end{aligned} \quad (5.99)$$

Theorem 5.8 *When m is in (5.97), the Erdős-Straus equation (5.99) holds.*

5.9 Section Summary

In the above sections, $m = g(x, d, y)$ and $\mathfrak{S}_2 = h(y)$ are integral polynomials, where $y = 2^{r+1}t$, $y = 2^{2+1}t$ or $y = 2^{2r+2}t$. If we change y to any positive integer, they still give a solution of the Erdős-Straus equation. In summary, we have the following table of solutions:

Table 5.6

m	$\mathfrak{S}_1$	$\mathfrak{S}_2$
$(d+1)2^{r+2}(2x+1)(24y+23) - y - 2^r(2x+1) - 1$	$2^{r+3}(6x+3)$	$24y+23$
$2^r(2x+1)[4(16y+15+d(24y+23))+1] - y - 1$	$2^{r+3}(2x+1)$	$24y+23$
$(d+1)2^{2r+3}(6x+1)(24y+7) - 2y - 2^{2r+2}x - (2^{2r+1}+1)/3$	$2^{2r+4}(6x+1)$	$24y+7$
$2^{2r+2}[2(6x+5)(16y+4+d(24y+7)) + 7x+6] - y - (2^{2r+1}+1)/3$	$2^{2r+4}(6x+5)$	$24y+7$
$(d+1)2^{2r+3}(6x+5)(8y+5) - y - 2^{2r}x - (5 \cdot 2^{2r+1}+2)/3$	$2^{2r+4}(6x+5)$	$24y+15$
$2^{2r+1}[2(6x+1)(16y+4+d(24y+7)) + 7x+1] - y + (2^{2r}-1)/3$	$2^{2r+3}(6x+1)$	$24y+7$
$(d+1)2^{2r+2}(6x+1)(8y+5) - y - 2^{2r+1}x - (2^{2r}+2)/3$	$2^{2r+3}(6x+1)$	$24y+15$
$(d+1)2^{2r+2}(6x+5)(24y+7) - y - 2^{2r+1}x - (3 \cdot 2^{2r}+1)/3$	$2^{2r+3}(6x+5)$	$24y+7$

References

- [1] K. Bradford, A note on the Erdős-Straus conjecture, *Integers* **21** (2021), Paper No. A24, 10pp.
- [2] M. Bright and D. Loughran, Brauer-Manin obstruction for Erdős-Straus surfaces, *Bull. Lond. Math. Soc.* **52** (2020), no. 4, 746–761.
- [3] C. Elscholtz and T. Tao, Counting the number of solutions to the equations on unit fractions, *J. Aust. Math. Soc.* **94** (2013), 50–105.
- [4] T. R. Hagedorn, A proof of a conjecture on Egyptian fractions, *Amer. Math. Monthly* **107** (2000), no. 1, 62–63.
- [5] J. Huang and R. Vaughan, Mean value theorems for binary Egyptian fractions, *J. Number Theory* **121** (2011), no. 9, 1641–1656.
- [6] J. Huang and R. Vaughan, Mean value theorems for binary Egyptian fractions II, *Acta Arith.* **115** (2012), no. 3, 287–296.
- [7] J. Huang and R. Vaughan, On the exceptional set for binary Egyptian fractions, *Bull. Lond. Math. Soc.* **45** (2013), no. 4, 861–874.
- [8] B. Howerton, The wild primes formula sheet, *personal communication*, 12 June 2026.
- [9] C. Jia, The estimate for mean values on prime numbers relative to $4/n = 1/n_1 + 1/n_2 + 1/n_3$, *Sci. China: Math.* **55** (3), (2012), 465–474.
- [10] C. Jia, Mean value from representation of a rational number as sum of two Egyptian fractions, *J. Number Theory* **132** (2012), no. 4, 701–713.

- [11] E. J. Ionascu and A. Wilson, On the Erdős-Straus conjecture, *Rev. Roumaine Math. Pures Appl.* **56** (2011), no. 1, 21–30.
- [12] I. Kotsireas, The Erdős-Straus conjecture on Egyptian fractions, *Paul Erdős and his mathematics (Budapest, 1999)*, 140–144, *János Nolgai Math. Soc. Budapest*, 1999.
- [13] D. Li, On the equation $4/n = 1/x + 1/y + 1/z$, *J. Number Theory* **13** (1981), no. 4, 485–494.
- [14] S. Maiti, A study on Erdős-Straus conjecture on Diophantine equation $\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$, *arXiv: 2010.00975v1 [math.GM]* 30 Dec 2020.
- [15] L. J. Mordell, *Diophantine Equations*, Academic Press, London/New York, 1969.
- [16] S. Prugsapitak, The Egyptian fraction of the form $\frac{1}{a} + \frac{1}{b} = \frac{q-1}{pq}$, *Int. J. Math. Comput. Sci.* **18** (2023), no. 4, 595–597.
- [17] L. Antonio Rosati, Sull’equazione diofantea $4/n = 1/x_1 + 1/x_2 + 1/x_3$, *Boll. Uni. Mat. Ital. (3)* **9** (1954), 59–63.
- [18] J. W. Sander, On $4/n = 1/x + 1/y + 1/z$ and Rosser’s sieve, *Acta Arith.* **59** (1991), no. 2, 183–204.
- [19] J. W. Sander, On $4/n = 1/x + 1/y + 1/z$ and Iwanniec’s half-dimensional sieve, *J. Number Theory* **46** (1994), no. 2, 123–136.
- [20] J. W. Sander, Egyptian fractions and the Erdős-Straus conjecture, *Nieuw Arch. Wisk (4)* **15** (1997), no. 1-2, 43–50.
- [21] J. D. Serna, Partial proof without words: shapping some cases of the Erdős-Straus conjecture, *College. Math. J.* **40** (2015), no. 3, 181.
- [22] S. Subburam and A. Togb’e, A note on Erdős-Straus conjecture, *Period Math. Hungar.* **72** (2016), no. 1, 43–49.
- [23] D. G. Terzi, On a conjecture by Erdős and Straus, *Nordisk Tidsk. Informationsbehandling (BIT)* **11** (1971), 212–216.
- [24] R. C. Vaughan, On a problem of Erdős, Straus and Schinzel, *Mathematika* **17** (1970), 193–198.
- [25] W. A. Webb, On $4/n = 1/x + 1/y + 1/z$, *Proc. Amer. Math. Soc.* **25** (1970), 578–584.
- [26] X. Xu, Congruence classes of supporting the Erdős-Straus conjecture I: tame solutions, *arXiv: 2605.2360[v] [math.NT]* May 2026.

E-Mail: xiaoping@math.ac.cn