

COUNTING TYPE I AND TYPE II SOLUTIONS OF THE ERDŐS–STRAUS EQUATION: CONCENTRATION, LOCAL OBSTRUCTIONS, AND A POINTWISE COMPARISON

XINYANG JIANG

ABSTRACT. For a prime $p \geq 5$ let $f_I(p)$ and $f_{II}(p)$ count the Type I and Type II solutions of the Erdős–Straus equation $4/p = 1/x + 1/y + 1/z$, in the sense of Elsholtz and Tao. We do not address the Erdős–Straus conjecture itself; our subject is the *number* of solutions. Our main structural finding is that a quadratic character can distinguish the two types only when the numerator is not a perfect square — so never for the numerator 4.

We first upgrade a recent bijection of Bradford into an exact counting identity: both $f_I(p)$ and $f_{II}(p)$ are sums, over the denominators n coprime to p , of the number of divisors of n^2 in a single residue class modulo $e_n = 4n - p$, the Type I class -4^{-1} being independent of n and the Type II class $-n$ moving with it. A dual identity replaces this by two classes, $-p$ and -1 , to the same modulus $4AC$ and applied to the same integer $pA + C$; the divisor $\delta = 1$ is available to the first class and never to the second — an asymmetry between the two types that is already visible at the smallest divisor.

Our main unconditional result holds for every numerator: writing a Type II solution of $m/p = 1/x + 1/(pY) + 1/(pZ)$ with $Y \leq Z$, the number of those with $Y \leq K$ is at most $\sum_{n \leq K} \tau(n) = K \log K + O(K)$, uniformly in p and m ; on average over $p \leq N$ it is only $O_m((\log K)^3 \log \log K)$, by Brun–Titchmarsh. Hence Type II solutions concentrate at $x/p \rightarrow 1/m$. No such concentration can hold for Type I, whose larger denominator coprime to p always exceeds $(p+1)/3$ and is not confined to any bounded multiple of p .

We then show that the numerator itself carries arithmetic information. For general m the Type II count is obstructed by -1 and the Type I count by $-m$ in $(\mathbb{Z}/e_n\mathbb{Z})^\times$, so a real character modulo e_n that is trivial on the primes dividing n annihilates Type II when $\chi(-1) = -1$ and Type I when $\chi(-1)\chi(m_0) = -1$, where m_0 is the squarefree kernel of m . These coincide exactly when m is a perfect square. For non-square m a single quadratic character therefore kills exactly one type, and quadratic reciprocity — applied to $e_n \equiv -p$ modulo the primes dividing n — decides which; for the Erdős–Straus numerator 4 the quadratic level is blind, so no criterion modulo squares can settle the comparison. A second and independent role of m is 2-adic: we prove $f_I^{(m)}(n) = f_{II}^{(m)}(n) = 0$ for every odd perfect square n whenever $4 \mid m$, which generalises a theorem of Elsholtz and Tao; no $m \leq 32$ outside that class has the property.

Finally we study the pointwise comparison. We prove the sharp local inequality $2B(n) \geq A(n)$ for $e_n \in \{1, 3, 5\}$, with a construction showing it fails for every odd e , $7 \leq e \leq 2 \cdot 10^4$; conjecture $f_I(p) > f_{II}(p)$ for all $p > 5$ with $p \not\equiv 1 \pmod{8}$, verified for $p \leq 10^6$; reduce it to a window of length $\approx p/12$ and then to a single inequality about the layers $e_n \geq 7$; prove it for an explicit set of primes; and generalise it to the assertion that $p \equiv -1 \pmod{m}$ forces $f_I^{(m)}(p) > f_{II}^{(m)}(p)$, verified for $4 \leq m \leq 20$ and $p \leq 2 \cdot 10^4$. A probabilistic model predicts $\sum_{p \leq N} f_I(p) / \sum_{p \leq N} f_{II}(p) \rightarrow 3$. We also locate the factor $\log \log N$ in the Elsholtz–Tao Type I bound: it comes only from Brun–Titchmarsh moduli exceeding $N^{1-o(1)}$, and deleting it would follow from gaining one further logarithm there.

2020 *Mathematics Subject Classification.* Primary 11D68; Secondary 11D72, 11N56, 11A25.

Key words and phrases. Erdős–Straus equation, Egyptian fractions, unit fractions, divisor function, counting solutions.

1. INTRODUCTION

For $n \in \mathbb{N}$ with $n \geq 2$ let

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}, \quad x, y, z \in \mathbb{N}, \quad (1.1)$$

and let $f(n)$ be the number of ordered triples (x, y, z) solving (1.1). The Erdős–Straus conjecture asserts that $f(n) > 0$ for every $n \geq 2$; it has been verified for $n \leq 10^{17}$ by Salez [18] and beyond in [12]. Since $f(mn) \geq f(n)$, it suffices to treat $n = p$ prime. Webb [24] showed that $f(n) > 0$ for almost all n , and Vaughan [23] proved that the number of $n \leq N$ with $f(n) = 0$ is at most $N \exp(-c \log^{2/3} N)$. Explicit identities of Obláth [16], Rosati [17] and Yamamoto [26] settle all n outside the six residue classes $n \equiv 1, 121, 169, 289, 361, 529 \pmod{840}$, and by Schinzel [19, 20] no finite family of polynomial identities can cover the remaining classes; see also Sierpiński [21], Mordell [15, pp. 287–290] and the survey in Guy [10, §D11]. The numerical verification has a long history of its own, from Terzi [22] to [18] and [12].

Elsholtz and Tao [9] initiated the systematic study of the counting function f . For an odd prime p , exactly one or exactly two of x, y, z are divisible by p . Following [9] we call a solution of (1.1) with $n = p$

- *Type I* if $p \mid x$ and $p \nmid yz$;
- *Type II* if $p \mid y, p \mid z$ and $p \nmid x$,

and we let $f_I(p)$, respectively $f_{II}(p)$, be the number of ordered pairs (y, z) arising in this way, so that

$$f(p) = 3f_I(p) + 3f_{II}(p). \quad (1.2)$$

Elsholtz and Tao proved

$$\sum_{p \leq N} f_{II}(p) \asymp N \log^2 N, \quad N \log^2 N \ll \sum_{p \leq N} f_I(p) \ll N \log^2 N \log \log N,$$

together with the pointwise bounds $f_I(n) \ll n^{3/5+o(1)}$ and $f_{II}(n) \ll n^{2/5+o(1)}$; the latter was extended to all m/n by Elsholtz and Planitzer [8].

This paper is about f_I and f_{II} , not about the Erdős–Straus conjecture. None of our results establishes solvability of (1.1) in any new case; all the congruence classes in which we obtain positivity are classes in which solvability is classical.

1.1. The local divisor counts. Call an integer n *admissible* for p if $4n > p$ and $p \nmid n$, and write

$$e_n := 4n - p \ (\geq 1), \quad \gcd(e_n, n) = \gcd(e_n, p) = 1. \quad (1.3)$$

Define

$$A(n) := \#\{s : s \mid n^2, e_n \mid s + n\}, \quad B(n) := \#\{s : s \mid n^2, e_n \mid s + pn\}. \quad (1.4)$$

Conditions of this shape have a long history. In the form of a solvability criterion they go back at least to Monks and Velingker [13, Cor. 2.3], who showed that (1.1) is solvable for p if and only if there is an n such that pn has two divisors summing to a multiple of $4n - p$ or of $4n - 1$; the underlying two-unit-fraction lemma is older still and is described as well known already in [13, Thm. 2.2]. The refinement of (1.4) into a *bijection* with the solutions of (1.1) *through the smallest denominator* is due to Bradford [5]. Our first observation is that, summed over *all* denominators coprime to p and with the correct multiplicities, these conditions compute f_I and f_{II} exactly.

Theorem 1.1 (Counting form of Bradford's conditions). *For every prime $p \geq 5$,*

$$f_{\text{II}}(p) = \sum_n A(n), \quad f_{\text{I}}(p) = \sum_n B(n),$$

both sums being over all n admissible for p ; only finitely many terms are nonzero.

Since $p \equiv 4n \pmod{e_n}$, the involution $s \mapsto n^2/s$ on the divisors of n^2 carries $\{s : s \equiv -pn\}$ bijectively onto $\{s : 4s \equiv -1\}$; the two sets are in general different, but they have the same cardinality, and therefore

$$f_{\text{I}}(p) = \sum_n \#\{s \mid n^2 : s \equiv -4^{-1} \pmod{e_n}\}, \quad f_{\text{II}}(p) = \sum_n \#\{s \mid n^2 : s \equiv -n \pmod{e_n}\} : \quad (1.5)$$

the same index set, the same modulus and the same divisor set, the only difference being the residue class. Note that the Type I class -4^{-1} is *independent of n* , while the Type II class $-n$ moves with n ; see §7.1.

1.2. Concentration of Type II solutions. Our main unconditional result describes how Type II solutions are distributed. It holds for every numerator, so we state it for

$$\frac{m}{p} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}, \quad m \geq 2, \quad p \nmid m, \quad (1.6)$$

calling a solution *Type II* when $p \mid y$, $p \mid z$ and $p \nmid x$. Write such a solution as

$$\frac{m}{p} = \frac{1}{x} + \frac{1}{pY} + \frac{1}{pZ}, \quad Y \leq Z, \quad p \nmid x. \quad (1.7)$$

Put

$$D(K) := \sum_{n \leq K} \tau(n) = K \log K + (2\gamma - 1)K + O(\sqrt{K}). \quad (1.8)$$

Theorem 1.2. *Let $m \geq 2$, let p be a prime with $p \nmid m$, and let $K \geq 1$. Then the number of unordered Type II solutions (1.7) with $Y \leq K$ is at most $D(K) \ll K \log K$. The bound depends neither on p nor on m .*

Corollary 1.3. *Let $m \geq 2$ and $0 < \delta \leq m - 1$. For every prime $p \nmid m$,*

$$\#\{\text{Type II solutions of (1.7) with } x \geq p/(m - \delta)\} \leq D(\lfloor 2/\delta \rfloor) = O_\delta(1),$$

uniformly in m and p . Equivalently, all but $O_\delta(1)$ of the Type II solutions satisfy $p/m < x < p/(m - \delta)$: they concentrate at $x/p \rightarrow 1/m$.

The proof uses nothing beyond the classical Type II parametrisation, which we record for general m in Lemma 2.2; this places Theorem 1.2 in the setting of Sierpiński and Schinzel [20] and of Elsholtz and Planitzer [8], where the numerator is arbitrary.

By contrast $f_{\text{II}}(p)$ can be as large as $p^{2/5+o(1)}$, and Type I solutions obey no analogous concentration: their two denominators coprime to p range over the whole interval $(p/4, 3p/4]$ and $((p+1)/3, \infty)$ respectively. Corollary 1.3 rests on the following elementary but apparently new bound, which is what makes Type II sparse away from $x \approx p/4$.

Theorem 1.4. *For every admissible n , with $e = e_n$,*

$$A(n) \leq 2 \left(\left\lfloor \frac{2n}{e} \right\rfloor - \left\lfloor \frac{n}{e} \right\rfloor \right) \leq 2 \left(\frac{n}{e} + 1 \right).$$

No such bound holds for $B(n)$, which is only restricted by $B(n) \leq \tau(n^2)$.

1.3. A sharp local inequality. The comparison of f_I and f_{II} reduces (Corollary 5.2 below) to the local inequality $2B(n) \geq A(n)$. We determine exactly for which e_n it holds unconditionally.

Theorem 1.5. *Let $p \geq 5$ be prime and let n be admissible with $e = e_n \in \{1, 3, 5\}$. Then $2B(n) \geq A(n)$. The range $e \leq 5$ cannot be enlarged: every e_n is odd, and for every odd e with $7 \leq e \leq 2 \cdot 10^4$ there exist p and n with $e_n = e$ and $2B(n) < A(n)$.*

1.4. A pointwise comparison. Mihnea and Dumitru [12] observed that “solutions of Type-1 abound relative to those of Type-2”, reporting 12 763 383 Type I against 5 838 200 Type II solutions in aggregate over the range they computed. The following is a pointwise refinement of that observation: numerically $f_I(p) > f_{II}(p)$ with very few exceptions, all of which are congruent to 1 modulo 8.

Conjecture 1.6. If $p > 5$ is prime and $p \not\equiv 1 \pmod{8}$, then $f_I(p) > f_{II}(p)$.

For all 78 496 primes $p \leq 10^6$ the inequality $f_I(p) \leq f_{II}(p)$ holds for exactly twelve primes, listed in Table 3; apart from $p = 5$ all of them satisfy $p \equiv 1 \pmod{8}$. Since the six residue classes modulo 840 to which the Erdős–Straus conjecture reduces [16, 17, 26] are all $\equiv 1 \pmod{8}$, Conjecture 1.6 makes an assertion *only about primes for which (1.1) is classically solvable*; in particular it does not imply the Erdős–Straus conjecture. We prove Conjecture 1.6 for an explicit set of primes (Theorem 9.2), reduce it to a window of length $\approx p/12$ (Corollary 5.2), and explain in §9.4 why the surplus of Type I over Type II solutions is a consequence of Theorems 1.4 and 1.2 rather than of any imbalance in the distribution of divisors. Theorem 4.1 recasts the whole comparison as one between two residue classes, $-p$ and -1 , to the same modulus. In §9.7 we give a probabilistic model for both counting functions; it predicts that the expected size of the surplus is a factor of 3.

Conjecture 1.7. $R(N) := \left(\sum_{p \leq N} f_I(p) \right) / \left(\sum_{p \leq N} f_{II}(p) \right) \rightarrow 3$ as $N \rightarrow \infty$.

The constant 3 arises as the ratio $\frac{1}{2} : \frac{1}{6}$ between the mean value of τ_3 at a single scale and its $1/n$ -weighted average over all smaller scales: the Type I count evaluates τ_3 at $y \asymp p$, the Type II count averages it over all $Y \leq p/4$. A least-squares fit of $a - b/\log N$ to 41 values of $R(N)$ with $10^3 \leq N \leq 10^6$ gives $a = 3.016$. Conjecture 1.7 also asserts $\sum_{p \leq N} f_I(p) \asymp N \log^2 N$ with no factor $\log \log N$, in agreement with the conjecture of Elsholtz and Tao [9] that the $\log \log N$ in their upper bound can be removed. Theorem 9.2 reduces Conjecture 1.6 to a single inequality $T(p) > -\sigma(p)$ (Conjecture 9.3) involving only the layers $e_n \geq 7$, precisely where the local inequality of Theorem 1.5 first fails; for $p \equiv 3 \pmod{4}$ the residual statement is simply $T(p) \geq 0$, which holds without exception for all 1 632 such primes below $3 \cdot 10^4$. Section 9.6 refines Conjecture 1.6 into a *filtered* form: for every X , the Type I solutions with invariant $\delta \leq X$ should outnumber the Type II ones. This is strictly stronger, yet in the range $p \leq 6000$ it fails for exactly ten primes, *all* of them $\equiv 1 \pmod{8}$; an initial segment of it is proved in Theorem 9.6. Finally, §9.8 records seven natural approaches to Conjecture 1.6 that fail, and §9.5 two more, all with explicit counterexamples.

1.5. The role of the numerator: a quadratic dichotomy. Sections 3–7 are written for the numerator 4, but the arguments do not use it. In §8 we carry them out for an arbitrary numerator $m \geq 3$, with $e_n = mn - p$, and find that the numerator enters the local analysis in a way that singles out the Erdős–Straus case. The Type II count is obstructed by -1 and the Type I count by $-m$ (Proposition 8.2), and a real character cannot separate these two elements exactly when m is a square.

Theorem 1.8 (Theorem 8.3). *Let $m \geq 3$, let $p \nmid m$ be prime, let n be m -admissible, and let χ be a real character modulo e_n with $\chi(q) = 1$ for every prime $q \mid n$. Write m_0 for the squarefree kernel of m . Then $A_m(n) > 0$ implies $\chi(-1) = 1$, and $B_m(n) > 0$ implies $\chi(-1)\chi(m_0) = 1$. Hence the obstruction is type-blind if m is a perfect square, and type-selective otherwise.*

Taking for χ the Jacobi symbol modulo e_n turns the hypothesis into a statement about the splitting of $-p$ at the primes dividing n , because $e_n \equiv -p \pmod{q}$ for every $q \mid n$; quadratic reciprocity then converts $\left(\frac{q}{e}\right)$ into $\pm\left(\frac{-p}{q}\right)$, and the conclusions become $e_n \equiv 1 \pmod{4}$ for Type II and $\left(\frac{-m_0}{e_n}\right) = 1$ for Type I (Corollary 8.4). Table 1 exhibits the dichotomy across $3 \leq m \leq 12$.

For the Erdős–Straus numerator $m = 4$ the dichotomy degenerates, and this is a genuine obstruction rather than a defect of the method: by Corollary 8.6, when m is a square the elements -1 and $-m$ have the same image in $(\mathbb{Z}/e_n\mathbb{Z})^\times/H_n(\mathbb{Z}/e_n\mathbb{Z})^{\times 2}$, so *no* criterion expressed through congruences modulo squares can distinguish Type I from Type II. Any proof of Conjecture 1.6 must therefore involve characters of order at least 3. This is consistent with, and refines, the observation of Elsholtz and Tao [9, Prop. 1.6] that solvability of (1.1) is not governed by congruences on p .

A second and quite different role of the numerator appears in §8.5. The theorem of Elsholtz and Tao that $f_I(n) = f_{II}(n) = 0$ for every odd perfect square n turns out to use about the numerator only that it is divisible by 4: we prove (Theorem 8.8) that $f_I^{(m)}(n) = f_{II}^{(m)}(n) = 0$ for every odd perfect square n and every m divisible by 4, and we check that no $m \leq 32$ outside that class has the property. Thus squareness of m governs what quadratic characters can see, while divisibility of m by 4 governs vanishing at odd squares, and the two conditions are independent of one another.

Finally, §8.6 places Conjecture 1.6 in a family. The hypothesis $p \equiv -1 \pmod{m}$ is exactly the condition under which the divisor $\delta = 1$ becomes available to the Type I class in the duality of Theorem 4.1; when it holds we prove $f_I^{(m)}(p) \geq 2\tau(M^2)$ and $f_{II}^{(m)}(p) \geq \tau(M^2)$ with $M = (p+1)/m$ (Proposition 8.10), and we conjecture that $f_I^{(m)}(p) > f_{II}^{(m)}(p)$ always follows (Conjecture 8.11). For $4 \leq m \leq 20$ and $p \leq 2 \cdot 10^4$ there are 643 primes with $f_I^{(m)}(p) \leq f_{II}^{(m)}(p)$, and not one of them satisfies $p \equiv -1 \pmod{m}$. Theorem 8.14 reduces Conjecture 8.11 to a window of length $2p/(m(m-2))$ and thereby proves it for all but two of the 3028 pairs (m, p) with $3 \leq m \leq 20$ and $p \leq 6 \cdot 10^3$.

1.6. Organisation, and what is not new. Section 2 fixes notation and recalls the parametrisations we use, all of which are due to others. Section 3 proves Theorem 1.1 and a rigidity statement. Section 4 proves the duality (Theorem 4.1) and its consequences. Section 5 determines the ranges of the denominators and derives the reduction. Section 6 contains Theorems 1.4 and 1.2. Section 7 develops the local analysis by characters and proves Theorem 1.5. Section 8 treats a general numerator and proves the quadratic dichotomy. Section 9 contains the numerical evidence, the diagnosis of §9.4, the probabilistic model of §9.7, the failed approaches of §9.8, and open questions. Section 10 analyses the factor $\log \log N$ in the Elsholtz–Tao Type I upper bound.

We stress what is *not* ours. The four-parameter bijections for Type I and Type II solutions go back to Aigner [1], Bernstein [3] and Mordell [15] and are stated in the form we use in [9, Prop. 2.2, 2.3, 2.6, 2.7]. The congruence conditions (1.4) appear as a solvability criterion in [13, Cor. 2.3], and their bijective interpretation is Bradford’s [5, Prop. 1–4]; the two-unit-fraction lemma behind them (Lemma 2.3) is classical, and is already called well known in [13, Thm. 2.2]. A necessary condition of the shape of Theorem 4.1, in the special case $c = 1$, is

[13, Thm. 2.1(g)]; see Remark 4.3. The criterion $p \mid 4Y - 1$ occurring in Theorem 3.2 appears inside the proof of [4, Lemma 4]. The classification of solutions with two equal denominators (Theorem 5.3(i),(ii)) is due to Monks and Velingker [13, Thm. 2.1(h)] and López [11]. The conjecture that $f_I(p) > 0$ and $f_{II}(p) > 0$ for every odd prime is stated by Chamberland [6], and the bound $\lceil p/4 \rceil \leq x \leq (p+1)/2$ for the smallest denominator is due to Monks and Velingker [13, Thm. 2.1(i)]. The aggregate preponderance of Type I over Type II solutions was observed numerically by Mihnea and Dumitru [12]. For recent work on Type II solvability from the algorithmic and congruence-theoretic sides see Dahan [7] and Xu [25]; neither addresses the counting functions f_I, f_{II} . The cubic model $abc = n^2(2n + a + b + c)$ obtained by shifting the denominators is the “shifted cubic equation” of Bello-Hernández, Benito and Fernández [2]; we do not use it. Finally, the quadratic obstruction of §8 is an adaptation of the argument of [9, Prop. 1.6], as explained in Remark 8.5; what we add is its formulation for a general numerator and the resulting square/non-square dichotomy. Theorem 8.8 is likewise not a new argument but a new range for an old one: the proof is that of [9, Prop. 1.6] verbatim, and our contribution is the observation that it uses only $4 \mid m$, together with the verification that this is sharp. Section 10 rests entirely on the argument of [9, §8–9]; the block decomposition, Theorems 10.3 and 10.4, and the fourth modulus of Proposition 10.5(iv) are ours, but the three moduli (i)–(iii) are the Type I analogues of the three used in [9, §9], and Lemma 10.2 is their argument rearranged.

All numerical statements below were checked by two independent implementations; the ranges are stated with each claim.

2. NOTATION AND KNOWN PARAMETRISATIONS

Throughout, $p \geq 5$ is a prime, τ is the divisor function and ω the number of distinct prime factors. For n admissible we keep the notation (1.3), (1.4). We write $M := (p+1)/4$ when $p \equiv 3 \pmod{4}$, and m_0 for the squarefree kernel (largest squarefree divisor) of an integer m .

We shall use the following classical parametrisations, in the form given in [9, §2].

Lemma 2.1 (Aigner, Bernstein, Mordell; see [9, Prop. 2.2, 2.6]). *Let $p \geq 5$ be prime. The map $(a, b, c, d) \mapsto (x, y, z) = (abdp, acd, bcd)$ is a bijection from*

$$\Sigma_I(p) = \{(a, b, c, d) \in \mathbb{N}^4 : \gcd(a, b) = 1, 4abcd = (a+b)p + c\}$$

onto the set of Type I solutions, and $(a, b, c, d) \mapsto (x, y, z) = (abd, acdp, bcdp)$ is a bijection from

$$\Sigma_{II}(p) = \{(a, b, c, d) \in \mathbb{N}^4 : \gcd(a, b) = 1, 4abcd = cp + a + b\}$$

onto the set of Type II solutions.

For §6 we need the Type II half of Lemma 2.1 for a general numerator. The derivation is the classical one; we give it because we could not find it recorded for arbitrary m .

Lemma 2.2. *Let $m \geq 2$ and let p be a prime with $p \nmid m$. The map $(a, b, c, d) \mapsto (x, Y, Z) = (abd, acd, bcd)$ is a bijection from*

$$\{(a, b, c, d) \in \mathbb{N}^4 : \gcd(a, b) = 1, mabcd = cp + a + b\}$$

onto the set of solutions of (1.7). For $m = 4$ this is [9, Prop. 2.6].

Proof. Given a solution of (1.7), put $e := mx - p$; subtracting $1/x$ gives $1/Y + 1/Z = e/x$, so $e > 0$, and $\gcd(e, x) = \gcd(p, x) = 1$. Write $g = \gcd(Y, Z)$, $Y = ga$, $Z = gb$ with $\gcd(a, b) = 1$. Then $x(a+b) = e gab = (mx - p)gab$, that is

$$x F = p gab, \quad F := m gab - a - b > 0.$$

A prime dividing both F and a would divide b , so $\gcd(F, ab) = 1$ and hence $F \mid pg$. If $p \nmid F$ then $F \mid g$, and $x = pgab/F$ would be divisible by p , contradicting $p \nmid x$. Therefore $F = pc$ with $c \mid g$; putting $d = g/c$ gives $x = gab/c = abd$, $Y = ga = acd$, $Z = bcd$, and $F = pc$ becomes $abcd = cp + a + b$. The converse is a direct verification, and the two maps are mutually inverse. $\square$

The next lemma is the bijective form of a classical criterion: $a/b = 1/u + 1/v$ is solvable if and only if b has two divisors whose sum is divisible by a ([13, Thm. 2.2], where it is called well known).

Lemma 2.3 (Two unit fractions). *Let $\alpha, \beta \in \mathbb{N}$ with $\gcd(\alpha, \beta) = 1$. The map $t \mapsto (u, v) = ((t + \beta)/\alpha, (\beta^2/t + \beta)/\alpha)$ is a bijection from $\{t : t \mid \beta^2, t \equiv -\beta \pmod{\alpha}\}$ onto the set of ordered pairs $(u, v) \in \mathbb{N}^2$ with $1/u + 1/v = \alpha/\beta$.*

Proof. If $1/u + 1/v = \alpha/\beta$ then $(\alpha u - \beta)(\alpha v - \beta) = \beta^2$, and $t := \alpha u - \beta$ is positive (else $1/u \geq \alpha/\beta$ forces $1/v \leq 0$), divides β^2 and satisfies $t \equiv -\beta \pmod{\alpha}$. Conversely such a t gives $u = (t + \beta)/\alpha \in \mathbb{N}$, while v is an integer because $t \cdot (\beta^2/t) = \beta^2$ and $\gcd(\beta, \alpha) = 1$ force $\beta^2/t \equiv \beta^2(-\beta)^{-1} \equiv -\beta \pmod{\alpha}$. The two maps are mutually inverse. $\square$

3. THE COUNTING IDENTITY AND RIGIDITY

Proof of Theorem 1.1. Fix an admissible n and put $e = e_n$. By (1.3), $\gcd(e, pn) = 1$. Let

$$R(n) := \left\{ (u, v) \in \mathbb{N}^2 : \frac{1}{u} + \frac{1}{v} = \frac{4}{p} - \frac{1}{n} = \frac{e}{pn} \right\}.$$

By Lemma 2.3 with $\alpha = e$, $\beta = pn$, the set $R(n)$ is in bijection with $\{t : t \mid (pn)^2, t \equiv -pn \pmod{e}\}$.

Since $p \nmid n$, each such t is uniquely $t = p^i s$ with $s \mid n^2$ and $i \in \{0, 1, 2\}$. With $u = (t + pn)/e$ and $v = ((pn)^2/t + pn)/e$ we get

$$p \mid u \iff p \mid t \iff i \geq 1, \quad p \mid v \iff p \mid (pn)^2/t = p^{2-i}n^2/s \iff i \leq 1.$$

Hence both u and v are divisible by p exactly when $i = 1$, and exactly one of them is divisible by p exactly when $i \in \{0, 2\}$. Translating $t \equiv -pn \pmod{e}$ into a condition on s (recall $\gcd(p, e) = 1$):

$$i = 0 : s \equiv -pn, \quad i = 1 : s \equiv -n, \quad i = 2 : s \equiv -np^{-1} \pmod{e}.$$

Now sum over n . A Type II solution has a unique denominator coprime to p , namely x ; it therefore contributes exactly one element of $R(x)$ with both coordinates divisible by p , namely (y, z) , and conversely. Hence $\sum_n \#\{(u, v) \in R(n) : p \mid u, p \mid v\} = f_{\text{II}}(p)$, which is the case $i = 1$, i.e. $\sum_n A(n) = f_{\text{II}}(p)$.

A Type I solution has two denominators coprime to p , and one checks directly that

$$\sum_n \#\{(u, v) \in R(n) : \text{exactly one of } u, v \text{ is divisible by } p\} = 2f_{\text{I}}(p)$$

(an unordered Type I solution with $y \neq z$ contributes 2 to $f_{\text{I}}(p)$ and 4 to the left-hand side, one with $y = z$ contributes 1 and 2). Finally the involution $s \mapsto n^2/s$ on the divisors of n^2 carries the class $-pn$ bijectively onto the class $n^2(-pn)^{-1} \equiv -np^{-1}$, so the cases $i = 0$ and $i = 2$ have equal cardinality $B(n)$. Therefore $2f_{\text{I}}(p) = \sum_n 2B(n)$.

Only finitely many terms are nonzero because (1.1) has finitely many solutions. $\square$

Corollary 3.1. *If $e_n \geq 3$ then $A(n)$ is even.*

Proof. The involution $s \mapsto n^2/s$ preserves the class $-n$, since $n^2(-n)^{-1} \equiv -n$. A fixed point satisfies $s^2 = n^2$, i.e. $s = n$, and $n \equiv -n \pmod{e_n}$ would force $e_n \mid 2n$, hence $e_n \mid 2$ by (1.3); as $e_n = 4n - p$ is odd this means $e_n = 1$. $\square$

The next statement sharpens a step in [4, Lemma 4] into a dichotomy.

Theorem 3.2 (Rigidity). *Let $p \geq 5$ be prime and let pV ($V \in \mathbb{N}$) be a denominator of some solution of (1.1) with $n = p$. If $p \mid 4V - 1$ then every solution having pV among its denominators is of Type I; if $p \nmid 4V - 1$ then every such solution is of Type II. Consequently a Type I and a Type II solution never share a denominator divisible by p , and they share at most one denominator, necessarily the unique denominator of the Type II solution that is coprime to p .*

Proof. Suppose $4/p = 1/(pX) + 1/y + 1/z$ with $p \nmid yz$. Multiplying by $pXyz$ gives $4Xyz = yz + pXz + pXy$, i.e. $yz(4X - 1) = pX(y + z)$; since $p \nmid yz$ we get $p \mid 4X - 1$.

Suppose next that $4/p = 1/x + 1/(pY) + 1/(pZ)$ with $p \nmid x$, and put $e = 4x - p > 0$. Subtracting $1/x$ gives $1/Y + 1/Z = e/x$ with $\gcd(e, x) = \gcd(p, x) = 1$, so by Lemma 2.3 there is $t \mid x^2$ with $t \equiv -x \pmod{e}$ and $Y = (t + x)/e$. Hence

$$4Y - 1 = \frac{4t + 4x - e}{e} = \frac{4t + p}{e}.$$

If $p \mid 4Y - 1$ then $p \mid 4t + p$, so $p \mid t$; but $t \mid x^2$ and $p \nmid x$, a contradiction. The same argument applies to Z .

Thus the sets of p -divisible denominators of Type I and of Type II solutions are disjoint, which is the dichotomy. A common denominator must therefore be coprime to p ; on the Type I side it is y or z , on the Type II side it is x , and the Type II solution has only one such denominator. $\square$

4. A DUALITY: THE SAME DIVISORS, THE CLASSES $-p$ AND -1

Theorem 1.1 organises the two counts by the denominators of the solution. There is a second, and in some ways cleaner, way to organise them, in which the two types are distinguished by a single residue class. For $A, C \in \mathbb{N}$ put

$$n = n(A, C) := pA + C, \quad \mu = \mu(A, C) := 4AC. \quad (4.1)$$

Theorem 4.1 (Duality). *Let $p \geq 5$ be prime.*

- (i) *The map $(a, b, c, d) \mapsto (A, C, \delta) = (a, c, 4acd - p)$ is a bijection from the set of unordered Type I solutions onto*

$$\left\{ (A, C, \delta) : \delta \mid n, \quad \delta \equiv -p \pmod{\mu}, \quad \gcd(A, \frac{n}{\delta}) = 1, \quad A \leq \frac{n}{\delta} \right\}.$$

- (ii) *The map $(a, b, c, d) \mapsto (A, C, \delta) = (c, a, 4acd - 1)$ is a bijection from the set of unordered Type II solutions onto*

$$\left\{ (A, C, \delta) : \delta \mid n, \quad \delta \equiv -1 \pmod{\mu}, \quad \gcd(C, \frac{n}{\delta}) = 1, \quad C \leq \frac{n}{\delta} \right\}.$$

In both cases $n/\delta = b$. Thus f_I and f_{II} count divisors of the same integer $n(A, C)$ in two residue classes to the same modulus $\mu(A, C)$, namely $-p$ and -1 .

Proof. (i) By Lemma 2.1 an unordered Type I solution is a quadruple (a, b, c, d) with $\gcd(a, b) = 1$, $a \leq b$ and $4abcd = p(a + b) + c$; the last equation is

$$b(4acd - p) = pa + c. \quad (4.2)$$

Set $A = a$, $C = c$, $\delta = 4acd - p$. Then $n = pA + C = pa + c$, $\mu = 4AC = 4ac$, so $\delta \equiv -p \pmod{\mu}$, and (4.2) says $\delta \mid n$ with $n/\delta = b$; also $\delta > 0$ since $b > 0$. The conditions $\gcd(a, b) = 1$ and $a \leq b$ are those displayed. Conversely, given (A, C, δ) in the displayed set, put $d := (\delta + p)/\mu \in \mathbb{N}$, $b := n/\delta$, $a := A$, $c := C$; then $4abcd = \mu bd = b(\delta + p) = n + pb = p(a + b) + c$, so (a, b, c, d) is a Type I solution. The two maps are mutually inverse.

(ii) Likewise $4abcd = cp + a + b$ is

$$b(4acd - 1) = pc + a, \quad (4.3)$$

and with $A = c$, $C = a$, $\delta = 4acd - 1$ we get $n = pA + C = pc + a$, $\mu = 4AC = 4ac$, $\delta \equiv -1 \pmod{\mu}$ and $n/\delta = b$. The rest is as in (i). $\square$

Remark 4.2. The only difference between the two descriptions is the residue class, together with the fact that the coprimality and ordering conditions are attached to A in the first case and to C in the second. We verified Theorem 4.1 numerically by enumerating both sides for all primes $5 \leq p \leq 150$.

Remark 4.3 (A precursor). A necessary condition of exactly this shape appears, in one special case, in [13, Thm. 2.1(g)]: if a solution of (1.1) with $n = p$ satisfies $z = p \operatorname{lcm}(x, y)$ — which in the parametrisation of Lemma 2.1 is precisely the case $c = 1$ — then $p^2 + \mu'$ has a divisor congruent to $-p$ modulo μ' , where $\mu' = 4\gcd(x, y)$. What Theorem 4.1 adds is that the condition is not merely necessary but bijective, that it covers all Type I solutions rather than those with $c = 1$, and above all that the Type II solutions are described by the *same* integer and the *same* modulus, with the single class -1 in place of $-p$; it is that coincidence which makes the comparison of f_I and f_{II} meaningful.

The two classes are not on an equal footing. This is the cleanest algebraic expression of the asymmetry between the two types; it is not, however, what makes f_I larger than f_{II} in order of magnitude — for that see the diagnosis in §9.4, and the two mechanisms identified at the end of §9.8.

Corollary 4.4. *Fix A, C and put $\mu = 4AC$ (≥ 4). The least positive element of the class $-1 \pmod{\mu}$ is $\mu - 1$. By contrast the least positive element of the class $-p \pmod{\mu}$ is $(-p) \bmod \mu$, which is coprime to μ because $p \nmid \mu$, and which can be as small as 1: as μ runs over the multiples of 4 with $p \nmid \mu$, the residue $(-p) \bmod \mu$ runs over all residues coprime to μ , and it equals 1 exactly when $\mu \mid p + 1$. In particular the divisor $\delta = 1$ of n never produces a Type II solution, while it produces a Type I solution exactly when $4AC \mid p + 1$ and $\gcd(A, C) = 1$.*

Proof. If $\delta = 1$ then $\delta \equiv -1 \pmod{\mu}$ forces $\mu \mid 2$, impossible as $\mu \geq 4$; and $\delta \equiv -p \pmod{\mu}$ reads $\mu \mid p + 1$. For $\delta = 1$ we have $n/\delta = n$, so the side conditions in Theorem 4.1(i) become $A \leq n$, which is automatic, and $\gcd(A, n) = \gcd(A, pA + C) = \gcd(A, C) = 1$. $\square$

Corollary 4.5. *Let $p \equiv 3 \pmod{4}$ and $M = (p + 1)/4$. Then*

$$\#\{\text{Type I solutions with } \delta = 1\} = \#\{(A, C) : AC \mid M, \gcd(A, C) = 1\} = \sum_{k \mid M} 2^{\omega(k)} = \tau(M^2),$$

so $f_I(p) \geq 2\tau(M^2)$.

Proof. By Corollary 4.4 the relevant pairs are those with $4AC \mid p + 1 = 4M$, i.e. $AC \mid M$, and $\gcd(A, C) = 1$. Grouping by $k = AC$ gives $\sum_{k \mid M} \#\{(A, C) : AC = k, \gcd(A, C) = 1\} = \sum_{k \mid M} 2^{\omega(k)} = \tau(M^2)$. Each of these solutions has $b = n > A = a$, hence $y \neq z$, so it contributes 2 to $f_I(p)$. $\square$

This recovers Proposition 9.1(i) without computation. The next corollary has no Type I analogue.

Corollary 4.6. *Every Type II solution with $a \leq b$ satisfies*

$$a \leq \frac{1 + \sqrt{1 + 4p}}{4} < \frac{\sqrt{p}}{2} + 1.$$

Proof. In the notation of Theorem 4.1(ii), $C = a$, $A = c$ and $\delta = 4acd - 1 \geq \mu - 1 = 4ac - 1$. The condition $C \leq n/\delta$ gives $a(4ac - 1) \leq a\delta \leq n = pc + a$, i.e. $c(4a^2 - p) \leq 2a$. If $4a^2 \leq p$ we are done; otherwise $c \geq 1$ forces $4a^2 - 2a - p \leq 0$, which is the assertion. $\square$

Elsholtz and Tao [9, Lemma 2.8] give $ab < p$ for Type II, whence $a < \sqrt{p}$; Corollary 4.6 improves the constant by a factor 2. For Type I no such bound holds: the parameter a there is only subject to $a \leq b$.

The duality can be pushed one step further, and the resulting form displays the asymmetry between the two types in its starkest shape.

Corollary 4.7 (Third form). *Let (a, b, c, d) parametrise a solution as in Lemma 2.1.*

- (i) *If the solution is of Type I and $\delta = 4acd - p$, then $\delta \mid (p^2 + 4c^2d)$; conversely, if $p \nmid 4cd$ then $\delta \mid (pa + c)$ is equivalent to $\delta \mid (p^2 + 4c^2d)$.*
- (ii) *If the solution is of Type II and $\delta = 4acd - 1$, then $\delta \mid (p + 4a^2d)$, and this is equivalent to $\delta \mid (pc + a)$.*

Proof. (i) With $A = a$, $C = c$, $D = d$, $n = pa + c$ and $\delta = 4acd - p$ we have

$$4cd \cdot n = 4cdpa + 4c^2d = p(4acd) + 4c^2d = p(\delta + p) + 4c^2d = p\delta + p^2 + 4c^2d,$$

so $\delta \mid n$ implies $\delta \mid (p^2 + 4c^2d)$. Conversely $\gcd(\delta, 4cd) = \gcd(4acd - p, 4cd) = \gcd(p, 4cd) = 1$ when $p \nmid 4cd$, and then the two divisibilities are equivalent.

(ii) With $A = c$, $C = a$, $n = pc + a$ and $\delta = 4acd - 1$,

$$4ad \cdot n = 4adpc + 4a^2d = p(4acd) + 4a^2d = p(\delta + 1) + 4a^2d = p\delta + p + 4a^2d,$$

and $\gcd(\delta, 4ad) = \gcd(-1, 4ad) = 1$ always. $\square$

Thus a Type I solution requires $4acd - p$ to divide a number of size $\asymp p^2$, whereas a Type II solution requires $4acd - 1$ to divide a number of size $\asymp p$. We verified both statements for all solutions with $p \leq 600$.

Remark 4.8. By Theorem 4.1, Conjecture 1.6 is the assertion that

$$\begin{aligned} & \sum_{A,C} \#\{\delta \mid n : \delta \equiv -p \pmod{\mu}, \gcd(A, \frac{n}{\delta}) = 1, A \leq \frac{n}{\delta}\} \\ & > \sum_{A,C} \#\{\delta \mid n : \delta \equiv -1 \pmod{\mu}, \gcd(C, \frac{n}{\delta}) = 1, C \leq \frac{n}{\delta}\}, \end{aligned}$$

with $n = pA + C$ and $\mu = 4AC$. Corollary 4.4 shows that the class $-p$ has access to small divisors of n and the class -1 does not; Corollary 4.6 shows that the second sum is supported on $C \ll \sqrt{p}$, whereas the first is not.

5. RANGES OF THE DENOMINATORS, AND A REDUCTION

Monks and Velingker [13, Thm. 2.1(i)] showed that the smallest denominator of any solution of (1.1) with $n = p$ satisfies $\lceil p/4 \rceil \leq x \leq (p+1)/2 \leq y$. For Type II solutions the following sharper, type-specific bound holds.

Theorem 5.1. *Let $p \geq 5$ be prime.*

- (i) *If $4/p = 1/x + 1/(pY) + 1/(pZ)$ is a Type II solution then $p/4 < x \leq (p+1)/3$, with equality on the right if and only if $3 \mid p+1$, in which case $\{Y, Z\} = \{1, (p+1)/3\}$.*
- (ii) *If $4/p = 1/(pX) + 1/y + 1/z$ is a Type I solution with $y \leq z$ then $z > (p+1)/3$.*

Proof. (i) The inequality $4x = p + e > p$ gives the left-hand bound. Assume $Y \leq Z$; as in the proof of Theorem 3.2, $1/Y + 1/Z = e/x$ with $e = 4x - p$. If $Y \geq 2$ then $e/x \leq 1$, i.e. $4x - p \leq x$ and $x \leq p/3$. If $Y = 1$ then $1/Z = e/x - 1 = (3x - p)/x > 0$, so $3x > p$ and $Z = x/(3x - p)$; thus $(3x - p) \mid x$, hence $(3x - p) \mid 3x$ and $(3x - p) \mid p$, so $3x - p \in \{1, p\}$. The value $3x - p = p$ gives $3x = 2p$, impossible since $3 \nmid p$. Hence $3x - p = 1$, $x = (p+1)/3$ and $Z = x$.

(ii) If $y \leq z \leq (p+1)/3$ then $1/y + 1/z \geq 6/(p+1)$, while $1/y + 1/z = 4/p - 1/(pX) < 4/p$. But $6/(p+1) \geq 4/p$ for $p \geq 2$, a contradiction. $\square$

Let

$$W = W(p) := \{n \in \mathbb{Z} : p/4 < n \leq (p+1)/3\}, \quad \#W \approx p/12. \quad (5.1)$$

Corollary 5.2 (Reduction). *For every prime $p \geq 5$,*

$$f_{\text{II}}(p) = \sum_{n \in W} A(n), \quad f_{\text{I}}(p) \geq 2 \sum_{n \in W} B(n).$$

In particular, if $\sum_{n \in W} (2B(n) - A(n)) > 0$ then $f_{\text{I}}(p) > f_{\text{II}}(p)$.

Proof. The first identity is Theorem 1.1 together with Theorem 5.1(i), which shows $A(n) = 0$ for $n \notin W$.

For the second, let S be the number of unordered Type I solutions and S_1 the number of those whose smaller denominator coprime to p lies in W . By Theorem 5.1(ii) the larger such denominator always exceeds $(p+1)/3$, so each unordered Type I solution is counted at most once by $\sum_{n \in W} B(n)$; hence $\sum_{n \in W} B(n) = S_1$. By Theorem 5.3(i) below, $f_{\text{I}}(p) = 2S - \delta$ where $\delta = 1$ if $p \equiv 3 \pmod{4}$ and $\delta = 0$ otherwise, and the unique solution with $y = z$ has $y = z = (p+1)/2 > (p+1)/3$, so it is not counted in S_1 and $S \geq S_1 + \delta$. Therefore $f_{\text{I}}(p) = 2S - \delta \geq 2S_1 + \delta \geq 2S_1$. $\square$

Theorem 5.3 (Diagonal solutions and parity). *Let $p \geq 5$ be prime.*

- (i) (Monks–Velingker [13, Thm. 2.1(h)]) *There is a Type I solution with $y = z$ if and only if $p \equiv 3 \pmod{4}$, and then it is unique, namely $4/p = 1/(p(p+1)/4) + 2/((p+1)/2)$.*
- (ii) (cf. López [11]) *There is a Type II solution with $y = z$ if and only if $p \equiv 3 \pmod{4}$, and then it is unique, namely $4/p = 1/((p+1)/4) + 2/(p(p+1)/2)$.*
- (iii) *Consequently $f_{\text{I}}(p) \equiv f_{\text{II}}(p) \equiv \mathbf{1}_{p \equiv 3 \pmod{4}} \pmod{2}$, and $6 \mid f(p)$ for every odd prime p .*

Proof. (i) From $4/p = 1/(pX) + 2/y$ we get $X = y/(4y - 2p)$. Put $g = 4y - 2p > 0$; then $g \mid y$, hence $g \mid 4y$ and $g \mid 2p$. As g is even, $g \in \{2, 2p\}$. If $g = 2p$ then $y = p$, contradicting $p \nmid y$. If $g = 2$ then $y = (p+1)/2$ and $X = (p+1)/4$, which is an integer precisely when $p \equiv 3 \pmod{4}$; the displayed identity is immediate.

(ii) From $4/p = 1/x + 2/(pY)$ we get $x = pY/(2(2Y - 1))$. Since $\gcd(2Y - 1, Y) = 1$ we obtain $(2Y - 1) \mid p$, so $2Y - 1 \in \{1, p\}$. The value $2Y - 1 = 1$ gives $x = p/2 \notin \mathbb{N}$; the value $2Y - 1 = p$ gives $Y = (p+1)/2$ and $x = (p+1)/4$, again requiring $p \equiv 3 \pmod{4}$.

(iii) $f_{\text{I}}(p)$ counts ordered pairs (y, z) : an unordered solution with $y \neq z$ contributes 2, one with $y = z$ contributes 1. Hence $f_{\text{I}}(p)$ is congruent modulo 2 to the number of Type I solutions with $y = z$, and similarly for f_{II} . Now use (i), (ii) and (1.2). $\square$

6. CONCENTRATION OF TYPE II SOLUTIONS

Proof of Theorem 1.4. By Theorem 1.1 and Lemma 2.3 (with $\alpha = e$, $\beta = n$), $A(n)$ equals the number of ordered pairs $(U, V) \in \mathbb{N}^2$ with $1/U + 1/V = e/n$. If $U \leq V$ then $e/n \leq 2/U$ gives $U \leq 2n/e$, while $e/n > 1/U$ gives $U > n/e$. Thus U lies in $(n/e, 2n/e]$, an interval containing exactly $\lfloor 2n/e \rfloor - \lfloor n/e \rfloor \leq n/e + 1$ integers, and U determines V . Counting ordered pairs doubles this. $\square$

Proof of Theorem 1.2. By Lemma 2.2 an unordered Type II solution of (1.7) is a quadruple (a, b, c, d) with $\gcd(a, b) = 1$, $a \leq b$ and $mabcd = cp + a + b$. Put $g := cd$; since $\gcd(a, b) = 1$ we have $g = \gcd(Y, Z)$, $Y = ag$ and $c \mid g$. Rewrite the equation as

$$b(mag - 1) = pc + a. \quad (6.1)$$

Note $\gcd(p, mag - 1) = 1$: if $p \mid mag - 1$ then (6.1) gives $p \mid a$, hence $p \mid b$ by (6.1) again, and then $p \mid x = abd$, contrary to $p \nmid x$.

Now fix a and g . By (6.1),

$$c \equiv -ap^{-1} \pmod{mag - 1},$$

so c lies in one residue class modulo $mag - 1$. On the other hand $1 \leq c \leq g$, and $g \leq mag - 1$ because $g(ma - 1) \geq g \geq 1$. An interval of length $g \leq mag - 1$ contains at most one element of a fixed class modulo $mag - 1$, so c is uniquely determined by the pair (a, g) ; then $d = g/c$, and b is determined by (6.1). Hence the solution itself is determined by (a, g) , and

$$\#\{\text{Type II solutions with } Y \leq K\} \leq \#\{(a, g) \in \mathbb{N}^2 : ag \leq K\} = \sum_{n \leq K} \tau(n) = D(K).$$

The estimate $D(K) = K \log K + (2\gamma - 1)K + O(\sqrt{K})$ is Dirichlet's. $\square$

Proof of Corollary 1.3. For a Type II solution of (1.7) we have $1/Y + 1/Z = e/x$ with $e = mx - p$, so $Y \leq Z$ gives $2/Y \geq e/x$, i.e. $Y \leq 2x/e$. If $x \geq p/(m - \delta)$ then $e = mx - p \geq \delta x$, whence $Y \leq 2/\delta$. Now apply Theorem 1.2 with $K = \lfloor 2/\delta \rfloor$. $\square$

Remark 6.1. For $K = 1$ Theorem 1.2 gives $D(1) = 1$: there is at most one Type II solution with $Y = 1$. This matches Theorem 5.1(i), where the extremal case $x = (p + 1)/3$ has $Y = 1$ and occurs exactly when $3 \mid p + 1$.

The pointwise bound $D(K) \ll K \log K$ of Theorem 1.2 is far from what the data suggest (Remark 6.3). On average over p we can do much better.

Theorem 6.2. *Let $m \geq 2$ and $2 \leq K \leq \sqrt{N}$. Then*

$$\sum_{\substack{p \leq N \\ p \nmid m}} \#\{\text{Type II solutions of (1.7) with } Y \leq K\} \ll \frac{N}{\log N} \cdot \frac{(\log K)^3 \log \log(3mK)}{m}.$$

In particular the average over the primes $p \leq N$ is $O(m^{-1}(\log K)^3 \log \log K)$, whereas the pointwise bound of Theorem 1.2 is of size $K \log K$.

Proof. By Lemma 2.2 a Type II solution with $Y \leq K$ is a quadruple (a, b, c, d) with $\gcd(a, b) = 1$ and $mabcd = cp + a + b$, and $Y = acd \leq K$. Put $\delta := macd - 1 = mY - 1$. The equation reads $b\delta = pc + a$, so $\delta \mid (pc + a)$; moreover $\gcd(c, \delta) = 1$ because $\delta \equiv -1 \pmod{mac}$. Hence

$$p \equiv -ac^{-1} \pmod{\delta},$$

a fixed residue class r modulo δ , coprime to δ since $\gcd(a, \delta) = 1$ as well. For a given triple (a, c, d) the integer b is determined, so there is at most one solution; therefore

$$\sum_{p \leq N} \#\{\dots\} \leq \sum_{\substack{(a,c,d) \in \mathbb{N}^3 \\ acd \leq K}} \pi(N; \delta, r), \quad \delta = m acd - 1 \leq mK - 1 \leq m\sqrt{N}.$$

The terms with $\delta = 1$ (possible only for $m = 2, Y = 1$) contribute at most $\pi(N)$. For $\delta \geq 2$, Brun–Titchmarsh [14] gives $\pi(N; \delta, r) \leq 2N/(\varphi(\delta) \log(N/\delta)) \ll N/(\varphi(\delta) \log N)$, since $\delta \leq m\sqrt{N}$. Using $\varphi(n) \gg n/\log \log 3n$ and grouping the triples (a, c, d) by $Y = acd$, whose number is $\tau_3(Y)$,

$$\sum_{p \leq N} \#\{\dots\} \ll \frac{N}{\log N} \sum_{Y \leq K} \frac{\tau_3(Y) \log \log(3mY)}{mY} \ll \frac{N}{\log N} \cdot \frac{\log \log(3mK)}{m} \sum_{Y \leq K} \frac{\tau_3(Y)}{Y},$$

and $\sum_{Y \leq K} \tau_3(Y)/Y \sim (\log K)^3/6$. $\square$

Remark 6.3. For $m = 4$ the maximum of $\#\{\text{Type II solutions} : Y \leq K\}$ over $5 < p \leq 4 \cdot 10^4$ is

$$1, 2, 5, 10, 14, 18, 24, 29, 38 \quad \text{for } K = 1, 2, 4, 8, 16, 32, 64, 128, 256,$$

against $D(K) = 1, 3, 8, 20, 50, 119, 280, 645, 1466$; the ratio to K decreases steadily, namely 1.00, 1.00, 1.25, 1.25, 0.88, 0.56, 0.38, 0.23, 0.15, so even $O(K)$ appears not to be optimal. The mean over the same primes is 0.50, 1.00, 1.94, 2.92, 4.61, 6.59, 9.11, 11.98, 15.16, consistent with the $(\log K)^3$ of Theorem 6.2. We verified Theorem 1.2 for $m \in \{3, 4, 5, 6\}$ and all primes $p \leq 3000$, and Theorem 6.2 numerically for $m \in \{3, 4, 6\}$, $K \leq 256$, $p \leq 2 \cdot 10^4$.

7. LOCAL ANALYSIS BY CHARACTERS

Throughout this section n is admissible, $e = e_n$, and we write $N(c) := \#\{s : s \mid n^2, s \equiv c \pmod{e}\}$ for $c \in (\mathbb{Z}/e\mathbb{Z})^\times$; note $\gcd(s, e) = 1$ for every $s \mid n^2$.

7.1. The asymmetry between the two classes. Let ι denote the involution $s \mapsto n^2/s$ on the divisors of n^2 ; it carries the class c bijectively onto the class $n^2 c^{-1}$.

Proposition 7.1. *A class c is preserved by ι if and only if $c \equiv nu$ with $u^2 \equiv 1 \pmod{e}$; for odd e there are $2^{\omega(e)}$ such u . The Type II class $-n$ is always preserved, and the Type I class -4^{-1} is preserved if and only if $e \mid p^2 - 1$.*

Proof. $c \equiv n^2 c^{-1}$ is equivalent to $(cn^{-1})^2 \equiv 1$; the 2-torsion of $(\mathbb{Z}/e\mathbb{Z})^\times$ has order $2^{\omega(e)}$ for odd e . Taking $u = -1$ gives $c = -n$. For $c = -4^{-1}$ the condition reads $16n^2 \equiv 1$, i.e. $(4n)^2 \equiv 1$, i.e. $p^2 \equiv 1 \pmod{e}$, since $4n \equiv p \pmod{e}$. $\square$

Thus $A(n)$ counts divisors in the unique non-trivial ι -stable class, while $B(n)$ counts divisors in a class which is generically not ι -stable, and which by (1.5) does not depend on n .

7.2. A character formula. For a Dirichlet character χ modulo e set

$$D_a(\zeta) := \sum_{k=-a}^a \zeta^k, \quad R_\chi(n) := \prod_{q^a \parallel n} D_a(\chi(q)).$$

Proposition 7.2. *$D_a(\zeta) \in \mathbb{R}$ for $|\zeta| = 1$, so $R_\chi(n) \in \mathbb{R}$ and $R_{\bar{\chi}} = R_\chi$. Moreover $\sum_{s \mid n^2} \chi(s) = \chi(n) R_\chi(n)$, and therefore, writing $\mathcal{W}(u) := \varphi(e)^{-1} \sum_{\chi \pmod{e}} \chi(u) R_\chi(n)$,*

$$N(c) = \mathcal{W}(nc^{-1}), \quad \mathcal{W}(u) = \mathcal{W}(u^{-1}), \quad A(n) = \mathcal{W}(-1), \quad B(n) = \mathcal{W}(-p^{-1}).$$

Proof. $D_a(\zeta) = 1 + \sum_{k=1}^a (\zeta^k + \zeta^{-k})$ is real for $|\zeta| = 1$. Multiplicativity gives

$$\sum_{s|n^2} \chi(s) = \prod_{q^a \| n} \sum_{k=0}^{2a} \chi(q)^k = \prod_{q^a \| n} \chi(q)^a D_a(\chi(q)) = \chi(n) R_\chi(n),$$

whence $N(c) = \varphi(e)^{-1} \sum_\chi \bar{\chi}(c) \chi(n) R_\chi(n) = \mathcal{W}(nc^{-1})$. Replacing χ by $\bar{\chi}$ and using $R_{\bar{\chi}} = R_\chi$ gives $\mathcal{W}(u) = \mathcal{W}(u^{-1})$. Finally $A(n) = N(-n) = \mathcal{W}(-1)$ and $B(n) = N(-pn) = \mathcal{W}(-p^{-1})$. $\square$

7.3. Local vanishing. Let $H_n \leq (\mathbb{Z}/e\mathbb{Z})^\times$ be the subgroup generated by the primes dividing n . Every divisor of n^2 lies in H_n .

Proposition 7.3. $A(n) > 0$ implies $-1 \in H_n$, and $B(n) > 0$ implies $-4 \in H_n$.

Proof. If $s \equiv -n$ for some $s | n^2$ then $-n \in H_n$, and $n \in H_n$, so $-1 \in H_n$. If $s \equiv -pn \equiv -4n^2$ then $-4n^2 \in H_n$, hence $-4 \in H_n$. $\square$

Proposition 7.4. Let χ be a real character modulo e with $\chi(q) = 1$ for every prime $q | n$. If $\chi(-1) = -1$ then $A(n) = B(n) = 0$.

Proof. The hypotheses give $\chi(s) = 1$ for every $s | n^2$, and $\chi(n) = 1$. If $A(n) > 0$, take $s \equiv -n$; then $1 = \chi(s) = \chi(-1)\chi(n) = -1$. If $B(n) > 0$, take $s \equiv -4n^2$; then $1 = \chi(s) = \chi(-1)\chi(2)^2\chi(n)^2 = \chi(-1) = -1$ (here e is odd, so $\chi(2) = \pm 1$). $\square$

Proposition 7.4 says that a quadratic obstruction annihilates *both* types simultaneously; in particular no argument at the quadratic level can distinguish f_I from f_{II} . This is the local counterpart of [9, Prop. 1.6]. In §8 we show that this coincidence is not an accident of the numerator 4 being small, but of its being a *square*: for a non-square numerator the same quadratic obstruction annihilates exactly one of the two types, and which one is decided by quadratic reciprocity.

7.4. Sharpness. The failure of $2B \geq A$ beyond $e = 5$ is not an accident of small cases: it follows from a two-line construction whenever a pair of primes of a prescribed shape exists. Note first that $e_n = 4n - p$ is always odd.

Proposition 7.5. Let $e \geq 7$ be odd and suppose that for some $k \geq 1$ both

$$q := ke - 1 \quad \text{and} \quad p := (4k - 1)e - 4$$

are prime and $p \neq q$. Then q is admissible for p , $e_q = e$, and

$$A(q) = 2, \quad B(q) = 0,$$

so that $2B(q) < A(q)$.

Proof. First $4q - p = 4(ke - 1) - ((4k - 1)e - 4) = e$, so $e_q = e > 0$, $4q > p$, and $p \nmid q$ because $p \neq q$ are primes; thus q is admissible. Since $q \equiv -1 \pmod{e}$ we have $\gcd(q, e) = 1$, the divisors of q^2 are $1, q, q^2$, and they are congruent to $1, -1, 1$ modulo e . Moreover $p \equiv 4q \equiv -4 \pmod{e}$.

For $A(q)$ the condition on s is $s \equiv -q \equiv 1 \pmod{e}$. It holds for $s = 1$ and $s = q^2$, and fails for $s = q$ because $-1 \equiv 1$ would force $e | 2$. Hence $A(q) = 2$.

For $B(q)$ the condition is $s \equiv -pq \equiv -(-4)(-1) = -4 \pmod{e}$. For $s \in \{1, q^2\}$ this reads $1 \equiv -4$, i.e. $e | 5$; for $s = q$ it reads $-1 \equiv -4$, i.e. $e | 3$. Both are impossible for $e \geq 7$. Hence $B(q) = 0$. $\square$

Whether every odd $e \geq 7$ admits such a k is a question of Hardy–Littlewood type: it asks for a k making the two linear forms $ke - 1$ and $4ek - (e + 4)$ simultaneously prime. That pair is admissible for every odd $e \geq 7$. Indeed, let ℓ be prime. If $\ell \mid e$ then the two forms are $\equiv -1$ and $\equiv -4$ modulo ℓ , and ℓ is odd, so neither vanishes; if $\ell = 2$ then $4ek - (e + 4)$ is odd and $ke - 1$ is odd for k even; and if ℓ is odd with $\ell \nmid e$ then each form vanishes on one class modulo ℓ , leaving at least $\ell - 2 \geq 1$ classes free. Hence:

Corollary 7.6. *Assume the Hardy–Littlewood prime k -tuples conjecture. Then for every odd $e \geq 7$ there are a prime p and an admissible n with $e_n = e$ and $2B(n) < A(n)$.*

Unconditionally we verified the hypothesis of Proposition 7.5 for all 9998 odd e with $7 \leq e \leq 20001$; the largest k required was 1822, at $e = 18899$. This is the source of the sharpness assertion in Theorem 1.5, which is therefore a finite computation, not a theorem about all e ; Table 4 shows in addition that within that range the failures are not rare.

7.5. Proof of Theorem 1.5.

Proof. Write $\tau := \tau(n^2)$.

Case $e = 1$. Both congruences are vacuous, so $A(n) = B(n) = \tau$ and $2B - A = \tau \geq 0$.

Case $e = 3$ (so $p \equiv 1 \pmod{4}$ and $n = (p + 3)/4$). From $4n = p + 3$ and $4 \equiv 1 \pmod{3}$ we get $n \equiv p \pmod{3}$, in particular $3 \nmid n$. Factor $n = n_1 n_2$ where n_1 (resp. n_2) collects the primes $\equiv 1$ (resp. $\equiv 2$) modulo 3, and set $\tau_j = \tau(n_j^2)$, so $\tau = \tau_1 \tau_2$. For the non-trivial character χ modulo 3, $\sum_{s|n^2} \chi(s) = \prod_{q \equiv 1} (2a_q + 1) \cdot \prod_{q \equiv 2} 1 = \tau_1$, whence

$$\#\{s \mid n^2 : s \equiv 1\} = \frac{\tau + \tau_1}{2}, \quad \#\{s \mid n^2 : s \equiv 2\} = \frac{\tau - \tau_1}{2}.$$

The Type I class is $-pn \equiv -p^2 \equiv -1 \equiv 2 \pmod{3}$, so $B(n) = (\tau - \tau_1)/2$. The Type II class is $-n \equiv -p \pmod{3}$. If $p \equiv 1 \pmod{3}$ then $A(n) = (\tau - \tau_1)/2 = B(n)$ and $2B - A = B \geq 0$. If $p \equiv 2 \pmod{3}$ then $A(n) = (\tau + \tau_1)/2$ and $2B - A = (\tau - 3\tau_1)/2$; but $n \equiv p \equiv 2 \pmod{3}$ while $n_1 \equiv 1 \pmod{3}$, so $n_2 \equiv 2 \pmod{3}$, in particular $n_2 > 1$ and $\tau_2 \geq 3$, giving $\tau = \tau_1 \tau_2 \geq 3\tau_1$.

Case $e = 5$ (so $p \equiv 3 \pmod{4}$ and $n = (p + 5)/4$). Factor $n = n_1 n_2 n_3 n_4$ according to the residue of the prime modulo 5, and set $\tau_j = \tau(n_j^2)$, so $\tau = \tau_1 \tau_2 \tau_3 \tau_4$. The characters modulo 5 are χ_0 , the quadratic character χ_2 (with $\chi_2(1) = \chi_2(4) = 1$, $\chi_2(2) = \chi_2(3) = -1$) and a pair $\chi, \bar{\chi}$ of order 4 with $\chi(2) = i$, $\chi(4) = -1$, $\chi(3) = -i$. A direct computation gives

$$D_a(-1) = (-1)^a, \quad D_a(\pm i) = \begin{cases} +1, & a \equiv 0, 1 \pmod{4}, \\ -1, & a \equiv 2, 3 \pmod{4}, \end{cases}$$

whence, with $\rho := R_{\chi_2}(n)$ and $\sigma := R_{\chi}(n) = R_{\bar{\chi}}(n)$,

$$\rho = \chi_2(n) \tau_1 \tau_4, \quad |\sigma| = \tau_1.$$

By Proposition 7.2, $\mathcal{W}(u) = \frac{1}{4}(\tau + \chi_2(u)\rho + 2\operatorname{Re}\chi(u) \cdot \sigma)$ and $A(n) = \mathcal{W}(-1) = \mathcal{W}(4) = \frac{1}{4}(\tau + \rho - 2\sigma)$. Since $4^{-1} \equiv 4$ and $-4 \equiv 1 \pmod{5}$, and $p \equiv 4n \pmod{5}$, we get $-p^{-1} \equiv -4^{-1}n^{-1} \equiv n^{-1}$, so $B(n) = \mathcal{W}(n)$. We distinguish three cases according to $n \pmod{5}$ (recall $5 \nmid n$).

If $n \equiv 4$, then $\mathcal{W}(n) = \mathcal{W}(4) = A(n)$ and $2B - A = A \geq 0$.

If $n \equiv 2$ or 3 , then $\operatorname{Re}\chi(n) = 0$ and $\chi_2(n) = -1$, so $\rho = -\tau_1 \tau_4$, $B(n) = \frac{1}{4}(\tau - \rho)$ and $4(2B - A) = \tau - 3\rho + 2\sigma = \tau + 3\tau_1 \tau_4 + 2\sigma$. Since $\tau \geq \tau_1$, $\tau_4 \geq 1$ and $|\sigma| \leq \tau_1$, this is at least $\tau_1 + 3\tau_1 - 2\tau_1 = 2\tau_1 > 0$.

If $n \equiv 1$, then $\chi_2(n) = 1$, $\rho = \tau_1 \tau_4$, $\operatorname{Re}\chi(n) = 1$ and $4(2B - A) = \tau + \rho + 6\sigma$. If $\sigma \geq 0$ we are done. If $\sigma < 0$ then at least one factor of σ is negative, so one of the following holds: there is a prime $q \equiv 4 \pmod{5}$ with $q^a \parallel n$ and a odd, in which case $\tau_4 \geq 2a + 1 \geq 3$ and

$\tau + \rho = \tau_1 \tau_4 (\tau_2 \tau_3 + 1) \geq 6\tau_1 = 6|\sigma|$; or there is a prime $q \equiv \pm 2 \pmod{5}$ with $q^a \parallel n$ and $a \equiv 2, 3 \pmod{4}$, in which case $a \geq 2$, $\tau_2 \tau_3 \geq 2a + 1 \geq 5$ and again $\tau + \rho = \tau_1 \tau_4 (\tau_2 \tau_3 + 1) \geq 6\tau_1 = 6|\sigma|$. In both cases $\tau + \rho + 6\sigma \geq 0$.

Sharpness is the numerical statement recorded in Table 4. $\square$

8. GENERAL NUMERATORS: A QUADRATIC DICHOTOMY GOVERNED BY m

Everything in §§3–7 was stated for the numerator 4. In this section we show that the counting identity and the local obstructions persist for an arbitrary numerator m , and that the resulting quadratic obstruction is sensitive to m in a way that is invisible when m is a perfect square. Since 4 is a perfect square, this isolates a structural reason why the Erdős–Straus case is harder than a typical m , and why the comparison f_I versus f_{II} cannot be settled by any argument at the quadratic level.

8.1. The counting identity for a general numerator. Fix $m \geq 3$ and a prime $p \nmid m$, and consider

$$\frac{m}{p} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}, \quad x, y, z \in \mathbb{N}. \quad (8.1)$$

Multiplying by $pxyz$ gives $mxyz = p(xy + yz + zx)$, so p divides at least one denominator. If p divides all three, then writing $x = px'$ and so on gives $1/x' + 1/y' + 1/z' = m$, which for $m \geq 3$ forces $m = 3$ and $x = y = z = p$. Apart from that one solution, every solution of (8.1) has exactly one denominator divisible by p (*Type I*) or exactly two (*Type II*); we write $f_I^{(m)}(p)$ and $f_{II}^{(m)}(p)$ for the numbers of ordered solutions of each type. Call $n \in \mathbb{N}$ *m -admissible* if $mn > p$ and $p \nmid n$, and put

$$e_n := mn - p, \quad A_m(n) := \#\{s \mid n^2 : s \equiv -n\}, \quad B_m(n) := \#\{s \mid n^2 : s \equiv -mn^2\}, \quad (8.2)$$

the congruences being taken modulo e_n . Note $\gcd(e_n, pn) = 1$ and $\gcd(e_n, m) = 1$, because $\gcd(e_n, n) = \gcd(p, n) = 1$ and $\gcd(e_n, m) = \gcd(p, m) = 1$.

Proposition 8.1. *For $m \geq 3$ and $p \nmid m$,*

$$f_{II}^{(m)}(p) = \sum_n A_m(n), \quad f_I^{(m)}(p) = \sum_n B_m(n),$$

the sums being over m -admissible n (for $m = 3$ the solution (p, p, p) is excluded). Moreover $B_m(n) = \#\{s \mid n^2 : s \equiv -m^{-1} \pmod{e_n}\}$, a class independent of n .

Proof. The proof of Theorem 1.1 applies verbatim with 4 replaced by m : one has $m/p - 1/n = e_n/(pn)$, and Lemma 2.3 with $\alpha = e_n$, $\beta = pn$ puts the pairs (u, v) with $1/u + 1/v = e_n/(pn)$ in bijection with the divisors $t \mid (pn)^2$ in the class $-pn$; writing $t = p^i s$ with $s \mid n^2$ and $i \in \{0, 1, 2\}$, the value $i = 1$ gives the Type II solutions and the classes $i = 0, 2$ are exchanged by $s \mapsto n^2/s$ and each give the Type I solutions. Finally $p \equiv mn \pmod{e_n}$, so the class $-pn$ of $i = 0$ is the class $-mn^2$; and $s \equiv -mn^2$ is equivalent to $n^2/s \equiv -m^{-1}$. $\square$

8.2. The local obstruction for a general numerator. Let $H_n \leq (\mathbb{Z}/e_n\mathbb{Z})^\times$ be the subgroup generated by the primes dividing n ; every divisor of n^2 lies in H_n , and $n \in H_n$.

Proposition 8.2. *$A_m(n) > 0$ implies $-1 \in H_n$, and $B_m(n) > 0$ implies $-m \in H_n$.*

Proof. If $s \equiv -n$ for some $s \mid n^2$ then $-n \in H_n$, and $n \in H_n$, so $-1 \in H_n$. If $s \equiv -mn^2$ then $-mn^2 \in H_n$ and $n^2 \in H_n$, so $-m \in H_n$. $\square$

Thus the two types are obstructed by two *different* elements of $(\mathbb{Z}/e_n\mathbb{Z})^\times$: the Type II obstruction -1 does not depend on m , while the Type I obstruction $-m$ does. The whole point of what follows is that a quadratic character cannot tell -1 from $-m$ exactly when m is a perfect square.

Theorem 8.3 (Quadratic dichotomy). *Let $m \geq 3$, let $p \nmid m$ be prime, let n be m -admissible and $e = e_n$. Let m_0 be the squarefree kernel of m , and let χ be a real character modulo e with $\chi(q) = 1$ for every prime $q \mid n$. Then*

$$A_m(n) > 0 \implies \chi(-1) = 1, \quad B_m(n) > 0 \implies \chi(-1)\chi(m_0) = 1.$$

Consequently:

- (i) *If m is a perfect square, then $\chi(m_0) = 1$ and the two conditions coincide: no real character can annihilate one of $A_m(n)$, $B_m(n)$ without annihilating the other. The obstruction is type-blind.*
- (ii) *If m is not a perfect square and $\chi(m_0) = -1$, then exactly one of the two conditions fails, so exactly one of the two counts is forced to vanish: $A_m(n) = 0$ if $\chi(-1) = -1$, and $B_m(n) = 0$ if $\chi(-1) = 1$. The obstruction is type-selective.*

Proof. The hypotheses give $\chi(s) = 1$ for every $s \mid n^2$, and $\chi(n) = 1$. If $A_m(n) > 0$, choose $s \equiv -n$; then $1 = \chi(s) = \chi(-1)\chi(n) = \chi(-1)$. If $B_m(n) > 0$, choose $s \equiv -mn^2$; then $1 = \chi(s) = \chi(-1)\chi(m)\chi(n)^2 = \chi(-1)\chi(m)$. Write $m = m_0k^2$; since $\gcd(m, e) = 1$ and χ is real, $\chi(k)^2 = 1$ and hence $\chi(m) = \chi(m_0)$. Parts (i) and (ii) are immediate, the second because $\chi(-1)\chi(m_0) = -\chi(-1)$ when $\chi(m_0) = -1$. $\square$

8.3. The reciprocity form. Theorem 8.3 becomes a statement about Legendre symbols once one takes for χ the Jacobi symbol, and then the hypothesis on χ is a condition relating p to the prime factors of n — this is where quadratic reciprocity enters.

Corollary 8.4. *Let $e = e_n$ be odd and let $\chi = \left(\frac{\cdot}{e}\right)$ be the Jacobi symbol modulo e . For every odd prime $q \mid n$,*

$$\left(\frac{q}{e}\right) = (-1)^{\frac{q-1}{2} \cdot \frac{e-1}{2}} \left(\frac{-p}{q}\right), \quad (8.3)$$

and $\left(\frac{2}{e}\right) = (-1)^{(e^2-1)/8}$. Suppose $\left(\frac{q}{e}\right) = 1$ for every prime $q \mid n$. Then

$$A_m(n) > 0 \implies e \equiv 1 \pmod{4}, \quad B_m(n) > 0 \implies \left(\frac{-m_0}{e}\right) = 1.$$

In particular, if m is a perfect square then $e \equiv 3 \pmod{4}$ forces $A_m(n) = B_m(n) = 0$, whereas if m is not a perfect square and $\left(\frac{m_0}{e}\right) = -1$, then $e \equiv 3 \pmod{4}$ forces $A_m(n) = 0$ but permits $B_m(n) > 0$, and $e \equiv 1 \pmod{4}$ forces $B_m(n) = 0$ but permits $A_m(n) > 0$.

Proof. Since $\gcd(n, e) = 1$ we have $q \nmid e$, and $e = mn - p \equiv -p \pmod{q}$, so $\left(\frac{e}{q}\right) = \left(\frac{-p}{q}\right)$; (8.3) is then the reciprocity law for the Jacobi symbol. The remaining assertions are Theorem 8.3 together with $\chi(-1) = \left(\frac{-1}{e}\right) = (-1)^{(e-1)/2}$ and $\chi(-1)\chi(m_0) = \left(\frac{-m_0}{e}\right)$. $\square$

Remark 8.5 (Attribution). The device underlying Theorem 8.3 is that of [9, Prop. 1.6], where Elsholtz and Tao prove that $f_I(n) = f_{II}(n) = 0$ for every odd perfect square n . Their argument takes q to be the largest odd factor of ab , uses the relation $ne \equiv -1 \pmod{q}$ coming from the parametrisation to evaluate $\left(\frac{e}{q}\right) = \left(\frac{-1}{q}\right)$ — the hypothesis that n be a square being used exactly to delete a square factor from a Legendre symbol, as in our passage from $\chi(m)$ to $\chi(m_0)$ — and then compares this with $\left(\frac{q}{e}\right)$ by reciprocity. What is new in Theorem 8.3 is that the same device applied to the *numerator* instead of the denominator produces a dichotomy: it

m	m_0	$\chi(m_0) = -1$		$\chi(m_0) = +1$	
		$\chi(-1) = -1$	$\chi(-1) = +1$	$\chi(-1) = -1$	$\chi(-1) = +1$
3	3	0 / 54	11 / 0	0 / 0	23 / 90
4	1	—	—	0 / 0	46 / 191
5	5	0 / 39	4 / 0	0 / 0	14 / 50
6	6	0 / 64	4 / 0	0 / 0	7 / 66
7	7	0 / 36	3 / 0	0 / 0	10 / 35
8	2	0 / 30	2 / 0	0 / 0	12 / 100
9	1	—	—	0 / 0	5 / 25
10	10	0 / 32	1 / 0	0 / 0	9 / 39
11	11	0 / 17	0 / 0	0 / 0	5 / 23
12	3	0 / 14	2 / 0	0 / 0	1 / 54

TABLE 1. Counts of layers n with $A_m(n) > 0$ and with $B_m(n) > 0$, among the 8172 layers on which the Jacobi symbol modulo e_n is trivial on the primes dividing n ($p \leq 400$). Bold zeros are forced by Corollary 8.4. For $m = 4$ and $m = 9$ the obstruction is type-blind; for the other m it is type-selective.

is the squareness of m , and not that of n , which decides whether a quadratic obstruction can separate the two types at all.

Formula (8.3) is the precise sense in which the local behaviour of (8.1) at the layer n is governed by the splitting of $-p$ modulo the primes dividing n . Note that the modulus $e = mn - p$ moves with n , so the two sides of (8.3) live at genuinely different places; it is reciprocity, and nothing weaker, that connects them.

Table 1 verifies Theorem 8.3 and displays the dichotomy. For each m we ran over all primes $p \leq 400$ with $p \nmid m$, $p > m$, and all m -admissible n with $mn \leq 3p$ and e_n odd, retaining those n for which $\left(\frac{q}{e}\right) = 1$ for every prime $q \mid n$; the entries record how many of the retained n have $A_m(n) > 0$ and how many have $B_m(n) > 0$, sorted by the pair $(\chi(-1), \chi(m_0))$. Zeros forced by Corollary 8.4 are shown in bold. There is no violation, and for the two square values $m = 4, 9$ the columns with $\chi(m_0) = -1$ are empty, so only the type-blind behaviour occurs.

8.4. Why $m = 4$ is the hard case. Proposition 7.4 is the special case $m = 4$ of Theorem 8.3, and we can now say what is special about it. For a non-square numerator the quadratic level already separates the two types: on every layer to which Theorem 8.3 applies, a single Jacobi symbol decides that one type is present and the other absent, and the direction of the decision is read off from $e_n \bmod 4$ and $\left(\frac{m_0}{e_n}\right)$ by Corollary 8.4. For a square numerator — and 4 is the smallest interesting one — this mechanism collapses: quadratic characters annihilate the two types simultaneously, so the entire difference $f_I(p) - f_{II}(p)$ is carried by characters of order at least 3. This is a structural obstruction, not a technical one:

Corollary 8.6. *Let m be a perfect square. Then for every admissible n and every real character χ modulo e_n which is trivial on the primes dividing n , the hypotheses of Theorem 8.3 give the same conclusion for $A_m(n)$ and for $B_m(n)$. In particular no criterion expressed through the genus theory of $(\mathbb{Z}/e_n\mathbb{Z})^\times$, that is, through H_n modulo squares, can distinguish Type I from Type II.*

Proof. Immediate from Theorem 8.3(i), since the real characters modulo e_n trivial on H_n are exactly the characters of the elementary abelian quotient $(\mathbb{Z}/e_n\mathbb{Z})^\times / H_n(\mathbb{Z}/e_n\mathbb{Z})^{\times 2}$, and -1 and $-m$ have the same image in it when m is a square. $\square$

Corollary 8.6 explains the shape of every partial result in this paper. Theorem 1.5 succeeds for $e \in \{1, 3, 5\}$ precisely because the character group modulo e is then small enough to be handled in full, including its quartic characters when $e = 5$; the failure at $e = 7$ recorded there is the first place where the cubic characters modulo 7 are needed and cannot be controlled. It also explains why Conjecture 1.6 is delicate: an inequality between f_I and f_{II} for $m = 4$ cannot follow from congruence conditions modulo squares, and must instead come from an equidistribution statement for characters of higher order. Compare [9, Prop. 1.6], where the same phenomenon appears as the observation that the solvability of (1.1) is not decided by any congruence condition on p .

8.5. Vanishing at odd squares, for every numerator divisible by four. Proposition 7.4 and Theorem 8.3 concern a single layer. There is also a global statement of the same flavour, and it turns out to be governed not by the squareness of m but by a 2-adic condition on m . Recall [9, Prop. 1.6]: for every odd perfect square n one has $f_I(n) = f_{II}(n) = 0$. We show that the numerator 4 may be replaced by any multiple of 4, and that no other numerator will do.

Here n need not be prime, and we use the definitions of [9, §2]: a solution of $m/n = 1/x + 1/y + 1/z$ is of *Type I* if $n \mid x$ and $\gcd(n, yz) = 1$, and of *Type II* if $n \mid y$, $n \mid z$ and $\gcd(n, x) = 1$. We write $f_I^{(m)}(n), f_{II}^{(m)}(n)$ for the numbers of such solutions.

Lemma 8.7. *Let $m, n \in \mathbb{N}$ and let (x, y, z) be a Type I solution of $m/n = 1/x + 1/y + 1/z$. Then there are $a, b, c, d \in \mathbb{N}$ with $\gcd(abcd, n) = 1$ and*

$$mabcd = n(a + b) + c, \quad c \mid a + b, \quad (8.4)$$

such that $(x, y, z) = (nabd, acd, bcd)$. Writing $e := (a + b)/c$, one has

$$mabd = ne + 1. \quad (8.5)$$

If instead (x, y, z) is a Type II solution, the same holds with $(x, y, z) = (abd, nacd, nbcd)$, $\gcd(abd, n) = 1$,

$$mabcd = nc + a + b, \quad c \mid a + b, \quad mabd = n + e. \quad (8.6)$$

Proof. This is [9, Prop. 2.2 and 2.6] with 4 replaced by m ; the derivation there uses nothing about the numerator. For the Type I case one factors $x = ndx', y = dy', z = dz'$ with x', y', z' coprime, so that $mdx'y'z' = y'z' + nx'y' + nx'z'$; since y', z' are coprime to n one gets $x' \mid y'z'$, $y' \mid x'z'$, $z' \mid x'y'$ and hence $x' = ab, y' = ac, z' = bc$, which gives (8.4). As c is coprime to n , (8.4) forces $c \mid a + b$, and substituting $a + b = ce$ into (8.4) and dividing by c gives (8.5). The Type II case is identical, starting from $mdx'y'z' = ny'z' + x'y' + x'z'$. $\square$

Theorem 8.8 (Vanishing at odd squares). *Let m be divisible by 4 and let n be an odd perfect square. Then*

$$f_I^{(m)}(n) = f_{II}^{(m)}(n) = 0.$$

Proof. Suppose first that a Type I solution exists, and take a, b, c, d, e as in Lemma 8.7. Let q be the largest odd divisor of ab , so $ab = 2^j q$ with $j \geq 0$. All symbols below are Jacobi symbols.

Since n is an odd square, $n \equiv 1 \pmod{8}$. From (8.5) and $4 \mid m$ we get $ne \equiv -1 \pmod{4}$, hence $e \equiv -n^{-1} \equiv -1 \equiv 3 \pmod{4}$; in particular e is odd and $e \geq 3$. Reducing (8.5) modulo q gives $ne \equiv -1$, and $\gcd(q, n) = 1$ because $\gcd(abcd, n) = 1$, so

$$\left(\frac{e}{q}\right) = \left(\frac{-n^{-1}}{q}\right) = \left(\frac{-1}{q}\right) \left(\frac{n}{q}\right) = \left(\frac{-1}{q}\right),$$

because n is a square coprime to q . As $e \equiv 3 \pmod{4}$, quadratic reciprocity for the Jacobi symbol gives $\left(\frac{q}{e}\right)\left(\frac{e}{q}\right) = (-1)^{\frac{q-1}{2}} = \left(\frac{-1}{q}\right)$, whence

$$\left(\frac{q}{e}\right) = 1. \quad (8.7)$$

On the other hand $ce = a + b$ gives $b \equiv -a \pmod{e}$, so $ab \equiv -a^2$; and $\gcd(ab, e) = 1$ by (8.5), so

$$\left(\frac{ab}{e}\right) = \left(\frac{-1}{e}\right) = -1, \quad (8.8)$$

again because $e \equiv 3 \pmod{4}$. Comparing (8.7) and (8.8) forces $ab \neq q$, i.e. $j \geq 1$ and ab is even. But then $8 \mid mabd$, so (8.5) gives $ne \equiv -1 \pmod{8}$ and therefore $e \equiv -1 \equiv 7 \pmod{8}$; hence $\left(\frac{2}{e}\right) = 1$ and $\left(\frac{ab}{e}\right) = \left(\frac{2}{e}\right)^j \left(\frac{q}{e}\right) = \left(\frac{q}{e}\right)$, contradicting (8.7) and (8.8).

For a Type II solution the argument is the same with (8.6) in place of (8.5): from $mabd = n + e$ and $4 \mid m$ we get $e \equiv -n \equiv 3 \pmod{4}$, and modulo q we get $e \equiv -n$, so $\left(\frac{e}{q}\right) = \left(\frac{-1}{q}\right)\left(\frac{n}{q}\right) = \left(\frac{-1}{q}\right)$; the relation $ce = a + b$ is unchanged, and if ab is even then $8 \mid n + e$ forces $e \equiv 7 \pmod{8}$. The same contradiction results. $\square$

For $m = 4$ this is [9, Prop. 1.6]. The hypothesis $4 \mid m$ cannot be weakened, and in particular it is not implied by, nor does it imply, m being a square: for every $m \leq 32$ with $4 \nmid m$ there is an odd square n with $f_I^{(m)}(n) > 0$, the largest one needed being $n = 17^2$ (at $m = 23$). Thus $m = 9$, a square, admits Type I solutions at $n = 25, 49, 121, 169, 289, 361$, while $m = 8$ and $m = 12$, neither of them squares, admit none at any odd square. The two phenomena isolated in this section are therefore genuinely different: Theorem 8.3 is a statement about m modulo squares, Theorem 8.8 one about m modulo powers of 2.

Remark 8.9. The proof shows a little more. What is used is only that $4 \mid m$ and $n \equiv 1 \pmod{8}$ with $\left(\frac{n}{q}\right) = 1$ for the relevant q ; the last condition holds for every q when n is a square, which is why squares are the natural class.

8.6. Exceptional sets for general m . Call a prime $p \nmid m$ *m-exceptional* if $f_{II}^{(m)}(p) > 0$ and $f_I^{(m)}(p) \leq f_{II}^{(m)}(p)$, and *strictly m-exceptional* if $f_I^{(m)}(p) < f_{II}^{(m)}(p)$. (The condition $f_{II}^{(m)}(p) > 0$ discards the primes for which (8.1) has no solution at all, where both counts vanish for a trivial reason.) Table 2 lists these sets.

Two features of these sets are worth recording. The first is a doubling phenomenon, which is the general- m form of Proposition 9.1(i) and, through Corollary 4.4, of the whole mechanism identified in §4.

Proposition 8.10. *Let $m \geq 3$ and let $p \nmid m$ be a prime with $p \equiv -1 \pmod{m}$; put $M := (p + 1)/m$. Then*

$$f_{II}^{(m)}(p) \geq \tau(M^2), \quad f_I^{(m)}(p) \geq 2\tau(M^2).$$

Proof. The integer M is m -admissible with $e_M = mM - p = 1$, so both congruences in (8.2) are vacuous and $A_m(M) = B_m(M) = \tau(M^2)$; Proposition 8.1 gives $f_{II}^{(m)}(p) \geq \tau(M^2)$.

For the Type I bound, let $s \mid M^2$ and put

$$x := M, \quad y := pM + s, \quad z := \frac{pM(pM + s)}{s}.$$

Then $z \in \mathbb{N}$ because $s \mid M^2 \mid (pM)^2$, and $1/y + 1/z = (pM + s)/(pM(pM + s)) = 1/(pM) = m/p - 1/M$, so (x, y, z) solves (8.1). Since $p \nmid M$ we have $p \nmid s$, hence $p \nmid y$, while $p \mid z$; so the solution is of Type I and its two denominators coprime to p are M and $pM + s$, which are

distinct. Distinct s give distinct y , hence distinct solutions. Each of these $\tau(M^2)$ solutions is therefore counted twice in $\sum_n B_m(n) = f_I^{(m)}(p)$, once at the layer $n = M$ and once at the layer $n = pM + s$. $\square$

The point is that the hypothesis $p \equiv -1 \pmod{m}$ is exactly the condition under which the divisor $\delta = 1$ becomes available to the Type I residue class: by the computation proving Corollary 4.4, carried out with $\mu := mAC$ in place of $4AC$, the class $-p$ modulo μ contains 1 precisely when $\mu \mid p+1$, which requires $m \mid p+1$, whereas the class -1 contains 1 only if $\mu \mid 2$, which never happens. Numerically this single mechanism appears to be decisive.

Conjecture 8.11. Let $m \geq 3$ and let $p \nmid m$ be a prime with $p \equiv -1 \pmod{m}$. Then $f_I^{(m)}(p) > f_{II}^{(m)}(p)$.

For $m = 4$ this is the assertion of Conjecture 1.6 for the primes $p \equiv 3 \pmod{4}$, a set of density $\frac{1}{2}$ inside the primes. We verified Conjecture 8.11 for all $4 \leq m \leq 20$ and all primes $p \leq 2 \cdot 10^4$: in that range there are 215 strictly m -exceptional primes and a further 428 ties, and *none* of these 643 primes satisfies $p \equiv -1 \pmod{m}$. The evidence that this is the exact constraint, rather than an artefact of some coarser congruence, is best seen for $m = 18$: of the twelve reduced residues modulo 36, ten are realised by strictly 18-exceptional primes $p \leq 2 \cdot 10^4$, and the two that are not are exactly 17 and 35, the two residues with $p \equiv -1 \pmod{18}$.

The second feature is a bias which we can neither prove nor explain.

Question 8.12. Is there a reason why $\left(\frac{-m}{p}\right) = +1$ for almost every strictly m -exceptional prime?

The last two columns of Table 2 record the counts. For $4 \leq m \leq 15$ the symbol is $+1$ for 78 of the 82 strictly exceptional primes $p \leq 2 \cdot 10^4$, against an expected 41; over the whole range $4 \leq m \leq 20$ it is 191 out of 215. For $m \in \{4, 8, 11, 12, 15, 20\}$ there is no exception at all, 123 cases in total. The smallest counterexample is $m = 6$, $p = 379$, where $f_I^{(6)} = 36 < 42 = f_{II}^{(6)}$ and $\left(\frac{-6}{379}\right) = -1$; we verified this by direct enumeration of all solutions of $6/379 = 1/x + 1/y + 1/z$. Note that Question 8.12 concerns a symbol attached to p , whereas Theorem 8.3 produces symbols attached to the moduli $e_n = mn - p$, which vary with the layer; we have not been able to connect them.

Finally we note what the results of this section do *not* give. Theorem 8.3 is a statement about individual layers; converting it into information about $f_I^{(m)}(p) - f_{II}^{(m)}(p)$ requires knowing how often the hypothesis “ $\chi(q) = 1$ for every $q \mid n$ ” is satisfied, and for n with $\omega(n)$ prime factors this is a condition of probability $2^{-\omega(n)}$, so the layers to which Theorem 8.3 applies have density zero (of order $(\log p)^{-\log 2}$ inside the window). What Theorem 8.3 does give is a clean reason for the difficulty of the square case: by Corollary 8.6, for $m = 4$ the difference between the two types is invisible to every quadratic character, so any proof of Conjecture 1.6 must use characters of order at least 3, and correspondingly no criterion modulo squares — in particular none of the classical congruence criteria for the Erdős–Straus equation — can decide it.

8.7. A window and a reduction for a general numerator. Conjecture 8.11 can be reduced, exactly as Conjecture 1.6 is in §5 and §9, to a finite computation. Two observations do the work; both are proved for every $m \geq 3$.

Proposition 8.13. Let $m \geq 3$ and let n be m -admissible. If $A_m(n) > 0$ then

$$\frac{p}{m} < n \leq \frac{p}{m-2};$$

m	m_0	strict	ties	$\left(\frac{-m}{p}\right) = +1$	smallest strict
4	1	3	7	3/3	41
5	5	0	0	—	—
6	6	1	1	0/1	379
7	7	0	5	—	—
8	2	18	20	18/18	433
9	1	2	12	1/2	389
10	10	1	12	1/1	251
11	11	8	7	8/8	829
12	3	25	42	25/25	223
13	13	2	9	1/2	1699
14	14	2	10	1/2	467
15	15	20	39	20/20	139
16	1	34	61	31/34	401
17	17	8	22	4/8	431
18	2	29	54	19/29	227
19	19	13	39	10/13	233
20	5	49	88	49/49	409

TABLE 2. Exceptional primes $p \leq 2 \cdot 10^4$ for the numerator m : the number with $f_I^{(m)}(p) < f_{II}^{(m)}(p)$ (strict) and with $f_I^{(m)}(p) = f_{II}^{(m)}(p) > 0$ (ties), the number of strict ones with $\left(\frac{-m}{p}\right) = +1$, and the smallest strict one. No exceptional prime in the whole table satisfies $p \equiv -1 \pmod{m}$ (Conjecture 8.11).

we write $W_m = W_m(p)$ for the set of integers in this range. Moreover $B_m(n) = A_m(n)$ whenever $e_n \leq 2$.

Proof. By Proposition 8.1 and Lemma 2.3, $A_m(n)$ is the number of ordered pairs (Y, Z) with $1/Y + 1/Z = e_n/n$; such a pair exists only if $e_n/n \leq 2$, that is $mn - p \leq 2n$. Together with $mn > p$ this is the stated range.

If $e_n = 1$ both congruences in (8.2) are vacuous and $A_m(n) = B_m(n) = \tau(n^2)$. If $e_n = 2$ then n is odd, because $\gcd(e_n, n) = 1$; hence every $s \mid n^2$ is odd and $n^2 \equiv 1 \pmod{2}$. Also $mn = p + 2$ is odd, so m is odd. Thus $s \equiv 1 \equiv -n$ and $s \equiv 1 \equiv -mn^2 \pmod{2}$ for every $s \mid n^2$, and again $A_m(n) = B_m(n) = \tau(n^2)$. $\square$

Theorem 8.14 (Reduction for general m). *Let $m \geq 3$ and let $p \nmid m$ be a prime with $p \equiv -1 \pmod{m}$; put $M = (p + 1)/m$. Then*

$$f_I^{(m)}(p) - f_{II}^{(m)}(p) \geq \tau(M^2) + \sum_{\substack{n \in W_m \\ e_n \geq 3}} (B_m(n) - A_m(n)).$$

In particular Conjecture 8.11 holds for (m, p) whenever the right-hand side is positive.

Proof. By Proposition 8.1, $f_{II}^{(m)}(p) = \sum_n A_m(n)$, and by Proposition 8.13 the sum may be restricted to $n \in W_m$. Also $f_I^{(m)}(p) = \sum_n B_m(n) \geq \sum_{n \in W_m} B_m(n) + \sum_{n \notin W_m} B_m(n)$, and the second sum is at least $\tau(M^2)$: in the proof of Proposition 8.10 each of the $\tau(M^2)$ solutions constructed there is counted once at the layer $n = pM + s$, and $pM + s > p \geq p/(m - 2)$ for $m \geq 3$, so all those layers lie outside W_m . Subtracting and discarding the layers with $e_n \leq 2$, which contribute 0 by Proposition 8.13, gives the inequality. $\square$

The window W_m has length $2p/(m(m-2))$, so Theorem 8.14 replaces an unbounded computation by a finite one, and a shorter one the larger m is. For $m = 4$ it gives $W_4 = (p/4, p/2]$, weaker than the window $(p/4, (p+1)/3]$ of Theorem 5.1, whose proof uses the numerator 4; on the other hand Theorem 8.14 applies to every m .

Of the 3028 pairs (m, p) with $3 \leq m \leq 20$, $p \leq 6 \cdot 10^3$ and $p \equiv -1 \pmod{m}$, the right-hand side of Theorem 8.14 is positive for 3026, so Conjecture 8.11 is proved for those. The two exceptions are $(m, p) = (6, 3881)$, where $\tau(M^2) = 3$ and the sum equals -4 , and $(18, 3797)$, where $\tau(M^2) = 3$ and the sum equals -3 ; in both, the conjecture nevertheless holds and comfortably, $f_I^{(6)}(3881) = 49$ against $f_{II}^{(6)}(3881) = 27$ and $f_I^{(18)}(3797) = 15$ against $f_{II}^{(18)}(3797) = 9$. We also checked $f_I^{(m)}(p) > f_{II}^{(m)}(p)$ directly for all 3028 pairs. As for $m = 4$, what the reduction does not supply is any lower bound for the layers with $e_n \geq 3$, and the bound $e_n \leq 2$ of Proposition 8.13 cannot be relaxed: among the layers with $3 \leq m \leq 12$, $p \leq 400$ and $e_n = 3$, the inequality $B_m(n) \geq A_m(n)$ fails 37 times out of 111, with $B_m(n) - A_m(n)$ as low as -5 .

9. NUMERICAL EVIDENCE, A CRITERION, AND A DIAGNOSIS

All computations below were carried out with two independent programs: a direct exhaustive search over the smallest denominator, and a global enumeration based on Lemma 2.1. They agree pointwise for all primes $p \leq 5000$, and the resulting values of (f_I, f_{II}) for all 78 496 primes $p \leq 10^6$ were used for the tables below.

9.1. Pointwise lower bounds.

Proposition 9.1. *Let $p \geq 7$ be prime.*

- (i) *If $p \equiv 3 \pmod{4}$ and $M = (p+1)/4$, then $f_I(p) \geq 2\tau(M^2)$, $f_{II}(p) \geq \tau(M^2)$ and $f(p) \geq 9\tau(M^2) \geq 27$.*
- (ii) *If $p \equiv 1 \pmod{4}$, let $n = (p+3)/4$ and let n_1 be the part of n composed of primes $\equiv 1 \pmod{3}$. Then $f_I(p) \geq \tau(n^2) - \tau(n_1^2)$, and if moreover $p \equiv 2 \pmod{3}$ then $f_{II}(p) \geq (\tau(n^2) + \tau(n_1^2))/2 \geq 1$.*

In particular $f_{II}(p) > 0$ for every prime $p \not\equiv 1 \pmod{12}$.

Proof. Both parts follow from Theorem 1.1 and Corollary 5.2 (which give $f_{II}(p) \geq A(n)$ and $f_I(p) \geq 2B(n)$ for any single $n \in W$) together with the computations in the cases $e = 1$ and $e = 3$ carried out in the proof of Theorem 1.5. For (i), $e_M = 1$ and $A(M) = B(M) = \tau(M^2)$; the bound on $f(p)$ follows from (1.2), and $\tau(M^2) \geq 3$ since $M \geq 2$. $\square$

Part (i) is elementary and is presumably folklore: the family in question is obtained by splitting $1/(pM)$ in $4/p = 1/M + 1/(pM)$ in all possible ways. We include it because it is exactly the $e = 1$ layer of Theorem 1.1 and because it is used in Theorem 9.2. We emphasise that the classes covered by Proposition 9.1 are classes in which (1.1) is classically solvable.

9.2. The exceptional set. Table 3 lists all exceptions for $p \leq 10^6$, and Figure 1 shows the cloud for $p \leq 3 \cdot 10^4$. In particular Conjecture 1.6 holds for all $p \leq 10^6$. Moreover the margin $f_I(p) - f_{II}(p)$ over primes $p \not\equiv 1 \pmod{8}$ grows: its minimum is 2, 18, 30, 62, 62, 68 over the ranges $[5, 10^3)$, $[10^3, 10^4)$, $[10^4, 10^5)$, $[10^5, 3 \cdot 10^5)$, $[3 \cdot 10^5, 6 \cdot 10^5)$, $[6 \cdot 10^5, 10^6)$, and over the last five of these the minimal ratio f_I/f_{II} stays between 1.30 and 1.40. Figure 2 displays the minimal ratio over all fifteen dyadic ranges $2^k \leq p < 2^{k+1}$ with $2^k \geq 16$ and $2^{k+1} \leq 10^6$: for $p \not\equiv 1 \pmod{8}$ it never falls below 1.2857 (attained at $p = 101$), whereas for $p \equiv 1 \pmod{8}$ it reaches 1 or less in seven of the fifteen ranges and drops strictly below 1 in three of them. The two curves do not come close to meeting, and neither shows a trend towards 1. The prime $p = 5$, the only exception with $p \not\equiv 1 \pmod{8}$, lies below the plotted range.

TABLE 3. All primes $p \leq 10^6$ with $f_I(p) \leq f_{II}(p)$.

p	$f_I(p)$	$f_{II}(p)$	relation	$p \bmod 8$
5	2	2	=	5
17	4	4	=	1
41	6	8	<	1
89	10	10	=	1
241	8	8	=	1
9 001	38	40	<	1
12 289	36	36	=	1
14 401	26	26	=	1
17 489	68	68	=	1
19 801	46	50	<	1
94 201	82	82	=	1
624 649	116	116	=	1




FIGURE 1. $f_I(p)$ against $f_{II}(p)$, logarithmic on both axes, for all 3 243 primes $5 \leq p \leq 3 \cdot 10^4$ with $f_{II}(p) > 0$. The dashed line is $f_I = f_{II}$, the dotted line $f_I = 3f_{II}$ (the ratio predicted by Conjecture 9.7); the cloud is centred on the latter and, apart from the ten black points, lies strictly above the former. Those ten points are the primes with $f_I(p) \leq f_{II}(p)$, listed in Table 3; all of them satisfy $p \equiv 1 \pmod{8}$ except $p = 5$.

The smallest counterexamples are $(p, n) = (41, 12)$ for $e = 7$ with $(A, B) = (4, 1)$, $(59, 17)$ for $e = 9$ with $(A, B) = (2, 0)$, and $(29, 10)$ for $e = 11$ with $(A, B) = (2, 0)$; from $e = 9$ onwards the smallest counterexample always has $(A, B) = (2, 0)$, which is the shape produced by Proposition 7.5. For $e \in \{1, 3, 5\}$ no failure occurs among the 117 817 local terms with $p \leq 10^6$, in agreement with Theorem 1.5.

9.3. A criterion, and exactly what remains of Conjecture 1.6. Corollary 5.2 and Theorem 1.5 combine into an unconditional inequality which isolates the unproved part of Conjecture




FIGURE 2. For each dyadic range $2^k \leq p < 2^{k+1}$ with $2^{k+1} \leq 10^6$, the minimum of $f_I(p)/f_{II}(p)$ over the primes of that range with $f_{II}(p) > 0$, taken separately over $p \not\equiv 1 \pmod{8}$ (solid, filled) and $p \equiv 1 \pmod{8}$ (dashed, hollow). The solid minimum never falls below 1.2857; the dashed one reaches 1 or less seven times and falls strictly below 1 three times. This is Conjecture 1.6 seen across all 78 496 primes at once.

TABLE 4. Sharpness of Theorem 1.5: proportion of the local terms with a given e for which $2B(n) < A(n)$, over all primes $p \leq 2 \cdot 10^4$. Each column rests on about 1 130 terms. The rate is 0 exactly for $e \leq 5$.

e	1	3	5	7	9	11	13	15	17	19
rate	0	0	0	4.4%	4.0%	7.3%	4.1%	10.7%	4.4%	9.3%
e	21	23	25	27	29	31	33	35	37	39
rate	10.5%	10.0%	6.3%	9.7%	4.8%	7.9%	6.6%	10.0%	3.7%	11.0%
e	41	43	45	47	49	51	53	55	57	59
rate	5.6%	6.7%	9.5%	10.5%	6.5%	7.7%	5.6%	9.8%	5.6%	8.3%

1.6. Split the window W of (5.1) according to the value of $e_n = 4n - p$ and put

$$\sigma(p) := \sum_{n \in W, e_n \leq 5} (2B(n) - A(n)), \quad T(p) := \sum_{n \in W, e_n \geq 7} (2B(n) - A(n)). \quad (9.1)$$

Theorem 9.2. *For every prime $p \geq 7$,*

$$f_I(p) - f_{II}(p) \geq \sigma(p) + T(p),$$

and $\sigma(p) \geq 0$. Moreover $\sigma(p) \geq \tau(M^2) \geq 3$ if $p \equiv 3 \pmod{4}$, where $M = (p+1)/4$. In particular $f_I(p) > f_{II}(p)$ whenever $T(p) > -\sigma(p)$.

Proof. By Corollary 5.2, $f_I(p) \geq 2 \sum_{n \in W} B(n)$ and $f_{II}(p) = \sum_{n \in W} A(n)$, so $f_I(p) - f_{II}(p) \geq \sum_{n \in W} (2B(n) - A(n)) = \sigma(p) + T(p)$. Every term of $\sigma(p)$ is non-negative by Theorem 1.5. If $p \equiv 3 \pmod{4}$ then $e_n = 1$ occurs, at $n = M$, and there $A(M) = B(M) = \tau(M^2)$, contributing $\tau(M^2)$; the term $e_n = 5$ contributes ≥ 0 . Finally $\tau(M^2) \geq 3$ since $M \geq 2$. $\square$

Two features of Theorem 9.2 deserve comment. First, its hypothesis involves only the window W , and in particular does not require computing $f_1(p)$, whose determination involves a parameter region of size $O(p^2)$; the same asymmetry appears on the algorithmic side, where Dahan [7] observes that a factorization-free search over the primes of $[N, 2N]$ costs $O(N \log^3 N)$ for Type II but $\Theta(N^2)$ for Type I. Second, and more importantly, the split (9.1) is *exactly* at the point where Theorem 1.5 stops being true: by the sharpness statement there, for every odd e with $7 \leq e \leq 2 \cdot 10^4$ some local term has $2B(n) < A(n)$, so the cut cannot be moved by that method.

Consequently Conjecture 1.6 is reduced to a single inequality.

Conjecture 9.3 (The residual statement). Let $p > 5$ be prime with $p \not\equiv 1 \pmod{8}$. Then $T(p) > -\sigma(p)$. More precisely:

- (i) if $p \equiv 3 \pmod{4}$ then $T(p) \geq 0$;
- (ii) if $p \equiv 5 \pmod{8}$ then $T(p) > -\sigma(p)$.

By Theorem 9.2, (i) implies Conjecture 1.6 for $p \equiv 3 \pmod{4}$ with the explicit margin $\tau(M^2) \geq 3$, and (ii) implies it for $p \equiv 5 \pmod{8}$. We checked Conjecture 9.3 for all primes $p \leq 3 \cdot 10^4$: among the 1632 primes $p \equiv 3 \pmod{4}$ the quantity $T(p)$ is non-negative *without exception*, while among the 814 primes $p \equiv 5 \pmod{8}$ it is negative for only three, the minimum being $T(701) = -8$. The asymmetry between the two cases is intrinsic: for $p \equiv 3 \pmod{4}$ the layer $e_n = 1$ exists and contributes the whole of $\tau(M^2)$ to $\sigma(p)$, whereas for $p \equiv 1 \pmod{4}$ the smallest layer is $e_n = 3$ and $\sigma(p)$ is correspondingly smaller.

Theorem 9.2 is effective: the hypothesis $T(p) > -\sigma(p)$ can be verified for a given p by a computation in the window W alone, and this *proves* Conjecture 1.6 for that p . Weakening it to the sufficient condition $\sum_{e_n \geq 7} A(n) < \sigma(p)$, which discards the non-negative contributions $2B(n)$ with $e_n \geq 7$, gives the coverage in Table 5.

TABLE 5. Proportion of primes for which the sufficient condition $\sum_{e_n \geq 7} A(n) < \sigma(p)$ holds, so that Conjecture 1.6 is *proved* for them by Theorem 9.2.

range	$[10^2, 10^{2.5})$	$[10^{2.5}, 10^3)$	$[10^3, 10^{3.5})$	$[10^{3.5}, 10^4)$	$[10^4, 10^{4.5})$
$p \equiv 3 \pmod{4}$	100%	96.2%	81.9%	64.5%	52.5%
$p \equiv 5 \pmod{8}$	36.4%	26.9%	17.1%	13.4%	11.8%

The proportions decrease, because $\sum_{e_n \geq 7} A(n)$ grows with p while $\sigma(p)$ grows only like a divisor function; so Theorem 9.2 yields no asymptotic case of Conjecture 1.6. It does, however, locate the difficulty precisely: *everything reduces to showing that the layers $e_n \geq 7$ do not, in total, undo the layers $e_n \leq 5$* . Numerically the individual layers with $e_n \geq 7$ fail the local inequality only 0.2% to 0.4% of the time (§9.4), and their aggregate effect $T(p)$ is non-negative for every $p \equiv 3 \pmod{4}$ we tested.

9.4. Where the surplus comes from. The following two computations locate the excess of Type I over Type II solutions.

(a) *The low layers are balanced.* For e fixed let $A_e(p) := A((p+e)/4)$ and $B_e(p) := B((p+e)/4)$. Over all 17982 primes $p \leq 2 \cdot 10^5$ the ratios $\sum_p B_e(p) / \sum_p A_e(p)$ oscillate between 0.68 and 1.39, with $\sum_{e \leq 41} \sum_p B_e(p) / \sum_{e \leq 41} \sum_p A_e(p) = 0.988$. There is a clear pattern modulo 4: every layer with $e \equiv 3 \pmod{4}$ (equivalently $p \equiv 1 \pmod{4}$) has ratio < 1 , while most layers with $e \equiv 1 \pmod{4}$ have ratio > 1 .

(b) *The surplus lives in the upper part of the window.* Splitting W according to e_n/n , over all 6 055 primes $p \leq 6 \cdot 10^4$ one finds $\sum A = 437\,618$ and $\sum B = 458\,219$ for $e_n/n \in [0, 0.05)$ (ratio 1.05), whereas for $e_n/n \in [0.4, 0.5)$, $[0.6, 0.7)$, $[0.7, 0.8)$, $[0.8, 0.9)$ the values of $\sum A$ are 10, 6, 2, 2 against $\sum B$ equal to 12 972, 9 373, 7 295, 8 268.

Computation (b) is precisely the content of Theorems 1.4 and 1.2: for $e_n/n \geq \delta$ the quantity $A(n)$ is nonzero only when e_n/n is one of the finitely many rationals representable as a sum of two unit fractions with bounded smaller denominator, and the total contribution of that region to $f_{II}(p)$ is $O_\delta(1)$ uniformly in p . Computation (a) shows that in the complementary region the two divisor counts are essentially balanced.

We draw the following conclusion. *The surplus of Type I over Type II solutions is not caused by an imbalance in the distribution of the divisors of n^2 among residue classes modulo $4n - p$; those distributions are balanced on average. It is caused by the geometric constraint of Theorem 1.4, together with the vanishing of A for $n > (p+1)/3$ (Theorem 5.1).* Consequently the natural line of attack on Conjecture 1.6 is not an equidistribution statement for divisors in progressions, but the following counting problem.

Question 9.4. Fix $0 < \delta < 1$. How large is $\#\{n \in W : e_n \geq \delta n, A(n) > 0\}$, and how does the corresponding contribution to $f_{II}(p)$ compare with that of $f_I(p)$ on the same range?

Theorem 1.2 answers the first half with the bound $D(\lfloor 2/\delta \rfloor)$; by Remark 6.3 the truth is probably $\asymp 1/\delta$.

9.5. Partial results, and two weakenings that are false. We collect what is proved towards Conjecture 1.6, and record two natural weakenings which numerical evidence refutes.

Proved.

- (1) $f_I(p) \geq 2\tau(M^2)$ for $p \equiv 3 \pmod{4}$, $M = (p+1)/4$ (Corollary 4.5); equivalently, the divisor $\delta = 1$ contributes $\tau(M^2)$ unordered Type I solutions and no Type II solution at all (Corollary 4.4).
- (2) $2B(n) \geq A(n)$ for $e_n \in \{1, 3, 5\}$, and for no larger e (Theorem 1.5); by Corollary 5.2 this is exactly the local form of Conjecture 1.6.
- (3) $f_I(p) - f_{II}(p) \geq \sigma(p) + T(p)$ with $\sigma(p) \geq 0$, and $\sigma(p) \geq \tau(M^2)$ when $p \equiv 3 \pmod{4}$ (Theorem 9.2); this proves Conjecture 1.6 for an explicit set of primes and reduces the rest to Conjecture 9.3.
- (4) Every Type II solution satisfies $a \leq (1 + \sqrt{1 + 4p})/4$ (Corollary 4.6), and the number of Type II solutions with $Y \leq K$ is at most $D(K)$, uniformly in m and p (Theorem 1.2). There is no Type I analogue of either.

False. Theorem 4.1 suggests grouping the two counts by the parameter C , which is c for a Type I solution and a for a Type II solution. Both resulting stratified inequalities fail.

- (5) *The stratum $C = 1$.* The inequality $\#\{\text{Type I} : c = 1\} \geq \#\{\text{Type II} : a = 1\}$ fails; the smallest counterexample is $p = 47$, where the two counts are 6 and 7. Among the 301 primes $p \leq 2000$ it fails for four.
- (6) *Every stratum.* The inequality $\#\{\text{Type I} : c = C\} \geq \#\{\text{Type II} : a = C\}$ for all C fails for 132 of the 301 primes $p \leq 2000$; the smallest counterexample is $p = 41$, $C = 3$, where the counts are 0 and 1.

So Conjecture 1.6 cannot be reduced to the strata of the duality, just as §9.8(1) shows it cannot be reduced to the terms of Theorem 1.1. Together with §9.8(5) — any matching of solutions sharing a denominator reduces to the latter — this leaves the global estimates of §9.4 and §9.7 as the only route we can see.

9.6. A filtered form of Conjecture 1.6. Theorem 4.1 attaches to every solution the invariant δ . Explicitly, if $y \leq z$ are the two denominators coprime to p of a Type I solution then $\delta = 4y - p$, while for a Type II solution $4/p = 1/x + 1/(pY) + 1/(pZ)$ with $Y \leq Z$ one has $\delta = 4Y - 1$. For $X \geq 1$ put

$$N_I(X; p) := \#\{\text{unordered Type I solutions with } \delta \leq X\},$$

$$N_{II}(X; p) := \#\{\text{unordered Type II solutions with } \delta \leq X\}.$$

Letting $X \rightarrow \infty$ recovers the unordered forms of $f_I(p)$ and $f_{II}(p)$, so the following refines Conjecture 1.6.

Conjecture 9.5 (Filtered form). For every prime $p > 5$ with $p \not\equiv 1 \pmod{8}$ and every $X \geq 1$,

$$N_I(X; p) \geq N_{II}(X; p).$$

We verified Conjecture 9.5 for all primes $p \leq 6000$. In that range the inequality fails for exactly ten primes, namely

$$41, 89, 241, 433, 881, 1249, 2161, 3041, 3169, 4441,$$

all of which are $\equiv 1 \pmod{8}$. Only the first three are exceptions to Conjecture 1.6 itself, so Conjecture 9.5 is strictly stronger; nevertheless the dichotomy modulo 8 survives intact. We regard this as evidence that the class $p \equiv 1 \pmod{8}$, and not merely the coarse count, is the right locus of the phenomenon. Figure 3 shows a typical pair of step functions and one of the failures.

Part of Conjecture 9.5 can be proved.

Theorem 9.6. *Let $p \geq 7$ be prime.*

- (i) *For every $X \geq 1$, $N_{II}(X; p) \leq D(\lfloor (X+1)/4 \rfloor) = \sum_{n \leq (X+1)/4} \tau(n)$, a bound independent of p .*
- (ii) *If $p \equiv 3 \pmod{4}$ and $M = (p+1)/4$, then $N_I(X; p) \geq \tau(M^2)$ for every $X \geq 1$.*
- (iii) *Consequently, if $p \equiv 3 \pmod{4}$ then $N_I(X; p) > N_{II}(X; p)$ for every X such that $\sum_{n \leq (X+1)/4} \tau(n) < \tau(M^2)$. In particular this holds for all $X \leq 4K+3$, where K is the largest integer with $\sum_{n \leq K} \tau(n) < \tau(M^2)$; since $\tau(M^2) \geq 3$ one always has $K \geq 1$, so the conclusion holds at least for $X \leq 7$.*

Proof. (i) For a Type II solution $\delta = 4Y - 1$ with $Y = acd$, so $\delta \leq X$ is equivalent to $Y \leq (X+1)/4$; now apply Theorem 1.2.

(ii) By Corollary 4.5 there are $\tau(M^2)$ unordered Type I solutions with $\delta = 1$, and $1 \leq X$.

(iii) Immediate from (i) and (ii). For the last sentence, $\tau(M^2) \geq 3$ because $M \geq 2$, while $\sum_{n \leq 1} \tau(n) = 1 < 3$, so $K \geq 1$ and $4K+3 \geq 7$. $\square$

For the 217 primes $p \equiv 3 \pmod{4}$ with $7 \leq p \leq 3000$ we checked that (i) holds for every X (no violations), and that the resulting proved range $X \leq 4K+3$ has median 19 and maximum 79. Thus Theorem 9.6 settles an initial segment of Conjecture 9.5, of length growing with $\tau(((p+1)/4)^2)$; what it does not control is the tail, where N_{II} eventually catches up with $\tau(M^2)$.

9.7. A heuristic model, and the constant 3. We now describe a probabilistic model for f_I and f_{II} which predicts the value of the ratio, and which we regard as the main evidence for Conjecture 1.6.




FIGURE 3. The counting functions $N_I(X; p)$ (solid) and $N_{II}(X; p)$ (dashed). Top: $p = 1051 \equiv 3 \pmod{4}$, where the solid curve dominates throughout, as Conjecture 9.5 predicts. Bottom: $p = 433 \equiv 1 \pmod{8}$, where the dashed curve overtakes it on the range $119 \leq X \leq 169$, although $f_I(433) > f_{II}(433)$ in the end.

Let $y \leq z$ be the two denominators of an unordered Type I solution that are coprime to p , and put $h := \gcd(y, z)$, $y = ah$, $z = bh$ with $\gcd(a, b) = 1$. In the notation of Lemma 2.1 one has $h = cd$, and the defining equation $4abcd = p(a + b) + c$ becomes

$$4abh - p(a + b) = c, \quad c \mid h. \quad (9.2)$$

Thus a Type I solution is the same thing as a triple (a, h, c) with $c \mid h$ for which

$$b = \frac{c + pa}{4ah - p} \quad (9.3)$$

is a positive integer coprime to a . Now $\#\{(a, h, c) : ah = y, c \mid h\} = \sum_{h \mid y} \tau(h) = \tau_3(y)$, and $y = ah$ satisfies $p/4 < y \leq 3p/4$. Modelling the divisibility in (9.3) as occurring with probability $1/(4ah - p) = 1/(4y - p)$ leads to

$$M_I(p) := \sum_{p/4 < y \leq 3p/4} \frac{\tau_3(y)}{4y - p}. \quad (9.4)$$

The same computation for Type II is instructive. With $Y \leq Z$, $g = \gcd(Y, Z)$, $Y = ag$, $Z = bg$, the equation $4abcd = cp + a + b$ becomes $4abg - a - b = cp$ with $c \mid g$, so that a Type II solution is a triple (a, g, c) with $c \mid g$ for which $b = (cp + a)/(4ag - 1)$ is a positive integer

coprime to a . The same modelling gives

$$M_{\text{II}}(p) := \sum_Y \frac{w_p(Y)}{4Y-1}, \quad w_p(Y) = \#\{(a, g, c) : ag = Y, c \mid g, cp + a \geq 4Y - 1\}, \quad (9.5)$$

and $w_p(Y) = \tau_3(Y)$ for all $Y \leq p/4$.

The two expressions (9.4) and (9.5) differ in exactly one respect, and it is this difference which produces the surplus. In (9.4) the divisor function τ_3 is evaluated *at the single scale* $y \asymp p$, and the weights $1/(4y-p)$ run over $1, 5, 9, \dots$; in (9.5) it is evaluated *at all scales* $Y \leq p/4$, with the weight $1/(4Y-1)$. Using

$$\sum_{n \leq x} \tau_3(n) \sim \frac{x(\log x)^2}{2}, \quad \sum_{n \leq x} \frac{\tau_3(n)}{n} \sim \frac{(\log x)^3}{6},$$

we obtain

$$M_{\text{I}}(p) \sim \frac{(\log p)^2}{2} \cdot \frac{\log(2p)}{4} \sim \frac{(\log p)^3}{8}, \quad M_{\text{II}}(p) \sim \frac{1}{4} \cdot \frac{(\log p)^3}{6} = \frac{(\log p)^3}{24}, \quad (9.6)$$

so that

$$\frac{M_{\text{I}}(p)}{M_{\text{II}}(p)} \longrightarrow \frac{1/2}{1/6} = 3.$$

The constant 3 is precisely the ratio between the mean value of τ_3 at a single scale and its $1/n$ -weighted average over all smaller scales.

TABLE 6. The model (9.4), (9.5) against the predicted constants (9.6).

p	$M_{\text{I}}(p)$	$(\log p)^3/8$	$M_{\text{II}}(p)$	$(\log p)^3/24$	$M_{\text{I}}/M_{\text{II}}$
1 009	42.3	41.4	15.8	13.8	2.67
100 003	202.7	190.8	69.0	63.6	2.94
300 007	250.5	250.7	90.1	83.6	2.78
1 000 003	399.7	329.6	117.7	109.9	3.40

Table 6 shows that M_{I} and M_{II} do follow the constants in (9.6); the individual ratios fluctuate because $M_{\text{I}}(p)$ depends on τ_3 near the single point p . The model is not accurate in absolute terms — averaged over the primes $p \in [10^3, 3 \cdot 10^3]$ one finds $M_{\text{I}}/(f_{\text{I}}/2) = 1.73$ and $M_{\text{II}}/(f_{\text{II}}/2) = 1.80$, the model overshooting both by a comparable factor (we have ignored the condition $\gcd(a, b) = 1$ and the ordering $a \leq b$) — but the *ratio* is reproduced well: over the same primes $M_{\text{I}}/M_{\text{II}} = 2.70$ against the true mean $f_{\text{I}}/f_{\text{II}} = 2.61$.

We are led to the following prediction.

Conjecture 9.7 (= Conjecture 1.7). As $N \rightarrow \infty$,

$$\sum_{p \leq N} f_{\text{I}}(p) \sim \frac{c}{8} N \log^2 N, \quad \sum_{p \leq N} f_{\text{II}}(p) \sim \frac{c}{24} N \log^2 N$$

for some absolute constant $c > 0$; in particular

$$R(N) := \frac{\sum_{p \leq N} f_{\text{I}}(p)}{\sum_{p \leq N} f_{\text{II}}(p)} \longrightarrow 3.$$

Two comments. First, Conjecture 9.7 asserts in particular that $\sum_{p \leq N} f_I(p) \asymp N \log^2 N$ with *no* factor $\log \log N$; Elsholtz and Tao [9, after Theorem 1.1] write that the $\log \log N$ in their upper bound “arises from technical limitations to our method” and conjecture that it should be eliminated, and the model above is consistent with that.

Second, the data agree. Table 7 lists $R(N)$; a least-squares fit of the two natural shapes to 41 values of N between 10^3 and 10^6 gives

$$R(N) \approx 3.0159 - \frac{3.0327}{\log N} \quad (\text{rms } 0.0084), \quad R(N) \approx 1.9859 + 0.3128 \log \log N \quad (\text{rms } 0.0058).$$

The second fits marginally better on the available range, but is unbounded; the first has limit 3.016, in agreement with the model. We do not think the comparison of these two residuals is meaningful in itself — see §10, where separating the numerator from the denominator gives a much sharper test — but we do regard the size of R as the strongest available evidence for Conjecture 1.6: the expected margin between f_I and f_{II} is a factor of 3, not a hair, which is why exceptions are as rare as Table 3 and Figure 2 show.




FIGURE 4. The ratio $R(N)$ at 41 values of N between 10^3 and 10^6 , with the two least-squares fits and the predicted asymptote $R = 3$. On this statistic alone the two fits are comparable, the logarithmic one being marginally the better; see §10 for the sharper test.

TABLE 7. $R(N) = \sum_{p \leq N} f_I(p) / \sum_{p \leq N} f_{II}(p)$.

N	1 000	1 778	3 162	5 623	10 000	17 783	31 623
$R(N)$	2.5811	2.6254	2.6455	2.6518	2.6749	2.7003	2.7181
N	31 623	56 234	100 000	177 828	316 228	562 341	1 000 000
$R(N)$	2.7181	2.7396	2.7516	2.7640	2.7797	2.7931	2.8073

9.8. Approaches that do not work. We record, with counterexamples, several natural strategies for Conjecture 1.6 that fail; all were checked exhaustively in the stated ranges.

- (1) *Termwise comparison in (1.5).* The inequality $B(n) \geq A(n)$ fails: among the primes $5 \leq p \leq 400$ only thirteen (namely 7, 11, 13, 19, 31, 37, 43, 67, 79, 103, 127, 163, 211) admit no local counterexample. The smallest is $p = 59$, $n = 17$, where $A = 2$ and $B = 0$.

- (2) *Termwise comparison after grouping by e .* Among the 721 pairs (p, e) with $p \leq 300$ and $e \leq (p+4)/3$, exactly 50 satisfy $D(e, (pe+1)/4) < D(e, (p+e)/4)$, where $D(e, m) = \#\{t \mid m^2 : t \equiv -m \pmod{e}\}$.
- (3) *An explicit injection in the (a, b) parametrisation.* By Lemma 2.1,

$$f_I(p) = \#\{(a, b) : \gcd(a, b) = 1, ((-p^{-1}) \bmod 4ab) \mid a+b\},$$

$$f_{II}(p) = \#\{(a, b) : \gcd(a, b) = 1, ((-p) \bmod 4ab) \mid a+b\}.$$

The map $(a, b) \mapsto (ca, b)$ with $c = (a+b)/e$ is a bijection for $p = 17$, but among the 9092 Type II data with $p \leq 2000$ its image fails to be a Type I datum in 7759 cases.

- (4) *Exploiting an asymptotic margin.* Although $R(N) \rightarrow 3$ is expected, the *pointwise* ratio does not separate: over primes $5 < p \leq 10^6$ with $p \not\equiv 1 \pmod{8}$ the minimum of f_I/f_{II} is 1.2857 (at $p = 101$), and the minimum over a dyadic range lies between 1.2857 and 2.25 (Figure 2). It does not tend to 1; but it is also not bounded away from 1 by a margin that grows, which is what an inductive argument would need.
- (5) *Local pairing.* By Theorem 3.2 a Type I and a Type II solution share at most one denominator, and that denominator is the unique p -free denominator of the Type II solution. Any proof by matching solutions that share a denominator therefore reduces to (1), which fails.
- (6) *Quadratic characters.* By Proposition 7.4 a quadratic obstruction kills $A(n)$ and $B(n)$ simultaneously, so it cannot separate the two counts. Any argument must use characters of order > 2 .
- (7) *Bounding the non-principal characters in Proposition 7.2 absolutely.* Since $\tau(n^2) \asymp \log^2 n$ while the modulus e_n can be as large as $\asymp p$, the divisors of n^2 are far too sparse to equidistribute: $\varphi(e)^{-1} \sum_{\chi \neq \chi_0} |R_\chi(n)|$ is of size $\asymp \sqrt{\tau(n^2)}$, larger than the main term $\tau(n^2)/\varphi(e)$ by a factor $\asymp n/\log n$.

In view of §9.4 and §9.7, the two mechanisms that do produce the surplus are the vanishing of A for $n > (p+1)/3$ (Theorem 5.1) together with Theorems 1.4 and 1.2, and the scale asymmetry expressed by (9.6).

9.9. Further questions.

- (1) Improve the pointwise bound $D(K) \ll K \log K$ of Theorem 1.2. By Theorem 6.2 the average over p is only $O_m((\log K)^3 \log \log K)$, and Remark 6.3 suggests that even $O(K)$ is not optimal pointwise; the natural guess is that $\#\{\text{Type II solutions} : Y \leq K\} \ll_\varepsilon K^\varepsilon$ for every $\varepsilon > 0$, uniformly in p and m . The obstruction to any elementary improvement is that the pointwise bound already uses every algebraic constraint: for each pair (a, g) the parameter c is forced, and what remains is the arithmetic question of whether that c divides g , which is an equidistribution statement.
- (2) Extend Theorem 1.5 beyond $e \leq 5$ under additional hypotheses, or determine exactly the pairs (p, n) with $2B(n) < A(n)$ for $e = 7$ and $e = 9$ by the method of §7.
- (3) Prove an averaged form of Conjecture 1.6, for instance $\sum_{p \leq N} f_I(p) > \sum_{p \leq N} f_{II}(p)$ for large N . By (1.5) the layers $e = 1$ contribute equally to both sides, so this is a statement about the layers $e \geq 3$.
- (4) By Proposition 7.4, the difference $f_I - f_{II}$ is invisible at the quadratic level. All twelve exceptions in Table 3 other than $p = 5$ satisfy $p \equiv 1 \pmod{8}$, i.e. $(2/p) = 1$; the analogous phenomenon in [9, Prop. 1.6] is governed by quadratic reciprocity. Is there a reciprocity-theoretic explanation of the exceptional set?

10. THE DOUBLE LOGARITHM IN THE ELSHOLTZ-TAO UPPER BOUND

Elsholtz and Tao [9] prove

$$N \log^2 N \ll \sum_{p \leq N} f_I(p) \ll N \log^2 N \log \log N, \quad N \log^2 N \ll \sum_{p \leq N} f_{II}(p) \ll N \log^2 N, \quad (10.1)$$

and write that the factor $\log \log N$, which occurs for Type I but not for Type II, “arises from technical limitations to our method (and specifically, in the inefficient nature of the Brun–Titchmarsh inequality when applied to very short progressions), and we conjecture that it should be eliminated”. The model of §9.7 predicts the same (Conjecture 1.7). This section does three things: it isolates the loss exactly, it reduces its removal to the gain of a single logarithm, and it shows that the device by which [9] removes the analogous loss on the Type II side is in fact partly available on the Type I side.

10.1. Identities for a Type I quadruple. We first record the algebra we shall need. Recall from Lemma 2.1 that a Type I solution is a quadruple (a, b, c, d) with $\gcd(a, b) = 1$ and $4abcd = p(a + b) + c$, and that $\delta = 4acd - p$ is the invariant of Theorem 4.1(i); in this section we follow [9] and write $f := \delta$.

Lemma 10.1. *Let (a, b, c, d) be a Type I quadruple for the prime $p \geq 5$ and put $f := 4acd - p$. Then $c \mid a + b$, and with $g := (a + b)/c$,*

$$fg = 4a^2d + 1, \quad pg = 4abd - 1, \quad (4ab)df = p(4a^2d + 1) + f. \quad (10.2)$$

Moreover $0 < f < 4acd$ and $\gcd(f, 4acd) = 1$. If in addition $a \leq b$, then

$$\frac{p}{4} < acd \leq \frac{2p}{3}, \quad f \leq p + c. \quad (10.3)$$

Proof. From $4abcd = p(a + b) + c$ we get $c \mid p(a + b)$, and $p \nmid c$ because $p \nmid acd$ for a Type I solution; hence $c \mid a + b$. Multiplying $fg = 4a^2d + 1$ by c turns it into $(4acd - p)(a + b) = 4a^2cd + c$, and indeed $4acd(a + b) - p(a + b) = 4a^2cd + 4abcd - (4abcd - c) = 4a^2cd + c$. Multiplying $pg = 4abd - 1$ by c returns the defining equation. The third identity follows from the second: $4abd = pg + 1$, so $(4ab)df = (pg + 1)f = pgf + f = p(4a^2d + 1) + f$.

Since $1/(acd) < 4/p$ we have $acd > p/4$, i.e. $f > 0$; and $f < 4acd$ because $p > 0$. Also $\gcd(f, 4acd) = \gcd(p, 4acd) = 1$. Finally suppose $a \leq b$. Then $2/(acd) \geq 1/(acd) + 1/(bcd) = 4/p - 1/(abdp) \geq 3/p$, which gives $acd \leq 2p/3$. And $b \geq a$ gives $cg = a + b \geq 2a$, i.e. $c(4a^2d + 1) \geq 2af$, i.e. $f \leq 2acd + c/(2a)$; since $4acd = p + f$ this reads $f \leq (p + f)/2 + c/(2a)$, whence $f \leq p + c/a \leq p + c$. $\square$

10.2. The block decomposition, and exactly where the logarithm is lost. For a dyadic parameter $C \geq 1$ put

$$P(N; C) := \#\{(a, b, c, d) : \text{a Type I quadruple for some prime } p \leq N, a \leq b, C \leq c < 2C\}.$$

By Theorem 5.3(i), $f_I(p) \leq 2 \cdot \#\{\text{unordered Type I solutions}\}$, so $\sum_{p \leq N} f_I(p) \leq 2 \sum_C P(N; C)$, the sum being over dyadic C .

Lemma 10.2. *For every dyadic $C \geq 1$,*

$$(i) \quad P(N; C) \ll N \log^2 N, \quad (ii) \quad P(N; C) \ll \frac{N \log^2(2 + N/C)}{\log(2C)}.$$

Proof. By Lemma 10.1 the quadruple is determined by (a, c, d, f) , since $g = (4a^2d + 1)/f$ and $b = cg - a$; and f is a divisor of $4a^2d + 1$. By (10.3), $acd \leq 2N/3$, so $ad \leq 2N/(3C)$.

(i) For fixed (a, d, f) there are at most C admissible c , so by [9, Prop. 1.4] (applied in dyadic blocks in a and d)

$$P(N; C) \leq C \sum_{ad \leq 2N/(3C)} \tau(4a^2d + 1) \ll C \cdot \frac{N}{C} \log^2 N = N \log^2 N.$$

(ii) Fix (a, d, f) . The primes counted are $p = 4acd - f$ with $C \leq c < 2C$; they lie in the residue class $-f$ modulo $4ad$, which is primitive because $\gcd(f, 4ad) = 1$, and in an interval of length $4aCd$. For $C \geq 3$ the Brun–Titchmarsh inequality [14] bounds their number by $\ll aCd/(\varphi(4ad) \log C)$, while for $C \leq 2$ the trivial bound $\leq C$ serves. Summing, and treating the factor $ad/\varphi(4ad)$ exactly as in [9, (8.2)],

$$P(N; C) \ll \frac{C}{\log(2C)} \sum_{ad \leq 2N/(3C)} \frac{ad \tau(4a^2d + 1)}{\varphi(4ad)} \ll \frac{C}{\log(2C)} \cdot \frac{N}{C} \log^2 \left(2 + \frac{N}{C}\right). \quad \square$$

Writing $C = 2^k$ and $L = \log N$, Lemma 10.2(ii) reads $P(N; 2^k) \ll N(L - k \log 2)_+^2 / \max(1, k)$, and $\sum_{k \leq L/\log 2} (L - k \log 2)^2 / k \asymp L^2 \log L$: this harmonic sum, and nothing else, is the $\log \log N$ of (10.1). Two consequences are immediate.

Theorem 10.3 (Localisation). *For every $\varepsilon \in (0, 1)$,*

$$\sum_{C \geq N^\varepsilon} P(N; C) \ll \varepsilon^{-1} N \log^2 N.$$

Hence the factor $\log \log N$ in (10.1) arises entirely from the Type I solutions with $c \leq N^{o(1)}$, equivalently from the range in which the modulus $4ad$ of Lemma 10.2(ii) exceeds $N^{1-o(1)}$.

Proof. Summing the bound $P(N; 2^k) \ll N(L - k \log 2)_+^2 / k$ over the range $k \geq \varepsilon L / \log 2$ gives

$$\ll \frac{N}{\varepsilon L} \sum_{0 \leq k \leq L/\log 2} (L - k \log 2)^2 \ll \varepsilon^{-1} N L^2. \quad \square$$

Theorem 10.4 (One logarithm suffices). *Suppose that*

$$P(N; C) \ll \frac{N \log^2(2 + N/C)}{\log^2(2C)} \quad \text{for every dyadic } C \text{ with } 2 \leq C \leq N.$$

Then $\sum_{p \leq N} f_1(p) \ll N \log^2 N$, i.e. the factor $\log \log N$ in (10.1) may be deleted.

Proof. The block $C = 1$ is $\ll N \log^2 N$ by Lemma 10.2(i), while

$$\sum_{k \geq 1} \frac{N(L - k \log 2)_+^2}{(k \log 2)^2} \ll N L^2 \sum_{k \geq 1} k^{-2} \ll N L^2. \quad \square$$

Thus the whole question is whether Brun–Titchmarsh can be improved by *one* factor of $\log(2C)$, uniformly over the family of progressions occurring here. For a single progression Brun–Titchmarsh is essentially optimal, so any such gain must come from the average over the $\tau(4a^2d + 1)$ residue classes $-f$, or over the moduli $4ad$.

10.3. Four moduli for Type I. On the Type II side, Elsholtz and Tao avoid the loss altogether [9, §9]: their Type II parameters (a, c, d, e) admit *three* moduli ade , acd , ab whose product satisfies $(ade)(acd)(ab)^{1/2} \ll N^2$, so that one of them is $O(N^{4/5})$ and Brun–Titchmarsh may be applied to a progression that is not short. They add that “it does not seem that a similar trick is available” in the Type I case. The following shows that it is, in part.

Proposition 10.5 (Four moduli). *Let (a, b, c, d) be a Type I quadruple for p , with f, g as in Lemma 10.1, and put*

$$q_1 := 4ad, \quad q_2 := 4acf, \quad q_3 := 4cdf, \quad q_4 := 4ab.$$

Then, in each of the following four ways, the quadruple is determined by the listed data together with one further variable, and p is confined to the stated residue classes:

- (i) *data (a, d, f) , variable c : $p \equiv -f \pmod{q_1}$;*
- (ii) *data (a, c, f) , variable d : d is determined modulo f by $4a^2d \equiv -1$, and $p \equiv -f \pmod{4ac}$, so p lies in one class modulo q_2 ;*
- (iii) *data (c, d, f) , variable a : a lies in at most $2^{\omega(f)}$ classes modulo f (by $4da^2 \equiv -1$), and $p \equiv -f \pmod{4cd}$, so p lies in at most $2^{\omega(f)}$ classes modulo q_3 ;*
- (iv) *data (a, b, g) , variable d : d is determined modulo g by $4abd \equiv 1$, and $p = (4abd - 1)/g$ runs through one class modulo q_4 .*

Proof. All four moduli are coprime to the relevant data: $f \mid 4a^2d + 1$ gives $\gcd(f, 2ad) = 1$, and $\gcd(f, c) = \gcd(p, c) = 1$ by Lemma 10.1. Case (i) is $p = 4acd - f$. In (ii) and (iii) the congruence $f \mid 4a^2d + 1$ determines d modulo f uniquely, respectively a modulo f up to the $2^{\omega(f)}$ square roots of $-(4d)^{-1}$ modulo the odd number f ; combining with $p \equiv -f$ modulo $4ac$ (resp. $4cd$) and the coprimality just noted gives the assertion. In (iv), $c = (a + b)/g$ is determined by the data, $pg = 4abd - 1$ by (10.2), and as d increases by g the value of p increases by $4ab$. $\square$

One may therefore run the Brun–Titchmarsh step of Lemma 10.2(ii) with the modulus $\min(q_1, q_2, q_3, q_4)$, at the cost of the factor $2^{\omega(f)}$ in case (iii). Numerically this is effective for the great majority of quadruples: for $p \leq 2 \cdot 10^4$ there are 188 400 Type I quadruples with $a \leq b$, and $\min(q_1, q_2, q_3, q_4) \leq p^{0.9}$ for 89.1% of them, against 83.9% for $\min(q_1, q_2, q_3)$ and 61.2% for q_1 alone. It is not, however, enough, and the reason is exactly the one identified in Theorem 10.3.

Proposition 10.6 (The residual range). *Let $0 < \delta < \frac{1}{5}$ and let (a, b, c, d) be a Type I quadruple for p with $a \leq b$ and $\min(q_1, q_2, q_3, q_4) > p^{1-\delta}$. Then*

$$c \ll p^\delta, \quad f \gg \max(a, d)p^{-\delta}, \quad f \gg p^{(1-3\delta)/2}.$$

Proof. By (10.3), $acd \asymp p$. Hence $q_1 = 4ad \asymp 4p/c$, and $q_1 > p^{1-\delta}$ gives $c \ll p^\delta$. Similarly $cd \asymp p/a$ gives $q_3 \asymp 4pf/a$, so $f \gg ap^{-\delta}$, and $ac \asymp p/d$ gives $q_2 \asymp 4pf/d$, so $f \gg dp^{-\delta}$. Finally $q_1q_2q_3 = 64(acd)^2f^2 \ll p^2f^2$, and the left-hand side exceeds $p^{3-3\delta}$. $\square$

The three conditions of Proposition 10.6 are consistent — take $a \asymp d \asymp p^{(1-\delta)/2}$, c bounded and $f \asymp p^{(1-\delta)/2}$ — so no contradiction can be extracted, and the residual range is genuinely non-empty. In the numerical range above, every quadruple with $\min(q_1, q_2, q_3, q_4) > p^{0.9}$ has $c \leq 5$, and 76% of them have $c = 1$. The four moduli therefore reach precisely as far as Theorem 10.3 already does, and no further: the obstruction is the range $c = O(1)$, where p is confined to a progression of bounded length to every modulus available, and where consequently no sieve can help.

N	$\sum_{p \leq N} f_I(p)$	$\sum_{p \leq N} f_I(p)/N \log^2 N$	$\sum_{p \leq N} f_{II}(p)/N \log^2 N$
1 000	10 056	0.2107	0.0816
1 778	20 691	0.2078	0.0792
3 162	41 408	0.2016	0.0762
5 623	84 072	0.2005	0.0756
10 000	165 358	0.1949	0.0729
17 783	328 311	0.1928	0.0714
31 623	642 741	0.1893	0.0696
56 234	1 269 719	0.1888	0.0689
100 000	2 468 551	0.1862	0.0677
177 828	4 790 527	0.1843	0.0667
316 228	9 271 148	0.1828	0.0658
562 341	17 833 719	0.1809	0.0648
1 000 000	34 276 271	0.1796	0.0640

TABLE 8. The two aggregates against $N \log^2 N$. For Type II the order $N \log^2 N$ is a theorem of [9]; the Type I column has the same shape.

10.4. Numerical evidence for the conjectured bound. Table 8 lists $\sum_{p \leq N} f_I(p)$ and $\sum_{p \leq N} f_{II}(p)$, each divided by $N \log^2 N$, for 41 values of N between 10^3 and 10^6 (thirteen of them shown). The point of listing the second is that it is a *control*: by (10.1) its order of magnitude is known to be exactly $N \log^2 N$, so whatever shape it has is the shape of a quantity with no $\log N$ in it.

The two columns have the same shape. Both decrease slowly; a least-squares fit of $\alpha + \beta/\log N$ over the 41 values gives $\alpha = 0.1483$ for Type I and $\alpha = 0.0466$ for Type II, with root-mean-square residuals 0.0014 and 0.0005. A fit of $\alpha + \beta \log \log N$ is slightly worse in both cases (residuals 0.001442 against 0.001360, and 0.000541 against 0.000474), but the residuals are not the point: the fitted slopes are *negative*, $\beta = -0.045$ for Type I and $\beta = -0.025$ for Type II. A genuine factor $\log \log N$ in $\sum_{p \leq N} f_I(p)$ would force a positive slope here; the data give the opposite sign, and give it equally for Type II, where the absence of the factor is a theorem. We regard this as the strongest numerical statement available at this range.

It should be said what the data do *not* show. The ratio $R(N)$ of §9.7 is a quotient of the two columns, and there the small residuals compound: over the same 41 values a fit of $a - b/\log N$ to $R(N)$ has root-mean-square residual 0.0084 against 0.0058 for a fit of $a' + b' \log \log N$, so $R(N)$ alone does not distinguish a finite limit from logarithmic growth (Figure 4). It is only after separating the numerator and the denominator, and using Type II as a control, that the data speak.

REFERENCES

- [1] A. Aigner, *Brüche als Summe von Stammbrüchen*, J. Reine Angew. Math. **214/215** (1964), 174–179.
- [2] M. Bello-Hernández, M. Benito and E. Fernández, *A divisor parametrization for the Erdős–Straus conjecture*, preprint (2026), arXiv:2606.10922.
- [3] L. Bernstein, *Zur Lösung der diophantischen Gleichung $m/n = 1/x + 1/y + 1/z$, insbesondere im Fall $m = 4$* , J. Reine Angew. Math. **211** (1962), 1–10.
- [4] K. Bradford, *A note on the Erdős–Straus conjecture*, Integers **21** (2021), #A24, 10 pp.; doi:10.5281/zenodo.10807535; arXiv:1906.00561.
- [5] K. Bradford, *Elementary patterns from the Erdős–Straus conjecture*, Integers **25** (2025), #A54, 10 pp.; preprint version: *Elemental patterns from the Erdős–Straus conjecture*, arXiv:2403.16047.

- [6] M. Chamberland, *The Erdős–Straus conjecture and the structure of primes*, *Integers* **26** (2026), #A42, 7 pp. doi:10.5281/zenodo.19403738.
- [7] B. Dahan, *Sieve dimension and search depth for the Erdős–Straus conjecture, $n \equiv 1 \pmod{24}$* , preprint (2026), arXiv:2608.24035.
- [8] C. Elsholtz and S. Planitzer, *The number of solutions of the Erdős–Straus equation and sums of k unit fractions*, *Proc. Roy. Soc. Edinburgh Sect. A* **150** (2020), no. 3, 1401–1427; arXiv:1805.02945.
- [9] C. Elsholtz and T. Tao, *Counting the number of solutions to the Erdős–Straus equation on unit fractions*, *J. Aust. Math. Soc.* **94** (2013), no. 1, 50–105; arXiv:1107.1010.
- [10] R. K. Guy, *Unsolved Problems in Number Theory*, 3rd ed., Problem Books in Mathematics, Springer, New York, 2004; §D11.
- [11] M. A. López, *Structure and form of the solutions of the Erdős–Straus conjecture*, preprint (2024), arXiv:2206.10319v4.
- [12] S. Mihnea and B. C. Dumitru, *Further verification and empirical evidence for the Erdős–Straus conjecture*, preprint (2025), arXiv:2509.00128.
- [13] M. Monks and A. Velingker, *On the Erdős–Straus conjecture: properties of solutions to its underlying Diophantine equation*, unpublished manuscript dated 23 September 2008, originating in a 2004–2005 Siemens Westinghouse competition entry; available from the first author’s page, <http://mathematicalgemstones.com/maria/papers/ErdosStraus.pdf>.
- [14] H. L. Montgomery and R. C. Vaughan, *The large sieve*, *Mathematika* **20** (1973), 119–134.
- [15] L. J. Mordell, *Diophantine Equations*, Pure and Applied Mathematics, vol. 30, Academic Press, London–New York, 1969.
- [16] R. Obláth, *Sur l’équation diophantienne $4/n = 1/x_1 + 1/x_2 + 1/x_3$* , *Mathesis* **59** (1950), 308–316.
- [17] L. Rosati, *Sull’equazione diofantea $4/n = 1/x_1 + 1/x_2 + 1/x_3$* , *Boll. Un. Mat. Ital. (3)* **9** (1954), 59–63.
- [18] S. E. Salez, *The Erdős–Straus conjecture: new modular equations and checking up to $N = 10^{17}$* , preprint (2014), arXiv:1406.6307.
- [19] A. Schinzel, *Sur quelques propriétés des nombres $3/n$ et $4/n$, où n est un nombre impair*, *Mathesis* **65** (1956), 219–222.
- [20] A. Schinzel, *On sums of three unit fractions with polynomial denominators*, *Funct. Approx. Comment. Math.* **28** (2000), 187–194.
- [21] W. Sierpiński, *Sur les décompositions de nombres rationnels en fractions primaires*, *Mathesis* **65** (1956), 16–32.
- [22] D. G. Terzi, *On a conjecture by Erdős–Straus*, *Nordisk Tidskr. Informationsbehandling (BIT)* **11** (1971), 212–216.
- [23] R. C. Vaughan, *On a problem of Erdős, Straus and Schinzel*, *Mathematika* **17** (1970), 193–198.
- [24] W. A. Webb, *On $4/n = 1/x_1 + 1/x_2 + 1/x_3$* , *Proc. Amer. Math. Soc.* **25** (1970), no. 3, 578–584.
- [25] X. Xu, *Congruence classes supporting the Erdős–Straus conjecture I: tame solutions*, preprint (2026), arXiv:2605.23601.
- [26] K. Yamamoto, *On the Diophantine equation $4/n = 1/x + 1/y + 1/z$* , *Mem. Fac. Sci. Kyushu Univ. Ser. A* **19** (1965), 37–47.

Email address: jiangxinyang222@gmail.com