

Congruence Classes of Supporting the Erdős-Straus Conjecture II: Wild Solutions ¹

Xiaoping Xu

HLM, Institute of Mathematics, Academy of Mathematics & System Sciences
Chinese Academy of Sciences, Beijing 100190, P.R. China
& School of Mathematics, University of Chinese Academy of Sciences,
Beijing 100049, P.R. China

Abstract

In 1948, Erdős and Straus formulated a conjecture : for any positive integer $n > 2$, there exist positive integers n_1, n_2 and n_3 such that

$$\frac{4}{n} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3},$$

which is still open. It is known that the conjecture holds if one can prove it for any prime $n \equiv 1 \pmod{24}$. If $n = 24m + 1$ and $n_1 \leq n_2, n_3$, then $n_1 = 6m + k$ with $1 \leq k \leq 12m$. A solution (n_1, n_2, n_3) of the above equation is called a *tame solution* if n_2 and n_3 are factors of $(6m + k)(24m + 1)$. We call $n = 24m + 1$ *wild* if it does not have any tame solution. Based on the information in our earlier work on tame solutions posed in arXiv, Howerton found that there are only fourteen wild primes of the form $n = 24m + 1 \leq 2.4 \times 10^{11}$. In this paper, we derive thirty-four families of wild solutions of the above equation, which contain the solvability of the fourteen wild primes. Together with our earlier tame polynomial solutions, numeric test shows that they cover all the primes of the form $24m + 1$.

Keywords: Erdős-Straus conjecture; Egyptian fraction; congruence class; tame solution; wild solution; wild prime.

1 Introduction

Ancient Egyptians used sums of unit fractions (whose numerators are 1) to express fractions due to their ways of distributing food. For instance, $5/8$ was interpreted by ancient Egyptians as distributing five pancakes fairly among eight people. They cut first four pancakes into halves and then cut the last one into eight equal pieces. So each person got the same share: $1/2 + 1/8$ pancakes. This amazingly interpreted the mathematical equation

$$\frac{5}{8} = \frac{1}{2} + \frac{1}{8}. \quad (1.1)$$

So an Egyptian fraction is a sum of distinct unit fractions. By repeatedly applying the simple fact

$$\frac{1}{k} = \frac{1}{k+1} + \frac{1}{k(k+1)}, \quad (1.2)$$

¹2010 Mathematical Subject Classification. Primary 11D68; Secondary 11D85, 11A67, 11B75

one can easily prove that any fraction is a Egyptian fraction. However, it is very difficult to determine if a fraction can be expressed as a sum of fixed number of unit fractions (cf. [3] for an excellent exposition and extensive references). For example, it is difficult to know if a fraction can be written as a sum of two unit fractions (e.g., cf. [6, 7, 8, 10, 16]). In 1948, Erdős and Straus formulated a conjecture : for any positive integer $n > 2$, there exist positive integers n_1, n_2 and n_3 such that

$$\frac{4}{n} = \frac{1}{n_1} + \frac{1}{n_2} + \frac{1}{n_3}, \quad (1.3)$$

which is still open up to now. For each n , the number of solution can go to infinity as n does. Elscholtz and Tao [3] found excellent bounds for it.

Mordell [15] proved that the conjecture holds for positive integers in 834 congruence classes modulo 840. Terzi [23] used computer to verify that the conjecture holds for positive integers in the congruence classes modulo 120120 except 198 classes. Kotsireas [12] verified the conjecture for every $n < 10^{10}$. There are other interesting partial results on the conjecture or related works (e.g., cf. [1, 2, 4, 11, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25]).

It is known that one only needs to prove the conjecture for any prime number n such that $n \equiv 1 \pmod{24}$. Suppose that $n = 24m + 1$ and $n_1 \leq n_2, n_3$. Then $n_1 = 6m + k$ with $1 \leq k \leq 12m$ and

$$\frac{4}{24m + 1} = \frac{1}{6m + k} + \frac{4k - 1}{(6m + k)(24m + 1)}. \quad (1.4)$$

If

$$\frac{1}{n_2} = \frac{\mathfrak{S}_1}{(6m + k)(24m + 1)} \quad \text{and} \quad \frac{1}{n_2} = \frac{\mathfrak{S}_2}{(6m + k)(24m + 1)} \quad (1.5)$$

for some positive integers $\mathfrak{S}_1$ and $\mathfrak{S}_2$ such that $\mathfrak{S}_1 + \mathfrak{S}_2 = 4k - 1$, we call the triple (n_1, n_2, n_3) a *tame solution* of the Erdős-Straus equation (1.3) for the positive integer $n = 24m + 1$. Moreover, we call $n = 24m + 1$ *wild* if it does not have any tame solution. Based on the information in our earlier work on tame solutions posed in arXiv, Howerton found that there are only fourteen wild primes of the form $n = 24m + 1 \leq 2.4 \times 10^{11}$.

By analysing the obtained wild solutions, we find a way of obtaining them as follows. In (1.4), we multiply a proper integer $\ell \geq 2$ both to the numerator and denominator of the fraction

$$\frac{4k - 1}{(6m + k)(24m + 1)} \implies \frac{\ell(4k - 1)}{\ell(6m + k)(24m + 1)} \quad (1.6)$$

and find positive integers $\mathfrak{S}'_1$ and $\mathfrak{S}'_2$ such that

$$\mathfrak{S}'_1 + \mathfrak{S}'_2 = \ell(4k - 1), \quad (1.7)$$

$$g.c.d(\ell, \mathfrak{S}'_1, \mathfrak{S}'_2) = 1 \quad (1.8)$$

and

$$\frac{1}{n_2} = \frac{\mathfrak{S}'_1}{\ell(6m + k)(24m + 1)} \quad \text{and} \quad \frac{1}{n_2} = \frac{\mathfrak{S}'_2}{\ell(6m + k)(24m + 1)} \quad (1.9)$$

are unit fractions. We call the integer ℓ a *multiplier* of the wild solution (n_1, n_2, n_3) . Moreover, we call $\mathfrak{S}'_1$ and $\mathfrak{S}'_2$ the *numerator summands* for (n_1, n_2, n_3) . According to our definitions, a tame prime of the form $24m + 1$ may have wild solutions, but a wild prime

of the form $24m + 1$ only has wild solutions. Based on the information in our earlier work on tame solutions posed in arXiv, Howerton [5] found that there are only fourteen wild primes of the form $n = 24m + 1 \leq 2.4 \times 10^{11}$. In this paper, we derive thirty-four families of wild solutions of the above equation according to (1.6)-(1.9), which contain the solvability of the fourteen wild primes. Together with our tame polynomial solutions in [26], numeric test shows that they cover all the primes of the form $24m + 1$. The paper is organized as follows.

In Section 2, we find a relatively simple wild solution for each of the fourteen wild primes of the form $24m + 1$. In Section 3, we solve the Erdős-Straus equation for $k \leq 5$ by the method described in (1.6)-(1.9). Moreover, we solve the equation for $k = 6$ in Section 4.

2 Solutions for the Fourteen Wild Primes

Howerton [5] found that there are only wild primes of the form $24m + 1$ with $m \leq 10^9$. They are

$$409, \quad 577, \quad 5569, \quad 9601, \quad 23929, \quad 83449, \quad 102001, \quad 329617, \quad 712321, \quad (2.1)$$

$$1134241, \quad 1724209, \quad 1726201, \quad 5212561, \quad 8813281. \quad (2.2)$$

In the form of $n = 24m + 1$, the corresponding m 's are as follows:

$$17, \quad 24, \quad 232, \quad 400, \quad 997, \quad 3477, \quad 4250, \quad 13734, \quad 29680, \quad (2.3)$$

$$47260, \quad 71842, \quad 71925, \quad 217190, \quad 367220. \quad (2.4)$$

In this section, we derive a relatively simple wild solution for each of them as the starting point of this paper.

Throughout this paper, we denote by $\mathbb{N}$ the set of nonnegative integers. Suppose $m \equiv 4 \pmod{5}$; that is, $m = 4 + 5c$ for some $c \in \mathbb{N}$. We have

$$\begin{aligned} \frac{4}{24m+1} &= \frac{1}{6m+1} + \frac{3}{(6m+1)(24m+1)} \\ &= \frac{1}{6m+1} + \frac{6}{2(30c+25)(24m+1)} \\ &= \frac{1}{6m+1} + \frac{5+1}{10(6c+5)(24m+1)} \\ &= \frac{1}{6m+1} + \frac{1}{2(6c+5)(24m+1)} + \frac{1}{10(6c+5)(24m+1)}. \end{aligned} \quad (2.5)$$

Taking $m = 24$ and $c = 4$, we get $24m + 1 = 577$. $6m + 1 = 145$ and

$$\frac{4}{577} = \frac{1}{145} + \frac{1}{58 \times 577} + \frac{1}{290 \times 577}. \quad (2.6)$$

When $m = 13734$, $24m + 1 = 329617$. $6m + 1 = 82405$, $c = 2746$ and

$$\frac{4}{329617} = \frac{1}{82405} + \frac{1}{32962 \times 329617} + \frac{1}{164810 \times 329617}. \quad (2.7)$$

Next we assume $m \equiv 3 \pmod{7}$; that is, $m = 3 + 7c$ for some $c \in \mathbb{N}$. We have

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+3} + \frac{11}{(6m+3)(24m+1)} \\
&= \frac{1}{6m+3} + \frac{22}{2(42c+21)(24m+1)} \\
&= \frac{1}{6m+3} + \frac{21+1}{42(2c+1)(24m+1)} \\
&= \frac{1}{6m+3} + \frac{1}{2(2c+1)(24m+1)} + \frac{1}{42(2c+1)(24m+1)}. \tag{2.8}
\end{aligned}$$

Taking $m = 17$ and $c = 2$, we get $24m + 1 = 409$, $6m + 3 = 105$ and

$$\frac{4}{409} = \frac{1}{105} + \frac{1}{10 \times 409} + \frac{1}{210 \times 409}. \tag{2.9}$$

When $m = 47260$ and $c = 6751$, we have $24m + 1 = 1134241$, $6m + 3 = 283563$ and

$$\frac{4}{1134241} = \frac{1}{283563} + \frac{1}{27006 \times 1134241} + \frac{1}{567126 \times 1134241}. \tag{2.10}$$

If $m = 367220$ and $c = 52031$, we have $24m + 1 = 8813281$, $6m + 3 = 2203323$ and

$$\frac{4}{8813281} = \frac{1}{2203323} + \frac{1}{104063 \times 8813281} + \frac{1}{2185323 \times 8813281}. \tag{2.11}$$

Suppose $m \equiv 11 \pmod{17}$; that is, $m = 11 + 17c$ for some $c \in \mathbb{N}$. We have

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+2} + \frac{7}{(6m+3)(24m+1)} \\
&= \frac{1}{6m+2} + \frac{35}{5(102c+68)(24m+1)} \\
&= \frac{1}{6m+2} + \frac{34+1}{5 \times 34(3c+2)(24m+1)} \\
&= \frac{1}{6m+2} + \frac{1}{5(3c+2)(24m+1)} + \frac{1}{170(3c+2)(24m+1)}. \tag{2.12}
\end{aligned}$$

Taking $m = 232$ and $c = 13$, we get $24m + 1 = 5569$, $6m + 2 = 1394$ and

$$\frac{4}{5569} = \frac{1}{1394} + \frac{1}{1141545} + \frac{1}{38815930}. \tag{2.13}$$

When $m = 997$, we have $c = 58$, $24m + 1 = 23929$, $6m + 2 = 5984$ and

$$\frac{4}{23929} = \frac{1}{5984} + \frac{1}{21057520} + \frac{1}{715955680}. \tag{2.14}$$

Let $m = 400$. Then $24m + 1 = 9601$ is a primes. Moreover, $6m + 5 = 2405 = 5 \times 13 \times 37$.

So

$$\begin{aligned}
\frac{4}{9601} &= \frac{1}{2405} + \frac{19}{2405 \times 9601} \\
&= \frac{1}{2405} + \frac{19}{(5 \times 13 \times 37) \times 9601} \\
&= \frac{1}{2405} + \frac{38}{2 \times (5 \times 13 \times 37) \times 9601} \\
&= \frac{1}{2405} + \frac{37+1}{2 \times (5 \times 13 \times 37) \times 9601} \\
&= \frac{1}{2405} + \frac{1}{2 \times (5 \times 13) \times 9601} + \frac{1}{2 \times (5 \times 13 \times 37) \times 9601} \\
&= \frac{1}{2405} + \frac{1}{1248130} + \frac{1}{46180810}.
\end{aligned} \tag{2.15}$$

In the case $m = 3477$, $24m + 1 = 83449$ is a primes. Moreover, $6m + 3 = 20865 = 3 \times 5 \times 13 \times 107$. Therefore,

$$\begin{aligned}
\frac{4}{83449} &= \frac{1}{20865} + \frac{11}{20865 \times 83449} \\
&= \frac{1}{20865} + \frac{11}{(3 \times 5 \times 13 \times 107) \times 83449} \\
&= \frac{1}{20865} + \frac{66}{6 \times (3 \times 65 \times 107) \times 83449} \\
&= \frac{1}{20865} + \frac{65+1}{6 \times (3 \times 65 \times 107) \times 83449} \\
&= \frac{1}{20865} + \frac{1}{6 \times (3 \times 107) \times 83449} + \frac{1}{6 \times (3 \times 65 \times 107) \times 83449} \\
&= \frac{1}{20865} + \frac{1}{160722774} + \frac{1}{10446980310}.
\end{aligned} \tag{2.16}$$

Assume $m = 4250$, $24m + 1 = 102001$ is a primes. Moreover, $6m + 2 = 25502 = 2 \times 41 \times 311$. Thus

$$\begin{aligned}
\frac{4}{102001} &= \frac{1}{25502} + \frac{7}{25502 \times 102001} \\
&= \frac{1}{25502} + \frac{7}{(2 \times 41 \times 311) \times 102001} \\
&= \frac{1}{25502} + \frac{42}{6 \times (2 \times 41 \times 311) \times 102001} \\
&= \frac{1}{25502} + \frac{41+1}{6 \times (2 \times 41 \times 311) \times 102001} \\
&= \frac{1}{25502} + \frac{1}{6 \times (2 \times 311) \times 102001} + \frac{1}{6 \times (2 \times 41 \times 311) \times 102001} \\
&= \frac{1}{25502} + \frac{1}{380667732} + \frac{1}{15607377012}.
\end{aligned} \tag{2.17}$$

We consider $m = 29680$, $24m + 1 = 712321$ is a primes. Moreover, $6m + 6 = 178086 =$

$2 \times 3 \times 67 \times 443$. Thus

$$\begin{aligned}
\frac{4}{712321} &= \frac{1}{178086} + \frac{23}{178086 \times 712321} \\
&= \frac{1}{178086} + \frac{23}{(2 \times 3 \times 67 \times 443) \times 712321} \\
&= \frac{1}{178086} + \frac{69}{3 \times (2 \times 3 \times 67 \times 443) \times 712321} \\
&= \frac{1}{178086} + \frac{67+2}{3 \times (2 \times 3 \times 67 \times 443) \times 712321} \\
&= \frac{1}{178086} + \frac{1}{3 \times (2 \times 3 \times 443) \times 712321} + \frac{1}{3 \times (3 \times 67 \times 443) \times 712321} \\
&= \frac{1}{178086} + \frac{1}{5680047654} + \frac{1}{190281596409}. \tag{2.18}
\end{aligned}$$

Now we assume $m \equiv 9 \pmod{29}$; equivalently, $m = 9 + 29c$ for some $c \in \mathbb{N}$. In this case,

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+4} + \frac{15}{2(3m+2)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{30}{4(3m+2)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{29+1}{116(3c+1)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{1}{4(3c+1)(24m+1)} + \frac{1}{116(3c+1)(24m+1)}. \tag{2.19}
\end{aligned}$$

Taking $m = 71842$, we get $c = 2477$ and $24m+1 = 1724209$ is a wild prime. By the above equation,

$$\frac{4}{1724209} = \frac{1}{431056} + \frac{1}{29728 \times 1724209} + \frac{1}{862112 \times 1724209}. \tag{2.20}$$

When $m = 217190$, we have $c = 7489$ and $24m+1 = 5212561$ is a wild prime. According to (2.18).,

$$\frac{4}{5212561} = \frac{1}{1303144} + \frac{1}{89872 \times 5212561} + \frac{1}{2606288 \times 5212561}. \tag{2.21}$$

Finally we consider $m = 71925$, $24m+1 = 1726201$ is a wild primes. Moreover, $6m+6 = 431556 = 4 \times 3 \times 35963$. In fact,

$$3 \times 35963 = 107889, \quad 23 \times 4691 = 107893 = 3 \times 35963 + 4. \tag{2.22}$$

Thus

$$\begin{aligned}
\frac{4}{1726201} &= \frac{1}{431556} + \frac{23}{431556 \times 1726201} \\
&= \frac{1}{431556} + \frac{4691 \times 23}{4691 \times 431556 \times 1726201} \\
&= \frac{1}{431556} + \frac{3 \times 35963 + 4}{4691 \times 4 \times 3 \times 35963 \times 1726201} \\
&= \frac{1}{431556} + \frac{1}{18764 \times 1726201} + \frac{1}{4691 \times 107889 \times 1726201}. \tag{2.23}
\end{aligned}$$

The above examples suggest us to solve the Erdős-Straus equation (1.3) for wild solutions under the conditions $k \leq 6$. In fact, we solve the equation by the method described in (1.6)-(1.9) and under the condition

$$\mathfrak{S}'_1[\ell(6m+k)], \quad \mathfrak{S}'_2[\ell(6m+k)]. \quad (2.24)$$

In the tame case, the above expression holds as long as we assume that $n = 24m + 1$ is a prime (cf. [26]). Computer shows that it does not hold in general wild case.

3 Wild Solutions with $k \leq 5$

In this section, we present the wild solutions for an integer $n = 24m + 1$ with $k = 1, 2, 3, 4, 5$, where

$$\frac{4}{n} = \frac{1}{6m+k} + \frac{4k-1}{(6m+k+1)(24m+1)}. \quad (3.1)$$

Since the case $k = 6$ is extremely complicated, we will solve it in next section.

3.1 Case $k = 1$

In this case, we consider

$$\begin{aligned} \frac{4}{n} &= \frac{1}{6m+1} + \frac{3}{(6m+1)(24m+1)} \\ &= \frac{1}{6m+1} + \frac{2\ell \times 3}{2\ell \times (6m+1)(24m+1)} \\ &= \frac{1}{6m+1} + \frac{(6\ell-1)+1}{2\ell(6m+1)(24m+1)}, \end{aligned} \quad (3.2)$$

where ℓ is a positive integer. Motivated from (2.5), we assume

$$6m+1 \equiv 0 \pmod{6\ell-1}. \quad (3.3)$$

Equivalently,

$$m+\ell \equiv 0 \pmod{6\ell-1}. \quad (3.4)$$

Thus

$$m \equiv 5\ell-1 \pmod{6\ell-1}. \quad (3.5)$$

When $\ell = 1$, we recover the condition $m \equiv 4 \pmod{5}$ for (2.5). Hence

$$m = 5\ell - 1 + c(6\ell - 1) \quad \text{for some } c \in \mathbb{N}. \quad (3.6)$$

Suppose that m is of the above form. Then

$$6m+1 = 30\ell - 6 + 6c(6\ell-1) + 1 = (6c+5)(6\ell-1). \quad (3.7)$$

According to (3.2),

$$\begin{aligned} \frac{4}{24m+1} &= \frac{1}{6m+1} + \frac{(6\ell-1)+1}{2\ell(6m+1)(24m+1)} \\ &= \frac{1}{6m+1} + \frac{(6\ell-1)+1}{2\ell(6c+5)(6\ell-1)(24m+1)} \\ &= \frac{1}{6m+1} + \frac{1}{2\ell(6c+5)(24m+1)} + \frac{1}{2\ell(6c+5)(6\ell-1)(24m+1)}. \end{aligned} \quad (3.8)$$

Theorem 3.1 *For any positive integer m of the form (3.6), we have the wild solution (3.8) of the Erdős-Straus equation.*

3.2 Case $k = 2$

In this case, we first consider

$$\begin{aligned}\frac{4}{n} &= \frac{1}{6m+2} + \frac{7}{(6m+2)(24m+1)} \\ &= \frac{1}{6m+2} + \frac{2\ell \times 7}{2\ell \times (6m+2)(24m+1)} \\ &= \frac{1}{6m+2} + \frac{(14\ell-1)+1}{4\ell(3m+1)(24m+1)},\end{aligned}\tag{3.9}$$

where ℓ is a positive integer. Motivated from (2.5), we assume

$$3m+1 \equiv 0 \pmod{14\ell-1}.\tag{3.10}$$

It is impossible if $\ell \equiv 2 \pmod{3}$. Assume $\ell = 3r$ with $0 < r \in \mathbb{N}$, Then the above equation becomes

$$3m+1 \equiv 0 \pmod{42r-1}.\tag{3.11}$$

Equivalently,

$$m+14r \equiv 0 \pmod{42r-1}.\tag{3.12}$$

Thus

$$m \equiv 28r-1 \pmod{42r-1}.\tag{3.13}$$

So

$$m = 28r-1 + c(42r-1) \quad \text{for some } c \in \mathbb{N}.\tag{3.14}$$

Suppose that m is of the above form. Then

$$3m+1 = 84r-3 + 3c(42r-1) + 1 = (3c+2)(42r-1).\tag{3.15}$$

According to (3.9),

$$\begin{aligned}\frac{4}{24m+1} &= \frac{1}{6m+2} + \frac{(14\ell-1)+1}{4\ell(3m+1)(24m+1)} \\ &= \frac{1}{6m+2} + \frac{(42r-1)+1}{12r(3c+2)(42r-1)(24m+1)} \\ &= \frac{1}{6m+2} + \frac{1}{12r(3c+2)(24m+1)} + \frac{1}{12r(3c+2)(42r-1)(24m+1)}.\end{aligned}\tag{3.16}$$

When $r = 1$ and $c = 103$, the above equation yields (2.17).

Next we assume $\ell = 3r+1$ with $r \in \mathbb{N}$, Then Equation (3.10) becomes

$$3m+1 \equiv 0 \pmod{42r+13}.\tag{3.17}$$

Equivalently,

$$m \equiv 14r + 4 \pmod{42r + 13}. \quad (3.18)$$

Hence

$$m = 14r + 4 + c(42r + 13) \quad \text{for some } c \in \mathbb{N}. \quad (3.19)$$

If m is of the above form, then

$$3m + 1 = 42r + 12 + 3c(42r + 13) + 1 = (3c + 1)(42r + 13). \quad (3.20)$$

By (3.9),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 2} + \frac{(14\ell - 1) + 1}{4\ell(3m + 1)(24m + 1)} \\ &= \frac{1}{6m + 2} + \frac{(42r + 13) + 1}{4(3r + 1)(3c + 1)(42r + 13)(24m + 1)} \\ &= \frac{1}{6m + 2} + \frac{1}{4(3r + 1)(3c + 1)(24m + 1)} \\ &\quad + \frac{1}{4(3r + 1)(3c + 1)(42r + 13)(24m + 1)}. \end{aligned} \quad (3.21)$$

Now we consider

$$\begin{aligned} \frac{4}{n} &= \frac{1}{6m + 2} + \frac{7}{(6m + 2)(24m + 1)} \\ &= \frac{1}{6m + 2} + \frac{(2\ell + 1) \times 7}{(2\ell + 1) \times (6m + 2)(24m + 1)} \\ &= \frac{1}{6m + 2} + \frac{(14\ell + 6) + 1}{2(2\ell + 1)(3m + 1)(24m + 1)}, \end{aligned} \quad (3.22)$$

where ℓ is a positive integer. Motivated from (2.5), we assume

$$3m + 1 \equiv 0 \pmod{7\ell + 3}. \quad (3.23)$$

It is impossible if $\ell \equiv 0 \pmod{3}$. Assume $\ell = 3r + 1$ with $r \in \mathbb{N}$, Then the above equation becomes

$$3m + 1 \equiv 0 \pmod{21r + 10}. \quad (3.24)$$

Equivalently,

$$m \equiv 7r + 3 \pmod{21r + 10}. \quad (3.25)$$

Therefore,

$$m = 7r + 3 + c(21r + 10) \quad \text{for some } c \in \mathbb{N}. \quad (3.26)$$

When m is of the above form,

$$3m + 1 = 21r + 9 + 3c(21r + 10) + 1 = (3c + 1)(21r + 10). \quad (3.27)$$

By (3.22),

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+2} + \frac{(14\ell+6)+1}{2(2\ell+1)(3m+1)(24m+1)} \\
&= \frac{1}{6m+2} + \frac{2(21r+10)+1}{6(2r+1)(3c+1)(21r+10)(24m+1)} \\
&= \frac{1}{6m+2} + \frac{1}{3(2r+1)(3c+1)(24m+1)} \\
&\quad + \frac{1}{6(2r+1)(3c+1)(21r+10)(24m+1)}.
\end{aligned} \tag{3.28}$$

Next we assume $\ell = 3r + 2$ with $r \in \mathbb{N}$, Then Equation (3.23) becomes

$$3m + 1 \equiv 0 \pmod{21r + 17}. \tag{3.29}$$

Equivalently,

$$m + 7r + 6 \equiv 0 \pmod{21r + 17}; \tag{3.30}$$

that is

$$m \equiv 14r + 11 \pmod{21r + 17}; \tag{3.31}$$

Hence

$$m = 14r + 11 + c(21r + 17) \quad \text{for some } c \in \mathbb{N}. \tag{3.32}$$

Suppose that m is of the above form. Then

$$3m + 1 = 42r + 33 + 3c(21r + 17) + 1 = (3c + 2)(21r + 17). \tag{3.33}$$

By (3.22),

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+2} + \frac{(14\ell+6)+1}{2(2\ell+1)(3m+1)(24m+1)} \\
&= \frac{1}{6m+2} + \frac{2(21r+17)+1}{2(6r+5)(3c+2)(21r+17)(24m+1)} \\
&= \frac{1}{6m+2} + \frac{1}{(6r+5)(3c+2)(24m+1)} \\
&\quad + \frac{1}{2(6r+5)(3c+2)(21r+17)(24m+1)}.
\end{aligned} \tag{3.34}$$

When $r = 0$, the above equation becomes (2.12).

Finally we consider

$$\begin{aligned}
\frac{4}{n} &= \frac{1}{6m+2} + \frac{7}{(6m+2)(24m+1)} \\
&= \frac{1}{6m+2} + \frac{(2\ell+1) \times 7}{(2\ell+1) \times (6m+2)(24m+1)} \\
&= \frac{1}{6m+2} + \frac{(14\ell+5)+2}{2(2\ell+1)(3m+1)(24m+1)},
\end{aligned} \tag{3.35}$$

where ℓ is a positive integer. Motivated from (2.15), we assume

$$3m + 1 \equiv 0 \pmod{14\ell + 5}. \quad (3.36)$$

It is impossible if $\ell \equiv 2 \pmod{3}$. Assume $\ell = 3r$ with $0 < r \in \mathbb{N}$, Then the above equation becomes

$$3m + 1 \equiv 0 \pmod{42r + 5}. \quad (3.37)$$

Equivalently,

$$m + 14r + 2 \equiv 0 \pmod{42r + 5}; \quad (3.38)$$

that is

$$m \equiv 28r + 3 \pmod{42r + 5}; \quad (3.39)$$

Thus

$$m = 28r + 3 + c(42r + 5) \quad \text{for some } c \in \mathbb{N}. \quad (3.40)$$

If m is of the above form, then

$$3m + 1 = 84r + 9 + 3c(42r + 5) + 1 = (3c + 2)(42r + 5). \quad (3.41)$$

By (3.35),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 2} + \frac{(14\ell + 5) + 2}{2(2\ell + 1)(3m + 1)(24m + 1)} \\ &= \frac{1}{6m + 2} + \frac{(42r + 5) + 2}{2(6r + 1)(3c + 2)(42r + 5)(24m + 1)} \\ &= \frac{1}{6m + 2} + \frac{1}{2(6r + 1)(3c + 2)(24m + 1)} \\ &\quad + \frac{1}{(6r + 1)(3c + 2)(42r + 5)(24m + 1)}. \end{aligned} \quad (3.42)$$

Next we assume $\ell = 3r + 1$ with $r \in \mathbb{N}$, Then Equation (3.36) becomes

$$3m + 1 \equiv 0 \pmod{42r + 19}. \quad (3.43)$$

Equivalently,

$$m \equiv 14r + 6 \pmod{42r + 19}. \quad (3.44)$$

In this way,

$$m = 14r + 6 + c(42r + 19) \quad \text{for some } c \in \mathbb{N}. \quad (3.45)$$

When m is of the above form,

$$3m + 1 = 42r + 18 + 3c(42r + 19) + 1 = (3c + 1)(42r + 19). \quad (3.46)$$

According to (3.35),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 2} + \frac{(14\ell + 5) + 2}{2(2\ell + 1)(3m + 1)(24m + 1)} \\ &= \frac{1}{6m + 2} + \frac{(42r + 19) + 2}{6(2r + 1)(3c + 1)(42r + 19)(24m + 1)} \\ &= \frac{1}{6m + 2} + \frac{1}{6(2r + 1)(3c + 1)(24m + 1)} \\ &\quad + \frac{1}{3(2r + 1)(3c + 1)(42r + 19)(24m + 1)}. \end{aligned} \quad (3.47)$$

Theorem 3.2 *For any positive integer m of the form (3.14), we have the wild solution (3.16) of the Erdős-Straus equation. If m is of the form (3.19), we have the wild solution (3.21) of the Erdős-Straus equation. When m is of the form (3.26), we have the wild solution (3.28) of the Erdős-Straus equation. Suppose that m is of the form (3.32), we have the wild solution (3.34) of the Erdős-Straus equation. Let m be of the form (3.40), we have the wild solution (3.42) of the Erdős-Straus equation. Assume that m is of the form (3.45), we have the wild solution (3.47) of the Erdős-Straus equation.*

3.3 Case $k = 3$

In this case, we first consider

$$\begin{aligned}\frac{4}{n} &= \frac{1}{6m+3} + \frac{11}{(6m+3)(24m+1)} \\ &= \frac{1}{6m+3} + \frac{2\ell \times 11}{6\ell \times (2m+1)(24m+1)} \\ &= \frac{1}{6m+3} + \frac{(22\ell-1)+1}{6\ell(2m+1)(24m+1)},\end{aligned}\tag{3.48}$$

where ℓ is a positive integer. Suppose $\ell \equiv 1 \pmod{3}$; that is, $\ell = 3r + 1$ for some $r \in \mathbb{N}$. Then

$$22\ell - 1 = 66r + 21 = 3(22r + 7).\tag{3.49}$$

By (2.16), we assume

$$2m + 1 \equiv 0 \pmod{22r + 7}.\tag{3.50}$$

Equivalently,

$$m \equiv 11r + 3 \pmod{22r + 7}.\tag{3.51}$$

Thus

$$m = 11r + 3 + c(22r + 7) \quad \text{for some } c \in \mathbb{N}.\tag{3.52}$$

Assume that m is of the above form. Then

$$2m + 1 = 22r + 6 + 2c(22r + 7) + 1 = (2c + 1)(22r + 7).\tag{3.53}$$

According to (3.48),

$$\begin{aligned}\frac{4}{24m+1} &= \frac{1}{6m+3} + \frac{(22\ell-1)+1}{6\ell(2m+1)(24m+1)} \\ &= \frac{1}{6m+3} + \frac{3(22r+7)+1}{6(3r+1)(2c+1)(22r+7)(24m+1)} \\ &= \frac{1}{6m+3} + \frac{1}{2(3r+1)(2c+1)(24m+1)} \\ &\quad + \frac{1}{6(3r+1)(2c+1)(22r+7)(24m+1)}.\end{aligned}\tag{3.54}$$

When $r = 0$, the above equation yields (2.8). A subtle point is that we have well used the factor “3” in $6m + 3$ of the denominator.

Next we assume $\ell \equiv 0 \pmod{3}$; that is, $\ell = 3r$ for some $0 < r \in \mathbb{N}$. Then

$$22\ell - 1 = 66r - 1. \quad (3.55)$$

Motivated from (2.103), we impose

$$2m + 1 \equiv 0 \pmod{66r - 1}. \quad (3.56)$$

Equivalently,

$$m \equiv 33r - 1 \pmod{66r - 1}. \quad (3.57)$$

Thus

$$m = 33r - 1 + c(66r - 1) \quad \text{for some } c \in \mathbb{N}. \quad (3.58)$$

When m is of the above form,

$$2m + 1 = 66r - 2 + 2c(66r - 1) + 1 = (2c + 1)(66r - 1). \quad (3.59)$$

By (3.48),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 3} + \frac{(22\ell - 1) + 1}{6\ell(2m + 1)(24m + 1)} \\ &= \frac{1}{6m + 3} + \frac{(66r - 1) + 1}{18r(2c + 1)(66r - 1)(24m + 1)} \\ &= \frac{1}{6m + 3} + \frac{1}{18r(2c + 1)(24m + 1)} \\ &\quad + \frac{1}{18r(2c + 1)(66r - 1)(24m + 1)}. \end{aligned} \quad (3.60)$$

When $r = 1$ and $c = 53$, the above equation yields (2.13).

Thirdly we assume $\ell \equiv 2 \pmod{3}$; that is, $\ell = 3r + 2$ for some $r \in \mathbb{N}$. Then

$$22\ell - 1 = 66r + 43. \quad (3.61)$$

Motivated from (2.13), we impose

$$2m + 1 \equiv 0 \pmod{66r + 43}. \quad (3.62)$$

Equivalently,

$$m \equiv 33r + 21 \pmod{66r + 43}. \quad (3.63)$$

Thus

$$m = 33r + 21 + c(66r + 43) \quad \text{for some } c \in \mathbb{N}. \quad (3.64)$$

If m is of the above form,

$$2m + 1 = 66r + 42 + 2c(66r + 43) + 1 = (2c + 1)(66r + 43). \quad (3.65)$$

Expression (3.48) yields

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 3} + \frac{(22\ell - 1) + 1}{6\ell(2m + 1)(24m + 1)} \\ &= \frac{1}{6m + 3} + \frac{(66r + 43) + 1}{6(3r + 2)(2c + 1)(66r + 43)(24m + 1)} \\ &= \frac{1}{6m + 3} + \frac{1}{6(3r + 2)(2c + 1)(24m + 1)} \\ &\quad + \frac{1}{6(3r + 2)(2c + 1)(66r + 43)(24m + 1)}. \end{aligned} \quad (3.66)$$

Now we consider

$$\begin{aligned}
\frac{4}{n} &= \frac{1}{6m+3} + \frac{11}{(6m+3)(24m+1)} \\
&= \frac{1}{6m+3} + \frac{2\ell \times 11}{6\ell \times (2m+1)(24m+1)} \\
&= \frac{1}{6m+3} + \frac{(22\ell-3)+3}{6\ell(2m+1)(24m+1)},
\end{aligned} \tag{3.67}$$

where ℓ is a positive integer. According to (1.8) with $\mathfrak{S}'_1 = 22\ell - 3$ and $\mathfrak{S}'_2 = 3$, $\ell \not\equiv 0 \pmod{3}$. Suppose $\ell \equiv 1 \pmod{3}$; that is, $\ell = 3r + 1$ for some $0 < r \in \mathbb{N}$. Then

$$22\ell - 3 = 66r + 19. \tag{3.68}$$

Motivated from (2.8), we assume

$$2m + 1 \equiv 0 \pmod{66r + 19}. \tag{3.69}$$

Equivalently,

$$m \equiv 33r + 9 \pmod{66r + 19}. \tag{3.70}$$

Thus

$$m = 33r + 9 + c(66r + 19) \quad \text{for some } c \in \mathbb{N}. \tag{3.71}$$

Assume that m is of the above form. Then

$$2m + 1 = 66r + 18 + 2c(66r + 19) + 1 = (2c + 1)(66r + 19). \tag{3.72}$$

According to (3.67),

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+3} + \frac{(22\ell-3)+3}{6\ell(2m+1)(24m+1)} \\
&= \frac{1}{6m+3} + \frac{(66r+19)+3}{6(3r+1)(2c+1)(66r+19)(24m+1)} \\
&= \frac{1}{6m+3} + \frac{1}{6(3r+1)(2c+1)(24m+1)} \\
&\quad + \frac{1}{2(3r+1)(2c+1)(66r+19)(24m+1)}.
\end{aligned} \tag{3.73}$$

Secondly we suppose $\ell \equiv 2 \pmod{3}$; that is, $\ell = 3r + 2$ for some $0 < r \in \mathbb{N}$. Then

$$22\ell - 3 = 66r + 41. \tag{3.74}$$

Motivated from (2.8), we assume

$$2m + 1 \equiv 0 \pmod{66r + 41}. \tag{3.75}$$

Equivalently,

$$m \equiv 33r + 20 \pmod{66r + 41}. \tag{3.76}$$

Thus

$$m = 33r + 20 + c(66r + 41) \quad \text{for some } c \in \mathbb{N}. \quad (3.77)$$

Let m be of the above form. Then

$$2m + 1 = 66r + 40 + 2c(66r + 41) + 1 = (2c + 1)(66r + 41). \quad (3.78)$$

By (3.67),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 3} + \frac{(22\ell - 3) + 3}{6\ell(2m + 1)(24m + 1)} \\ &= \frac{1}{6m + 3} + \frac{(66r + 41) + 3}{6(3r + 1)(2c + 1)(66r + 41)(24m + 1)} \\ &= \frac{1}{6m + 3} + \frac{1}{6(3r + 1)(2c + 1)(24m + 1)} \\ &\quad + \frac{1}{2(3r + 1)(2c + 1)(66r + 41)(24m + 1)}. \end{aligned} \quad (3.79)$$

Theorem 3.3 *For any positive integer m of the form (3.52), we have the wild solution (3.54) of the Erdős-Straus equation. If m is of the form (3.58), we have the wild solution (3.60) of the Erdős-Straus equation. When m is of the form (3.64), we have the wild solution (3.66) of the Erdős-Straus equation. Suppose that m is of the form (3.71), we have the wild solution (3.73) of the Erdős-Straus equation. Let m be of the form (3.77), we have the wild solution (3.79) of the Erdős-Straus equation.*

3.4 Case $k = 4$

In this case, we first consider

$$\begin{aligned} \frac{4}{n} &= \frac{1}{6m + 4} + \frac{15}{(6m + 4)(24m + 1)} \\ &= \frac{1}{6m + 4} + \frac{2\ell \times 15}{2\ell \times 2(3m + 2)(24m + 1)} \\ &= \frac{1}{6m + 4} + \frac{(30\ell - 1) + 1}{4\ell(3m + 2)(24m + 1)}, \end{aligned} \quad (3.80)$$

where ℓ is a positive integer. Motivated from (2.8), we assume

$$3m + 2 \equiv 0 \pmod{30\ell - 1}. \quad (3.81)$$

Equivalently,

$$m \equiv 10\ell - 1 \pmod{30\ell - 1}. \quad (3.82)$$

Thus

$$m = 10\ell - 1 + c(30\ell - 1) \quad \text{for some } c \in \mathbb{N}. \quad (3.83)$$

Assume that m is of the above form. Then

$$3m + 2 = 30\ell - 3 + 3c(30\ell - 1) + 2 = (3c + 1)(30\ell - 1). \quad (3.84)$$

According to (3.80),

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+4} + \frac{(30\ell-1)+1}{4\ell(3m+2)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{(30\ell-1)+1}{4\ell(3c+1)(30\ell-1)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{1}{4\ell(3c+1)(24m+1)} \\
&\quad + \frac{1}{4\ell(3c+1)(30\ell-1)(24m+1)}.
\end{aligned} \tag{3.85}$$

Taking $\ell = 1$, we get (2.19).

Next we consider

$$\begin{aligned}
\frac{4}{n} &= \frac{1}{6m+4} + \frac{15}{(6m+4)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{(2\ell+1) \times 15}{(2\ell+1) \times 2(3m+2)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{(30\ell+14)+1}{2(2\ell+1)(3m+2)(24m+1)},
\end{aligned} \tag{3.86}$$

where ℓ is a positive integer. Motivated from (2.8), we assume

$$3m+2 \equiv 0 \pmod{15\ell+7}. \tag{3.87}$$

Equivalently,

$$m \equiv 10\ell+4 \pmod{15\ell+7}. \tag{3.88}$$

Thus

$$m = 10\ell+4 + c(15\ell+7) \quad \text{for some } c \in \mathbb{N}. \tag{3.89}$$

When m is of the above form,

$$3m+2 = 30\ell+12 + 3c(15\ell+7) + 2 = (3c+2)(15\ell+7). \tag{3.90}$$

By (3.86),

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+4} + \frac{(30\ell+14)+1}{2(2\ell+1)(3m+2)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{2(15\ell+7)+1}{2(2\ell+1)(3c+2)(15\ell+7)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{1}{(2\ell+1)(3c+2)(24m+1)} \\
&\quad + \frac{1}{2(2\ell+1)(3c+2)(15\ell+7)(24m+1)}.
\end{aligned} \tag{3.91}$$

Finally we consider

$$\begin{aligned}
\frac{4}{n} &= \frac{1}{6m+4} + \frac{15}{(6m+4)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{(2\ell+1) \times 15}{(2\ell+1) \times 2(3m+2)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{(30\ell+13)+2}{2(2\ell+1)(3m+2)(24m+1)},
\end{aligned} \tag{3.92}$$

where ℓ is a positive integer. Motivated from (2.15), we assume

$$3m+2 \equiv 0 \pmod{30\ell+13}. \tag{3.93}$$

Equivalently,

$$m \equiv 20\ell+8 \pmod{30\ell+13}. \tag{3.94}$$

Thus

$$m = 20\ell+8 + c(30\ell+13) \quad \text{for some } c \in \mathbb{N}. \tag{3.95}$$

If m is of the above form,

$$3m+2 = 60\ell+24 + 3c(30\ell+13) + 2 = (3c+2)(30\ell+13). \tag{3.96}$$

Expression (3.92) gives

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+4} + \frac{(30\ell+13)+2}{2(2\ell+1)(3m+2)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{(30\ell+13)+2}{2(2\ell+1)(3c+2)(30\ell+13)(24m+1)} \\
&= \frac{1}{6m+4} + \frac{1}{2(2\ell+1)(3c+2)(24m+1)} \\
&\quad + \frac{1}{(2\ell+1)(3c+2)(30\ell+13)(24m+1)}.
\end{aligned} \tag{3.97}$$

Theorem 3.4 *For any positive integer m of the form (3.83), we have the wild solution (3.85) of the Erdős-Straus equation. If m is of the form (3.89), we have the wild solution (3.91) of the Erdős-Straus equation. When m is of the form (3.95), we have the wild solution (3.97) of the Erdős-Straus equation.*

3.5 Case $k = 5$

In this case, we consider

$$\begin{aligned}
\frac{4}{n} &= \frac{1}{6m+5} + \frac{19}{(6m+5)(24m+1)} \\
&= \frac{1}{6m+5} + \frac{2\ell \times 19}{2\ell \times (6m+5)(24m+1)} \\
&= \frac{1}{6m+5} + \frac{(38\ell-1)+1}{2\ell(6m+5)(24m+1)},
\end{aligned} \tag{3.98}$$

where ℓ is a positive integer. As the earlier cases, we assume

$$6m + 5 \equiv 0 \pmod{38\ell - 1}. \quad (3.99)$$

It is impossible if $\ell \equiv 2 \pmod{3}$. Assume $\ell = 3r$ with $0 < r \in \mathbb{N}$, Then the above equation becomes

$$6m + 5 \equiv 0 \pmod{114r - 1}. \quad (3.100)$$

Equivalently,

$$m + 95r \equiv 0 \pmod{114r - 1}. \quad (3.101)$$

Thus

$$m \equiv 19r - 1 \pmod{114r - 1}. \quad (3.102)$$

So

$$m = 19r - 1 + c(114r - 1) \quad \text{for some } c \in \mathbb{N}. \quad (3.103)$$

When m is of the above form,

$$6m + 5 = 114r - 6 + 6c(114r - 1) + 5 = (6c + 1)(114r - 1). \quad (3.104)$$

According to (3.98),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 5} + \frac{(38\ell - 1) + 1}{2\ell(6m + 5)(24m + 1)} \\ &= \frac{1}{6m + 5} + \frac{(114r - 1) + 1}{6r(6c + 1)(114r - 1)(24m + 1)} \\ &= \frac{1}{6m + 5} + \frac{1}{6r(6c + 1)(24m + 1)} \\ &\quad + \frac{1}{6r(6c + 1)(114r - 1)(24m + 1)}. \end{aligned} \quad (3.105)$$

Suppose $\ell \equiv 1 \pmod{3}$; that is, $\ell = 3r + 1$ for some $r \in \mathbb{N}$. Then

$$38\ell - 1 = 114r + 37. \quad (3.106)$$

Equation (3.99) becomes

$$6m + 5 \equiv 0 \pmod{114r + 37}. \quad (3.107)$$

Equivalently,

$$m \equiv 95r + 30 \pmod{114r + 37}. \quad (3.108)$$

So

$$m = 95r + 30 + c(114r + 37) \quad \text{for some } c \in \mathbb{N}. \quad (3.109)$$

Let m be of the above form,

$$6m + 5 = 570r + 180 + 6c(114r + 37) + 5 = (6c + 5)(114r + 37). \quad (3.110)$$

According to (3.98),

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+5} + \frac{(38\ell-1)+1}{2\ell(6m+5)(24m+1)} \\
&= \frac{1}{6m+5} + \frac{(114r+37)+1}{2(3r+1)(6c+5)(114r+37)(24m+1)} \\
&= \frac{1}{6m+5} + \frac{1}{2(3r+1)(6c+5)(24m+1)} \\
&\quad + \frac{1}{2(3r+1)(6c+5)(114r+37)(24m+1)}. \tag{3.111}
\end{aligned}$$

When $r = 0$ and $c = 10$, the above equation gives (2.15).

Theorem 3.5 *For any positive integer m of the form (3.103), we have the wild solution (3.105) of the Erdős-Straus equation. If m is of the form (3.109), we have the wild solution (3.111) of the Erdős-Straus equation.*

4 Wild Solutions with $k = 6$

In this section, we solve the Erös-Straus equation for $n = 24m + 1$ with $k = 6$ in (3.1). We divide it into the cases of even multiplier and odd multiplier, respectively.

4.1 Case of even multiplier

In this subcase, we first consider

$$\begin{aligned}
\frac{4}{n} &= \frac{1}{6m+6} + \frac{23}{(6m+6)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{2\ell \times 23}{2\ell \times 6(m+1)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{(46\ell-1)+1}{12\ell(m+1)(24m+1)}, \tag{4.1}
\end{aligned}$$

where ℓ is a positive integer. Suppose $\ell \equiv 1 \pmod{3}$; that is, $\ell = 3r + 1$ for some $r \in \mathbb{N}$. Then

$$46\ell - 1 = 138r + 45 = 3(46r + 15). \tag{4.2}$$

According to (2.16), we impose

$$m + 1 \equiv 0 \pmod{46r + 15}. \tag{4.3}$$

Equivalently,

$$m \equiv 46r + 14 \pmod{46r + 15}. \tag{4.4}$$

Thus

$$m = 46r + 14 + c(46r + 15) \quad \text{for some } c \in \mathbb{N}. \tag{4.5}$$

Assume that m is of the above form. Then

$$m + 1 = 46r + 14 + c(46r + 15) + 1 = (c + 1)(46r + 15). \tag{4.6}$$

According to (4.1),

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+6} + \frac{(46\ell-1)+1}{12\ell(m+1)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{3(46r+15)+1}{12(3r+1)(c+1)(46r+15)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{1}{4(3r+1)(c+1)(24m+1)} \\
&\quad + \frac{1}{12(3r+1)(c+1)(46r+15)(24m+1)}. \tag{4.7}
\end{aligned}$$

A subtle point is that we have well used the factor “3” in $6m+6$ of the denominator.

Next we assume $\ell \equiv 0 \pmod{3}$; that is, $\ell = 3r$ for some $0 < r \in \mathbb{N}$. Then

$$46\ell - 1 = 138r - 1. \tag{4.8}$$

We assume

$$m+1 \equiv 0 \pmod{138r-1}. \tag{4.9}$$

Equivalently,

$$m \equiv 138r - 2 \pmod{138r-1}. \tag{4.10}$$

Thus

$$m = 138r - 2 + c(138r - 1) \quad \text{for some } c \in \mathbb{N}. \tag{4.11}$$

When m is of the above form,

$$m+1 = 138r - 2 + c(138r - 1) + 1 = (c+1)(138r - 1). \tag{4.12}$$

By (4.1),

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+6} + \frac{(46\ell-1)+1}{12\ell(m+1)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{(138r-1)+1}{36r(c+1)(138r-1)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{1}{36r(c+1)(24m+1)} \\
&\quad + \frac{1}{36r(c+1)(138r-1)(24m+1)}. \tag{4.13}
\end{aligned}$$

Thirdly we assume $\ell \equiv 2 \pmod{3}$; that is, $\ell = 3r + 2$ for some $r \in \mathbb{N}$. Then

$$46\ell - 1 = 138r + 91. \tag{4.14}$$

We impose

$$m+1 \equiv 0 \pmod{138r+91}. \tag{4.15}$$

Equivalently,

$$m \equiv 138r + 90 \pmod{138r+91}. \tag{4.16}$$

Thus

$$m = 138r + 90 + c(138r + 91) \quad \text{for some } c \in \mathbb{N}. \quad (4.17)$$

If m is of the above form, then

$$m + 1 = 138r + 90 + c(138r + 91) + 1 = (c + 1)(138r + 91). \quad (4.18)$$

Expression (4.1) yields

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 6} + \frac{(46\ell - 1) + 1}{12\ell(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(138r + 91) + 1}{12(3r + 2)(c + 1)(138r + 91)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{12(3r + 2)(c + 1)(24m + 1)} \\ &\quad + \frac{1}{12(3r + 2)(c + 1)(138r + 91)(24m + 1)}. \end{aligned} \quad (4.19)$$

Now we consider

$$\begin{aligned} \frac{4}{n} &= \frac{1}{6m + 6} + \frac{23}{(6m + 6)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{2\ell \times 23}{2\ell \times 6(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(46\ell - 3) + 3}{12\ell(m + 1)(24m + 1)}, \end{aligned} \quad (4.20)$$

where ℓ is a positive integer. According to (1.8) with $\mathfrak{S}'_1 = 46\ell - 3$ and $\mathfrak{S}'_2 = 3$, $\ell \not\equiv 0 \pmod{3}$. Suppose $\ell \equiv 1 \pmod{3}$; that is, $\ell = 3r + 1$ for some $r \in \mathbb{N}$. Then

$$46\ell - 3 = 138r + 43. \quad (4.21)$$

We assume

$$m + 1 \equiv 0 \pmod{138r + 43}. \quad (4.22)$$

Equivalently,

$$m \equiv 138r + 42 \pmod{138r + 43}. \quad (4.23)$$

Thus

$$m = 138r + 42 + c(138r + 43) \quad \text{for some } c \in \mathbb{N}. \quad (4.24)$$

When m is of the above form,

$$m + 1 = 138r + 42 + c(138r + 43) + 1 = (c + 1)(138r + 43). \quad (4.25)$$

Expression (4.20) gives

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 6} + \frac{(46\ell - 3) + 3}{12\ell(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(138r + 43) + 3}{12(3r + 1)(c + 1)(138r + 43)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{12(3r + 1)(c + 1)(24m + 1)} \\ &\quad + \frac{1}{4(3r + 1)(c + 1)(138r + 43)(24m + 1)}. \end{aligned} \quad (4.26)$$

We assume $\ell \equiv 2 \pmod{3}$; that is, $\ell = 3r + 2$ for some $r \in \mathbb{N}$. Then

$$46\ell - 3 = 138r + 89. \quad (4.27)$$

We impose

$$m + 1 \equiv 0 \pmod{138r + 89}. \quad (4.28)$$

Equivalently,

$$m \equiv 138r + 88 \pmod{138r + 89}. \quad (4.29)$$

Thus

$$m = 138r + 89 + c(138r + 89) \quad \text{for some } c \in \mathbb{N}. \quad (4.30)$$

If m is of the above form, then

$$m + 1 = 138r + 88 + c(138r + 89) + 1 = (c + 1)(138r + 89). \quad (4.31)$$

Expression (4.20) yields

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 6} + \frac{(46\ell - 3) + 3}{12\ell(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(138r + 89) + 3}{12(3r + 2)(c + 1)(138r + 89)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{12(3r + 2)(c + 1)(24m + 1)} \\ &\quad + \frac{1}{4(3r + 2)(c + 1)(138r + 89)(24m + 1)}. \end{aligned} \quad (4.32)$$

Theorem 4.1 *For any positive integer m of the form (4.5), we have the wild solution (4.7) of the Erdős-Straus equation. If m is of the form (4.11), we have the wild solution (4.13) of the Erdős-Straus equation. When m is of the form (4.17), we have the wild solution (4.19) of the Erdős-Straus equation. Suppose that m is of the form (4.24), and then we have the wild solution (4.26) of the Erdős-Straus equation. Let m be of the form (4.30), and then we have the wild solution (4.32) of the Erdős-Straus equation.*

4.2 Case of odd multiplier

In this case, we first consider

$$\begin{aligned} \frac{4}{n} &= \frac{1}{6m + 6} + \frac{23}{(6m + 6)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(2\ell + 1) \times 23}{(2\ell + 1) \times 6(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{2(23\ell + 11) + 1}{6(2\ell + 1)(m + 1)(24m + 1)}, \end{aligned} \quad (4.33)$$

where ℓ is a positive integer. Suppose $\ell \equiv 2 \pmod{3}$; that is, $\ell = 3r + 2$ for some $r \in \mathbb{N}$. Then

$$23\ell + 11 = 69r + 57 = 3(23r + 19). \quad (4.34)$$

By (2.16), we impose

$$m + 1 \equiv 0 \pmod{23r + 19}. \quad (4.35)$$

Equivalently,

$$m \equiv 23r + 18 \pmod{23r + 19}. \quad (4.36)$$

Thus

$$m = 23r + 18 + c(23r + 19) \quad \text{for some } c \in \mathbb{N}. \quad (4.37)$$

Assume that m is of the above form. Then

$$m + 1 = 23r + 18 + c(23r + 19) + 1 = (c + 1)(23r + 19). \quad (4.38)$$

According to (4.33),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 6} + \frac{2(23\ell + 11) + 1}{6(2\ell + 1)(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{6(23r + 19) + 1}{6(6r + 5)(c + 1)(23r + 19)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{(6r + 5)(c + 1)(24m + 1)} \\ &\quad + \frac{1}{6(6r + 5)(c + 1)(23r + 19)(24m + 1)}. \end{aligned} \quad (4.39)$$

A subtle point is that we have well used the factor “6” in $6m + 6$ of the denominator.

Next we assume $\ell \equiv 0 \pmod{3}$; that is, $\ell = 3r$ for some $0 < r \in \mathbb{N}$. Then

$$23\ell + 11 = 69r + 11. \quad (4.40)$$

We assume

$$m + 1 \equiv 0 \pmod{69r + 11}. \quad (4.41)$$

Equivalently,

$$m \equiv 69r + 10 \pmod{69r + 11}. \quad (4.42)$$

Thus

$$m = 69r + 10 + c(69r + 11) \quad \text{for some } c \in \mathbb{N}. \quad (4.43)$$

When m is of the above form,

$$m + 1 = 69r + 10 + c(69r + 11) + 1 = (c + 1)(69r + 11). \quad (4.44)$$

By (4.33),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 6} + \frac{2(23\ell + 11) + 1}{6(2\ell + 1)(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{2(69r + 11) + 1}{6(6r + 1)(c + 1)(69r + 11)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{3(6r + 1)(c + 1)(24m + 1)} \\ &\quad + \frac{1}{6(6r + 1)(c + 1)(69r + 11)(24m + 1)}. \end{aligned} \quad (4.45)$$

Let $\ell \equiv 1 \pmod{3}$; that is, $\ell = 3r + 1$ for some $0 < r \in \mathbb{N}$. Then

$$23\ell + 11 = 69r + 34. \quad (4.46)$$

We assume

$$m + 1 \equiv 0 \pmod{69r + 34}. \quad (4.47)$$

Equivalently,

$$m \equiv 69r + 33 \pmod{69r + 34}. \quad (4.48)$$

Thus

$$m = 69r + 33 + c(69r + 34) \quad \text{for some } c \in \mathbb{N}. \quad (4.49)$$

If m be of the above form, then

$$m + 1 = 69r + 33 + c(69r + 34) + 1 = (c + 1)(69r + 34). \quad (4.50)$$

By (4.43),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 6} + \frac{2(23\ell + 11) + 1}{6(2\ell + 1)(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{2(69r + 34) + 1}{18(2r + 1)(c + 1)(69r + 34)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{9(2r + 1)(c + 1)(24m + 1)} \\ &\quad + \frac{1}{18(2r + 1)(c + 1)(69r + 34)(24m + 1)}. \end{aligned} \quad (4.51)$$

Furthermore, we consider

$$\begin{aligned} \frac{4}{n} &= \frac{1}{6m + 6} + \frac{23}{(6m + 6)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(2\ell + 1) \times 23}{(2\ell + 1) \times 6(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(46\ell + 21) + 2}{6(2\ell + 1)(m + 1)(24m + 1)}, \end{aligned} \quad (4.52)$$

where ℓ is a positive integer. Suppose $\ell \equiv 0 \pmod{3}$; that is, $\ell = 3r$ for some $0 < r \in \mathbb{N}$. Then

$$46\ell + 21 = 3(46r + 7). \quad (4.53)$$

By (2.16), we impose

$$m + 1 \equiv 0 \pmod{46r + 7}. \quad (4.54)$$

Equivalently,

$$m \equiv 46r + 6 \pmod{46r + 7}. \quad (4.55)$$

Thus

$$m = 46r + 6 + c(46r + 7) \quad \text{for some } c \in \mathbb{N}. \quad (4.56)$$

Assume that m is of the above form. Then

$$m + 1 = 46r + 7 + c(46r + 7) + 1 = (c + 1)(46r + 7). \quad (4.57)$$

According to (4.52),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 6} + \frac{(46\ell + 21) + 2}{6(2\ell + 1)(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{3(46r + 7) + 2}{6(6r + 1)(c + 1)(46r + 7)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{2(6r + 1)(c + 1)(24m + 1)} \\ &\quad + \frac{1}{3(6r + 1)(c + 1)(46r + 7)(24m + 1)}. \end{aligned} \quad (4.58)$$

A subtle point is that we have well used the factor “3” in $6m + 6$ of the denominator.

Next we assume $\ell \equiv 1 \pmod{3}$; that is, $\ell = 3r + 1$ for some $r \in \mathbb{N}$. Then

$$46\ell + 21 = 138r + 67. \quad (4.59)$$

So we impose

$$m + 1 \equiv 0 \pmod{138r + 67}. \quad (4.60)$$

Equivalently,

$$m \equiv 138r + 66 \pmod{138r + 67}. \quad (4.61)$$

Thus

$$m = 138r + 66 + c(138r + 67) \quad \text{for some } c \in \mathbb{N}. \quad (4.62)$$

Let m be of the above form. Then

$$m + 1 = 138r + 66 + c(138r + 67) + 1 = (c + 1)(138r + 67). \quad (4.63)$$

By (4.52),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 6} + \frac{(46\ell + 21) + 2}{6(2\ell + 1)(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(138r + 67) + 2}{18(2r + 1)(c + 1)(138r + 67)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{18(2r + 1)(c + 1)(24m + 1)} \\ &\quad + \frac{1}{9(2r + 1)(c + 1)(138r + 67)(24m + 1)}. \end{aligned} \quad (4.64)$$

When $r = 0$ and $c = 442$, the above equation becomes (2.18).

Let $\ell \equiv 2 \pmod{3}$; that is, $\ell = 3r + 2$ for some $r \in \mathbb{N}$. Then

$$46\ell + 21 = 138r + 113. \quad (4.65)$$

So we impose

$$m + 1 \equiv 0 \pmod{138r + 113}. \quad (4.66)$$

Equivalently,

$$m \equiv 138r + 112 \pmod{138r + 113}. \quad (4.67)$$

Hence

$$m = 138r + 112 + c(138r + 113) \quad \text{for some } c \in \mathbb{N}. \quad (4.68)$$

Assume that m is of the above form. Then

$$m + 1 = 138r + 112 + c(138r + 113) + 1 = (c + 1)(138r + 113). \quad (4.69)$$

Expression (4.52) yields

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 6} + \frac{(46\ell + 21) + 2}{6(2\ell + 1)(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(138r + 113) + 2}{6(6r + 5)(c + 1)(138r + 113)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{6(6r + 5)(c + 1)(24m + 1)} \\ &\quad + \frac{1}{3(6r + 5)(c + 1)(138r + 113)(24m + 1)}. \end{aligned} \quad (4.70)$$

In addition, we consider

$$\begin{aligned} \frac{4}{n} &= \frac{1}{6m + 6} + \frac{23}{(6m + 6)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(2\ell + 1) \times 23}{(2\ell + 1) \times 6(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{2(23\ell + 10) + 3}{6(2\ell + 1)(m + 1)(24m + 1)}, \end{aligned} \quad (4.71)$$

where ℓ is a positive integer. According to (1.8) with $\mathfrak{I}'_1 = 2(23\ell + 10) + 3$ and $\mathfrak{I}'_2 = 3$, $\ell \not\equiv 1 \pmod{3}$. Suppose $\ell \equiv 0 \pmod{3}$; that is, $\ell = 3r$ for some $0 < r \in \mathbb{N}$. Then

$$23\ell + 10 = 69r + 10. \quad (4.72)$$

So we impose

$$m + 1 \equiv 0 \pmod{69r + 10}. \quad (4.73)$$

Equivalently,

$$m \equiv 69r + 9 \pmod{69r + 10}. \quad (4.74)$$

Thus

$$m = 69r + 9 + c(69r + 10) \quad \text{for some } c \in \mathbb{N}. \quad (4.75)$$

Assume that m is of the above form. Then

$$m + 1 = 69r + 9 + c(69r + 10) + 1 = (c + 1)(69r + 10). \quad (4.76)$$

According to (4.71),

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+6} + \frac{2(23\ell+10)+3}{6(2\ell+1)(m+1)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{2(69r+10)+3}{6(6r+1)(c+1)(69r+10)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{1}{3(6r+1)(c+1)(24m+1)} \\
&\quad + \frac{1}{2(6r+1)(c+1)(69r+10)(24m+1)}. \tag{4.77}
\end{aligned}$$

A subtle point is that we have well used the factor “3” in $6m+6$ of the denominator.

Let $\ell \equiv 2 \pmod{3}$; that is, $\ell = 3r + 2$ for some $0 < r \in \mathbb{N}$. Then

$$23\ell + 10 = 69r + 56. \tag{4.78}$$

So we impose

$$m + 1 \equiv 0 \pmod{69r + 56}. \tag{4.79}$$

Equivalently,

$$m \equiv 69r + 55 \pmod{69r + 56}. \tag{4.80}$$

Thus

$$m = 69r + 55 + c(69r + 56) \quad \text{for some } c \in \mathbb{N}. \tag{4.81}$$

When m is of the above form,

$$m + 1 = 69r + 55 + c(69r + 56) + 1 = (c + 1)(69r + 56). \tag{4.82}$$

By (4.71),

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+6} + \frac{2(23\ell+10)+3}{6(2\ell+1)(m+1)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{2(69r+56)+3}{6(6r+5)(c+1)(69r+56)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{1}{3(6r+5)(c+1)(24m+1)} \\
&\quad + \frac{1}{2(6r+5)(c+1)(69r+56)(24m+1)}. \tag{4.83}
\end{aligned}$$

Next we assume that $m = 2m_1 + 1$ with $m_1 \in \mathbb{N}$. We consider

$$\begin{aligned}
\frac{4}{n} &= \frac{1}{6m+6} + \frac{23}{(6m+6)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{(2\ell+1) \times 23}{(2\ell+1) \times 6(m+1)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{(46\ell+19)+4}{12(2\ell+1)(m_1+1)(24m+1)}, \tag{4.84}
\end{aligned}$$

where ℓ is a positive integer. First we suppose

$$(46\ell + 19) \mid [3(m_1 + 1)]. \quad (4.85)$$

Assume $\ell = 3s + 2$. Then

$$46\ell + 19 = 138s + 111 = 3(46s + 37) \implies (46s + 37) \mid (m_1 + 1). \quad (4.86)$$

So

$$m_1 = t(46s + 37) + 46s + 36 = 36 + 37t + 46s(t + 1). \quad (4.87)$$

In particular,

$$m = 73 + 74t + 92s(t + 1). \quad (4.88)$$

By (4.84),

$$\begin{aligned} \frac{4}{n} &= \frac{1}{6m + 6} + \frac{(46\ell + 19) + 4}{12(2\ell + 1)(m_1 + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{3(46s + 37) + 4}{12(6s + 5)(t + 1)(46s + 37)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{4(6s + 5)(t + 1)(24m + 1)} \\ &\quad + \frac{1}{3(6s + 5)(t + 1)(46s + 37)(24m + 1)}. \end{aligned} \quad (4.89)$$

When $m = 71925$, $n = 24m + 1 = 1726201$ is a prime. Taking $s = 781$ and $t = 0$, we have $2\ell + 1 = 4691$ and

$$\frac{1}{1726201} = \frac{1}{431556} + \frac{1}{32390435564} + \frac{1}{873642925641099}, \quad (4.90)$$

which coincides with (2.23).

Assume $3 \nmid (46\ell + 19)$. Then

$$(46\ell + 19) \mid (m_1 + 1) \implies m_1 = t(46\ell + 19) + 46\ell + 18 = 18 + 19t + 46\ell(t + 1). \quad (4.91)$$

In particular,

$$m = 37 + 38t + 92\ell(t + 1). \quad (4.92)$$

By (4.84),

$$\begin{aligned} \frac{4}{n} &= \frac{1}{6m + 6} + \frac{(46\ell + 19) + 4}{12(2\ell + 1)(m_1 + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(46\ell + 19) + 4}{12(2\ell + 1)(t + 1)(46\ell + 19)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{12(2\ell + 1)(t + 1)(24m + 1)} \\ &\quad + \frac{1}{3(2\ell + 1)(t + 1)(46\ell + 19)(24m + 1)}. \end{aligned} \quad (4.93)$$

Finally we consider

$$\begin{aligned}
\frac{4}{n} &= \frac{1}{6m+6} + \frac{23}{(6m+6)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{(2\ell+1) \times 23}{(2\ell+1) \times 6(m+1)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{(46\ell+17)+6}{6(2\ell+1)(m+1)(24m+1)},
\end{aligned} \tag{4.94}$$

where ℓ is a positive integer. According to (1.8) with $\mathfrak{S}'_1 = 46\ell + 17$ and $\mathfrak{S}'_2 = 3$, $\ell \not\equiv 1 \pmod{3}$. Suppose $\ell \equiv 0 \pmod{3}$; that is, $\ell = 3r$ for some $0 < r \in \mathbb{N}$. Then

$$46\ell + 17 = 138r + 17. \tag{4.95}$$

By (2.13), we impose

$$m + 1 \equiv 0 \pmod{138r + 17}. \tag{4.96}$$

Equivalently,

$$m \equiv 138r + 16 \pmod{138r + 17}. \tag{4.97}$$

Thus

$$m = 138r + 16 + c(138r + 17) \quad \text{for some } c \in \mathbb{N}. \tag{4.98}$$

Assume that m is of the above form. Then

$$m + 1 = 138r + 16 + c(138r + 17) + 1 = (c + 1)(138r + 17). \tag{4.99}$$

According to (4.84),

$$\begin{aligned}
\frac{4}{24m+1} &= \frac{1}{6m+6} + \frac{(46\ell+17)+6}{6(2\ell+1)(m+1)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{(138r+17)+6}{6(6r+1)(c+1)(138r+17)(24m+1)} \\
&= \frac{1}{6m+6} + \frac{1}{6(6r+1)(c+1)(24m+1)} \\
&\quad + \frac{1}{(6r+1)(c+1)(138r+17)(24m+1)}.
\end{aligned} \tag{4.100}$$

Let $\ell \equiv 2 \pmod{3}$; that is, $\ell = 3r + 2$ for some $0 < r \in \mathbb{N}$. Then

$$46\ell + 17 = 138r + 109. \tag{4.101}$$

By (2.13), we impose

$$m + 1 \equiv 0 \pmod{138r + 109}. \tag{4.102}$$

Equivalently,

$$m \equiv 138r + 108 \pmod{138r + 109}. \tag{4.103}$$

Thus

$$m = 138r + 108 + c(138r + 109) \quad \text{for some } c \in \mathbb{N}. \tag{4.104}$$

Assume that m is of the above form. Then

$$m + 1 = 138r + 108 + c(138r + 109) + 1 = (c + 1)(138r + 109). \quad (4.105)$$

By (4.84),

$$\begin{aligned} \frac{4}{24m + 1} &= \frac{1}{6m + 6} + \frac{(46\ell + 17) + 6}{6(2\ell + 1)(m + 1)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{(138r + 109) + 6}{6(6r + 5)(c + 1)(138r + 109)(24m + 1)} \\ &= \frac{1}{6m + 6} + \frac{1}{6(6r + 5)(c + 1)(24m + 1)} \\ &\quad + \frac{1}{(6r + 5)(c + 1)(138r + 109)(24m + 1)}. \end{aligned} \quad (4.106)$$

Theorem 4.2 *For any positive integer m of the form (4.37), we have the wild solution (4.39) of the Erdős-Straus equation. If m is of the form (4.43), we have the wild solution (4.45) of the Erdős-Straus equation. When m is of the form (4.49), we have the wild solution (4.51) of the Erdős-Straus equation. Suppose that m is of the form (4.56), and then we have the wild solution (4.58) of the Erdős-Straus equation. Let m be of the form (4.62), and then we have the wild solution (4.64) of the Erdős-Straus equation.*

For any positive integer m of the form (4.88), we have the wild solution (4.89) of the Erdős-Straus equation. For any positive integer m of the form (4.92), we have the wild solution (4.93) of the Erdős-Straus equation. For any positive integer m of the form (4.68), we have the wild solution (4.70) of the Erdős-Straus equation. If m is of the form (4.75), we have the wild solution (4.77) of the Erdős-Straus equation. When m is of the form (4.81), we have the wild solution (4.83) of the Erdős-Straus equation. Suppose that m is of the form (4.98), and then we have the wild solution (4.100) of the Erdős-Straus equation. Let m be of the form (4.104), and then we have the wild solution (4.1066) of the Erdős-Straus equation.

References

- [1] K. Bradford, A note on the Erdős-Straus conjecture, *Integers* **21** (2021), Paper No. A24, 10pp.
- [2] M. Bright and D. Loughran, Brauer-Manin obstruction for Erdős-Straus surfaces, *Bull. Lond. Math. Soc.* **52** (2020), no. 4, 746–761.
- [3] C. Elscholtz and T. Tao, Counting the number of solutions to the equations on unit fractions, *J. Aust. Math. Soc.* **94** (2013), 50–105.
- [4] T. R. Hagedon, A proof of a conjecture on Egyptian fractions, *Amer. Math. Monthly* **107** (2000), no. 1, 62–63.
- [5] B. Howerton, The wild primes formula sheet, *personal communication*, 12 June 2026.

- [6] J. Huang and R. Vaughan, Mean value theorems for binary Egyptian fractions, *J. Number Theory* **121** (2011), no. 9, 1641–1656.
- [7] J. Huang and R. Vaughan, Mean value theorems for binary Egyptian fractions II, *Acta Arith.* **115** (2012), no. 3, 287–296.
- [8] J. Huang and R. Vaughan, On the exceptional set for binary Egyptian fractions, *Bull. Lond. Math. Soc.* **45** (2013), no. 4, 861–874.
- [9] C. Jia, The estimate for mean values on prime numbers relative to $4/n = 1/n_1 + 1/n_2 + 1/n_3$, *Sci. China: Math.* **55** (3), (2012), 465–474.
- [10] C. Jia, Mean value from representation of a rational number as sum of two Egyptian fractions, *J. Number Theory* **132** (2012), no. 4, 701–713.
- [11] E. J. Ionascu and A. Wilson, On the Erdős-Straus conjecture, *Rev. Roumaine Math. Pures Appl.* **56** (2011), no. 1, 21–30.
- [12] I. Kotsireas, The Erdős-Straus conjecture on Egyptian fractions, *Paul Erdős and his mathematics (Budapest, 1999)*, 140–144, *János Nolgai Math. Soc. Budapest*, 1999.
- [13] D. Li, On the equation $4/n = 1/x + 1/y + 1/z$, *J. Number Theory* **13** (1981), no. 4, 485–494.
- [14] S. Maiti, A study on Erdős-Straus conjecture on Diophantine equation $\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$, *arXiv: 2010.00975v1 [math.GM]* 30 Dec 2020.
- [15] L. J. Mordell, *Diophantine Equations*, Academic Press, London/New York, 1969.
- [16] S. Prugsapitak, The Egyptian fraction of the form $\frac{1}{a} + \frac{1}{b} = \frac{q-1}{pq}$, *Int. J. Math. Comput. Sci.* **18** (2023), no. 4, 595–597.
- [17] L. Antonio Rosati, Sull’equazione diofantea $4/n = 1/x_1 + 1/x_2 + 1/x_3$, *Boll. Uni. Mat. Ital. (3)* **9** (1954), 59–63.
- [18] J. W. Sander, On $4/n = 1/x + 1/y + 1/z$ and Rosser’s sieve, *Acta Arith.* **59** (1991), no. 2, 183–204.
- [19] J. W. Sander, On $4/n = 1/x + 1/y + 1/z$ and Iwanniec’s half-dimensional sieve, *J. Number Theory* **46** (1994), no. 2, 123–136.
- [20] J. W. Sander, Egyptian fractions and the Erdős-Straus conjecture, *Nieuw Arch. Wisk (4)* **15** (1997), no. 1-2, 43–50.
- [21] J. D. Serna, Partial proof without words: shapping some cases of the Erdős-Straus conjecture, *College. Math. J.* **40** (2015), no. 3, 181.
- [22] S. Subburam and A. Togb’e, A note on Erdős-Straus conjecture, *Period Math. Hungar.* **72** (2016), no. 1, 43–49.
- [23] D. G. Terzi, On a conjecture by Erdős and Straus, *Nordisk Tidsk. Informationsbehandling (BIT)* **11** (1971), 212–216.

- [24] R. C. Vaughan, On a problem of Erdős, Straus and Schinzel, *Mathematika* **17** (1970), 193–198.
- [25] W. A. Webb, On $4/n = 1/x + 1/y + 1/z$, *Proc. Amer. Math. Soc.* **25** (1970), 578–584.
- [26] X. Xu, Congruence classes of supporting the Erdős-Straus conjecture I: tame solutions, *arXiv: 2605.2360[v2]* [math.NT] 19 Sep 2026

E-Mail: xiaoping@math.ac.cn