

Partial Resolution of the Erdős–Straus, Sierpiński, and Generalized Erdős–Straus Conjectures Using New Analytical Formulas

Philemon Urbain MBALLA

philemonmballa@gmail.com

philemon-urbain.mballa@etu.u-paris.fr

Abstract

This article proposes a unified analytical approach leading to a partial resolution of the Erdős–Straus, Sierpiński conjectures, and their generalization. We introduce an equivalent reformulation of these conjectures while constructing two new explicit analytical formulas. The first formula, which is a special case of the second, is based on a divisibility condition, whereas the second, more general formula, relies on the existence of a perfect square, which we conjecture to always hold. Under these conditions, the formulas verify the conjectures even for very large numerical values. Moreover, our method reduces the problem to the search for a suitable perfect square, thereby opening the way to a complete proof of these conjectures. In conclusion, we present open questions and conjectures to the mathematical community regarding the generalization of these formulas.

1 Introduction

The Erdős–Straus conjecture, formulated in 1948 by Paul Erdős and Ernst G. Straus, states that for every natural number $n \geq 2$, there exist $x, y, z \in \mathbb{N}^*$ such that

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}. \quad (1)$$

Louis Mordell demonstrated that if n is not congruent to $1, 11^2, 13^2, 17^2, 19^2$, or 23^2 modulo 840, then the conjecture holds. Furthermore, numerical verifications have confirmed its validity up to $n = 10^{17}$.

For decades, mathematicians have been intrigued by the following question: **Does an explicit formula exist relating x, y, z , and n that systematically verifies the conjecture?**

In this article, we propose two new analytical formulas obtained through an in-depth study of the Diophantine equation (1). The first relies on a natural divisibility condition, while the second is based on the existence of a perfect square. These results provide a partial proof of the conjecture and represent a significant advancement in its study.

The Polish mathematician Waław Sierpiński also investigated this conjecture and posed a similar question: **Is a similar decomposition possible if 4 is replaced by 5?** He

conjectured that for every natural number $n \geq 2$, there exist $x, y, z \in \mathbb{N}^*$ such that

$$\frac{5}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}. \quad (2)$$

In this article, we will demonstrate that our formulas also verify this conjecture under the same conditions as before.

Finally, following the formulation of the Erdős–Straus conjecture, several mathematicians explored the possibility of generalizing this Egyptian fraction representation. This led to the generalized Erdős–Straus conjecture, which may be more precisely formulated as follows: for every integer $a \geq 5$, there exists an integer $N_0 \in \mathbb{N}$ with $N_0 \geq 2$ (possibly depending on a , since it is easy to see that N_0 cannot be strictly smaller than $a/3$) such that for all $n \geq N_0$, there exist $x, y, z \in \mathbb{N}^*$ satisfying

$$\frac{a}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}. \quad (3)$$

Moreover, this result is obtained using the same analytical formulas developed for the Erdős–Straus and Sierpiński conjectures.

In this article, we present a detailed analytical approach leading to these formulas, along with numerical tests supporting our results.

2 The Erdős–Straus Conjecture

One of the methods that is known nowadays is that if n is even integer, meaning $n = 2k$ with $k \in \mathbb{N}^*$. In this case, it is well established that:

$$\frac{4}{n} = \frac{1}{\frac{n}{2}} + \frac{1}{n} + \frac{1}{n}. \quad (4)$$

This result is one of the most trivial among those obtained since the conjecture was stated, as it shows that all even numbers satisfy the conjecture.

In our new approach, I will demonstrate how this result can be naturally recovered when n is even. Before introducing the general method that allowed me to establish two formulas verifying the conjecture under a certain condition, I will first illustrate my process by analytically testing the conjecture for some values of n .

Our method is based on the structural analysis of equation (1). It is a simple yet powerful and general approach, applicable to even numbers, odd numbers, and prime numbers alike.

—

2.1 Case $n = 2$ (Even Number)

The fundamental equation becomes:

$$\frac{4}{2} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}.$$

I choose not to simplify $\frac{4}{2}$ immediately in order to maintain the general structure of the method. However, ****one can simplify this fraction directly****, and by following the same steps that I will present, the same result is always obtained.

The equation can then be rewritten as:

$$\frac{4}{2} - \frac{1}{x} = \frac{1}{y} + \frac{1}{z},$$

which reformulates as:

$$\frac{4x - 2}{2x} = \frac{y + z}{yz}.$$

Since we are searching for solutions in $\mathbb{N}^*$, it is necessary that $4x - 2$ be strictly positive, that is, $4x - 2 > 0$, which gives $x \geq \lfloor \frac{1}{2} \rfloor + 1 = 1$.

For the rest of the document, the notation $\lfloor x \rfloor$ will always denote the integer part of x to avoid any confusion.

We then obtain the equation:

$$yz(4x - 2) = 2x(y + z).$$

At this stage, I apply a method commonly used in geometry of curves and surfaces, called parameterization.

2.2 Parameterization Technique

Let $a, c \in \mathbb{R}$, and $b, d \in \mathbb{R}^*$ such that:

$$ab = cd.$$

Then, for $t \in \mathbb{R}$, we can set:

$$a = dt, \quad c = bt.$$

This parametric representation holds for $t \in \mathbb{R}$ and allows rewriting $(dt)b = (bt)d$, which verifies the initial equation. This technique will be crucial for my results.

2.3 Application to Our Equation

We then set:

$$y + z = (4x - 2)t, \quad yz = 2xt.$$

This parameterization is valid for $t \in \mathbb{Q}_+^*$.

According to the sum and product method, y and z are solutions of the quadratic equation:

$$V^2 - ((4x - 2)t)V + 2xt = 0.$$

Its discriminant is given by:

$$\Delta = t^2(4x - 2)^2 - 8xt.$$

We imposed $x \geq 1$. Choosing $x = 1$, we obtain:

$$\Delta = 4t^2 - 8t = t(4t - 8).$$

For y and z to be integers, Δ must be a perfect square, and $4x - 2$ must have the same parity as $\sqrt{\Delta}$.

Taking $t = 2$ (a trivial value for which $\Delta = 0$), we obtain a double root:

$$y = z = \frac{(4 \times 1 - 2) \times 2}{2} = 2.$$

Thus, for $n = 2$, we find $x = 1, y = 2, z = 2$, verifying:

$$\frac{4}{2} = \frac{1}{1} + \frac{1}{2} + \frac{1}{2}.$$

This confirms the validity of the conjecture for $n = 2$.

2.4 Analysis of the Method

if we had directly applied the known formula for even numbers given by equation (4) , we would have found:

$$x = \frac{2}{2} = 1, \quad y = z = 2.$$

Thus, we obtain exactly the same values.

One of the key advantages of our analytical approach is its applicability not only to even numbers but also to odd numbers and prime numbers. Unlike classical methods—which have relied primarily on brute-force techniques that merely test the existence of solutions without offering insight into the problem’s structure—our method derives solutions directly from the inherent logical structure of the underlying Diophantine equation. This provides a general, rigorous framework for addressing the Erdős–Straus, Sierpiński, and generalized Erdős–Straus conjectures. Moreover, the elegance and clarity of our approach make it a valuable pedagogical tool for teaching methods of solving Diophantine equations, as it not only yields the solutions but also deepens our understanding of the arithmetic properties involved.

Of course, testing the conjecture for specific values of n does not constitute a general proof. My goal here is to demonstrate the efficiency of this method in solving equation (1) analytically.

Next, we will examine the cases $n = 3$, $n = 17$, and finally $n = 841$ (one of the numbers that Mordell had not proven), before moving on to generalization. Since the main challenge concerns odd numbers, I will not dwell much on even numbers. However, you can reproduce this reasoning for any even n , and it will work the same way.

2.5 Case $n = 3$ (odd and prime Number)

Applying the same procedure, we impose:

$$x \geq \left\lfloor \frac{3}{4} \right\rfloor + 1 = 1.$$

We then obtain the system:

$$\begin{cases} y + z = (4x - 3)t, \\ yz = 3xt, \end{cases}$$

for $t \in \mathbb{Q}_+^*$.

The values of y and z satisfy the quadratic equation:

$$V^2 - ((4x - 3)t)V + 3xt = 0.$$

The discriminant is given by:

$$\Delta = t^2(4x - 3)^2 - 12xt.$$

We choose $x \geq \left\lfloor \frac{4}{3} \right\rfloor + 1 = 2$.

Taking $x = 2$, we obtain:

$$\Delta = 25t^2 - 24t = t(25t - 24).$$

A trivial solution ensuring that Δ is a perfect square is $t = 1$, yielding $\Delta = 1$.

Thus, we find:

$$y = \frac{(4 \times 2 - 3) - 1}{2} = 2, \quad z = \frac{(4 \times 2 - 3) + 1}{2} = 3.$$

For $n = 3$, we have $x = 2$, $y = 2$, and $z = 3$, verifying:

$$\frac{4}{3} = \frac{1}{2} + \frac{1}{2} + \frac{1}{3}.$$

By selecting a different x , we can determine corresponding values of t that yield integer solutions. For instance, choosing $x = 4$ results in:

$$\Delta = 169t^2 - 48t = t(169t - 48).$$

For $t = 1$, we find a perfect square $11^2 = 121$, yielding $y = 1$ and $z = 12$.

—

2.6 Case $n = 17$ (odd and prime Number)

We impose:

$$x \geq \left\lfloor \frac{17}{4} \right\rfloor + 1 = 5.$$

The system obtained is:

$$\begin{cases} y + z = (4x - 17)t, \\ yz = 17xt, \end{cases}$$

for $t \in \mathbb{Q}_+^*$.

y and z satisfy the quadratic equation:

$$V^2 - ((4x - 17)t)V + 17xt = 0.$$

The discriminant is:

$$\Delta = t^2(4x - 17)^2 - 68xt.$$

To minimize $(4x - 17)^2$ and simplify calculations, we choose the smallest $x \geq 5$, i.e., $x = 5$.

This gives:

$$\Delta = 9t^2 - 340t = t(9t - 340).$$

We require Δ to be a perfect square, meaning $9t - 340 \geq 0$, so $t \geq \frac{340}{9} \approx 37.77$. Using a Python script to scan t in the range $[38, 1000]$, we find the perfect square values of Δ :

$$t = 68 \Rightarrow \sqrt{\Delta} = 136, \quad t = 180 \Rightarrow \sqrt{\Delta} = 480.$$

For $t = 68$, we obtain:

$$y = 34, \quad z = 170.$$

Thus, we confirm:

$$\frac{4}{17} = \frac{1}{5} + \frac{1}{34} + \frac{1}{170}.$$

The reader may compute y and z for $t = 180$ using the same approach.

2.7 Case $n = 841$ (number not proven by Mordell)

Following the same procedure, we impose:

$$x \geq \left\lfloor \frac{841}{4} \right\rfloor + 1 = 211.$$

The system obtained is:

$$\begin{cases} y + z = (4x - 841)t, \\ yz = 841xt, \end{cases}$$

for all $t \in \mathbb{Q}_+^*$.

y and z satisfy the quadratic equation:

$$V^2 - ((4x - 841)t)V + 841xt = 0.$$

The discriminant is:

$$\Delta = t^2(4x - 841)^2 - 3364xt.$$

Choosing $x = 211$, we obtain:

$$\Delta = 9t^2 - 709804t = t(9t - 709804).$$

For Δ to be a perfect square, we require $9t - 709804 \geq 0$, so $t \geq \frac{709804}{9} \approx 78867.11$. Scanning for t in the range $[78868, 500000]$, we find:

$$t = 185600, \quad \sqrt{\Delta} = 422240.$$

Thus, we compute:

$$y = 67280, \quad z = 489520.$$

Verifying:

$$\frac{4}{841} = \frac{1}{211} + \frac{1}{67280} + \frac{1}{489520}.$$

Both sides yield the same numerical value.

NB. In most cases, we restrict the search for t to $\mathbb{N}^*$. However, numerical simulations indicate that allowing t to range over non-integer positive rationals $\mathbb{Q}_+^* \setminus \mathbb{N}$ becomes necessary for $a \geq 5$ when n is small, namely in a neighborhood of the threshold N_0 , since solutions remain sparse in this region. Beyond a certain threshold, we observe an abundance of integer values of t for which the discriminant is a perfect square. Moreover, t may also naturally take values in $\mathbb{Q}_+^*$ in the context of Formula (1) as soon as the divisibility condition is satisfied.

3 Conclusion

This analytical approach effectively solves equation (1) and finds solutions satisfying the conjecture for even, odd, and prime numbers.

With this method, one can manually prove the conjecture for many numbers not previously proven by Mordell.

However, verifying the conjecture for specific values does not constitute a general proof. We will now proceed to the generalization and present the results obtained.

4 Generalization of the Erdős-Straus Conjecture

We want to prove that for all n in $\mathbb{N}^*$ with $n \geq 2$, there exist x, y, z in $\mathbb{N}^*$ such that

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$$

We have called this equation (1).

My proposed proof follows the same structure as before, as you may have noticed, the value of n does not change the structure of the proof. I will use exactly the same procedure as previously.

Equation (1) is equivalent to:

$$\frac{4}{n} - \frac{1}{x} = \frac{y+z}{yz}$$

which is also equivalent to

$$\frac{4x-n}{nx} = \frac{y+z}{yz}$$

For the right-hand side to be strictly positive, it is necessary that $4x-n > 0$, i.e., $x > \frac{n}{4}$, so we have $x \geq \left\lceil \frac{n}{4} \right\rceil$.

We still have:

$$\frac{4x-n}{nx} = \frac{y+z}{yz}$$

which holds if and only if

$$yz(4x-n) = nx(y+z)$$

Similarly, using parameterization, we can express:

$$\begin{cases} y+z = (4x-n)(2t) \\ yz = nx(2t) \end{cases}$$

for $t \in \mathbb{Q}_+^*$. If we replace $y+z$ and yz with these expressions, the equation $yz(4x-n) = nx(y+z)$ will always hold.

Later, I will explain why I parameterized using $2t$ in this general case instead of using t as I did for specific values of n . (Of course, we are allowed to parameterize using any multiple of $t \in \mathbb{Q}_+^*$, as the equation will always hold.)

Now, let us use sums and products. From the system above, y and z satisfy the quadratic equation in V :

$$V^2 - (\text{sum})V + \text{product} = 0$$

which gives:

$$V^2 - ((4x - n)(2t))V + 2ntx = 0.$$

The discriminant is given by:

$$\Delta = 4t^2(4x - n)^2 - 8ntx. \quad (5)$$

You may recall that I promised to retrieve equation (4) for even values of n . If we set $n = 2x$, meaning n is even since x is in $\mathbb{N}^*$, the discriminant becomes:

$$\Delta = 4t^2(4x - 2x)^2 - 16x^2t.$$

$$= 16x^2t^2 - 16x^2t.$$

An obvious choice for t so that Δ is a perfect square is $t = 1$, which gives $\Delta = 0$. Thus,

$$y = z = \frac{2(4x - 2x)}{2} = 2x$$

so $y = z = n = 2x$, which implies $x = \frac{n}{2}$, and we obtain:

$$\frac{4}{n} = \frac{1}{\frac{n}{2}} + \frac{1}{n} + \frac{1}{n}$$

which retrieves equation (4).

This was a side note, as our goal is to generalize. So, we return to our quadratic equation in V in its general form, with $n \in \mathbb{N}^*$ such that $n \geq 2$ (as required by the conjecture).

We still have our discriminant computed in (5):

To obtain integer solutions for this equation, it is necessary that Δ be a perfect square. I struggled to find an obvious perfect square for Δ . It seems to me that the polynomial given in (5) is rich, but unfortunately, after some quick and unsuccessful attempts to find a classic perfect square of Δ without imposing any condition on n (since we are in the process of generalization), the idea I had was to use the fact that the discriminant Δ holds for $t \in \mathbb{Q}_+^*$ arbitrary and see what can be deduced to obtain a perfect square in order to bypass the difficulty. Let's see how I proceed Let's revisit the compact form of Δ in (5):

5 Compact Form of Δ

$$\Delta = 4t^2(4x - n)^2 - 8ntx$$

This simplifies to:

$$\Delta = 4t(t(4x - n)^2 - 2nx).$$

This is valid for $t \in \mathbb{Q}_+^*$ arbitrary because, as mentioned earlier, the parametrization is valid for $t \in \mathbb{Q}_+^*$ arbitrary. You also observed how this information allowed us to prove

the conjecture (manually for some n) by providing specific values of t that help us obtain a perfect square. We will use this information in this generalization.

We saw that $\Delta > 0$ for

$$t > \frac{2nx}{(4x - n)^2}.$$

Recalling the expression for Δ ,

$$\Delta = 4t(t(4x - n)^2 - 2nx),$$

and since $\Delta > 0$ for

$$t > \frac{2nx}{(4x - n)^2}.$$

for the x and t which respect this condition, it is clear that there exist x and t such that $t(4x - n)^2 - 2nx$ is in $\mathbb{Q}_+^*$. We can then write:

$$t(4x - n)^2 - 2nx = k,$$

with $k \in \mathbb{Q}_+^*$. Thus,

$$\Delta = 4t(t(4x - n)^2 - 2nx) = 4tk,$$

which holds for $t \in \mathbb{Q}_+^*$ arbitrary. So, for $t = k$ (This is why we excluded the case $\Delta = 0$, otherwise $k = (t(4x - n)^2 - 2nx)$ can be zero and we cannot set $t = k$ because from the start, during the parameterization, t is non-zero). , so we have:

$$\Delta = 4tk = 4k^2 = (2k)^2.$$

We have thus obtained a perfect square (rational or integer perfect square). Let's now see under what conditions this transformation holds.

6 Conditions for Obtaining a Perfect Square

To obtain this perfect square, we wrote $t(4x - n)^2 - 2nx = k$, and then set $t = k$, simplifying the equation to:

$$k(4x - n)^2 - 2nx = k,$$

or:

$$k((4x - n)^2 - 1) = 2nx.$$

This implies that

$$k = \frac{2nx}{(4x - n - 1)(4x - n + 1)}, \quad k \in \mathbb{Q}_+^*.$$

At this stage, no additional condition is required, since the above expression naturally defines a positive rational value of k without any restriction.

Since, $\Delta = 4k^2 = (2k)^2$, so:

$$V = \frac{(4x - n)(2k) \pm 2k}{2}.$$

there is a simplification of 2, This is why I had previously parameterized by $2t$, and we just replaced t with k to calculate the roots of this polynomial, and this was possible because we set $t = k$ to obtain a perfect square.

we have $y = k(4x - n - 1)$ and $z = k(4x - n + 1)$. Now, from earlier, we know that:

$$k = \frac{2nx}{(4x - n - 1)(4x - n + 1)}$$

Substituting k into the expressions for y and z , we get:

$$y = \frac{2nx}{(4x - n - 1)(4x - n + 1)}(4x - n - 1) = \frac{2nx}{4x - n + 1}$$

and

$$z = \frac{2nx}{(4x - n - 1)(4x - n + 1)}(4x - n + 1) = \frac{2nx}{4x - n - 1}$$

A natural condition for y and z to be integers is that both $4x - n - 1$ and $4x - n + 1$ divide $2nx$.

Next, we find the conditions for x such that both y and z are positives. Since $2nx > 0$, we only need to ensure that the denominators are strictly positive.

Case for y :

$y > 0$ if:

$$4x - n + 1 > 0 \quad \text{which implies} \quad x > \frac{n - 1}{4}$$

Thus, for y , we need $x \geq \left\lceil \frac{n-1}{4} \right\rceil + 1$.

Case for z :

$z > 0$ if:

$$4x - n - 1 > 0 \quad \text{which implies} \quad x > \frac{n + 1}{4}$$

Thus, for z , we need $x \geq \left\lceil \frac{n+1}{4} \right\rceil + 1$.

Therefore, the overall condition is:

$$x \geq \max \left\{ \left\lceil \frac{n-1}{4} \right\rceil + 1, \left\lceil \frac{n+1}{4} \right\rceil + 1 \right\}$$

which gives:

$$x \geq \left\lceil \frac{n+1}{4} \right\rceil + 1$$

Thus, $x \in \left[\left\lceil \frac{n+1}{4} \right\rceil + 1, +\infty \right[$.

Now, let us check for which values of x , y , and z are in $\mathbb{N}^*$. We recall the following property: if $a \in \mathbb{N}^*$ and $b \in \mathbb{N}$, and if a divides b , then $a \leq b$.

Therefore, $y \in \mathbb{N}^*$ if and only if $4x - n + 1$ divides $2nx$, which implies:

$$4x - n + 1 \leq 2nx$$

$$4x - 2nx \leq n - 1$$

$$x(4 - 2n) \leq n - 1$$

For $n > 2$, $4 - 2n < 0$, so:

$$x \geq \frac{n-1}{2n-4}, \quad x \in \mathbb{N}$$

Similarly, we can show that $z \in \mathbb{N}^*$ if:

$$x \geq \frac{n+1}{2n-4}$$

Thus, we have the following conditions:

$$x \geq \left\lceil \frac{n+1}{4} \right\rceil + 1$$

$$x \geq \frac{n-1}{2n-4}$$

$$x \geq \frac{n+1}{2n-4}$$

The two latter values of x are not valid for $n = 2$. Therefore, the correct choice that satisfies all conditions is:

$$x \geq \left\lceil \frac{n+1}{4} \right\rceil + 1$$

In other words, $x \in \left[\left\lceil \frac{n+1}{4} \right\rceil + 1, +\infty \right[$.

As we will see in the numerical checks, under these divisibility conditions, this formula will allow us to decompose the fraction $\frac{4}{n}$ into three unit fractions for those n satisfying these conditions up to infinity (since we have shown that the values of x for which $4x - n - 1$ and $4x - n + 1$ divide $2nx$ extend to infinity). This is good news because we always hope that for any given n , there is an infinite number of choices for x that satisfy the divisibility condition, allowing us to prove the conjecture for that specific n .

Before proceeding with this, let's first verify whether, under these divisibility conditions, the Erdős–Strauss conjecture is satisfied for:

$$y = \frac{2nx}{4x - n + 1} \quad \text{and} \quad z = \frac{2nx}{4x - n - 1}$$

We need to check if:

$$\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = \frac{4}{n}$$

Substituting the expressions for y and z :

$$\frac{1}{x} + \frac{1}{\frac{2nx}{4x-n+1}} + \frac{1}{\frac{2nx}{4x-n-1}} = \frac{1}{x} + \frac{4x-n+1}{2nx} + \frac{4x-n-1}{2nx}$$

Simplifying:

$$\frac{1}{x} + \frac{(4x-n+1) + (4x-n-1)}{2nx} = \frac{1}{x} + \frac{8x-2n}{2nx} = \frac{1}{x} + \frac{4x-n}{nx} = \frac{4}{n}$$

Thus, x , y , and z satisfy the Erdős–Strauss conjecture under the established divisibility conditions.

In summary, we have proven that for all $n \geq 2$, If there exists $x_0 \in \mathbb{N}^*$ with $x_0 \in \left[\left[\frac{n+1}{4}\right] + 1, +\infty\right[$. such that $4x_0 - n - 1$ and $4x_0 - n + 1$ divide $2nx_0$, then there exist $y = \frac{2nx_0}{4x_0 - n - 1}$ and $z = \frac{2nx_0}{4x_0 - n + 1}$ in $\mathbb{N}^*$ such that:

$$\frac{4}{n} = \frac{1}{x_0} + \frac{1}{y} + \frac{1}{z}$$

we therefore have the following theorem:

Theorem 1. *Let $n \geq 2$ be an integer. Suppose there exists $x_0 \in \mathbb{N}^*$ with*

$$x_0 \in \left[\left[\frac{n+1}{4}\right] + 1, +\infty\right)$$

such that both integers $4x_0 - n - 1$ and $4x_0 - n + 1$ divide $2nx_0$. Then there exist $y, z \in \mathbb{N}^$ given by*

$$y = \frac{2nx_0}{4x_0 - n - 1} \quad \text{and} \quad z = \frac{2nx_0}{4x_0 - n + 1},$$

such that

$$\frac{4}{n} = \frac{1}{x_0} + \frac{1}{y} + \frac{1}{z}.$$

Below, I will provide numerical calculations that, depending on the range searched for x_0 , cover a some n . Based on this formula verifying the conjecture, you will also be able to perform calculations using more powerful computational resources to identify very large x_0 and n . Of course, with numerical computations, it is impossible to reach $+\infty$, but what assures us of the validity of the results is the analytical result we have just established. Every time we find such an x_0 , the conjecture holds for the corresponding n . My question to the mathematical community for a complete proof of the conjecture is as follows: How can we explicitly find x_0 ? If it is not possible to obtain an explicit expression for x_0 (which may depend on n), can we prove the existence of x_0 by induction on n (whether classical or strong)? I remain available for any suggestions or collaborations to address this question and definitively resolve the conjecture.

7 Second Formula: Condition of the Existence of a Perfect Square

While the first formula is derived from a divisibility condition, the second one is more fundamental in nature, as it provides a genuinely more general framework. Indeed, it is based on the existence of a perfect square and subsumes the first formula as a particular case. We now introduce this second formula.

Let us reconsider the discriminant Δ of our quadratic equation:

$$\Delta = 4t^2(4x - n)^2 - 8ntx = 4(t^2(4x - n)^2 - 2ntx).$$

We have already seen that $\Delta \geq 0$ for $t > \frac{2nx}{(4x-n)^2}$ with $t \in \mathbb{N}^*$. Therefore, for such values of t , we define

$t^2(4x - n)^2 - 2ntx = q^2$, with $q \in \mathbb{N}$ (as mentioned previously, here and in most cases we take t and q to be integers in order to automatically guarantee that, once the perfect square condition is satisfied, the resulting solutions are integral. Our parametrization

allows t to take arbitrary values without affecting the structure of the equation, and it is therefore up to us to choose suitable values of t ensuring that the corresponding values of y and z are integers).

Thus, we have:

$$\Delta = 4q^2 = (2q)^2.$$

We will now prove that the conjecture is true under the condition that this perfect square exists.

Let $y = t(4x - n) - q$ and $z = t(4x - n) + q$. We need to show that, with this formula, the conjecture holds for those q such that

$$t^2(4x - n)^2 - 2ntx = q^2, \quad q \in \mathbb{N}.$$

It is sufficient to check if

$$\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = \frac{1}{x} + \frac{1}{t(4x - n) - q} + \frac{1}{t(4x - n) + q}.$$

We can simplify this expression:

$$\frac{1}{x} + \frac{1}{t(4x - n) - q} + \frac{1}{t(4x - n) + q} = \frac{1}{x} + \frac{2t(4x - n)}{t^2(4x - n)^2 - q^2}.$$

Now, since $q^2 = t^2(4x - n)^2 - 2ntx$, we have:

$$\frac{1}{x} + \frac{2t(4x - n)}{t^2(4x - n)^2 - t^2(4x - n)^2 + 2ntx} = \frac{1}{x} + \frac{2t(4x - n)}{2nxt}.$$

Simplifying further:

$$= \frac{1}{x} + \frac{4x - n}{nx} = \frac{n - n + 4x}{nx} = \frac{4}{n}.$$

Thus, the second formula allows us to state the following theorem.

Theorem 2. *For any $n \in \mathbb{N}$ with $n \geq 2$, if there exists $(x_0, t_0) \in (\mathbb{N}^*)^2$ such that*

$$t_0^2(4x_0 - n)^2 - 2nt_0x_0 = q^2 \in \mathbb{N},$$

then there exist

$$y = t_0(4x_0 - n) - q \quad \text{and} \quad z = t_0(4x_0 - n) + q,$$

with $(y, z) \in \mathbb{N}^2$, such that

$$\frac{4}{n} = \frac{1}{x_0} + \frac{1}{y} + \frac{1}{z}.$$

This also proves the conjecture under this condition. I will also present below the numerical values of n found within a range where we search for (x_0, t_0) through numerical calculations.

I also pose the following question to the mathematical community: how can we find an explicit formula for (x_0, t_0) that works for all n without any conditions? In other words, how can we prove that the polynomial $t^2(4x - n)^2 - 2ntx$ or $4t^2(4x - n)^2 - 8ntx$ always admits a perfect square for all $n \in \mathbb{N}$, with $n \geq 2$, in order to complete the proof of the conjecture?

I remain available for suggestions or collaborations to address this question.

This is the contribution of this article to the Erdős-Strauss conjecture.

8 Sierpiński Conjecture

The Sierpiński conjecture is similar to the Erdős-Straus conjecture, except that 4 is replaced by 5. It states that for any $n \in \mathbb{N}$ with $n \geq 2$, there exist $x, y, z \in \mathbb{N}^*$ such that:

$$\frac{5}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}. \quad (7)$$

A trivial case arises when n is divisible by 3. In this case, one can simply take $x = \frac{n}{3}$, $y = z = n$, and the conjecture is verified for all n divisible by 3.

The good news is that with this new approach, everything stated for the Erdős-Straus conjecture remains valid for the Sierpiński conjecture by simply replacing 4 with 5. This demonstrates the power and generality of this method.

To keep the article concise, I will not manually test specific values of n as I did for the Erdős-Straus conjecture. However, the reasoning remains the same, and the reader may verify this independently.

8.1 Transforming the Equation

Equation (7) is equivalent to:

$$\frac{5}{n} - \frac{1}{x} = \frac{y+z}{yz}, \quad (4)$$

which can be rewritten as:

$$\frac{5x-n}{nx} = \frac{y+z}{yz}. \quad (5)$$

For the right-hand side to be strictly positive, it is necessary that $5x - n > 0$, i.e., $x > \frac{n}{5}$, leading to $x \geq \left\lfloor \frac{n}{5} \right\rfloor + 1$.

Using parameterization, we set:

$$y+z = (5x-n)(2t), \quad yz = nx(2t), \quad t \in \mathbb{Q}_+^*. \quad (6)$$

From the above, y and z satisfy the quadratic equation:

$$V^2 - (5x-n)(2t)V + 2ntx = 0. \quad (7)$$

The discriminant of this equation is:

$$\Delta = 4t^2(5x-n)^2 - 8ntx. \quad (8)$$

8.2 Special Case: n Divisible by 3

If $n = 3x$, then:

$$\begin{aligned} \Delta &= 4t^2(5x-3x)^2 - 24x^2t, \\ &= 16x^2t^2 - 24x^2t. \end{aligned}$$

A simple choice of t making Δ a perfect square is $t = 2$, yielding $\Delta = 16x^2 = (4x)^2$. Thus, solving for y and z :

$$y = \frac{4(5x - 3x) - 4x}{2} = 2x = \frac{2n}{3},$$

$$z = 2n.$$

Hence, we confirm:

$$\frac{5}{n} = \frac{1}{n/3} + \frac{1}{2n/3} + \frac{1}{2n}. \quad (9)$$

This case was just an example; our goal is to generalize the approach.

8.3 First Formula for the Sierpiński Conjecture

Solving for t such that $\Delta > 0$, we obtain:

$$t > \frac{2nx}{(5x - n)^2}.$$

Since:

$$\Delta = 4t^2(5x - n)^2 - 8ntx = 4t(t(5x - n)^2 - 2nx), \quad (10)$$

we can set $t = k = k(5x - n)^2 - 2nx$. Thus, we find:

$$\Delta = 4k^2 = (2k)^2, \quad (11)$$

leading to:

$$y = k(5x - n - 1), \quad z = k(5x - n + 1). \quad (12)$$

Replacing k by its expression:

$$k = \frac{2nx}{(5x - n - 1)(5x - n + 1)}, \quad (13)$$

We obtain:

$$y = \frac{2nx}{5x - n + 1},$$

$$z = \frac{2nx}{5x - n - 1}.$$

Thus, $y, z \in \mathbb{N}^*$ if and only if both $5x - n - 1$ and $5x - n + 1$ divide $2nx$. It follows that:

$$\frac{5}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}. \quad (14)$$

8.4 Conclusion

All details provided for the Erdős-Straus conjecture remain valid here, so I will not repeat them. I will also include numerical computations generating certain n values that satisfy these conditions.

Similarly, for the second approach:

$$\Delta = 4(t^2(5x - n)^2 - 2ntx), \quad (15)$$

for values of x satisfying $\Delta > 0$, we set

$$t^2(5x - n)^2 - 2ntx = q^2, \quad q \in \mathbb{N}.$$

, leading to:

$$\Delta = 4q^2 = (2q)^2. \quad (16)$$

We will now prove that the conjecture is true under the condition that this perfect square exists.

We have:

$$y = \frac{2t(5x - n) - 2q}{2} = t(5x - n) - q, \quad z = t(5x - n) + q.$$

Similarly, we can see that:

$$\frac{5}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}.$$

By substituting q^2 with $t^2(5x - n)^2 - 2ntx$, we obtain the desired result.

We may therefore conclude this section with the following theorem.

Theorem 3. *For any integer $n \geq 2$, if there exists $(x_0, t_0) \in \mathbb{N}^{*2}$ such that*

$$t_0^2(5x_0 - n)^2 - 2nt_0x_0 = q^2 \in \mathbb{N}, \quad (17)$$

then the quantities

$$y = t_0(5x_0 - n) - q \quad \text{and} \quad z = t_0(5x_0 - n) + q$$

satisfy

$$\frac{5}{n} = \frac{1}{x_0} + \frac{1}{y} + \frac{1}{z},$$

with $(y, z) \in \mathbb{N}^$.*

I invite the mathematical community to find a general explicit formula for (x_0, t_0) that works for all n , thereby fully resolving the Sierpiński conjecture.

9 Generalization of the Erdős-Straus and Sierpiński Conjectures

We recall that the generalized of the Erdős-Straus and Sierpiński Conjectures states that for every integer $a \geq 5$, there exists an integer $N_0 \in \mathbb{N}$ with $N_0 \geq 2$ such that for all $n \geq N_0$, there exist $x, y, z \in \mathbb{N}^*$ satisfying

$$\frac{a}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}. \quad (18)$$

It should be noted that since the inception of this conjecture, there has not been significant research on it, and thus no concrete leads have emerged. In this paper, we will attempt to partially prove it as I did with the two conjectures above, and we will reduce the difficulty to verifying two conditions to completely settle all three conjectures. Here, we will use the same approach as in the previous two conjectures.

It is evident that the Erdős-Straus conjecture and the Sierpiński conjecture are special cases of the generalized Erdős-Straus and Sierpiński conjecture. Therefore, all results established here remain valid for the two conjectures discussed earlier.

9.1 Trivial Case

As before, let us start with the trivial case: If $(a - 2)$ divides n , then we can simply take $x = \frac{n}{a-2}$ and $y = z = n$, which proves the conjecture in this particular case.

9.2 Equivalence of the Conjecture

Theorem 4. *Let $a \geq 4$ be a fixed integer. There exists an integer $N_0 \in \mathbb{N}$ with $N_0 \geq 2$ such that, for all $n \geq N_0$, the generalized Erdős-Straus and Sierpiński conjecture holds if and only if there exist $x \in \mathbb{N}^*$ and $t \in \mathbb{Q}_+^*$ such that y and z are solutions of the quadratic equation in V :*

$$V^2 - (ax - n)(2t)V + 2nxt = 0.$$

That is, we will establish that for all $n \geq N_0$, the identity

$$\frac{a}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$$

is satisfied if and only if y and z are roots of the quadratic equation above.

9.3 Preliminary Result

Proposition 1 (Vieta's Formulas). *Let X_1 and X_2 be two real (or complex) numbers, and consider the quadratic equation*

$$X^2 - SX + P = 0,$$

where S and P are real (or complex) parameters.

Then the following statements are equivalent:

1. X_1 and X_2 are solutions of the quadratic equation.
2. X_1 and X_2 satisfy the relations

$$X_1 + X_2 = S \quad \text{and} \quad X_1 X_2 = P.$$

This property is a direct consequence of Vieta's formulas.

9.4 Proof of the Theorem (Equivalence)

Let $a \in \mathbb{N}$ with $a \geq 4$, $n \in \mathbb{N}$ with $n \geq N_0$, and Suppose that there exist integers $x, y, z \in \mathbb{N}^*$ such that:

$$\frac{a}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$$

which is equivalent to:

$$\frac{a}{n} - \frac{1}{x} = \frac{1}{y} + \frac{1}{z}$$

which is also equivalent to:

$$\frac{ax - n}{nx} = \frac{y + z}{yz}.$$

For $\frac{y+z}{yz} > 0$, it is necessary that $ax - n > 0$, i.e., $x \geq \left\lceil \frac{n}{a} \right\rceil + 1$. This last equation is equivalent to:

$$yz(ax - n) = nx(y + z).$$

Indeed, for any $a, c \in \mathbb{R}$ and $b, d \in \mathbb{R}^*$, we have:

$$\frac{a}{b} = \frac{c}{d} \iff ad = bc.$$

(This is a fundamental property of fractions.)

Thus, our last equation is equivalent to the following system:

$$(S) \quad \begin{cases} y + z = (ax - n)(2t), \\ yz = nx(2t), \end{cases}$$

for $t \in \mathbb{Q}_+^*$.

By substituting $y + z$ with $(ax - n)(2t)$ and yz with $nx(2t)$ in $yz(ax - n) = nx(y + z)$, the equation remains valid, which shows that this equation implies the system (S).

9.5 Reciprocal Proof

Conversely, let us show that if:

$$\begin{cases} y + z = (ax - n)(2t), \\ yz = nx(2t), \end{cases}$$

for $t \in \mathbb{Q}_+^*$, then it follows that $yz(ax - n) = nx(y + z)$.

Indeed, since $y + z = (ax - n)(2t)$, we obtain:

$$2t = \frac{y + z}{ax - n},$$

noting that $ax - n \neq 0$ since we imposed $x \geq \left\lceil \frac{n}{a} \right\rceil + 1$ to ensure $ax - n > 0$.

On the other hand, since $yz = nx(2t)$, we also have:

$$2t = \frac{yz}{nx}.$$

By the transitivity of equality in $\mathbb{R}$, we deduce:

$$\frac{y+z}{ax-n} = \frac{yz}{nx},$$

which implies:

$$nx(y+z) = yz(ax-n).$$

This proves that system (S) implies $nx(y+z) = yz(ax-n)$.

From the previous paragraph, we have justified the equivalence between $nx(y+z) = yz(ax-n)$ and the system (S) .

Finally, by applying Vieta's formulas given in the above proposition, stated earlier, we conclude that the system (S) is equivalent to stating that y and z are solutions of the following quadratic equation in V :

$$V^2 - (y+z)V + yz = 0,$$

which simplifies to:

$$V^2 - (ax-n)(2t)V + 2nxt = 0.$$

This completes the proof of the equivalence.

10 Summary of the generalized Erdős-Straus and Sierpiński conjecture

We have shown that the generalized Erdős-Straus and Sierpiński conjecture (which includes the classical Erdős-Straus conjecture and the classical Sierpiński conjecture) is true if and only if the quadratic equation:

$$V^2 - (ax-n)(2t)V + 2nxt = 0 \tag{19}$$

always has solutions in $\mathbb{N}^*$ for any $a \in \mathbb{N}$ with $a \geq 4$ and any $n \in \mathbb{N}$ with $n \geq N0$. This reduces the difficulty of proving the conjecture to finding integer solutions (in $\mathbb{N}^*$) for a second-degree equation, whose discriminant is given by:

$$\Delta = 4t^2(ax-n)^2 - 8ntx. \tag{20}$$

Rewriting,

$$\Delta = 4(t^2(ax-n)^2 - 2ntx). \tag{21}$$

As in previous cases, we propose a partial resolution of the conjecture. We seek to solve the equation:

$$V^2 - (ax-n)(2t)V + 2nxt = 0, \tag{22}$$

with discriminant:

$$\Delta = 4t^2(ax-n)^2 - 8ntx. \tag{23}$$

Solving $\Delta > 0$, we obtain the solution range:

$$t > \frac{2nx}{(ax-n)^2}. \tag{24}$$

similary by doing the same manipulations for the t for which $\Delta > 0$

Since:

$$\Delta = 4t^2(ax - n)^2 - 8ntx = 4t(t(ax - n)^2 - 2nx), \quad (25)$$

we set:

$$t = k = t(ax - n)^2 - 2nx, \quad (26)$$

which leads to:

$$k = \frac{2nx}{(ax - n)^2 - 1}, \quad (27)$$

where $k \in \mathbb{Q}_+^*$.

Since:

$$\Delta = 4k^2 = (2k)^2, \quad (28)$$

we obtain:

$$V = \frac{(ax - n)(2k) \pm 2k}{2}. \quad (29)$$

Thus,

$$y = k(ax - n - 1), \quad z = k(ax - n + 1). \quad (30)$$

Replacing k by its expression:

$$y = \frac{2nx}{ax - n + 1}, \quad z = \frac{2nx}{ax - n - 1}. \quad (31)$$

The quantities y and z are integers if and only if $2nx$ is divisible by both $ax - n - 1$ and $ax - n + 1$. in this case x, y, z satisfy:

$$\frac{a}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}. \quad (32)$$

As previously, the condition:

$$x \geq \left\lceil \frac{n+1}{a} \right\rceil + 1 \quad (33)$$

must hold. We will provide numerical computations at the end of this paper to generate certain (a, n) pairs that satisfy this condition.

Similarly, for the second formula:

$$\Delta = 4(t^2(ax - n)^2 - 2ntx), \quad (34)$$

for $\Delta > 0$, we can set:

$$t^2(ax - n)^2 - 2ntx = q^2, \quad q \in \mathbb{N}.$$

so that:

$$\Delta = 4q^2 = (2q)^2. \quad (35)$$

We will now prove that the conjecture is true under the condition that this perfect square exists. We obtain:

$$y = \frac{2t(ax - n) - 2q}{2} = t(ax - n) - q, \quad z = t(ax - n) + q. \quad (36)$$

Verifying,

$$\frac{a}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}. \quad (37)$$

Replacing q^2 by $t^2(ax - n)^2 - 2ntx$.

we will also provide numerical computations of some (a, n) pairs satisfying this condition.

Summary:

we have demonstrated that for all $a \in \mathbb{N}$ with $a \geq 4$ and for all $n \in \mathbb{N}$ with $n \geq 2$, if there exists $x_0 \in \mathbb{N}^*$ such that $ax_0 - n - 1$ and $ax_0 - n + 1$ divide $2nx_0$, then there exist y and z in $\mathbb{N}^*$ such that:

$$y = \frac{2nx_0}{ax_0 - n - 1}, \quad z = \frac{2nx_0}{ax_0 - n + 1}$$

and we have:

$$\frac{a}{n} = \frac{1}{x_0} + \frac{1}{y} + \frac{1}{z}.$$

partially proving the conjecture under this divisibility condition.

Similarly, the second formula allows us to state that for any $a \in \mathbb{N}$ with $a \geq 4$ and for any $n \in \mathbb{N}$ with $n \geq N_0$, if there exist $(x_0, t_0) \in \mathbb{N}^{*2}$ such that:

$$t_0^2(ax_0 - n)^2 - 2nt_0x_0 = q^2, \quad q \in \mathbb{N}.$$

then there exist $y, z \in \mathbb{N}^*$ given by:

$$y = t_0(ax_0 - n) - q, \quad z = t_0(ax_0 - n) + q$$

such that:

$$\frac{a}{n} = \frac{1}{x_0} + \frac{1}{y} + \frac{1}{z}.$$

This also proves the conjecture under this condition.

I once again pose the same questions as for the classical Erdős-Straus conjecture and Sierpiński's conjecture to the mathematical community in order to definitively resolve the generalized Erdős-Straus and Sierpiński's conjecture.

11 Comparison with Mordell's Approach

Mordell's method provided a partial resolution of the Erdős-Straus conjecture by showing that for all integers n such that

$$n \not\equiv 1, 11^2, 13^2, 17^2, 19^2, 23^2 \pmod{840},$$

the conjecture holds. However, for the exceptional cases left unresolved by Mordell, our second formula proves to be a powerful tool for generating solutions.

Through numerical testing, we observed that a direct implementation of the second formula, applied to the values of n not covered by Mordell's result in the range $2 \leq n \leq 10\,000$, successfully produces solutions for all such n . In this interval, this corresponds to a numerical success rate of 100%.

Remark: This observation is purely experimental and does not constitute a proof. However, it strongly suggests that our approach captures the full underlying structure of the original Erdős-Straus equation. This is very likely due to the equivalence established earlier between the original Diophantine equation and our associated quadratic parametric equation, from which integer solutions are obtained directly via the analysis of its discriminant.

Moreover, since the Erdős–Straus conjecture has already been verified by other authors for all $n \leq 10^{17}$, it follows from the proven equivalence that our quadratic equation must necessarily admit integer solutions at least up to this bound in the classical case $a = 4$. From this perspective, it is therefore not surprising that our formula successfully produces solutions for the values of n left unresolved by Mordell, in particular for all n in the interval $[2, 10\,000]$.

For completeness and reproducibility, we provide in the appendix the numerical codes used to test the cases not covered by Mordell’s result, as well as to explore all values of n in user-specified intervals. These codes apply more generally to all fixed integers $a \geq 4$, including the classical Erdős–Straus case ($a = 4$), the Sierpiński case ($a = 5$), and their natural generalizations ($a \geq 6$).

12 Testing Second Formula for Mordell’s Exceptional Cases between $[2, 5000]$

Mordell’s exceptional cases in the given range are:

[121, 169, 289, 361, 529, 841, 961, 1009, 1129, 1201, 1369, 1681, 1801, 1849, 1969,
2041, 2209, 2521, 2641, 2689, 2809, 2881, 3049, 3361, 3481, 3529, 3649,
3721, 3889, 4201, 4321, 4369, 4489, 4561, 4729].

Total exceptional cases: 35

n	x	t	y	z
121	31	1078	1694	4774
169	44	325	1690	2860
289	75	510	2550	8670
361	92	1368	8664	10488
529	133	15778	42826	51842
841	220	319	6380	18502
961	248	496	15376	15376
1009	260	1009	10090	52468
1129	2964	104	312	2230904
1201	9548	775	310	57335740
1369	348	1813	38332	45066
1681	423	11808	121032	138744
1801	35720	342	456	96497580
1849	468	3483	60372	99846
1969	493	215818	633302	661606
2041	516	4239	73476	121518
2209	564	1128	53016	53016
2521	18908	163	652	23833534
2641	665	9730	184870	184870
2689	9412	181	724	12654434
2809	710	4240	112360	150520
2881	724	18963	242004	326886
3049	75768	275	770	165011880
3361	73525	34	850	19769402
3481	885	1770	104430	104430
3529	27056	342	912	71610468
3649	915	55432	569244	650260
3721	943	2806	115046	171166
3889	40836	83	996	26468534
4201	1056	16804	369688	403296
4321	1085	25984	483952	503440
4369	1104	4369	200974	209712
4489	1139	2278	152626	152626
4561	15964	307	1228	36405902
4729	244728	66	1188	128590968

The above table shows that the second formula successfully produces valid integer solutions for all Mordell exceptional values of n in the interval $[2, 5000]$. Consequently, a numerical success rate of 100% is achieved on this entire set of previously unresolved cases.

13 Testing Second Formula for Mordell's Exceptional Cases between [5000,10000]

Mordell's exceptional cases in the given range are:

[5041, 5161, 5209, 5329, 5401, 5569, 5881, 6001, 6049, 6169, 6241, 6409, 6721,
6841, 6889, 7009, 7081, 7249, 7561, 7681, 7729, 7849, 7921, 8089,
8401, 8521, 8569, 8689, 8761, 8929, 9241, 9361, 9409, 9529, 9601, 9769].

Total exceptional cases: 36.

n	x	t	y	z
5041	1264	56800	806560	897440
5161	1314	1588	115924	185796
5209	96888	275	1320	210287330
5329	1460	73	21316	53290
5401	1353	120786	1328646	1328646
5569	1112412	41	1394	364413084
5881	92628	83	1494	60527252
6001	1520	3177	180030	321936
6049	1518	34716	798468	798468
6169	1544	388864	2680384	2763712
6241	1580	3160	249640	249640
6409	1612	13600	512720	548080
6721	1683	186966	2056626	2056626
6841	32923	328	1804	81900452
6889	1743	3486	289338	289338
7009	1755	203424	2186808	2288520
7081	1815	1056	96360	281688
7249	1815	217470	2392170	2392170
7561	368600	190	1900	557396920
7681	47047	286	2002	103248002
7729	1935	247328	2658776	2782440
7849	1974	14028	659316	659316
7921	2136	89	47526	63368
8089	139536	342	2052	376235568
8401	2108	36856	1142536	1142536
8521	332320	268	2144	707924680
8569	2145	303810	3341910	3341910
8689	2184	17378	729876	903656
8761	2208	8761	403006	841056
8929	426921	88	2244	298978636
9241	181935	104	2340	149445452
9361	2343	362526	3987786	3987786
9409	2425	970	94090	470450
9529	2398	11728	639176	838552
9601	327639	62	2418	161316002
9769	2460	9769	586140	801058

The above table shows that the second formula successfully produces valid integer solutions for all Mordell exceptional values of n in the interval $[5000, 10000]$. Consequently, a numerical success rate of 100% is achieved on this entire set of previously unresolved cases.

Taken together, the two preceding tables show that our second formula captures 100% of the integers left unresolved by Mordell for all values of n in the interval $[2, 10000]$.

Any reader wishing to verify the correctness of these numerical simulations may proceed as follows: for a given value of n and the corresponding triple (x, y, z) provided in the tables, it suffices to check the identity

$$4xyz = n(xy + xz + yz),$$

which is strictly equivalent to the original Erdős–Straus equation

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}.$$

For large values of n , it is strongly preferable to verify solutions using the polynomial identity $4xyz = n(xy + xz + yz)$ rather than the fractional equation itself. Indeed, the latter involves floating-point arithmetic and may suffer from numerical round-off errors, potentially preventing exact equality from being detected by the machine. In contrast, the polynomial form relies solely on integer arithmetic and therefore provides an exact, unambiguous mathematical verification.

In the following tables, we provide the numerical values of some n , x_0 , y , and z that satisfy the conditions of each formula.

The value L (for “Left”) represents the ratio a/n , while R (for “Right”) is computed as $R = \frac{1}{x_0} + \frac{1}{y} + \frac{1}{z}$ (we use the fractional form here because the values of n tested in the tables below are relatively small, and therefore no floating-point precision issues arise).

The last column verifies whether $L = R$:

- Rows labeled **"True"** indicate equality, meaning the conjecture holds for the given values of n , x_0 , y , and z .
- Rows labeled **"False"** indicate that the conjecture does not hold for those specific values.

If a value of n is not displayed in the table, this does **not** mean that the number does not satisfy the formula or the conjecture. It simply indicates that the algorithm did not find an x_0 and/or a t_0 within the specific search range that meets one of the formula’s conditions for that n . This is due to the fact that there are infinitely many possible choices for x_0 .

According to the equivalence established below between the original equation

$$\frac{a}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$$

and the parametric quadratic equation

$$V^2 - 2t(ax - n)V + 2nxt = 0,$$

the case $a = 4$ (classical Erdős–Straus conjecture) reassures us that this quadratic equation admits integer solutions at least up to $n = 10^{17}$, which is the current numerical verification threshold for the Erdős–Straus conjecture.

For $a = 5$ (classical Sierpiński case), we also know today that very high numerical verifications have been carried out. I personally performed quick numerical checks for some values of $a \in \{6, 7, 8\}$ up to reasonable thresholds (which I do not officially report in this article, as these were rapid tests), and the equation consistently appears to admit integer solutions via the study of its discriminant

$$\Delta = 4t^2(ax - n)^2 - 8nxt = 4(t^2(ax - n)^2 - 2nxt),$$

which seems always to contain a perfect square, thus providing integer solutions to the quadratic equation, and by equivalence, to the generalized Erdős–Straus equation. Since no proof is currently known, these observations lead us to state the following conjectures.

Conjecture 1 (strong version, integer t). For every integer $n \geq 2$, there exists $(x, t) \in \mathbb{N}^{*2}$ such that

$$t^2(4x - n)^2 - 2nxt = q^2, \quad q \in \mathbb{N}.$$

Conjecture 2 (weak version, rational t , not necessarily integer). For every integer $n \geq 2$, there exists $(x, t) \in \mathbb{N}^* \times \mathbb{Q}_+^*$ such that

$$t^2(4x - n)^2 - 2nxt = q^2, \quad q \in \mathbb{Q}_+.$$

Conjecture 3 (weak version, rational t , not necessarily integer). For every fixed integer $a \geq 5$, there exists $N_0 \geq 2$ in $\mathbb{N}^*$ (with N_0 not strictly smaller than $a/3$) such that for all $n \geq N_0$, there exists

$$(x, t) \in \mathbb{N}^* \times \mathbb{Q}_+^* \quad \text{with} \quad t^2(ax - n)^2 - 2nxt = q^2, \quad q \in \mathbb{Q}_+.$$

Conjecture 4 (strong version, integer t). For every fixed integer $a \geq 5$, there exists $N_1 \geq N_0$ (N_0 is the one given in Conjecture 3), with N_1 not strictly smaller than $a/3$, such that for all $n \geq N_1$, there exists

$$(x, t) \in \mathbb{N}^{*2} \quad \text{such that} \quad t^2(ax - n)^2 - 2nxt = q^2, \quad q \in \mathbb{N}.$$

Conjecture 1 is equivalent to the classical Erdős–Straus conjecture. Indeed, under Conjecture 1, the integer q satisfies

$$q^2 = t^2(4x - n)^2 - 2nxt,$$

and the variables y and z are given by

$$y = t(4x - n) - q, \quad z = t(4x - n) + q.$$

It is immediate that $z \in \mathbb{N}^*$, since $4x - n > 0$ because

$$x \geq \left\lfloor \frac{n}{4} \right\rfloor + 1 > \frac{n}{4},$$

and with $t \in \mathbb{N}^*$, $x \in \mathbb{N}^*$, $n \in \mathbb{N}^*$ and $q \in \mathbb{N}$, we have $z \in \mathbb{N}^*$.

Let us now show that $y \in \mathbb{N}^*$. From

$$q^2 = t^2(4x - n)^2 - 2nxt < t^2(4x - n)^2,$$

since $-2nxt < 0$ for $n, x, t > 0$, and since the square root function is increasing on $\mathbb{R}_+^*$, we obtain

$$q < t(4x - n).$$

Hence $t(4x - n) - q > 0$, and since t, x, n, q are integers, it follows that

$$y = t(4x - n) - q \in \mathbb{N}^*.$$

We have therefore rigorously shown that whenever the condition

$$t^2(4x - n)^2 - 2nxt = q^2, \quad q \in \mathbb{N},$$

is satisfied (Conjecture 1), one can automatically construct $y, z \in \mathbb{N}^*$ solving the quadratic equation

$$V^2 - 2t(4x - n)V + 2nxt = 0,$$

which, by the equivalence theorem proved earlier, is equivalent to the original Erdős–Straus equation

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}.$$

Thus, Conjecture 1 is rigorously equivalent to the classical Erdős–Straus conjecture.

Using the same reasoning, one also proves that Conjecture 4 is equivalent to the generalized Sierpiński conjecture ($a \geq 5$) for all $n \geq N_1$, where N_1 is the minimal threshold beyond which the expression

$$t^2(ax - n)^2 - 2nxt$$

always admits an integer perfect square (under Conjecture 4).

Conjecture 2 is stated for researchers interested in working over $\mathbb{Q}_+^*$, either to enlarge the search space for t , or because it may be useful in other research contexts beyond Egyptian fractions.

Conjecture 3 is particularly useful for numerical verification of the generalized Sierpiński conjecture ($a \geq 5$) for values of n in the finite interval $[N_0, N_1]$, which is often not too large according to our simulations, since Conjecture 4 addresses the problem for all $n \geq N_1$.

Finally, why do we exclude values $n < a/3$? Since $x, y, z \in \mathbb{N}^*$, we have

$$\frac{1}{x} + \frac{1}{y} + \frac{1}{z} \leq 3,$$

which implies $a/n \leq 3$, and hence $n \geq a/3$. Therefore, the decomposition of a/n as a sum of three Egyptian fractions is impossible for all integers $n < a/3$. Consequently, Conjecture 4 cannot hold for such values of n , as it is equivalent to the generalized Sierpiński conjecture.

Similarly, even if Conjecture 3 produces rational perfect squares for $n < a/3$, these cannot yield integer solutions to the quadratic equation

$$V^2 - 2t(ax - n)V + 2nxt = 0,$$

since, by the equivalence theorem, this quadratic equation admits no integer solutions when $n < a/3$, because the equivalent equation

$$\frac{a}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$$

has no integer solutions in that range.

Table 1: Results with the first formula of the Erdős-Straus conjecture test for n from 2 to 90

n	x	y	z	L	R	L = R (verification)
3	2	3	2	1.3333	1.3333	True
5	2	10	5	0.8000	0.8000	True
6	2	24	8	0.6667	0.6667	True
9	4	12	9	0.4444	0.4444	True
10	3	60	20	0.4000	0.4000	True
13	4	52	26	0.3077	0.3077	True
14	5	28	20	0.2857	0.2857	True
18	5	180	60	0.2222	0.2222	True
19	6	57	38	0.2105	0.2105	True
20	6	80	48	0.2000	0.2000	True
21	6	126	63	0.1905	0.1905	True
22	6	264	88	0.1818	0.1818	True
27	8	108	72	0.1481	0.1481	True
28	9	72	56	0.1429	0.1429	True
29	8	232	116	0.1379	0.1379	True
30	8	480	160	0.1333	0.1333	True
34	9	612	204	0.1176	0.1176	True
35	12	70	60	0.1143	0.1143	True
36	10	240	144	0.1111	0.1111	True
37	10	370	185	0.1081	0.1081	True
39	16	52	48	0.1026	0.1026	True
41	12	164	123	0.0976	0.0976	True
42	11	924	308	0.0952	0.0952	True
43	12	258	172	0.0930	0.0930	True
45	12	540	270	0.0889	0.0889	True
46	12	1104	368	0.0870	0.0870	True
49	16	112	98	0.0816	0.0816	True
50	14	280	200	0.0800	0.0800	True
51	14	357	238	0.0784	0.0784	True
53	14	742	371	0.0755	0.0755	True
54	14	1512	504	0.0741	0.0741	True
55	16	220	176	0.0727	0.0727	True
56	15	560	336	0.0714	0.0714	True
57	16	304	228	0.0702	0.0702	True
58	15	1740	580	0.0690	0.0690	True
60	16	640	384	0.0667	0.0667	True
61	16	976	488	0.0656	0.0656	True
66	17	2244	748	0.0606	0.0606	True
67	18	603	402	0.0597	0.0597	True
69	18	1242	621	0.0580	0.0580	True
70	18	2520	840	0.0571	0.0571	True
71	20	355	284	0.0563	0.0563	True

n	x	y	z	L	R	L = R (verification)
75	20	750	500	0.0533	0.0533	True
77	20	1540	770	0.0519	0.0519	True
78	20	3120	1040	0.0513	0.0513	True
80	21	1120	672	0.0500	0.0500	True
82	21	3444	1148	0.0488	0.0488	True
85	22	1870	935	0.0471	0.0471	True
89	24	712	534	0.0449	0.0449	True
90	23	4140	1380	0.0444	0.0444	True

Table 2: Results of Erdős–Straus Conjecture with the second formula testing for n from 2 to 50

n	x	y	z	L	R	L = R (verification)
2	1	2	2	2.0	2.0	True
3	1	6	6	1.3333	1.3333	True
4	2	4	4	1.0	1.0	True
5	2	4	20	0.8	0.8	True
6	2	12	12	0.6667	0.6667	True
7	2	28	28	0.5714	0.5714	True
8	3	12	12	0.5	0.5	True
9	3	18	18	0.4444	0.4444	True
10	3	30	30	0.4	0.4	True
11	3	66	66	0.3636	0.3636	True
12	4	24	24	0.3333	0.3333	True
13	4	26	52	0.3077	0.3077	True
14	4	56	56	0.2857	0.2857	True
15	4	120	120	0.2667	0.2667	True
16	5	40	40	0.25	0.25	True
17	5	34	170	0.2353	0.2353	True
18	5	90	90	0.2222	0.2222	True
19	5	190	190	0.2105	0.2105	True
20	6	60	60	0.2	0.2	True
21	6	84	84	0.1905	0.1905	True
22	6	132	132	0.1818	0.1818	True
23	6	276	276	0.1739	0.1739	True
24	7	84	84	0.1667	0.1667	True
25	7	100	140	0.16	0.16	True
26	7	182	182	0.1538	0.1538	True
27	7	378	378	0.1481	0.1481	True
28	8	112	112	0.1429	0.1429	True
29	8	116	232	0.1379	0.1379	True
30	8	240	240	0.1333	0.1333	True
31	8	496	496	0.129	0.129	True
32	9	144	144	0.125	0.125	True
33	9	198	198	0.1212	0.1212	True
34	9	306	306	0.1176	0.1176	True

n	x	y	z	L	R	L = R (verification)
35	9	630	630	0.1143	0.1143	True
36	10	180	180	0.1111	0.1111	True
37	10	148	740	0.1081	0.1081	True
38	10	380	380	0.1053	0.1053	True
39	10	780	780	0.1026	0.1026	True
40	11	220	220	0.1	0.1	True
41	12	82	492	0.0976	0.0976	True
42	11	462	462	0.0952	0.0952	True
43	11	946	946	0.093	0.093	True
44	12	264	264	0.0909	0.0909	True
45	12	360	360	0.0889	0.0889	True
46	12	552	552	0.087	0.087	True
47	12	1128	1128	0.0851	0.0851	True
48	13	312	312	0.0833	0.0833	True
49	14	196	196	0.0816	0.0816	True
50	13	650	650	0.08	0.08	True

Table 3: Results of Erdős–Straus Conjecture with the second formula testing for n from 51 to 100

n	x	y	z	L	R	L = R (verification)
51	13	1326	1326	0.0784	0.0784	True
52	14	364	364	0.0769	0.0769	True
53	16	80	2120	0.0755	0.0755	True
54	14	756	756	0.0741	0.0741	True
55	14	1540	1540	0.0727	0.0727	True
56	15	420	420	0.0714	0.0714	True
57	15	570	570	0.0702	0.0702	True
58	15	870	870	0.069	0.069	True
59	15	1770	1770	0.0678	0.0678	True
60	16	480	480	0.0667	0.0667	True
61	16	488	976	0.0656	0.0656	True
62	16	992	992	0.0645	0.0645	True
63	16	2016	2016	0.0635	0.0635	True
64	17	544	544	0.0625	0.0625	True
65	17	650	850	0.0615	0.0615	True
66	17	1122	1122	0.0606	0.0606	True
67	17	2278	2278	0.0597	0.0597	True
68	18	612	612	0.0588	0.0588	True
69	18	828	828	0.058	0.058	True
70	18	1260	1260	0.0571	0.0571	True
71	18	2556	2556	0.0563	0.0563	True
72	19	684	684	0.0556	0.0556	True
73	20	292	730	0.0548	0.0548	True
74	19	1406	1406	0.0541	0.0541	True
75	19	2850	2850	0.0533	0.0533	True

n	x	y	z	L	R	L = R (verification)
76	20	760	760	0.0526	0.0526	True
77	20	980	1078	0.0519	0.0519	True
78	20	1560	1560	0.0513	0.0513	True
79	20	3160	3160	0.0506	0.0506	True
80	21	840	840	0.05	0.05	True
81	21	1134	1134	0.0494	0.0494	True
82	21	1722	1722	0.0488	0.0488	True
83	21	3486	3486	0.0482	0.0482	True
84	22	924	924	0.0476	0.0476	True
85	23	340	1564	0.0471	0.0471	True
86	22	1892	1892	0.0465	0.0465	True
87	22	3828	3828	0.046	0.046	True
88	23	1012	1012	0.0455	0.0455	True
89	24	534	712	0.0449	0.0449	True
90	23	2070	2070	0.0444	0.0444	True
91	23	4186	4186	0.044	0.044	True
92	24	1104	1104	0.0435	0.0435	True
93	24	1488	1488	0.043	0.043	True
94	24	2256	2256	0.0426	0.0426	True
95	24	4560	4560	0.0421	0.0421	True
96	25	1200	1200	0.0417	0.0417	True
97	28	194	2716	0.0412	0.0412	True
98	25	2450	2450	0.0408	0.0408	True
99	25	4950	4950	0.0404	0.0404	True
100	26	1300	1300	0.04	0.04	True

Table 4: Results of Erdős–Straus Conjecture with second formula testing for n from 101 to 150

n	x	y	z	L	R	L = R (verification)
101	38	76	7676	0.0396	0.0396	True
102	26	2652	2652	0.0392	0.0392	True
103	26	5356	5356	0.0388	0.0388	True
104	27	1404	1404	0.0385	0.0385	True
105	27	1890	1890	0.0381	0.0381	True
106	27	2862	2862	0.0377	0.0377	True
107	27	5778	5778	0.0374	0.0374	True
108	28	1512	1512	0.037	0.037	True
109	28	1526	3052	0.0367	0.0367	True
110	28	3080	3080	0.0364	0.0364	True
111	28	6216	6216	0.036	0.036	True
112	29	1624	1624	0.0357	0.0357	True
113	132	36	22374	0.0354	0.0354	True
114	29	3306	3306	0.0351	0.0351	True
115	29	6670	6670	0.0348	0.0348	True
116	30	1740	1740	0.0345	0.0345	True

n	x	y	z	L	R	L = R (verification)
117	30	2340	2340	0.0342	0.0342	True
118	30	3540	3540	0.0339	0.0339	True
119	30	7140	7140	0.0336	0.0336	True
120	31	1860	1860	0.0333	0.0333	True
121	33	726	726	0.0331	0.0331	True
122	31	3782	3782	0.0328	0.0328	True
123	31	7626	7626	0.0325	0.0325	True
124	32	1984	1984	0.0323	0.0323	True
125	32	2400	3000	0.032	0.032	True
126	32	4032	4032	0.0317	0.0317	True
127	32	8128	8128	0.0315	0.0315	True
128	33	2112	2112	0.0312	0.0312	True
129	33	2838	2838	0.031	0.031	True
130	33	4290	4290	0.0308	0.0308	True
131	33	8646	8646	0.0305	0.0305	True
132	34	2244	2244	0.0303	0.0303	True
133	34	2856	3192	0.0301	0.0301	True
134	34	4556	4556	0.0299	0.0299	True
135	34	9180	9180	0.0296	0.0296	True
136	35	2380	2380	0.0294	0.0294	True
137	35	2740	3836	0.0292	0.0292	True
138	35	4830	4830	0.029	0.029	True
139	35	9730	9730	0.0288	0.0288	True
140	36	2520	2520	0.0286	0.0286	True
141	36	3384	3384	0.0284	0.0284	True
142	36	5112	5112	0.0282	0.0282	True
143	36	10296	10296	0.028	0.028	True
144	37	2664	2664	0.0278	0.0278	True
145	37	3190	4070	0.0276	0.0276	True
146	37	5402	5402	0.0274	0.0274	True
147	37	10878	10878	0.0272	0.0272	True
148	38	2812	2812	0.027	0.027	True
149	70	80	16688	0.0268	0.0268	True
150	38	5700	5700	0.0267	0.0267	True

Table 5: Results of Erdős–Straus Conjecture with second formula testing for n from 150 to 200

n	x	y	z	L	R	L = R (verification)
151	38	11476	11476	0.0265	0.0265	True
152	39	2964	2964	0.0263	0.0263	True
153	39	3978	3978	0.0261	0.0261	True
154	39	6006	6006	0.026	0.026	True
155	39	12090	12090	0.0258	0.0258	True
156	40	3120	3120	0.0256	0.0256	True
157	40	3768	4710	0.0255	0.0255	True

n	x	y	z	L	R	L = R (verification)
158	40	6320	6320	0.0253	0.0253	True
159	40	12720	12720	0.0252	0.0252	True
160	41	3280	3280	0.025	0.025	True
161	42	1932	1932	0.0248	0.0248	True
162	41	6642	6642	0.0247	0.0247	True
163	41	13366	13366	0.0245	0.0245	True
164	42	3444	3444	0.0244	0.0244	True
165	42	4620	4620	0.0242	0.0242	True
166	42	6972	6972	0.0241	0.0241	True
167	42	14028	14028	0.024	0.024	True
168	43	3612	3612	0.0238	0.0238	True
169	44	1690	2860	0.0237	0.0237	True
170	43	7310	7310	0.0235	0.0235	True
171	43	14706	14706	0.0234	0.0234	True
172	44	3784	3784	0.0233	0.0233	True
173	45	1730	3114	0.0231	0.0231	True
174	44	7656	7656	0.023	0.023	True
175	44	15400	15400	0.0229	0.0229	True
176	45	3960	3960	0.0227	0.0227	True
177	45	5310	5310	0.0226	0.0226	True
178	45	8010	8010	0.0225	0.0225	True
179	45	16110	16110	0.0223	0.0223	True
180	46	4140	4140	0.0222	0.0222	True
181	48	1086	2896	0.0221	0.0221	True
182	46	8372	8372	0.022	0.022	True
183	46	16836	16836	0.0219	0.0219	True
184	47	4324	4324	0.0217	0.0217	True
185	47	5180	6580	0.0216	0.0216	True
186	47	8742	8742	0.0215	0.0215	True
187	47	17578	17578	0.0214	0.0214	True
188	48	4512	4512	0.0213	0.0213	True
189	48	6048	6048	0.0212	0.0212	True
190	48	9120	9120	0.0211	0.0211	True
191	48	18336	18336	0.0209	0.0209	True
192	49	4704	4704	0.0208	0.0208	True
193	676	52	65234	0.0207	0.0207	True
194	49	9506	9506	0.0206	0.0206	True
195	49	19110	19110	0.0205	0.0205	True
196	50	4900	4900	0.0204	0.0204	True
197	148	74	29156	0.0203	0.0203	True
198	50	9900	9900	0.0202	0.0202	True
199	50	19900	19900	0.0201	0.0201	True
200	51	5100	5100	0.02	0.02	True

Table 6: Results of Sierpiński Conjecture with second formula testing for n from 2 to 45

n	x	t	y	z	F	q
2	1	$\frac{1}{2}$	1	2	$\frac{1}{4}$	$\frac{1}{2}$
3	1	2	2	6	4	2
4	1	8	8	8	0	0
5	2	$\frac{4}{5}$	4	4	0	0
6	2	2	4	12	16	4
7	5	2	2	70	1156	34
8	2	8	16	16	0	0
9	2	36	36	36	0	0
10	3	$\frac{5}{2}$	10	15	$\frac{25}{4}$	$\frac{5}{2}$
11	5	8	4	220	11664	108
12	3	8	24	24	0	0
13	3	26	26	78	676	26
14	3	84	84	84	0	0
15	4	5	20	30	25	5
16	4	8	32	32	0	0
17	4	17	34	68	289	17
18	4	36	72	72	0	0
19	4	152	152	152	0	0
20	5	8	40	40	0	0
21	5	14	42	70	196	14
22	5	44	44	220	7744	88
23	5	98	70	322	15876	126
24	5	240	240	240	0	0
25	6	12	60	60	0	0
26	6	26	52	156	2704	52
27	6	36	108	108	0	0
28	6	84	168	168	0	0
29	6	348	348	348	0	0
30	7	20	60	140	1600	40
31	7	62	62	434	34596	186
32	7	50	140	160	100	10
33	7	128	176	336	6400	80
34	7	476	476	476	0	0
35	8	35	70	280	11025	105
36	8	36	144	144	0	0
37	8	74	148	296	5476	74
38	8	152	304	304	0	0
39	8	624	624	624	0	0

n	x	t	y	z	F	q
40	9	30	120	180	900	30
41	12	169	26	6396	10144225	3185
42	9	84	252	252	0	0
43	9	258	258	774	66564	258
44	9	792	792	792	0	0
45	10	36	180	180	0	0

Table 7: Results of Sierpiński Conjecture with second formula testing for n from 46 to 89

n	x	t	y	z	F	q
46	10	98	140	644	63504	252
47	10	188	188	940	141376	376
48	10	240	480	480	0	0
49	10	980	980	980	0	0
50	11	44	220	220	0	0
51	12	17	102	204	2601	51
52	11	128	352	416	1024	32
53	11	512	352	1696	451584	672
54	11	1188	1188	1188	0	0
55	12	55	220	330	3025	55
56	12	84	336	336	0	0
57	12	152	456	456	0	0
58	12	348	696	696	0	0
59	12	1416	1416	1416	0	0
60	13	65	260	390	4225	65
61	15	242	66	6710	11035684	3322
62	14	62	124	868	138384	372
63	13	416	728	936	10816	104
64	13	1664	1664	1664	0	0
65	14	80	280	520	14400	120
66	14	128	352	672	25600	160
67	15	134	134	2010	879844	938
68	14	476	952	952	0	0
69	14	1932	1932	1932	0	0
70	15	84	420	420	0	0
71	15	142	426	710	20164	142
72	15	240	720	720	0	0
73	15	584	876	1460	85264	292
74	15	2220	2220	2220	0	0
75	16	96	480	480	0	0

n	x	t	y	z	F	q
76	16	152	608	608	0	0
77	16	275	770	880	3025	55
78	16	624	1248	1248	0	0
79	16	2528	2528	2528	0	0
80	17	125	400	850	50625	225
81	18	36	324	324	0	0
82	24	169	52	12792	40576900	6370
83	20	125	100	4150	4100625	2025
84	17	2856	2856	2856	0	0
85	18	240	360	2040	705600	840
86	18	258	516	1548	266256	516
87	18	348	1044	1044	0	0
88	18	792	1584	1584	0	0
89	18	3204	3204	3204	0	0

Table 8: Results of the generalized conjecture ($a = 6$) with second formula, testing for n from 2 to 45

n	x	t	y	z	F	q
2	1	1/4	1	1	0	0
3	1	2/3	2	2	0	0
4	1	2	4	4	0	0
5	1	10	10	10	0	0
6	2	2/3	4	4	0	0
7	3	2	2	42	400	20
8	2	2	8	8	0	0
9	2	4	12	12	0	0
10	2	10	20	20	0	0
11	2	44	44	44	0	0
12	3	2	12	12	0	0
13	3	32	8	312	23104	152
14	3	7	14	42	196	14
15	3	10	30	30	0	0
16	3	24	48	48	0	0
17	3	102	102	102	0	0
18	4	4	24	24	0	0
19	4	32	16	304	20736	144
20	4	10	40	40	0	0
21	4	21	42	84	441	21
22	4	44	88	88	0	0
23	4	184	184	184	0	0

n	x	t	y	z	F	q
24	5	12	24	120	2304	48
25	5	10	50	50	0	0
26	6	32	16	624	92416	304
27	5	30	90	90	0	0
28	5	70	140	140	0	0
29	5	290	290	290	0	0
30	6	10	60	60	0	0
31	39	2	6	806	160000	400
32	6	24	96	96	0	0
33	6	44	132	132	0	0
34	6	102	204	204	0	0
35	6	420	420	420	0	0
36	7	14	84	84	0	0
37	27	32	8	7992	15936064	3992
38	7	76	76	532	51984	228
39	8	54	36	936	202500	450
40	7	140	280	280	0	0
41	7	574	574	574	0	0
42	8	21	84	168	1764	42
43	8	43	86	344	16641	129
44	8	44	176	176	0	0
45	8	80	240	240	0	0

Table 9: Results of the generalized conjecture ($a = 11$)
with first formula testing for n from 500 to 625

n	x	y	z	L	R	L = R
501	46	11523	7682	0.021956088	0.021956088	True
503	46	23138	11569	0.021868787	0.021868787	True
504	46	46368	15456	0.021825397	0.021825397	True
510	49	1785	1666	0.021568627	0.021568627	True
513	50	1425	1350	0.021442495	0.021442495	True
514	47	24158	12079	0.021400778	0.021400778	True
521	48	8336	6252	0.021113244	0.021113244	True
523	48	12552	8368	0.021032505	0.021032505	True
525	48	25200	12600	0.020952381	0.020952381	True
526	48	50496	16832	0.020912548	0.020912548	True
528	51	1683	1584	0.020833333	0.020833333	True
531	49	7434	5782	0.020715631	0.020715631	True
534	49	13083	8722	0.020599251	0.020599251	True
536	49	26264	13132	0.020522388	0.020522388	True

n	x	y	z	L	R	L = R
537	49	52626	17542	0.020484171	0.020484171	True
545	51	3706	3270	0.020183486	0.020183486	True
546	50	18200	10920	0.020146520	0.020146520	True
547	50	27350	13675	0.020109689	0.020109689	True
550	51	5610	4675	0.020000000	0.020000000	True
556	51	14178	9452	0.019784173	0.019784173	True
558	51	28458	14229	0.019713262	0.019713262	True
559	51	57018	19006	0.019677996	0.019677996	True
567	52	14742	9828	0.019400353	0.019400353	True
569	52	29588	14794	0.019332162	0.019332162	True
570	52	59280	19760	0.019298246	0.019298246	True
575	54	3450	3105	0.019130435	0.019130435	True
576	53	10176	7632	0.019097222	0.019097222	True
580	53	30740	15370	0.018965517	0.018965517	True
581	54	5229	4482	0.018932874	0.018932874	True
584	55	3212	2920	0.018835616	0.018835616	True
589	54	15903	10602	0.018675722	0.018675722	True
590	54	21240	12744	0.018644068	0.018644068	True
591	54	31914	15957	0.018612521	0.018612521	True
592	54	63936	21312	0.018581081	0.018581081	True
594	55	6534	5445	0.018518519	0.018518519	True
596	55	8195	6556	0.018456376	0.018456376	True
600	55	16500	11000	0.018333333	0.018333333	True
601	56	4808	4207	0.018302829	0.018302829	True
602	55	33110	16555	0.018272425	0.018272425	True
603	55	66330	22110	0.018242123	0.018242123	True
609	56	11368	8526	0.018062397	0.018062397	True
610	56	13664	9760	0.018032787	0.018032787	True
612	65	780	765	0.017973856	0.017973856	True
613	56	34328	17164	0.017944535	0.017944535	True
620	57	11780	8835	0.017741935	0.017741935	True
622	57	17727	11818	0.017684887	0.017684887	True
624	57	35568	17784	0.017628205	0.017628205	True

14 References

References

- [1] **Wikipédia**, *Conjecture d'Erdős-Straus*, Disponible à : https://fr.m.wikipedia.org/wiki/Conjecture_d'Erd%C5%91s-Straus
- [2] **P. Erdős**, *On a diophantine equation*, Mat. Lapok, vol. 1, pp. 192-210, 1950.
- [3] **L. A. Rosati**, *Sull'equazione diofantea $4/n = 1/x_1 + 1/x_2 + 1/x_3$* , Boll. Un. Mat. Ital., vol. 9, pp. 59-63, 1954.
- [4] **M. A. Lopez**, *A Complete Congruence System for the Erdős-Straus Conjecture*, arXiv, 2024. Disponible à : <https://arxiv.org/abs/2404.01508>
- [5] **I. V. Morozov**, *Approaches to the Erdős-Straus Conjecture*, CUNY Academic Works, 2022. Disponible à : https://academicworks.cuny.edu/ny_pubs/1065/
- [6] **M. Gionfriddo**, *A short proof of Erdős-Straus conjecture for every $n \equiv 13 \pmod{24}$* , ResearchGate, 2021.
- [7] **Wikipédia**, *Fraction égyptienne*, Disponible à : https://fr.wikipedia.org/wiki/Fraction_%C3%A9gyptienne

Appendix: Python Code for Testing the Generalized Erdős-Straus Conjecture (First Formula)

The following code allows testing the generalized Erdős-Straus conjecture numerically for different values of a and n . It provides direct output of the results without generating a CSV file.

```
1 # Python implementation of the first formula for the Generalized Erdos-
   Straus Conjecture
2
3
4 def generalized_erdos_straus_conjecture(a, n_min, n_max):
5     """
6     Tests the generalized Erdos-Straus conjecture for a >= 4.
7     .
8     This function prints results directly instead of saving them to a
       file.
9
10    :param a: Integer, the numerator in the equation a/n = 1/x + 1/y +
       1/z.
11    :param n_min: Integer, the starting value of n.
12    :param n_max: Integer, the ending value of n.
13    """
14    epsilon = 1e-9 # Tolerance for floating-point comparison
15    for n in range(n_min, n_max + 1):
16        solution_found = False
17        L = a / n # Left-hand side of the equation
18        for x in range(n // a + 1, 10 * n): # Searching for x > n/a
19            lambda_value = a * x - n - 1
```

```

20     if lambda_value <= 0:
21         continue
22     y = (2 * n * x) // lambda_value
23     z = (2 * n * x) // (lambda_value + 2)
24
25     # Ensure y and z are integers
26     if (2 * n * x) % lambda_value == 0 and (2 * n * x) % (
27         lambda_value + 2) == 0:
28         R = 1/x + 1/y + 1/z # Right-hand side of the equation
29         is_true = abs(L - R) < epsilon # Floating-point
30         comparison with tolerance
31         print(f"For n = {n}, x = {x}, y = {y}, z = {z}: L = {L
32             :.4f}, R = {R:.4f}, Conjecture holds: {is_true}")
33         solution_found = True
34         break
35
36 # Example usage:
37 a = 4 #(for the classic Erdos_Strauss Conjecture)
38 n_min = 2
39 n_max = 100
40 generalized_erdos_straus_conjecture(a, n_min, n_max)

```

Appendix: Python Code for Testing the Generalized Erdős-Straus Conjecture (Second Formula)

The following code tests the generalized Erdős-Straus conjecture using the second formula independently. It prints results directly based on the existence of a perfect square.

```
1 import math
2 import time
3 from fractions import Fraction
4
5 print("VERIFICATION OF OUR CONJECTURE -- CASE a = 4")
6 print("="*70)
7
8 def verify_our_conjecture(n_min=2, n_max=500,
9                           max_x_factor=300,
10                          max_t_range=500,
11                          max_q=10):
12
13     captured = 0
14     failed_n_list = []
15
16     start_time = time.time()
17     a = 4# You can change the value of a to test another conjecture
18
19     for n in range(n_min, n_max + 1):
20         found = False
21
22         x_start = n // a + 1
23         x_limit = max_x_factor * n
24
25         for x in range(x_start, x_limit + 1):
26             d = a * x - n
27             if d == 0:
28                 continue
29
30             denom = d * d
31             t_min = max(1, (2 * n * x) // denom)
32
33             # =====
34             # PHASE 1 : integer t
35             # =====
36             for t in range(t_min, t_min + max_t_range + 1):
37                 F = t*t*d*d - 2*t*x*n
38                 if F < 0:
39                     continue
40
41                 q = math.isqrt(F)
42                 if q*q == F:
43                     y = t*d - q
44                     z = t*d + q
45                     if y > 0 and z > 0 and y == int(y) and z == int(z):
46                         if a*x*y*z == n*(y*z + x*z + x*y):
47                             captured += 1
48                             found = True
49                             print(f"n={n:4d} | x={x}, t={t} [INTEGER],
50                                   y={y}, z={z}, F={F}, q={q}")
51                             break
```

```

51         if found:
52             break
53
54     # =====
55     # PHASE 2 : rational t
56     # =====
57     if not found:
58         for x in range(x_start, x_limit + 1):
59             d = a * x - n
60             if d == 0:
61                 continue
62
63             for q_den in range(2, max_q+1):
64                 for p in range(1, max_q*30):
65                     t = Fraction(p, q_den)
66                     F = t*t*d*d - 2*t*x*n
67                     if F < 0:
68                         continue
69
70                     F_num = F.numerator
71                     F_den = F.denominator
72                     if math.isqrt(F_num)**2 == F_num and math.isqrt(
73                         F_den)**2 == F_den:
74                         grat = Fraction(math.isqrt(F_num), math.
75                             isqrt(F_den))
76                         y = t*d - grat
77                         z = t*d + grat
78                         # === require y and z to be integers ===
79                         if y.denominator == 1 and z.denominator ==
80                             1 and y > 0 and z > 0:
81                             if a*x*int(y)*int(z) == n*(int(y)*int(z)
82                                 + x*int(z) + x*int(y)):
83                                 captured += 1
84                                 found = True
85                                 print(f"n={n:4d} | x={x}, t={t} [
86                                     RATIONAL], y={y}, z={z}, F={F},
87                                     q={grat}")
88                                 break
89
90                 if found:
91                     break
92
93             if found:
94                 break
95
96     if not found:
97         print(f"n={n:4d} : X FAILED")
98
99     failed_n_list.append(n)
100
101     elapsed_time = time.time() - start_time
102     total = n_max - n_min + 1
103     success_rate = 100 * captured / total
104
105     print("\n" + "="*70)
106     print("FINAL REPORT a = 4")
107     print("="*70)
108     print(f"Tested interval : n in [{n_min}, {n_max}]")
109
110     print(f"Success : {captured}/{total} ({success_rate:.2f}%)")

```

```

103     print(f"Time : {elapsed_time:.2f}s")
104
105     if failed_n_list:
106         print("Detected failures :", failed_n_list)
107     else:
108         print(" COMPLETE SUCCESS")
109
110     return
111
112
113
114 # DIRECT EXECUTION
115
116
117 verify_our_conjecture(n_min=2, n_max=500)#You can vary the check
    interval for a fixed a

```

Appendix: Python Code for Testing the Classic Erdos-Straus Conjecture for the Values Not Proven by Mordell (Second Formula)

The following code tests the generalized Erdős-Straus conjecture using the second formula specifically for values of n where Mordell's method was inconclusive.

```
1
2 import math
3 import time
4
5 print("VERIFICATION OF ERDOS-STRAUS CONJECTURE - a = 4 (Our Method)")
6
7 print("Testing n in quadratic residues modulo 840")
8 print("-" * 70)
9
10 # Parameters
11 a = 4
12 MOD = 840
13 RESIDUES = {(r * r) % MOD for r in [1, 11, 13, 17, 19, 23]}
14
15 def generate_admissible_n(n_min, n_max):
16     """Generate list of admissible n (quadratic residues mod 840) in [
17         n_min, n_max]"""
18     admissible_list = []
19     for n in range(n_min, n_max + 1):
20         if (n % MOD) in RESIDUES:
21             admissible_list.append(n)
22     return admissible_list
23
24 def verify_with_our_method_for_list(n_list, max_x_factor=300,
25     max_t_range=500):
26     """
27     Verify the conjecture using our method for a given list of n.
28     Returns: solutions dict, failures list, elapsed time
29     """
30     captured = 0
31     failed_n_list = []
32     solutions = {}
33
34     start_time = time.time()
35     total_n = len(n_list)
36
37     print(f"\nTesting {total_n} admissible n")
38     print(f"Parameters: max_x_factor={max_x_factor}, max_t_range={
39         max_t_range}")
40     print("-" * 70)
41
42     for idx, n in enumerate(n_list, 1):
43         found = False
44
45         # Search for x
46         x_start = n // a + 1
47         x_limit = max_x_factor * n
```

```

47     for x in range(x_start, x_limit + 1):
48         if found:
49             break
50
51         d = a * x - n
52         if d == 0:
53             continue
54
55         # Compute t_min to ensure F >= 0
56         denom = d * d
57         t_min = max(1, (2 * n * x) // denom)
58
59         # Search for t
60         for t in range(t_min, t_min + max_t_range + 1):
61             F = t * t * d * d - 2 * t * x * n
62
63             if F < 0:
64                 continue
65
66             q = math.isqrt(F)
67             if q * q == F:
68                 y = t * d - q
69                 z = t * d + q
70
71                 if y > 0 and z > 0:
72                     left = a * x * y * z
73                     right = n * (y*z + x*z + x*y)
74
75                     if left == right:
76                         solutions[n] = (x, t, y, z)
77                         captured += 1
78                         found = True
79
80                     # Print solution
81                     print(f"n={n:4d} : Ok x={x:5d}, t={t:5d}, y
82                           ={y:6d}, z={z:6d}")
83                     break
84
85         if not found:
86             failed_n_list.append(n)
87             print(f"n={n:4d} : FAILED")
88
89     elapsed_time = time.time() - start_time
90
91     return solutions, failed_n_list, elapsed_time
92
93 def intensive_search(failed_list, max_x_factor=1000, max_t_range=2000):
94     """Intensive search for failed n"""
95
96     print(f"\n" + "="*70)
97     print(f"INTENSIVE SEARCH FOR {len(failed_list)} FAILED N")
98     print("="*70)
99
100     a = 4
101     new_solutions = {}
102
103     for idx, n in enumerate(failed_list, 1):

```

```

104     print(f"Failed {idx}/{len(failed_list)} - n={n} : ", end="",
105           flush=True)
106
107     found = False
108     x_start = n // a + 1
109     x_limit = max_x_factor * n
110
111     for x in range(x_start, min(x_limit, 50000) + 1):
112         if found:
113             break
114
115         d = a * x - n
116         if d <= 0:
117             continue
118
119         t_min = max(1, (2 * n * x) // (d * d))
120         t_max = t_min + max_t_range
121
122         for t in range(t_min, min(t_max, t_min + 5000) + 1):
123             F = t * t * d * d - 2 * t * x * n
124
125             if F < 0:
126                 continue
127
128             q = math.isqrt(F)
129             if q * q == F:
130                 y = t * d - q
131                 z = t * d + q
132
133                 if y > 0 and z > 0:
134                     left = a * x * y * z
135                     right = n * (y*z + x*z + x*y)
136
137                     if left == right:
138                         new_solutions[n] = (x, t, y, z)
139                         found = True
140                         print(f"ok (x={x}, t={t}, y={y}, z={z})")
141                         break
142
143         if not found:
144             print("FAILED")
145
146     return new_solutions
147
148 # =====
149 # MAIN EXECUTION
150 # =====
151 if __name__ == "__main__":
152     # MODIFY THESE VALUES TO TEST DIFFERENT INTERVALS
153     n_min = 5000 # YOU CAN CHANGE THIS
154     n_max = 10000 # YOU CAN CHANGE THIS
155
156     # Step 1: Generate admissible n in the interval
157     print(f"\nGenerating admissible n in [{n_min}, {n_max}]")
158     print("-" * 70)
159
160     admissible_list = generate_admissible_n(n_min, n_max)

```

```

161
162 if len(admissible_list) == 0:
163     print("No admissible n in this interval!")
164     exit()
165
166 print(f"Total admissible n: {len(admissible_list)}")
167 print(f"\nList of admissible n ({len(admissible_list)} numbers):")
168 # Print admissible n in groups of 20 for readability
169 for i in range(0, len(admissible_list), 20):
170     print(f"    {admissible_list[i:i+20]}")
171
172 # Step 2: Verify using our method
173 solutions, failed, elapsed = verify_with_our_method_for_list(
174     n_list=admissible_list,
175     max_x_factor=300,
176     max_t_range=500
177 )
178
179 # Step 3: Final report
180 print(f"\n" + "="*70)
181 print("FINAL REPORT")
182 print("="*70)
183
184 success_rate = 100 * len(solutions) / len(admissible_list)
185
186 print(f"Interval tested          : [{n_min}, {n_max}]")
187 print(f"Admissible n tested        : {len(admissible_list)}")
188 print(f"Execution time             : {elapsed:.2f} seconds")
189 print(f"Solutions found            : {len(solutions)}/{len(
190     admissible_list)} ({success_rate:.2f}%)")
191 print(f"Failures                   : {len(failed)}")
192
193 if failed:
194     print(f"\nList of failures ({len(failed)} numbers):")
195     for i in range(0, len(failed), 20):
196         print(f"    {failed[i:i+20]}")
197
198 # Step 4: Intensive search for failures
199 new_solutions = intensive_search(
200     failed_list=failed,
201     max_x_factor=1000,
202     max_t_range=2000
203 )
204
205 if new_solutions:
206     print(f"\n" + "="*70)
207     print(f"INTENSIVE SEARCH RESULTS")
208     print("="*70)
209
210     print(f"New solutions found: {len(new_solutions)}")
211     for n in sorted(new_solutions.keys()):
212         x, t, y, z = new_solutions[n]
213         print(f"    n={n:4d} : x={x:6d}, t={t:6d}, y={y:8d}, z={z
214             :8d}")
215
216     # Update statistics
217     solutions.update(new_solutions)
218     remaining_failures = [n for n in failed if n not in

```

```

217         new_solutions]
218     new_total = len(solutions)
219     new_rate = 100 * new_total / len(admissible_list)
220
221     print(f"\nUpdated totals:")
222     print(f"    Total solutions : {new_total}/{len(
223         admissible_list)} ({new_rate:.2f}%)"
224     print(f"    Remaining failures : {len(remaining_failures)}")
225
226     if remaining_failures:
227         print(f"\nRemaining failures ({len(remaining_failures)}
228             numbers):")
229         for i in range(0, len(remaining_failures), 20):
230             print(f"    {remaining_failures[i:i+20]}")
231     else:
232         print(f"\nNo new solutions found despite intensive search")
233     else:
234         print(f"\n COMPLETE SUCCESS! All admissible n have a solution."
235             )
236
237     print(f"\n" + "="*70)
238     print("END OF PROGRAM")
239     print("="*70)
240
241     # Return the admissible list as well (useful for further analysis)
242     admissible_list_result = admissible_list

```

Users can execute this script in Python to test numerical values and verify the conjecture for different cases.