

Uniqueness of the Rank-5 Projection in the Quantum Gauge Theory of Time

Pasquale Camelia*

Revised June 2026

Abstract

We prove that $r = 5$ is the unique admissible rank within the Majorana–Fibonacci–Heegner projective class of the QGT framework. The result is conditional on four structural inputs: Noetherian balance, second-order Fibonacci autosimilar closure, phasorial realisation of the closure capacity by the admissible catalogue $\mathcal{N}_{\text{adm}} = \{2, 3, 5, 7, 11\}$, and operational metric stability. The proof is a two-sided exclusion. The lower bound does not identify the scalar curvature $|\lambda_+ - \lambda_-|^2 = 5$ with either a spacetime dimension or a Hilbert-space dimension. Rather, this curvature fixes a Fibonacci closure capacity. The rank-five lower bound is obtained only after that capacity is realised by five inequivalent rank-one phasorial sectors. The upper bound follows from the metric threshold $w_4 > \alpha > w_5$. Thus $r = 5$ is not a free truncation parameter inside this admissibility class.

Keywords. Non-invertible projection; rank uniqueness; autosimilar closure; boundary metric; Majorana spinor; finite-rank operator.

The QGT programme studies physical structure generated by a Noetherian balance under restricted accessibility, within the Majorana–Fibonacci–Heegner projective class [1, 20, 19]. In this setting the boundary is represented by a finite-rank non-invertible projection

$$\Pi : \mathcal{H}_{\text{bulk}} \longrightarrow \mathcal{H}_{\text{obs}}, \quad \ker \Pi \neq 0.$$

The admissible finite rank is constrained by two independent mechanisms. First, the Fibonacci closure capacity is realised phasorially by the admissible catalogue $\mathcal{N}_{\text{adm}} = \{2, 3, 5, 7, 11\}$, giving the lower bound $r \geq 5$. Second, operational metric stability gives the upper bound $r \leq 5$ through the bilateral threshold $w_4 > \alpha > w_5$, where α is derived independently from the Pincherle scale increment and Fibonacci curvature [21]. This note proves that these two bounds leave only $r = 5$.

Lemma 1 (Phasorial realisation of the closure capacity). *Let*

$$W := |\lambda_+ - \lambda_-|^2 = 5$$

be the Fibonacci closure capacity of the Majorana–Fibonacci admissibility class. This scalar quantity is not itself a dimension, and it is not the dimension of the pre-geometric domain $M_{\text{pre}}^5 = X_{\text{bdy}} \times S_{\chi}^1(\tau)$. The compact factor $S_{\chi}^1(\tau)$ is an internal phasorial fibre, not an additional observable spacetime direction. The rank-five lower bound is obtained only after the closure capacity is realised by five inequivalent phasorial sectors.

Let

$$\mathcal{N}_{\text{adm}} = \{2, 3, 5, 7, 11\}.$$

*Independent Researcher, Italy. pasquale.camelia@gmail.com. ORCID: [0009-0006-4779-4647](https://orcid.org/0009-0006-4779-4647). © 2026 Pasquale Camelia. Licensed under [CC BY-NC 4.0](https://creativecommons.org/licenses/by-nc/4.0/)

For $n \in \{2, 3, 7, 11\}$, the imaginary quadratic field $\mathbb{Q}(\sqrt{-n})$ has class-number-one arithmetic in the relevant phasorial channel, so its primitive norm form defines one stable rank-one phasorial sector. The remaining element $n = 5$ is the first lock-in channel where class-one rigidity breaks and the projection pole is met. Thus the admissible catalogue consists of four Heegner-stable channels plus one lock-in channel.

For each $n \in \mathcal{N}_{\text{adm}}$, let $s_n(\tau)$ be the corresponding normalised phasor and let

$$\Pi_n^{(1)} := |s_n\rangle\langle s_n|.$$

The five projectors $\{\Pi_n^{(1)} : n \in \mathcal{N}_{\text{adm}}\}$ define five inequivalent rank-one phasorial sectors. Collecting them as a sector decomposition,

$$\mathcal{V}_{\text{ph}} = \bigoplus_{n \in \mathcal{N}_{\text{adm}}} \mathbb{C} s_n, \quad \dim \mathcal{V}_{\text{ph}} = |\mathcal{N}_{\text{adm}}| = 5,$$

where the direct sum counts inequivalent phasorial channels as distinct rank-one superselection sectors of the boundary representation, not as five arbitrary vectors in a single undifferentiated function space. Therefore any boundary representation that resolves the complete admissible phasorial catalogue must satisfy

$$r \geq \dim \mathcal{V}_{\text{ph}} = |\mathcal{N}_{\text{adm}}| = 5.$$

Remark 2 (Rank is not domain dimension). The equality between the number 5 in the rank and the notation M_{pre}^5 is not dimensional. The domain

$$M_{\text{pre}}^5 = X_{\text{bdy}} \times S_\chi^1(\tau)$$

contains a compact internal phasorial fibre over the boundary sector. The projection rank $r = 5$ counts the number of inequivalent rank-one phasorial sectors that the boundary representation must resolve. It is therefore a statement about the observable sector space $\mathcal{H}_{\text{obs}}$, not about the number of spacetime dimensions.

Theorem 3 (Uniqueness of the rank-five projection). *Let $\Pi_r : \mathcal{H}_{\text{bulk}} \rightarrow \mathcal{H}_{\text{obs}}$ be a finite-rank non-invertible boundary projection in the Majorana–Fibonacci–Heegner admissibility class, satisfying:*

- (I) *Noetherian balance under restricted accessibility;*
- (II) *second-order Fibonacci autosimilar closure $v_{k+2} = v_{k+1} + v_k$;*
- (III) *phasorial realisation of the closure capacity by the admissible catalogue $\mathcal{N}_{\text{adm}} = \{2, 3, 5, 7, 11\}$;*
- (IV) *operational stability of the induced boundary metric.*

Then $r = 5$ is the unique admissible rank.

Proof. Lower bound. By Lemma 1, the scalar curvature

$$|\lambda_+ - \lambda_-|^2 = 5$$

is used only as the closure capacity of the admissibility class. The boundary lower bound is supplied by the phasorial realisation of that capacity: the admissible catalogue $\mathcal{N}_{\text{adm}} = \{2, 3, 5, 7, 11\}$ consists of five inequivalent rank-one phasorial sectors. These sectors are counted in the observable representation space; they are not coordinates of M_{pre}^5 . A boundary projection resolving the complete catalogue must therefore have

$$r \geq \dim \mathcal{V}_{\text{ph}} = |\mathcal{N}_{\text{adm}}| = 5.$$

For $r < 5$, at least one admissible phasorial sector has no representative in the retained boundary space. The Noetherian balance and the second-order autosimilar closure cannot then be represented on the full admissible catalogue. Hence $r \geq 5$.

Upper bound. In the SVD representative of the Fibonacci-autosimilar projection, the singular values are $\sigma_k = \varphi^{-k}$, and the transition weights are

$$w_k = \sigma_k \sigma_{k+1} = \varphi^{-(2k+1)}.$$

The threshold α is derived independently from the Pincherle scale increment and Fibonacci curvature, not from the rank choice. The bilateral inequality

$$w_4 > \alpha > w_5$$

then separates resolvable from sub-threshold transition cells. If $r > 5$, the retained space includes the transition $k = 5$, whose weight satisfies $w_5 < \alpha$. The induced boundary metric would assign positive distinguishability to a direction below the operational resolution threshold, violating metric stability. Hence $r \leq 5$.

Combining the two bounds gives $r = 5$. □

Corollary 4. *Under the hypotheses of Theorem 3, in the SVD representative of [21]: the retained singular values are $\sigma_k = \varphi^{-k}$, $k = 0, \dots, 4$; the boundary metric is $d_{\partial}^2 = \sum_{k=0}^4 \varphi^{-2k} |c_k|^2$; the threshold α satisfies $w_4 > \alpha > w_5$; in the QGT angular readout convention, $\mathcal{N}_{\text{adm}} = \{2, 3, 5, 7, 11\}$.*

Scope. The theorem is conditional on the Majorana–Fibonacci–Heegner admissibility class. It does not re-derive the Majorana condition, the Fibonacci autosimilar structure, or the phasorial catalogue from the bare conservation law alone. Its result is narrower: once those structural inputs are fixed, the finite-rank boundary projection cannot choose its rank freely. The complete admissible catalogue requires at least five rank-one phasorial sectors, while metric stability excludes all retained directions beyond the fifth. Thus $r = 5$ is unique inside the class.

Downstream QGT readouts that use $r = 5$ remain subject to their own declared hypotheses and open gates.

Declarations. All results are analytic. The author declares no competing interests and received no funding.

References

- [1] E. Noether, *Invariante Variationsprobleme*, Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse (1918), 235–257.
- [2] J. B. Conway, *A Course in Functional Analysis*, 2nd ed., Springer, New York, 1990.
- [3] M. Reed and B. Simon, *Methods of Modern Mathematical Physics I: Functional Analysis*, revised and enlarged ed., Academic Press, New York, 1980.
- [4] C. Eckart and G. Young, *The approximation of one matrix by another of lower rank*, Psychometrika **1** (1936), 211–218.
- [5] G. H. Golub and C. F. Van Loan, *Matrix Computations*, 4th ed., Johns Hopkins University Press, Baltimore, 2013.
- [6] G. W. Stewart, *On the early history of the singular value decomposition*, SIAM Review **35** (1993), 551–566.

- [7] S. Vajda, *Fibonacci and Lucas Numbers, and the Golden Section*, Ellis Horwood, Chichester, 1989.
- [8] T. Koshy, *Fibonacci and Lucas Numbers with Applications*, Wiley, New York, 2001.
- [9] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, 6th ed., Oxford University Press, Oxford, 2008.
- [10] K. Heegner, *Diophantische Analysis und Modulfunktionen*, Mathematische Zeitschrift **56** (1952), 227–253.
- [11] A. Baker, *Linear forms in the logarithms of algebraic numbers*, Mathematika **13** (1966), 204–216.
- [12] H. M. Stark, *A complete determination of the complex quadratic fields of class-number one*, Michigan Mathematical Journal **14** (1967), 1–27.
- [13] D. A. Cox, *Primes of the Form $x^2 + ny^2$: Fermat, Class Field Theory, and Complex Multiplication*, Wiley, New York, 1989.
- [14] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, 2nd ed., Springer, New York, 1990.
- [15] J. Bertrand, P. Bertrand, and J.-P. Ovarlez, *The Mellin Transform*, in A. D. Poularikas (ed.), *The Transforms and Applications Handbook*, 2nd ed., CRC Press, Boca Raton, 2000.
- [16] P. Flajolet, X. Gourdon, and P. Dumas, *Mellin transforms and asymptotics: harmonic sums*, Theoretical Computer Science **144** (1995), 3–58.
- [17] P. L. Duren, *Theory of H^p Spaces*, Academic Press, New York, 1970.
- [18] F. W. King, *Hilbert Transforms*, Vols. 1–2, Cambridge University Press, Cambridge, 2009.
- [19] P. Camelia, *The Quantum Gauge Theory of Time: The Majorana Qubit and the Structure of Physical Time*, First edition, independently published via Amazon KDP, 2026.
- [20] P. Camelia, *A Quantum Gauge Theory of Time: Foundations from a Noetherian Balance*, Preprint (2026). https://pasqualecamelia.github.io/papers/Camelia_QGT_Foundations_of_Physics.pdf
- [21] P. Camelia, *Projection-Induced SVD and Boundary Metric: A Rank-Five Fibonacci-Autosimilar Characterization*, Preprint (2026). https://pasqualecamelia.github.io/papers/Camelia_2026_ProjectionSVD_BoundaryMetric.pdf