

1. Functions of multivariables.

2. Partial derivatives

(a) Tangent plane to the surface of a given function

(b) Maxima & minima.

EE1203: Vector Calculus

Aneesh Sobhanan

Department of Electrical Engineering

IIT Hyderabad, India

16th January, 2025




भारतीय शैक्षणिक विज्ञान संस्थान
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad

Functions of several variables and derivatives.

* Calculus: → Studying functions!

1. Functions of 1 variable: $f(x) = \sin(x)$

Graph of a function $f(x)$



2. Function of 2-variables:

Given (x, y) in space → get a number $f(x, y)$

Example: $f(x, y) = x^2 + y^2$ Domain $\in \mathbb{R}^2$;
Range $\in \mathbb{R}$

$f(x,y) = \sqrt{y}$; only when $y \geq 0$ (y should not be negative)


$f(x,y) = \frac{1}{x+y}$; only defined when $x+y \neq 0$.

Physical example: $f(x,y) =$ temperature at point (x,y)
or 3 or more variables.


* Here we might again restrict in 2 or 3 variables:
But tools we learn is common even if we have n^1 variables.

Representation of function of 2 variables.

→ Graph: $Z = f(x,y)$



Example: $f(x, y) = 1 - x^2 - y^2$



in yz plane ($x=0$)

$$\Rightarrow z = 1 - y^2.$$

in xz plane: ($y=0$)

$$z = 1 - x^2.$$

Where it hits the xy plane?


$$\rightarrow \text{at } z=0$$

$$\Rightarrow 1 - x^2 - y^2 = 0$$
$$x^2 + y^2 = 1 \quad (\text{Circle of } r=1)$$

Example: Try to visualize (may be use a computer code)

$$\underline{z = y^2 - x^2}$$

Contour plot: Another elegant method to represent functions of 2-variables.




\equiv Elevation of $z = f(x,y)$
onto xy plane.

\Rightarrow Simpler to read the
map in contour plot.

Each of
these is
contour
plot is called
level curves!

- * Show all the points where $f(x,y) = C$ where $C = \text{Constant}$.
- * Usually C is chosen at regular intervals.
- * Equivalent of cutting the graph into horizontal planes

How is the contour plot useful?



It tells us many insights!

- * If x increases,
→ f increases (here)
 - * If y increases,
* f increases.

(1) If x decreases, } or y decreases } f decreases

Partial derivatives:

function of 1 variable :



Rate of change of $f(x)$ w.r.t change in x

$$f'(x) = \frac{df}{dx} = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}.$$

* Please review your basics on
Continuity, differentiability
Concepts.


How to extend this idea for Functions of 2 variables.

Partial derivatives:-

$$\frac{\partial f}{\partial x}(x_0, y_0) = \lim_{\Delta x \rightarrow 0} \frac{f(x_0 + \Delta x, y_0) - f(x_0, y_0)}{\Delta x}$$

$$\frac{\partial f}{\partial y}(x_0, y_0) = \lim_{\Delta y \rightarrow 0} \frac{f(x_0, y_0 + \Delta y) - f(x_0, y_0)}{\Delta y}$$

How to visualize this concept?



How to compute?

To find $\frac{\partial f}{\partial x} = f_x$

* treat 'y' as constant
& 'x' as variable &
Vice versa.

Usefulness of partial derivatives!

* minimization / maximization problems involving functions of multivariables.

Approximation formula:

If we change $x \rightarrow x + \Delta x$

$y \rightarrow y + \Delta y$.

How to justify this formula?

$Z = f(x, y)$: then

$$\Delta Z \approx f_x \Delta x + f_y \Delta y.$$

↓
neglecting higher orders

one of the powerful formulae in partial derivatives


Justification to the above formula:

Approximate formula in 1 variable case

If at $x_0 \rightarrow f(x) = f(x_0)$

Value at nearby point : $f(x) \approx f(x_0) + f'(x_0)(x - x_0)$

Ref: Vector calculus 4th Ed; by Susan Jane Colley.



* f_x, f_y are slopes of 2 tangent lines.


$$f(a, b) = z_0.$$

If $\frac{\partial f}{\partial x}(a, b) = p$;

$$\Rightarrow L_x = \begin{cases} z = z_0 + p(x-a) \\ y = b \end{cases}$$

If $\frac{\partial f}{\partial y}(a, b) = q$

$$\Rightarrow L_y = \begin{cases} z = z_0 + q(y-b) \\ x = a \end{cases}$$



The tangent plane at $(a, b, f(a, b))$ contains the lines
 tangent to the curves formed between
 $z = f(x, y) \text{ & } x=a$; $z = f(x, y) \text{ & } y=b$.

L_1 & L_2 are both tangent to the surface $Z = f(x,y)$

Together they determine a plane

$$Z = Z_0 + p(x-a) + q(y-b)$$

-- Egn of tangent plane!

If we fix $y=b$

and vary ' x ' we get

first tangent line,

fix $x=a$ and

Vary ' y ' \rightarrow the second tangent line. ✓

* This tangent plane is near to the graph / surface $f(x,y)$ (as it is only an approximation egn).

.. Can you think of another way to define the above egn using lines & planes in vector calculus.

Tangent plane Eqn. to the surface $z = f(x,y)$ at point $(a,b, f(a,b))$

$L_x \rightarrow$ tangent line to the traces of surface in plane $y=b$;
 $L_y \rightarrow$ " plane $x=a$.

If $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}$ exist at (a,b) , and is continuous,

Equation of tangent plane is; $A(x-a) + B(y-b) + C(z - f(a,b)) = 0$.

if $\vec{n} = (A, B, C)$ (normal vector)

Can we find this normal vector?


Tangent plane contains both $L_x \& L_y \Rightarrow$ If we know \vec{V}_x, \vec{V}_y parallel to L_x, L_y respectively

$$\Rightarrow \vec{n} = \vec{V}_x \times \vec{V}_y$$

$$= \begin{vmatrix} i & j & k \\ 1 & 0 & f_x(a,b) \\ 0 & 1 & f_y(a,b) \end{vmatrix}$$

$$= -\hat{i} f_x(a,b) - \hat{j} f_y(a,b) + \hat{k}$$

$$\Rightarrow f_x(a,b)(x-a) + f_y(y-b) - z + f(a,b) = 0.$$



Maxima & minima:

* Optimization problems.

→ Find min/max of a function in 2 variables $f(x, y)$

* At a local min or max $f_x = 0$ and $f_y = 0$.

* From Approx formula; $\Delta z = f_x \Delta x + f_y \Delta y$.


→ when $f_x = f_y = 0$ $\Delta z = 0$

⇒ The function does not change anymore

⇒ local max or local min.

⇒ Or it also means the tangent plane is horizontal to the graph of the function $z=f(x, y)$

Ex:



EE1203 Jan- Feb 2025
aneesh@ee.jith.ac.in

EE1203 Jan- Feb 2025
aneesh@ee.jith.ac.in

EE1203 Jan- Feb 2025
aneesh@ee.jith.ac.in

EE1203 Jan- Feb 2025
aneesh@ee.jith.ac.in