



Cambridge International Examinations

Cambridge International Advanced Level

CANDIDATE NAME									
CENTRE NUMBER					CANDIDATE NUMBER				
MATHEMATICS	;							97	09/52
Paper 5 Mecha	nics 2 (l	M2)					May	//Jun	e 2017
						1	hour	15 m	inutes
Candidates ans	wer on tl	ne Quest	ion Pa	iper.					
Additional Materials: List of Formulae (MF9)									

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

Where a numerical value for the acceleration due to gravity is needed, use 10 m s^{-2} .

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.


The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 50.



BLANK PAGE

on a	rticle P of mass 0.2 kg moves with speed $4 \mathrm{ms^{-1}}$ and angular speed $5 \mathrm{rads^{-1}}$ in a horizontal circ smooth surface. P is attached to one end of a light elastic string of natural length 0.6 m. The end of the string is attached to the point on the surface which is the centre of the circular motion.
(i)	Find the radius of this circle.
(ii)	Find the modulus of elasticity of the string.



The ends of two light inextensible strings of length 0.7 m are attached to a particle P. The other ends of the strings are attached to two fixed points A and B which lie in the same vertical line with A above B. The particle P moves in a horizontal circle which has its centre at the mid-point of AB. Both strings are inclined at 60° to the vertical. The tension in the string attached to A is $6 \, \text{N}$ and the tension in the string attached to B is $4 \, \text{N}$ (see diagram).

Find the mass of P .	[2]

(ii)	Calculate the speed of P .	[3]

An open box in the shape of a cube with edges of length 0.2 m is placed with its base horizontal and

Calculate the height of the centre of mass of the box above its base.	
	•••••

The box is now fitted with a thin uniform square lid of weight $3\,N$ and with edges of length $0.2\,m$. The lid is attached to the box by a hinge of length $0.2\,m$ and weight $2\,N$. The lid of the box is held partly open.

(including the lid and hinge) is 0.12 m above the base of the box.	
	,
	•••••
	•••••


desc	mall object of mass 0.4 kg is released from rest at a point 8 m above the ground. The object ends vertically and when its downwards displacement from its initial position is x m the object velocity v m s ⁻¹ . While the object is moving, a force of magnitude $0.2v^2$ N opposes the motion.
(i)	Show that $v \frac{\mathrm{d}v}{\mathrm{d}x} = 10 - 0.5v^2$. [2]
(ii)	Express v in terms of x . [4]

(iii)	Find the increase in the value of v during the final 4 m of the descent of the object. [2]

A particle of mass 0.3 kg is attached to one end of a light elastic string of natural length 0.8 m and

(Calculate the greatest speed of the particle during its descent.	
		•••••
•		
•		•••••
•		•••••
•		
•		
•		

•••••
•••••
•••••



The end A of a non-uniform rod AB of length 0.6 m and weight 8 N rests on a rough horizontal plane, with AB inclined at 60° to the horizontal. Equilibrium is maintained by a force of magnitude 3 N applied to the rod at B. This force acts at 30° above the horizontal in the vertical plane containing the rod (see diagram).

(i)	Find the distance of the centre of mass of the rod from A .	[2]
		•••••
		•••••
		•••••
		•••••

The 3 N force is removed, and the rod is held in equilibrium by a force of magnitude P N applied at B, acting in the vertical plane containing the rod, at an angle of 30° below the horizontal.

(ii)	Calculate <i>P</i> .	[2]
		••••••
In o	one of the two situations described, the rod AB is in limiting equilibrium.	
(iii)	Find the coefficient of friction at A .	[4]
		•••••

and vertically upwards displacements of P from O are x m and y m respectively. The equation of rajectory of P is $y = 2x - \frac{25x^2}{V^2}$.		
i) Write down the value of $\tan \theta$, whe	re θ is the angle of projection of P .	
hen $t = 4$, P passes through the point.	A where $x = y = a$.	
i) Calculate V and a .		

(iii)	Find the direction of motion of P when it passes through A . [3]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.