

Properties of curves and applications of differentiation

Kh notes

Contents

1 Properties of curves	2
1.1 Review Questions	2
1.1.1 Some fundamental derivatives:	2
1.1.2 Rules of differentiation:	2
1.2 Tangents	3
1.2.1 Horizontal Tangents	4
1.2.2 Ex13A	5
1.3 Normals	14
1.3.1 Ex 13B	14
1.4 Increasing and Decreasing	19
1.4.1 Basic sign diagram example	19
1.4.2 Sign diagram example with discontinuities	21
1.4.3 Ex 13 C	21
1.5 Stationary Points	26
1.5.1 Turning points (minima, maxima)	26
1.5.2 Stationary points of inflection	27
1.5.3 Greatest or least values (testing boundaries)	28
1.5.4 Ex 13D	29
1.6 Shape	36
1.6.1 Ex 13E	36
1.7 Inflection Points	39
1.7.1 Ex 13F	39
1.8 Understanding functions and their derivatives	44
2 Applications of differentiation	45

Chapter: 1 Properties of curves

1.1 Review Questions

1.1.1 Some fundamental derivatives:

Function	Derivative
$f(x) = x^n$	$f'(x) = nx^{n-1} \quad (n \in \mathbb{R})$
$f(x) = e^x$	$f'(x) = e^x$
$f(x) = \ln x$	$f'(x) = \frac{1}{x}$
$f(x) = \sqrt{x}$	$f'(x) = \frac{1}{2\sqrt{x}}$
$f(x) = \sin x$	$f'(x) = \cos x$
$f(x) = \cos x$	$f'(x) = -\sin x$
$f(x) = \tan x$	$f'(x) = \sec^2 x$

1.1.2 Rules of differentiation:

Chain Rule:

$$y = g(u_{(x)})$$
$$\frac{dy}{dx} = g'(u_{(x)})u'_{(x)}$$

Product Rule:

$$y = u_{(x)}v_{(x)}$$
$$\frac{dy}{dx} = u_{(x)}v'_{(x)} + u'_{(x)}v_{(x)}$$

Quotient Rule:


$$y = \frac{u_{(x)}}{v_{(x)}}$$
$$\frac{dy}{dx} = \frac{u'_{(x)}v_{(x)} - u_{(x)}v'_{(x)}}{[v_{(x)}]^2}$$

1.2 Tangents

- The tangent to a curve at a point A is the best approximating straight line to the curve at point A.
- (Leibniz definition) Tangent to the curve $y = f(x)$ at the point $(a, f(a))$ is the line through the infinitely close pair of points either side of $f(a)$

$$\frac{y - f(a)}{x - a} = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

- It is a single point of contact with the curve (although it may intersect the curve at some other point)



For the function $y = f(x)$, and some $x = a$

- $(a, f(a))$ is on the curve
- $f'(a)$ is the gradient of the curve at $x = a$


$$\begin{aligned} \frac{y - f(a)}{x - a} &= f'(a) \\ \Rightarrow y &= f'(a)(x - a) + f(a) \end{aligned}$$

Is the equation of the tangent line

1.2.1 Horizontal Tangents

Horizontal tangents have a gradient of 0.

These will be very important later, when we investigate stationary points in more detail.



In general, we will need to find where the stationary points are:

- Find $f'(x)$
 - Solve $f'(x) = 0$
 - Substitute the solution(s) into $f(x)$ to find the constant terms for the horizontal line equation
-

1.2.2 Ex13A

- Tangent equations from root and polynomial functions
- Horizontal tangents
- Natural log and exponent questions
- Concept questions

3 worked examples

- Polynomial
- Log or exponential
- Conceptual

Find the equation of the tangent to:

a) $y = x - 2x^2 + 3$ at $x=2$

$$\frac{dy}{dx} = 1 - 4x$$

when $x=2$

$$\frac{dy}{dx} = 1 - 4(2) = -7$$

and $y = (2) - 2(2)^2 + 3$
 $= -3$

$$\frac{dy}{dx} = -7 \quad \text{Point } (2, -3)$$

$$y - (-3) = -7(x - 2)$$

$$y + 3 = -7x + 14$$

$$y = -7x + 11$$

d) $y = \frac{4}{\sqrt{x}}$ at $(1, 4)$

$$y = 4x^{-\frac{1}{2}}$$

$$\frac{dy}{dx} = 4 \cdot -\frac{1}{2}x^{-\frac{3}{2}}$$

$$= -2x^{-\frac{3}{2}} = \frac{-2}{x^{\frac{3}{2}}}$$

$x = 1$

$$\frac{dy}{dx} = \frac{-2}{1} = -2$$

Point $(1, 4)$

$$y - 4 = -2(x - 1)$$

$$y - 4 = -2x + 2$$

$$y = -2x + 6$$

(Horizontal Tangents)

3. c)

Find the equation of horizontal tangents to

$$y = \sqrt{x} + \frac{1}{\sqrt{x}}$$

$$\Rightarrow y = x^{\frac{1}{2}} + x^{-\frac{1}{2}}$$

$$\frac{dy}{dx} = \frac{1}{2}x^{-\frac{1}{2}} + -\frac{1}{2}x^{-\frac{3}{2}}$$

(factorise using lowest power)

$$\frac{dy}{dx} = \frac{1}{2}x^{-\frac{3}{2}}(x^{\frac{1}{2}} - 1)$$

Set $\frac{dy}{dx} = 0$

$$\frac{1}{2}x^{-\frac{3}{2}}(x^{\frac{1}{2}} - 1) = 0$$

$$x^{-\frac{3}{2}} \neq 0 \quad x^{\frac{1}{2}} - 1 = 0$$

Find y

$$y = \sqrt{1} + \frac{1}{\sqrt{1}} = 1 + 1$$

$$= 2$$

Horizontal tangent at $(1, 2)$

Equation: $y = 2$

$$\frac{dy}{dx} = \frac{1}{2x^{\frac{1}{2}}} - \frac{1}{2x^{\frac{3}{2}}}$$

$$= \frac{x - 1}{2x^{\frac{3}{2}}}$$

$$\frac{x - 1}{2x^{\frac{3}{2}}} = 0 \Rightarrow x - 1 = 0$$

5. Find another tangent to

$$y = 1 - 3x + 12x^2 - 8x^3$$

which is parallel to the tangent at $(1, 2)$

$$\frac{dy}{dx} = -3 + 24x - 24x^2$$

when $x=1$

$$\begin{aligned}\frac{dy}{dx} &= -3 + 24(1) - 24(1)^2 \\ &= -3\end{aligned}$$

so gradient is -3

Find all x values where gradient $= -3$

$$-3 + 24x - 24x^2 = -3$$

$$24x - 24x^2 = 0$$

$$x - x^2 = 0$$

$$x(1-x) = 0$$

$$x=0 \quad \underbrace{x=1}_{a} \quad) \text{ already have}$$

Find equation of tangent at $x=0$


$$y = 1$$

$$y - 1 = -3(x - 0)$$

$$y - 1 = -3x, \quad y = -3x + 1$$

other tangent $y - 2 = -3(x - 1)$

$$y = -3x + 5$$



6. $y = x^2 + ax + b$

when $x=1$ tangent

$$is 2x+y=6$$

$$y = -2x + 6$$

$$\frac{dy}{dx} = -2 \quad y = 4$$

$(1, 4)$ point on curve.

-2 gradient at that point.

$$\frac{dy}{dx} = 2x + a$$

$$-2 = 2(1) + a$$

$$a = -4$$

$$y = x^2 - 4x + b$$


$$4 = (1)^2 - 4(1) + b$$

$$4 = -3 + b$$

$$b = 7.$$

$$\underline{y = x^2 - 4x + 7}$$

7. Find a and b



What do we know?

- when $x=4 \frac{dy}{dx} = 1$
- $(4, 4+2) = (4, 6)$ is on $y = a\sqrt{x} + b$

① Using $(4, 6)$

$$\begin{aligned} 6 &= a\sqrt{4} + b(4) \\ 6 &= 2a + 4b \\ \Rightarrow 3 &= a + 2b \end{aligned}$$

$$② \frac{dy}{dx} = \frac{a}{2\sqrt{x}} + b$$

$$1 = \frac{a}{2\sqrt{4}} + b$$

$$1 = \frac{a}{4} + b$$

$$4 = a + 4b$$

Equation

$$\begin{aligned} 3 &= a + 2b & ① \\ 4 &= a + 4b & ② \end{aligned}$$

$$② - ① \quad 1 = 2b$$

$$b = \frac{1}{2}$$

$$\begin{aligned} \text{using } ① \quad a &= 3 - 2b \\ &= 3 - 2\left(\frac{1}{2}\right) \\ &= 2 \end{aligned}$$

$$y = 2\sqrt{x} + \frac{x}{2}$$

8. Show equation of

tangent to $y = 2x^2 - 1$

at $x=a$ is

$$4ax - y = 2a^2 + 1$$

Find point:

$$x = a$$

$$y = 2a^2 - 1$$

$$(a, 2a^2 - 1)$$

$$\frac{dy}{dx} = 4x$$

$$= 4a$$

$$y - (2a^2 - 1) = 4a(x - a)$$

$$y - 2a^2 + 1 = 4ax - 4a^2$$

$$(-y) \quad -2a^2 + 1 = 4ax - 4a^2 - y$$

$$(+4a^2) \quad 2a^2 + 1 = 4ax - y$$

$$10) \frac{dy}{dx} = \ln \sqrt{x}$$

$$y = \ln(x^{\frac{1}{2}})$$

$$y = \frac{1}{2} \ln(x)$$

$$\frac{dy}{dx} = \frac{1}{2x}$$

$$y = -1$$

$$\frac{1}{2} \ln(x) = -1$$

$$\ln(x) = 2$$

$$x = e^2$$

$$(e^{-2}, -1)$$

$$\frac{dy}{dx} = \frac{1}{2e^{-2}}$$

$$y + 1 = \frac{1}{2e^{-2}}(x - e^{-2})$$

$$y = \frac{x}{2e^{-2}} - \frac{1}{2} - 1$$

$$y = \frac{e^2}{2}x - \frac{3}{2}$$

Using $\ln x$ and e^x

Find the equation of the tangent to $y = \ln x$ where $y = -1$

• Find x

$$\ln x = -1 \quad (\log_e x = 1)$$

$$x = e^{-1}$$

$$\frac{dy}{dx} = \frac{1}{x} = x^{-1}$$

$$x = e^{-1}, \quad \frac{dy}{dx} = (e^{-1})^{-1} = e^1 = e$$

Point $(e^{-1}, -1)$


$$\frac{dy}{dx} = e$$

$$y + 1 = e(x - e^{-1})$$

$$y + 1 = ex - e^1 e^{-1}$$

$$y + 1 = ex - 1$$

$$y = ex - 2$$



$$12. f(x) = \ln(x(x-2))$$

Domain of log function > 0

$$\Rightarrow x(x-2) > 0$$

$$\text{Solve } x(x-2) = 0$$

$$\begin{array}{c} x=0 \\ x=2 \end{array}$$

Domain $x < 0 \cup x > 2$

$$\{x \mid x < 0 \cup x > 2\}$$

$$f'(x) = \frac{1}{x(x-2)} \cdot \frac{d}{dx}(x^2 - 2x)$$

$$= \frac{1}{x(x-2)} \cdot 2x - 2$$

$$= \frac{2(x-1)}{x(x-2)}$$

$$= \frac{2x-2}{x^2-2x}$$

$$= \frac{2x-2}{x(x-2)}$$

$$= \frac{2x+x-2}{x(x-2)}$$

$$= \frac{1}{x} + \frac{1}{x-2}$$

$$x=3, f'(x) = \frac{1}{3} + \frac{1}{1} = \frac{4}{3}$$

$$x=3, f(3) = \ln(3(3-2)) = \ln 3$$

$$y - \ln 3 = \frac{4}{3}(x-3)$$

$$y = \frac{4}{3}x - 4 + \ln 3$$

$$13. y = x^2 e^x$$

$$y = uv \quad \frac{dy}{dx} = uv' + u'v$$

(product rule)

$$\begin{aligned} \frac{dy}{dx} &= x^2 e^x + 2x e^x \\ &= x e^x (x+2) \end{aligned}$$

$$x=1$$

$$\frac{dy}{dx} = e(3) = 3e$$

$$x=1, y=e$$

$$\text{Gradient: } 3e$$

$$\text{Point: } (1, e)$$

$$y - e = 3e(x-1)$$

$$y = 3ex - 2e \quad \text{equation of tangent}$$

Find intercepts.

$$x=0 \quad y = -2e$$

$$y=0 \quad x = \frac{2e}{3e}$$

$$x = \frac{2}{3}$$

Intercepts

$$(0, -2e) \quad y\text{-intercept}$$

$$\left(\frac{2}{3}, 0\right) \quad x\text{-intercept}$$

$$14. \quad y = 3xe^{\frac{x}{2}}$$

$$\begin{aligned} \frac{dy}{dx} &= 3x(e^{\frac{x}{2}})' + (3x)' e^{\frac{x}{2}} \\ &= 3x e^{\frac{x}{2}} \cdot \frac{1}{2} + 3e^{\frac{x}{2}} \\ &= 3e^{\frac{x}{2}} \left(\frac{x}{2} + 1 \right) \end{aligned}$$

When $x = -1$

$$\begin{aligned} \frac{dy}{dx} &= 3e^{-\frac{1}{2}} \left(-\frac{1}{2} + 1 \right) \\ &= \frac{3}{2} e^{-\frac{1}{2}} \end{aligned}$$

$$x = -1, \quad y = -3e^{-\frac{1}{2}}$$

$$y + 3e^{-\frac{1}{2}} = \frac{3e^{-\frac{1}{2}}}{2}(x + 1)$$

$$\begin{aligned} y &= \frac{3e^{-\frac{1}{2}}}{2}x + \frac{3e^{-\frac{1}{2}}}{2} - 3e^{-\frac{1}{2}} \\ &= \frac{3e^{-\frac{1}{2}}}{2}x - \frac{3e^{-\frac{1}{2}}}{2} \end{aligned}$$

$$\text{if } x = 0 \quad y = -\frac{3e^{-\frac{1}{2}}}{2}$$

$$\text{if } y = 0 \quad x = 1$$

$$\frac{3e^{-\frac{1}{2}}}{2} \sqrt{\frac{1}{A}} = \frac{3e^{-\frac{1}{2}}}{4}$$

$$A = \frac{3}{4\sqrt{e}} \text{ units}^2$$

15. Find the equation of the tangent to $y = \frac{1}{\sin 2x}$ at the point where

$$x = \frac{\pi}{4}$$

$$u = 1, u' = 0 \quad v = \sin 2x, v' = 2\cos(2x) \quad \left(\frac{u}{v}\right)' = \frac{u'v - uv'}{v^2}$$

$$\begin{aligned} \frac{dy}{dx} &= -\frac{1 \cdot 2\cos(2x)}{\sin^2(2x)} = -\frac{2\cos(2x)}{\sin^2(2x)} \\ &= -2\cot(2x)\csc(x) \end{aligned}$$


$$\text{when } x = \frac{\pi}{4} \quad \frac{dy}{dx} = -2\cot\left(\frac{\pi}{2}\right) = 0$$

$$\text{when } x = \frac{\pi}{4}, \quad y = \frac{1}{\sin\left(\frac{\pi}{2}\right)} = 1$$

Horizontal tangent

$$y = 1$$

27.



$$\text{Common Points } (b, ab^2)$$

$$\text{a) } (b, \ln b)$$

$$\Rightarrow ab^2 = \ln b$$

Common Tangent

$$\frac{dy}{dx} = 2ax \quad , \quad \frac{dy}{dx} = \frac{1}{x}$$

$$x=b \quad 2ab = \frac{1}{b} \Rightarrow 2ab^2 = 1$$

$$\text{b) } \begin{aligned} ab^2 &= \ln b & \textcircled{1} \\ 2ab^2 &= 1 & \textcircled{2} \end{aligned}$$

$$\begin{aligned} 2ab^2 &= 2\ln b \\ 2ab^2 &= 1 \end{aligned}$$

$$\begin{aligned} 2\ln b &= 1 \\ \ln b &= \frac{1}{2} \end{aligned}$$

$$b = e^{\frac{1}{2}} = \sqrt{e}$$

$$\text{Point } (e^{\frac{1}{2}}, \ln e^{\frac{1}{2}}) = (e^{\frac{1}{2}}, \frac{1}{2})$$

$$\text{c) } a = \frac{1}{2b^2}$$

$$a = \frac{1}{2e}$$

$$\text{d) } a = \frac{1}{2e}, b = \sqrt{e}$$

$$\text{Point } (e^{\frac{1}{2}}, \frac{1}{2})$$

$$\text{Gradient } \frac{dy}{dx} = \frac{1}{b} = \frac{1}{\sqrt{e}} = e^{-\frac{1}{2}}$$

$$y - \frac{1}{2} = e^{-\frac{1}{2}}(x - e^{\frac{1}{2}})$$

$$y - \frac{1}{2} = e^{-\frac{1}{2}}x - 1$$

$$y = 2\sqrt{e}x - \frac{1}{2}$$

$$28. \quad P(x) = ax^2$$

$$P'(x) = 2ax$$

$$x=s \quad (s, as^2) \quad P'(s) = 2as$$

$$y - as^2 = 2as(x - s)$$

$$\begin{aligned} y &= 2asx - 2as^2 + as^2 \\ y &= 2asx - as^2 \end{aligned}$$

$$\begin{aligned} x=t \\ y &= 2atx - at^2 \end{aligned}$$

$$2asx - as^2 = 2atx - at^2$$

$$x(2as - 2at) = as^2 - at^2$$

$$x = \frac{a(s^2 - t^2)}{2a(s-t)}$$

$$= \frac{(s+t)(s-t)}{2(s-t)}$$

$$= \frac{s+t}{2}$$

If perpendicular

$$2as \cdot 2at = -1$$

$$y = 2atx - at^2$$

$$x_c = \frac{s+t}{2}$$

$$y = \frac{2at(s+t)}{2} - at^2$$

$$= ast + at^2 - at^2$$

$$= ast$$

$$t = \frac{-1}{4a^2s}$$

$$y = ast$$


$$= as \cdot \frac{-1}{4a^2s}$$

$$= -\frac{1}{4a}$$

1.3 Normals

The product of the gradients of perpendicular lines = -1

There are various proofs, one example below.




For the function $y = f(x)$, and some $x = a$

- $(a, f(a))$ is on the curve
- $-\frac{1}{f'(a)}$ is the gradient of the **normal to the curve** at $x = a$

$$\frac{y - f(a)}{x - a} = -\frac{1}{f'(a)}$$

$$\Rightarrow y = -\frac{1}{f'(a)}(x - a) + f(a)$$

Is the equation of the tangent line



1.3.1 Ex 13B

13B

1. d)
Find the equation of the normal to:

$$y = 8\sqrt{x} - \frac{1}{x^2} \text{ at } x=1$$

Point $x=1$

$$y = 8\sqrt{1} - \frac{1}{1^2} = 7 \quad (1, 7)$$

Gradient:

$$\begin{aligned} y &= 8x^{\frac{1}{2}} - x^{-2} \\ \frac{dy}{dx} &= 8 \cdot \frac{1}{2} x^{-\frac{1}{2}} - -2x^{-3} \\ &= \frac{4}{x^{\frac{1}{2}}} + \frac{2}{x^3} \\ &= \frac{4x^{\frac{5}{2}} + 2}{x^3} \\ \frac{dy}{dx} \perp &= \frac{-x^3}{4x^{\frac{5}{2}} + 2} \end{aligned}$$

$$x=1$$

$$\frac{dy}{dx} \perp = \frac{-1}{4+2} = -\frac{1}{6}$$

$$(1, 7) \text{ and } -\frac{1}{6}$$

$$y-7 = -\frac{1}{6}(x-1)$$

$$y-7 = -\frac{x}{6} + \frac{1}{6}$$

$$y = -\frac{x}{6} + \frac{43}{6}$$

or

$$6y + x = 43$$

$$\text{or } x + 6y - 43 = 0 \quad \circ$$

3. Find the equation of
c) the normal to
 $y = e^{2x-1}$ at $x=1$

Point $(1, e^1)$

$$\begin{aligned} \frac{dy}{dx} &= e^{2x-1} \cdot 2 \\ &= 2e^{2x-1} \end{aligned}$$

$$\frac{dy}{dx} \perp = \frac{-1}{2e^{2x-1}} = -\frac{e^{1-2x}}{2}$$

$$x=1$$

$$\frac{dy}{dx} \perp = -\frac{1}{2e}$$

$$y - e = -\frac{1}{2e}(x-1)$$

$$y = -\frac{x}{2e} + \frac{1}{2e} + e$$

$$y = -\frac{xe^{-1}}{2} + \frac{1+2e^2}{2e}$$

or

$$2ey - 2e^2 = -x + 1$$

$$2ey + x = 1 + 2e^2$$

$$x + 2ey = 1 + 2e^2$$

5. Find the points where the normal to

$$y = x^3 - 2x^2 + 1$$

at $x=1$ meets the curve again.

Find equation of normal at $x=1$ and then set equal to y , find intersection points.

$$\frac{dy}{dx} = 3x^2 - 4x$$

$$\frac{dy}{dx} \perp = \frac{-1}{3x^2 - 4x}$$

$$x=1 \quad \frac{dy}{dx} \perp = \frac{-1}{3-4} = 1$$

$$x=1, \quad y = 1 - 2 + 1 = 0$$

$$y - 0 = 1(x-1)$$

$$y = x-1 \quad (\text{normal})$$

$$\text{set } x^3 - 2x^2 + 1 = x-1$$

$$\Rightarrow x^3 - 2x^2 - x + 2 = 0$$

We already have one solution

$$x=1 \Rightarrow x-1 \text{ is a}$$

factor

$$(x-1)(x^2 - x - 2) = 0$$

From deduce

$$\begin{array}{r} x^2 - x - 2 \\ \hline x-1) x^3 - 2x^2 - x + 2 \\ x^2 - x^2 \\ \hline -x^2 - x \\ -x^2 + x \\ \hline -2x + 2 \\ -2x + 2 \\ \hline 0 \end{array}$$

$$(x-1)(x^2 - x - 2)^0$$

$$\Rightarrow (x-1)(x-2)(x+1) = 0$$

Solution $x=1, x=2, x=-1$

so new points

$$x=2 \quad y = (2)^3 - 2(2)^2 + 1$$


$$= 1 \quad (2, 1)$$

$$x = -1$$


$$y = (-1)^3 - 2(-1)^2 + 1$$

$$= -2 \quad (-1, -2)$$

points are $(2, 1), (-1, -2)$



6. Find the equation of the normal to $f(x) = \cos(x)$ which passes through the origin.



let $x = a$
 $\Rightarrow (a, \cos(a))$ on the curve

$$\frac{dy}{dx} = -\sin(x)$$

$$\frac{dy}{dx} \perp = \frac{1}{\sin(x)}$$

$$\frac{dy}{dx} \perp = \frac{1}{\sin(a)}$$

$$\frac{\cos(a) - 0}{a - 0} = \frac{1}{\sin(a)}$$

$$\begin{aligned}\sin(a)\cos(a) &= a \\ 2\sin(a)\cos(a) &= 2a\end{aligned}$$

$$\sin(2a) = 2a$$

Solve $\sin(x) = x$?

We know $\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$

$$\begin{aligned}\Rightarrow \sin(x) &= x \\ \text{when } x &= 0 \\ a &= 0\end{aligned}$$


Point $(0, 1)$

Normal line is vertical

$$\Rightarrow x = 0$$

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$

Proof:



$$\sin(x) < x < \tan(x)$$


$$\Rightarrow \frac{1}{\sin(x)} > \frac{1}{x} > \frac{\cos(x)}{\sin(x)}$$

$$1 > \frac{\sin(x)}{x} > \cos(x)$$

$$\lim_{x \rightarrow 0} \Rightarrow 1 > \frac{\sin(x)}{x} > 1$$

$$\Rightarrow \lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$

7. Find the equation to $y = \sqrt{x}$
from the external point $(4, 0)$



$$y = \sqrt{x}, \frac{dy}{dx} = \frac{1}{2\sqrt{x}}, \frac{dy}{dx} \perp = -2\sqrt{x}$$

let the co ordinate on the
curve be (a, \sqrt{a})

$$\text{so } \frac{dy}{dx} \perp = -2\sqrt{a}$$

$$\text{and } \frac{\sqrt{a} - 0}{a - 4} = -2\sqrt{a}$$

$$\sqrt{a} = -2\sqrt{a}(a-4)$$

$$\sqrt{a} = -2a\sqrt{a} + 8\sqrt{a}$$

$$2a\sqrt{a} - 7\sqrt{a} = 0$$

$$\sqrt{a}(2a-7) = 0$$

$$\sqrt{a} = 0 \quad 2a-7 = 0$$

$$\xrightarrow{\text{reject}} \quad a = \frac{7}{2}$$

reject, because
derivative is not
defined at the
end point of a
curve

$$a = \frac{7}{2}$$

$$\text{Point } \left(\frac{7}{2}, \sqrt{\frac{7}{2}}\right)$$

$$\frac{dy}{dx} \perp = -2\sqrt{\frac{7}{2}}$$

$$y - \sqrt{\frac{7}{2}} = -2\sqrt{\frac{7}{2}}(x - \frac{7}{2})$$

$$y = -2\sqrt{\frac{7}{2}}x + 7\sqrt{\frac{7}{2}} + \sqrt{\frac{7}{2}}$$

$$y = -2\sqrt{\frac{7}{2}}x + 8\sqrt{\frac{7}{2}}$$

$$y = -\sqrt{\frac{4 \times 7}{2}}x + \sqrt{\frac{7 \times 64}{2}}$$

$$= -\sqrt{14}x + \sqrt{7 \times 32}$$

$$= -\sqrt{14}x + \sqrt{16 \times 7 \times 2}$$

$$= -\sqrt{14}x + 4\sqrt{14}$$

1.4 Increasing and Decreasing

When analysing functions, we are often interested in the intervals across which the function is increasing or decreasing.

- If the gradient is entirely positive on an interval, the function is increasing on that interval.
- If the gradient is entirely negative on an interval, the function is decreasing on that interval.


If a function changes from increasing to decreasing, it must pass through a stationary point (or cross a region where the function is not differentiable)

Hence we need to divide up the domain into intervals bounded by stationary points and points where the function is not differentiable.


Finding a single gradient on each interval will tell us whether the function is increasing or decreasing.

We can communicate all of this using a **sign diagram**.

1.4.1 Basic sign diagram example



Sign diagram: $f'(x)$ either calculated or observed from a graph



Worked Example:

Find the intervals where $f(x)$ is increasing, for the function: $f(x) = -4x^3 + 15x^2 + 18x + 3$

$$f'(x) = -12x^2 + 30x + 18$$

set $f'(x) = 0$ (to find stationary points)

$$-12x^2 + 30x + 18 = 0$$

$$-6(2x+1)(x-3) = 0$$

$$x = -\frac{1}{2} \quad x = 3$$

$$-6(2x^2 - 5x - 3) = 0$$

$$-6[2x^2 - 6x + x - 3] = 0$$

$$-6[2x(x-3) + (x-3)] = 0$$



Test any
point in
each interval
(choose easy
points)

Test $x = -1$

$$\begin{aligned} f'(-1) &= -12(-1)^2 + 30(-1) + 18 \\ &= -24 \end{aligned}$$


Test $x = 0$

$$f'(0) = +18$$

Test $x = 4$

$$\begin{aligned} f'(4) &= -12(4)^2 + 30(4) + 18 \\ &= -54 \end{aligned}$$

Completed Diagram



1.4.2 Sign diagram example with discontinuities


Below is the graph of

$$y = \frac{x-3}{x^2-4}$$


Because division by 0 is undefined, the graph has vertical asymptotes when $x^2 - 4 = 0$
(i.e when $x = 2$ and $x = -2$)

The derivative is $\frac{dy}{dx} = \frac{-x^2 + 6x - 4}{(x^2 - 4)^2}$ and using a graphics calculator to solve $-x^2 + 6x - 4 = 0$,

we have stationary points at $x \approx 0.764$ and $x \approx 5.236$




Sign diagram: $f'(x)$ either calculated or observed from a graph




1.4.3 Ex 13 C

13c


1. c)



d)



f)



$$e. f(x) = \frac{2}{\sqrt{x}} = 2x^{-\frac{1}{2}}$$

$$f'(x) = 2 - \frac{1}{2}x^{-\frac{3}{2}}$$

$$= -\frac{1}{x^{3/2}} = -\frac{1}{x\sqrt{x}}$$

Domain: $\{x | x > 0\}$

$\text{if } x < 0 \quad \sqrt{x} \text{ undefined}$
 $\text{and } \frac{1}{0} \text{ is undefined.}$

$$\text{Test: } f'(1) = -1$$

$$\begin{array}{c} \ominus \\ \oplus \end{array} \longrightarrow f'(x)$$

Function Decreasing on $(0, \infty)$

$$5. f(x) = x^3 - 3x^2 + 5x + 2$$

$$f'(x) = 3x^2 - 6x + 5$$


$$f'(x) = 0 \Rightarrow \text{no solution}$$

$$\Delta = (-6)^2 - 4 \times 3 \times 5 \\ = 36 - 60 = -24$$

$f'(x) > 0$ for all x

$$\begin{array}{c} \oplus \\ \oplus \end{array} \longrightarrow f'(x)$$

Function always increasing



$$9. f(x) = \frac{-x^2 + 4x - 7}{x-1}$$

$$\left(\frac{u}{v}\right)' = \frac{u'v - uv'}{v^2}$$

$$u = -x^2 + 4x - 7, \quad u' = -2x + 4$$

$$v = x-1, \quad v' = 1$$

$$f'(x) = \frac{(-2x+4)(x-1) - (-x^2+4x-7)1}{(x-1)^2}$$

$$= \frac{-2x^2 + 6x - 4 + x^2 - 4x + 7}{(x-1)^2}$$

$$= \frac{-x^2 + 2x + 3}{(x-1)^2} = -\frac{(x^2 - 2x - 3)}{(x-1)^2}$$

$$= -\frac{(x-3)(x+1)}{(x-1)^2}$$

$$f'(x) = 0 \text{ when } x = 3, x = -1$$

also $f(x)$ undefined when
 $x-1=0, x=1$

$$\begin{array}{c} - \\ -1 \quad 1 \quad 3 \end{array}$$

$$\text{Test } f'(-2) = -\frac{(-5)(-1)}{(-2-1)^2}$$

$$f'(-2) < 0$$

$$\text{Test } f'(0) = -\frac{(-3)(+1)}{(-1)^2}$$

$$f'(0) > 0$$



$$\text{Test } f'(2) = -\frac{(-1)(3)}{(2-1)^2}$$

$$f'(2) > 0$$

$$\text{Test } f'(4) = -\frac{(1)(5)}{(4-1)^2}$$

$$f'(4) < 0$$

$$\begin{array}{c} - \quad + \quad + \quad - \\ \leftarrow \quad \quad \quad \quad \rightarrow \end{array} \frac{dy}{dx}$$



II. c)

$$f(x) = 3 + e^{-x}$$

$$f'(x) = -e^{-x}$$

$$f'(x) \neq 0$$

Domain $\{x \mid x \in \mathbb{R}\}$

$$f'(0) = -e^0 = -1$$

$\xleftarrow{\hspace{1cm}}$ $\xrightarrow{\hspace{1cm}}$
Decreasing everywhere

d) $f(x) = xe^x$

$$(uv)' = uv' + u'v$$

$$f'(x) = xe^x + 1e^x$$

$$= e^x(x+1)$$

$$e^x(x+1) = 0$$

$$e^x \neq 0 \quad x+1=0$$

$$x = -1$$


$$\text{Test } f'(0) = e^0(0+1) = 1$$

$$f'(0) > 0$$

$$\text{Test } f'(-2)$$

$$f'(-2) = e^{-2}(-2+1)$$

$$f'(-2) = -e^{-2} < 0$$



$$f) f(x) = x^3 \ln x$$

Domain $\{x \mid x > 0\}$

$$(uv)' = uv' + u'v$$

$$f'(x) = x^3 \cdot \frac{1}{x} + 3x^2 \ln x$$

$$= x^2 + 3x^2 \ln x$$

$$f'(x) = 0$$

$$\Rightarrow x^2 = 0, \quad x = 0$$

(outside domain)

$$1 + 3 \ln x = 0$$

$$3 \ln x = -1$$

$$\ln x = -\frac{1}{3}$$

$$x = e^{-\frac{1}{3}} \quad x \approx 0.717$$



Test $x = 0.5$

$$f'(0.5) = (0.5)^2 (1 + 3 \ln 0.5)$$

$$= (0.5)^2 (-1.079)$$


$$f'(0.5) < 0$$

Test $x = 1$

$$f'(1) = 1^2 (1 + 3 \ln 1)$$

$$= 1^2 (1)$$

$$f'(1) > 0$$



$$k) f(x) = \ln(x^2 + 4)$$
$$x^2 + 4 > 0 \quad \forall x \in \mathbb{R}$$

\Rightarrow Domain $\{x \mid x \in \mathbb{R}\}$



$$(g(u))' = g'(u) \cdot u'$$

$$\begin{aligned}f'(x) &= \frac{1}{x^2 + 4} \cdot 2x \\&= \frac{2x}{x^2 + 4}\end{aligned}$$

$$f'(x) = 0 \Rightarrow x = 0$$

$$\text{Test } x = -1 \quad f'(-1) = \frac{-2}{5} < 0$$

$$\text{Test } x = 1 \quad f'(1) = \frac{2}{5} > 0$$




1.5 Stationary Points


1.5.1 Turning points (minima, maxima)

From our 'standard' graph we can identify a couple of key features.

$$f(x) = x^3 - 1.5x^2 - 18x + 1$$



If we do not have the graph we can classify the stationary points using a sign diagram.




1.5.2 Stationary points of inflection

Stationary points are not necessarily local maxima or minima.


They may be **inflection** points.

This is where the shape of the curve is changing, but the gradient is not changing sign.



Constructing a sign diagram:

$$\begin{aligned}
 f(x) &= 2x^4 - 6x^3 - 15 \\
 f'(x) &= 8x^3 - 18x^2 \\
 &= 2x^2(4x - 9) \\
 \text{Set } f'(x) &= 0 \\
 \Rightarrow 2x^2(4x - 9) &= 0 \\
 x = 0 \text{ and } x &= \frac{9}{4}
 \end{aligned}$$



If the sign of the gradient repeats across a stationary point, we will have an inflection point.

Note: Inflection points can be at stationary points, or not. Consequently we classify the one above as a **stationary inflection point**

Please note: Summary chart in the textbook.

Stationary point where $f'(a) = 0$	Sign diagram of $f'(x)$ near $x = a$	Shape of curve near $x = a$
local maximum	$\leftarrow + \frac{1}{a} - \frac{f'(x)}{x}$	
local minimum	$\leftarrow - \frac{1}{a} + \frac{f'(x)}{x}$	
stationary inflection	$\leftarrow + \frac{1}{a} + \frac{f'(x)}{x}$ $\leftarrow - \frac{1}{a} - \frac{f'(x)}{x}$	


1.5.3 Greatest or least values (testing boundaries)

If we are given a function which is bounded on an interval, and are finding the greatest/least values on that interval: we need to

- find the values of the stationary points
- find the values at the end points of the function (as these could be greater/less than the stationary point values).

Following the example from the book using

$$y = x^3 - 6x^2 + 5 \text{ On the interval } -2 \leq x \leq 5$$




Example 13

Find the greatest and least value of $y = x^3 - 6x^2 + 5$ on the interval $-2 \leq x \leq 5$.

$$\begin{aligned} \text{Now } \frac{dy}{dx} &= 3x^2 - 12x \\ &= 3x(x - 4) \end{aligned}$$

$$\therefore \frac{dy}{dx} = 0 \text{ when } x = 0 \text{ or } 4.$$

The sign diagram of $\frac{dy}{dx}$ is:



\therefore there is a local maximum at $x = 0$, and a local minimum at $x = 4$.

Critical value (x)	y
-2 (end point)	-27
0 (local maximum)	5
4 (local minimum)	-27
5 (end point)	-20

If an interval is given, we must also check the value of the function at the end points.



The greatest of these values is 5 when $x = 0$.

The least of these values is -27 when $x = -2$ and when $x = 4$.

14 Find the greatest and least value of

1.5.4 Ex 13D

13 D

S.

Find and classify stationary points

g) $f(x) = x - \sqrt{x}$


Domain $\{x \mid x \geq 0, x \in \mathbb{R}\}$

$$f'(x) = 1 - \frac{1}{2\sqrt{x}}$$

$$1 - \frac{1}{2\sqrt{x}} = 0 \Rightarrow 2\sqrt{x} = 1$$


$$\sqrt{x} = \frac{1}{2}$$

$$x = \frac{1}{4}$$



$$\text{Test } f'(1) = 1 - \frac{1}{2} = \frac{1}{2} \quad f'(1) > 0$$

$$\text{Test } f'\left(\frac{1}{16}\right) = 1 - \frac{1}{\frac{2}{4}} = 1 - 2 = -1 \quad f'\left(\frac{1}{16}\right) < 0$$



Local minimum at $x = \frac{1}{4}$

$$f\left(\frac{1}{4}\right) = \frac{1}{4} - \sqrt{\frac{1}{4}} = -\frac{1}{4}$$
$$\left(\frac{1}{4}, -\frac{1}{4}\right)$$

Also note: $f(x) = 0$

when $x - \sqrt{x} = 0$

$$\sqrt{x}(\sqrt{x} - 1) = 0$$

$$x = 0 \quad x = 1$$



x -intercepts $(0, 0) \quad (1, 0)$

for $\forall x, x > 1$

$$x > \sqrt{x}$$

$$\Rightarrow \lim_{x \rightarrow \infty} x - \sqrt{x} = \infty$$

but increasing quite slowly



7. d)

$$y = e^{-x}(x+2)$$

$$\frac{dy}{dx} = e^{-x} \cdot 1 + -e^{-x}(x+2)$$

$$= e^{-x} - e^{-x}(x+2)$$

$$= e^{-x}(1 - (x+2))$$

$$= e^{-x}(-x-1)$$

$$= -e^{-x}(x+1)$$

$$\frac{dy}{dx} = 0 \quad x = -1$$

$$x = -2 \quad \frac{dy}{dx} = -e^2(-2+1) \\ > 0$$

$$x = 0 \quad \frac{dy}{dx}(0) = -e^0(0+1) \\ < 0$$

$$+ \quad | \quad - \\ -1$$



$$y(-1) = e^1(1) = e$$

$$\text{set } y = 0 \Rightarrow x = -2$$

$$x = 0 \quad y = 2$$

$$\lim_{x \rightarrow -\infty} e^{-x}(x+2) = -\infty$$

$$\lim_{x \rightarrow \infty} e^{-x}(x+2) = 0$$



$$10. \quad y = \frac{e^{ax}}{bx}$$

has stationary point at
 $(\frac{1}{3}, \frac{e}{2})$

Find a, b
 Classify the stationary point.

$$\frac{e}{2} = \frac{e^{\frac{a}{3}}}{b}$$

$$\frac{b}{3} = \frac{a}{3}$$

$$\frac{be}{b} = e^{\frac{a}{3}}$$

$$b = 6e^{\frac{a}{3}-1}$$

$$u = e^{ax} \quad u' = ae^{ax}$$

$$v = bx \quad v' = b$$

$$\frac{dy}{dx} = \frac{ae^{ax}(bx) - be^{ax}}{(bx)^2}$$

$$= \frac{be^{ax}(ax-1)}{(bx)^2}$$

$$= \frac{e^{ax}(ax-1)}{b^2x^2}$$

$$\frac{dy}{dx}(\frac{1}{3}) = 0$$

$$\Rightarrow \left(\frac{a}{3} - 1 \right) = 0$$

$$a = 3$$

$$b = 6e^{\frac{3}{3}-1}$$

$$= 6$$

$$y = \frac{e^{3x}}{6x}$$

$$\frac{dy}{dx} = \frac{e^{3x}(3x-1)}{6x^2}$$

Function is undefined at $x=0$



Test: $x = -1$


$$\frac{dy}{dx}(-1) = \frac{e^{-3}(-4)}{6(-1)^2} < 0$$

Test: $x = \frac{1}{4}$

$$\frac{dy}{dx}(\frac{1}{4}) = \frac{e^{3/4}(\frac{3}{4}-1)}{6(\frac{1}{4})^2} < 0$$

Test: $x = 1$

$$\frac{dy}{dx}(1) = \frac{e^3(3-1)}{6(1)^2} > 0$$





$$\lim_{x \rightarrow -\infty} \frac{e^x}{x} = -\infty$$

$$\lim_{x \rightarrow 0^+} \frac{e^x}{x} = +\infty$$

$$\lim_{x \rightarrow 0^-} \frac{e^x}{x} = -\infty$$

$$\lim_{x \rightarrow +\infty} \frac{e^x}{x} = +\infty$$



12. f)

$$f(x) = \sin(2x) + 2\cos(x)$$

$$f'(x) = 2\cos(2x) - 2\sin(x)$$

$$f'(x) = 0$$

$$\Rightarrow 2\cos(2x) - 2\sin(x) = 0$$

Trig identity:

$$\cos(2x) = 1 - 2\sin^2(x)$$

(chapter 9: page 238)

$$2(1 - 2\sin^2(x)) - 2\sin(x) = 0$$

$$-4\sin^2(x) - 2\sin(x) + 2 = 0$$

$$-2(2\sin^2(x) + \sin(x) - 1) = 0$$

$$2y^2 + y - 1 \quad \begin{matrix} -2 \\ +2 \end{matrix} \quad \begin{matrix} \times \\ -1 \end{matrix}$$

$$2y^2 + 2y - y - 1 \quad \begin{matrix} +1 \\ \end{matrix} \quad \begin{matrix} \oplus \\ \end{matrix}$$

$$2y(y+1) - 1(y+1) = (2y-1)(y+1)$$

$$-2(2\sin(x)-1)(\sin(x)-1) = 0$$

$$\sin(x) = \frac{1}{2} \Rightarrow \sin^{-1}\left(\frac{1}{2}\right) = \frac{\pi}{6}$$

Domain $0 \leq x \leq 2\pi$

$$\frac{\pi}{6} + 2k\pi \Rightarrow \frac{\pi}{6} \text{ in } D$$

$$\pi - \frac{\pi}{6} + 2k\pi \Rightarrow \frac{5\pi}{6} \text{ in } D$$

$$\sin(x) = -1 \Rightarrow \sin^{-1}(-1) = -\frac{\pi}{2}$$

$$-\frac{\pi}{2} + 2k\pi \Rightarrow \frac{3\pi}{2} \text{ in } D$$

Solutions: $\frac{\pi}{6}, \frac{5\pi}{6}, \frac{3\pi}{2}$

$$f'(x) = 2\cos(2x) - 2\sin(x)$$




Test $x=0$, Test $x=\frac{\pi}{2}$
 $f'(0)=2 > 0$ $f'(\frac{\pi}{2})=-4 < 0$

Test $x=\pi$

$$f'(\pi)=2 > 0$$

Test $x=2\pi$

$$f'(2\pi)=2 > 0$$



Local maximum at

$$x = \frac{\pi}{6}, f\left(\frac{\pi}{6}\right) = \sin\left(\frac{\pi}{3}\right) + 2\cos\left(\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2} + \frac{2\sqrt{3}}{2} = \frac{3\sqrt{3}}{2} \approx 2.598$$


Local minimum at $x = \frac{5\pi}{6}$

$$f\left(\frac{5\pi}{6}\right) = \sin\left(\frac{5\pi}{3}\right) - 2\cos\left(\frac{5\pi}{6}\right) = -\frac{\sqrt{3}}{2} - \frac{2\sqrt{3}}{2} = -\frac{3\sqrt{3}}{2}$$

Stationary Inflection point at $x = \frac{3\pi}{2}$

$$f\left(\frac{3\pi}{2}\right) = \sin\left(\frac{3\pi}{2}\right) - 2\cos\left(\frac{3\pi}{2}\right) = 0$$

$$\left(\frac{3\pi}{2}, 0\right)$$



16.

$$\begin{aligned}
 f(x) &= \sin(x) \cos(2x) \\
 f'(x) &= \sin(x)(-2\sin(2x)) \\
 &\quad + \cos(x)\cos(2x) \\
 &= -2\sin(x)\sin(2x) + \cos(x)\cos(2x) \\
 &= -2\sin(x)(2\sin(2x)\cos(x)) \\
 &\quad + \cos(x)(2\cos^2(x)-1) \\
 &= -4\sin^2(x)\cos(x) \\
 &\quad + 2\cos^3(x) - \cos(x) \\
 &= -4(1-\cos^2(x))\cos(x) \\
 &\quad + 2\cos^3(x) - \cos(x) \\
 &= -4\cos(x) + 4\cos^3(x) \\
 &\quad + 2\cos^3(x) - \cos(x) \\
 &= 6\cos^3(x) - 5\cos(x)
 \end{aligned}$$

$$f'(x) = 0$$

$$\Rightarrow 6\cos^3(x) - 5\cos(x) = 0$$

$$\cos(x)(6\cos^2(x) - 5) = 0$$

$$\cos(x) = 0$$

$$\cos(x) = \pm \sqrt{\frac{5}{6}}$$

$$\text{Domain } \{x \mid 0 \leq x \leq \pi\}$$

which is Domain of $\cos^{-1}(y)$

$$\cos(x) = 0 \Rightarrow x = \frac{\pi}{2} \approx 1.571$$

$$\cos(x) = \pm \sqrt{\frac{5}{6}}$$


$$x = \cos^{-1}\left(\pm \sqrt{\frac{5}{6}}\right) \approx 0.421$$

$$\cos(x) = -\sqrt{\frac{5}{6}}$$

$$x = \cos^{-1}\left(-\sqrt{\frac{5}{6}}\right)$$

$$\approx 2.721$$

$$(\text{or } \pi - 0.421 = 2.721)$$




$$\text{Test } f'(0) = 1 > 0$$

$$\text{Test } f'(1) = -1.755 < 0$$

$$\text{Test } f'(2) = 1.648 > 0$$


$$\text{Test } f'(\pi) = -1 < 0$$



$$f(0.421) = 0.272$$

$$f\left(\frac{\pi}{2}\right) = -1$$

$$f(2.721) = 0.272$$



18. Prove $\frac{\ln(x)}{x} \leq \frac{1}{e}$

for all $x > 0$

Create a sign diagram and hopefully identify a maximum value for $\frac{\ln x}{x}$

let $f(x) = \frac{\ln(x)}{x}$ $u = \ln x$ $u' = \frac{1}{x}$

$$f'(x) = \frac{\frac{1}{x} \cdot x - \ln(x) \cdot 1}{x^2}$$

$$= \frac{1 - \ln(x)}{x^2}$$

$$f'(x) = 0 \Rightarrow 1 - \ln x = 0$$

$$\ln x = 1$$


$$x = e$$

Test $x = 1$

$$f'(1) = \frac{1-0}{1^2} > 0$$

Test $x = e^2$

$$f'(e^2) = \frac{1 - 2\ln e}{e^4} = \frac{-1}{e^4} < 0$$



Global maximum at $x = e$

$$f(e) = \frac{\ln e}{e} = \frac{1}{e}$$

$$(e, \frac{1}{e})$$

So $\frac{1}{e}$ is largest possible value of $\frac{\ln x}{x}$

$$\Rightarrow \frac{\ln x}{x} \leq \frac{1}{e}$$

19. $f(x) = x - \ln(x)$ $D \{ x > 0 \}$

a) show only local minimum.

b) prove $\ln(x) < x - 1 \quad \forall x > 0$

$$f'(x) = 1 - \frac{1}{x}$$

$$f'(x) = 0$$

$$1 - \frac{1}{x} = 0 \quad x = 1$$



$$f'\left(\frac{1}{2}\right) = 1 - 2 = -1 < 0$$

$$f'(2) = 1 - \frac{1}{2} = \frac{1}{2} > 0$$

local minimum at $(1, 1)$


$$\Rightarrow x - \ln(x) \geq 1 \quad \forall x > 0$$

$$x - 1 \geq \ln(x)$$

$$\ln(x) \leq x - 1$$

1.6 Shape

Looking at the shape of the graph below, we can see that over a certain interval, the curve is turning downwards **it is concave down**, and on then it changes to be turning upwards **concave up**.




If the curve is concave down the **gradient** is decreasing, so we consider the second derivative

Leibnitz	or	Newton
$\frac{d}{dx} \left(\frac{dy}{dx} \right) < 0$		$(f'(x))' < 0$
$\Rightarrow \frac{d^2y}{dx^2} < 0$		$\Rightarrow f''(x) < 0$

Similarly, if the curve is concave up the **gradient** is increasing, so

$$\frac{d^2y}{dx^2} > 0 \quad \text{or} \quad f''(x) > 0$$

We can construct a sign diagram for (in this case) $f''(x) = 6x - 3$, and $f''(x) = 0$ when $x = \frac{1}{2}$



We can also observe that there is an **non-stationary inflection point** at $x = \frac{1}{2}$, $f\left(\frac{1}{2}\right) = -\frac{37}{4}$

1.6.1 Ex 13E

13 E

3 b

Determine the shape of

$$y = -2(x-3)(x+1)$$

$$\begin{aligned} \frac{dy}{dx} &= -2[(x-3)1 + 1(x+1)] \\ &= -2(2x-2) \\ &= -4x+4 \end{aligned}$$

$$\frac{d^2y}{dx^2} = -4 \quad \frac{d^2y}{dx^2} < 0 \text{ always}$$

$$\xleftarrow{\text{---}} \xrightarrow{\frac{d^2y}{dx^2}}$$

function is concave down always.

4 f

$$f(x) = x^4 - 12x^3$$

$$f'(x) = 4x^3 - 36x^2$$

$$4x^3 - 36x^2 = 0$$

$$4x^2(x-9) = 0$$

$$x=0 \quad x=9$$

$$\begin{array}{c} - \\ \hline - & - & + \\ \hline 0 & & 9 \end{array}$$

$$f''(x) = 12x^2 - 72x$$

$$f''(x) = 0$$

$$12x^2 - 72x = 0$$

$$12x(x-6) = 0$$

$$x=0 \quad x=6$$

$$\begin{array}{c} + \\ \hline + & - & + \\ \hline 0 & & 6 \end{array}$$



$f(x)$ is decreasing for $x < 9$

$f(x)$ is increasing for $x > 9$

$f(x)$ is concave up $x < 0$

concave down $0 < x < 6$

concave up $x > 6$



$$7. f(x) = \frac{e^x}{x}$$

a) When $x=0$, $f(x)$ is undefined
 \Rightarrow y -intercept does not exist.

for x -intercepts $\frac{e^x}{x} = 0$

but $e^x \neq 0$, so no x -intercepts

b) as $x \rightarrow \infty$

e^x gets very large

$\frac{1}{x}$ gets very small

but e^x is an exponential

$$\Rightarrow \lim_{x \rightarrow \infty} \frac{e^x}{x} = +\infty$$

and $\lim_{x \rightarrow -\infty} \frac{e^x}{x} = 0$

c) $f(x) = \frac{e^x}{x}$


$$f'(x) = \frac{xe^x - e^x \cdot 1}{x^2}$$

$$= \frac{e^x(x-1)}{x^2}$$

$$f'(x) = 0$$

$$\Rightarrow e^x(x-1) = 0$$

$x=1$ is a stationary pt $(1, e)$



$(1, e)$ is a stationary point
 and is a local minimum.

$$f'(x) = \frac{xe^x - e^x}{x^2}$$

$$u = xe^x - e^x$$

$$u' = xe^x + e^x - e^x$$

$$= xe^x$$

$$v = x^2$$

$$v' = 2x$$

$$f''(x) = \frac{xe^x(x^2) - (xe^x - e^x)2x}{x^4}$$

$$= \frac{x^3e^x - 2x^2e^x + 2xe^x}{x^4}$$

$$= \frac{xe^x(x^2 - 2x + 2)}{x^4}$$

$$xe^x(x^2 - 2x + 2) = 0$$

$$x=0 \quad \text{no sol.}$$

so no inflection points.

but need to look at
 either side of undefined

$$\begin{array}{c|cc} - & + & \\ \hline & & f''(x) \end{array}$$


$$\begin{array}{ll} \text{Concave down} & 0 \\ x < 0 & \end{array} \quad \begin{array}{ll} \text{Concave up} & 0 \\ x > 0 & \end{array}$$

tangent at $x=-1$

$$f'(-1) = -\frac{2}{e} \quad f(-1) = -\frac{1}{e}$$

$$y + \frac{1}{e} = -\frac{2}{e}(x+1)$$


$$y = -\frac{2x}{e} - \frac{3}{e} \quad \text{or} \quad 2x + ey = -3$$



1.7 Inflection Points

At an inflection point:

- the tangent line crosses the curve
- the shape of the curve is changing from concave up to concave down (or vice versa).



We can identify inflection points by

- Solving $f''(x) = 0$
- Then using a sign diagram to verify that the points we have found are actually inflection points.
The sign diagram must **change signs** across the 0 for the points to be inflection points.

1.7.1 Ex 13F

Find stationary and inflection points for
 $y = x^3(5-x)$

$$y = 5x^3 - x^4$$

$$\frac{dy}{dx} = 15x^2 - 4x^3$$

$$15x^2 - 4x^3 = 0$$

$$x^2(15 - 4x) = 0$$

$$x=0 \quad x = \frac{15}{4} = 3.75$$

Stationary points at $(0, 0)$ and $(3.75, 65.918)$

$$\frac{d^2y}{dx^2} = 30x - 12x^2$$

$$= 6x(5 - 2x)$$

If $x=0$

$$\frac{d^2y}{dx^2} = 0$$

If $x = 3.75$


$$\frac{d^2y}{dx^2} = -56.25 \quad \curvearrowleft$$

unknown at this stage.

$(3.75, 65.918)$ is a local max.

Sign diagram for $\frac{d^2y}{dx^2}$ $6x(5-2x)=0$


$$x=0 \quad x=2.5$$



Change of sign at
 $x=0$

Change of sign at
 2.5

So $(0,0)$ is a stationary inflection point so $(2.5, 39.0625)$ is a non-stationary inflection point



Find stationary and inflection points for:

$$f(x) = \frac{x^6}{30} + 6.4x$$

$$f'(x) = \frac{6x^5}{30} + 6.4 = \frac{x^5}{5} + 6.4$$

$$\frac{x^5}{5} + 6.4 = 0$$

$$x^5 = -32$$

$$x = \sqrt[5]{-32} = -2$$

Stationary point at $(-2, -10\frac{2}{3})$

$$f''(x) = \frac{5x^4}{5} = x^4$$

Sign diagram
not required.

$$f''(-2) = 16 > 0$$

Stationary point is a local minimum.

Set $f''(x) = 0$

$$\Rightarrow x = 0$$

could be an inflection point


but $\begin{array}{c} + \\ \hline + & + \\ \hline 0 \end{array} f''(x)$


Does not change sign

so $(0, 0)$ is not an inflection point.

A local minimum at $(-2, -10\frac{2}{3})$

No inflection points.





1.8 Understanding functions and their derivatives

Chapter: 2 Applications of differentiation

$$\text{The equation is: } 9a - 4 = 14 + 3a$$

$$\text{Subtract } 3a: \quad 6a - 4 = 14$$

$$\text{Subtract } 4: \quad 6a = 18$$

$$\text{Divide by } 6: \quad a = 3$$

$$A\widehat{B}C$$

$$A\widehat{B}CC$$

$$A\dot{B}C$$

$$N\tilde{a}$$

$$X \sim \mathcal{N}(\mu, \sigma^2)$$

