


Sequences and Series

Kh notes

Contents

1	Introduction (Number Sequences)	2
2	Arithmetic Sequences	3
2.1	Ex 5B.1	3
2.2	Ex 5B.2	6
3	Geometric Sequences	6
3.1	Ex 5C	6
4	Growth and Decay	6
4.1	Ex 5D	6
5	Financial Mathematics	6
5.1	Compound Interest	6
5.1.1	Ex 5E.1	7
5.2	Inflation	7
5.2.1	Ex 5E.2	7
5.3	Real Value of an Investment	7
5.3.1	Ex 5E.3	7
5.4	Depreciation	7
5.4.1	Ex 5E.4	7
5.5	Using Financial Models	7
5.5.1	Ex 5E.5	7
6	Series	7
6.1	Sigma Notation	7
6.1.1	Ex 5F	7
6.2	Arithmetic Series	7
6.2.1	Ex 5G	7
6.3	Finite Geometric Series	7
6.3.1	Ex 5H	7
6.4	Infinite Geometric Series	7
6.4.1	Ex 5I	7

1 Introduction (Number Sequences)



Grains of rice problem.

How do we define this problem mathematically?

- Create a number sequence for each square: $1, 2, 4, 8, 16, \dots, 2^{62}, 2^{63}$
- Consider relations between the number in each square: $s_2 = 2s_1$, or $s_9 = 2^8$
- Create a formula for each square: $s_n = 2^{n-1}$
- Communicate the idea of summing : $1 + 2 + 4 + 8 + 16, + \dots +, 2^{62} + 2^{63} = \sum_{i=1}^{64} 2^{i-1}$
- Actually solve the problem $1 + 2 + 4 + 8 + 16, + \dots +, 2^{62} + 2^{63} = ?$

Consider:

$$\begin{aligned} 1 + 1 + 2 + 4 + 8 &= 2 + 2 + 4 + 8 \\ &= 2(1 + 1 + 2 + 4) \\ &= 2(2 + 2 + 4) \\ &= 2(2(1 + 1 + 2)) \\ &= 2(2(2 + 2)) \\ &= 2(2(2(1 + 1))) \\ &= 2(2(2(2))) = 2^4 \end{aligned}$$

2 Arithmetic Sequences


An example of an arithmetic sequence is:

$$14, 17, 20, 23, \dots$$

An arithmetic sequence has a common difference between the terms (in this case 3)

- We need to be able to describe the sequence:
The sequence has a first term of 14 and increases by 3 each time
- We define u_n to be the n^{th} term of the sequence
- Unless otherwise stated u_1 is the first term.
- We define d as the common difference ($d = u_{n+1} - u_n$).
- $\{u_n\}$ represents the whole sequence.

Note: the sequence is linear (so the difference is the gradient).



The middle of any three consecutive terms is the arithmetic mean of the other two terms.

General term formula:

$$u_n = u_1 + (n - 1)d$$

(this can then be simplified to a ' $y = mx + c$ ' form)

2.1 Ex 5B.1

- Showing and proving a sequence is arithmetic,
- Finding terms
- Working with k
- Inserting terms between

Example Questions: 5B.1

7. A sequence is defined by

$$u_n = \frac{71 - 7n}{2}$$

a) prove the sequence is arithmetic

Show $u_{n+1} - u_n = d$

$$\begin{aligned} u_{n+1} - u_n &= \frac{71 - 7(n+1)}{2} - \frac{71 - 7n}{2} \\ &= \frac{71 - 7n - 7 - 71 + 7n}{2} \\ &= \frac{-7}{2} = -3.5 \end{aligned}$$

So $u_{n+1} - u_n = -3.5$ for all $n \in \mathbb{Z}^+$

b) $u_n = \frac{71}{2} - \frac{7n}{2}$

$$u_1 = \frac{71}{2} - \frac{7}{2} = \frac{64}{2} = 32$$

$$d = -\frac{7}{2}$$

c) $u_{75} = \frac{71 - 7(75)}{2} = -227$

d) $u_n < -200$

$$\frac{71 - 7n}{2} < -200$$

$$71 - 7n < -400$$

$$7n > 71 + 400$$

$$7n > 471$$

$$n > 67 \frac{2}{7}$$

for $n \geq 68$ $u_n < -200$

$$\begin{aligned} \text{check } u_{68} &= \frac{71 - 7(68)}{2} \\ &= -202.5 \end{aligned}$$

12. ^{d)} Find k given the consecutive arithmetic terms:

$$\begin{array}{ccc} 2k+18 & , & -2-k & , & 2k+2 \\ u_1 & & u_2 & & u_3 \end{array}$$

$$u_2 - u_1 = u_3 - u_2$$

$$-2-k - (2k+18) = 2k+2 - (-2-k)$$

$$-2-k-2k-18 = 2k+2+2+k$$

$$-20-3k = 3k+4$$

$$6k = -24$$

$$k = -4$$

check:

$$10, 2, -6$$

$$d = -8$$

Other notes:

Could have:

$$\frac{2k+2+2k+18}{2} = -2-k$$

$$4k+20 = -4-2k$$

$$6k = -24$$

or

$$u_2 - u_1 = u_3 - u_2$$

$$2u_2 - u_1 = u_3$$

$$2(-2-k) - (2k+18) = 2k+2$$

$$-4-2k-2k-18 = 2k+2$$

$$-22-4k = 2k+2$$

$$6k = -24$$

13. Find the general term u_n for the arithmetic sequence
 $u_5 = -2$ and $u_{12} = -12\frac{1}{2}$

$$u_5 = a + 4d$$

$$u_{12} = a + 11d$$

$$a + 4d = -2 \quad (1)$$

$$a + 11d = -\frac{25}{2} \quad (2)$$

$$\frac{a + 4d = -2}{a + 11d = -\frac{25}{2}} \quad (1) - (2)$$

$$-7d = +\frac{21}{2}$$

$$d = \frac{+21}{2} \times -\frac{1}{7} = -\frac{3}{2}$$

from (1) $a = -2 - 4d$
 $= -2 - 4\left(-\frac{3}{2}\right)$
 $= -2 + 6 = 4$

$$u_n = 4 - (n-1)\frac{3}{2}$$

Test: $u_5 = 4 - (5-1)\frac{3}{2}$

$$= 4 - 6 = -2$$

$$u_{12} = 4 - (12-1)\frac{3}{2}$$

$$= 4 - \frac{33}{2}$$

$$= \frac{8-33}{2}$$

$$= -\frac{25}{2} = -12\frac{1}{2}$$

15. Insert six numbers between -1 and 32 so all 8 are in an arithmetic sequence

$$a = -1$$

$$u_8 = -1 + 7d$$

$$-1 + 7d = 32$$

$$7d = 33$$

$$d = \frac{33}{7}$$

$$u_n = -1 + (n-1)\frac{33}{7}$$

or $d = \frac{u_8 - u_1}{7} = \frac{33}{7}$

2.2 Ex 5B.2

- Approximations using arithmetic sequences

3 Geometric Sequences

3.1 Ex 5C

4 Growth and Decay

Starter Questions:

1. A school had 1200 students and a year later this has increased by 8% . How many students are now in the school?
2. Mary buys a car for \$40,000 and in one year its price has decreased by 12%. What is the value of it now?
3. The population of Sydney is currently 5.2 million. If it increases at a rate of 1.25% annually, what will the population be after 3 years?

4.1 Ex 5D

5 Financial Mathematics

5.1 Compound Interest

$$u_n = u_0(1 + i)^n$$

u_0 Initial Investment (Principal)

i Interest rate per compounding period

n Number of periods

u_n The final value of the investment

5.1.1 Ex 5E.1

5.2 Inflation

5.2.1 Ex 5E.2

5.3 Real Value of an Investment

5.3.1 Ex 5E.3

5.4 Depreciation

5.4.1 Ex 5E.4

5.5 Using Financial Models

5.5.1 Ex 5E.5

6 Series

6.1 Sigma Notation

6.1.1 Ex 5F

6.2 Arithmetic Series

6.2.1 Ex 5G

6.3 Finite Geometric Series

6.3.1 Ex 5H

6.4 Infinite Geometric Series

6.4.1 Ex 5I