

Properties of curves and applications of differentiation

Kh notes

Contents

1 Properties of curves	2
1.1 Review Questions	2
1.1.1 Some fundamental derivatives:	2
1.1.2 Rules of differentiation:	2
1.2 Tangents	3
1.2.1 Horizontal Tangents	4
1.2.2 Ex13A	5
1.3 Normals	14
1.3.1 Ex 13B	14
1.4 Increasing and Decreasing	19
1.4.1 Basic Sign Diagram example	19
1.4.2 Sign Diagram example with discontinuites	20
1.5 Stationary Points	21
1.5.1 Turning points (minima, maxima)	21
1.5.2 Stationary points of inflection	21
1.6 Shape	21
1.7 Inflection Points	21
1.8 Understanding functions and their derivatives	21
2 Applications of differentiation	22

Chapter: 1 Properties of curves

1.1 Review Questions

1.1.1 Some fundamental derivatives:

Function	Derivative
$f(x) = x^n$	$f'(x) = nx^{n-1} \quad (n \in \mathbb{R})$
$f(x) = e^x$	$f'(x) = e^x$
$f(x) = \ln x$	$f'(x) = \frac{1}{x}$
$f(x) = \sqrt{x}$	$f'(x) = \frac{1}{2\sqrt{x}}$
$f(x) = \sin x$	$f'(x) = \cos x$
$f(x) = \cos x$	$f'(x) = -\sin x$
$f(x) = \tan x$	$f'(x) = \sec^2 x$

1.1.2 Rules of differentiation:

Chain Rule:

$$y = g(u_{(x)})$$
$$\frac{dy}{dx} = g'(u_{(x)})u'_{(x)}$$

Product Rule:

$$y = u_{(x)}v_{(x)}$$
$$\frac{dy}{dx} = u_{(x)}v'_{(x)} + u'_{(x)}v_{(x)}$$

Quotient Rule:


$$y = \frac{u_{(x)}}{v_{(x)}}$$
$$\frac{dy}{dx} = \frac{u'_{(x)}v_{(x)} - u_{(x)}v'_{(x)}}{[v_{(x)}]^2}$$

1.2 Tangents

- The tangent to a curve at a point A is the best approximating straight line to the curve at point A.
- (Leibniz definition) Tangent to the curve $y = f(x)$ at the point $(a, f(a))$ is the line through the infinitely close pair of points either side of $f(a)$

$$\frac{y - f(a)}{x - a} = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$$

- It is a single point of contact with the curve (although it may intersect the curve at some other point)



For the function $y = f(x)$, and some $x = a$

- $(a, f(a))$ is on the curve
- $f'(a)$ is the gradient of the curve at $x = a$


$$\begin{aligned} \frac{y - f(a)}{x - a} &= f'(a) \\ \Rightarrow y &= f'(a)(x - a) + f(a) \end{aligned}$$

Is the equation of the tangent line

1.2.1 Horizontal Tangents

Horizontal tangents have a gradient of 0.

These will be very important later, when we investigate stationary points in more detail.



In general, we will need to find where the stationary points are:

- Find $f'(x)$
 - Solve $f'(x) = 0$
 - Substitute the solution(s) into $f(x)$ to find the constant terms for the horizontal line equation
-

1.2.2 Ex13A

- Tangent equations from root and polynomial functions
- Horizontal tangents
- Natural log and exponent questions
- Concept questions

4 worked examples

Find the equation of the tangent to:

a) $y = x - 2x^2 + 3$ at $x=2$

$$\frac{dy}{dx} = 1 - 4x$$

when $x=2$

$$\frac{dy}{dx} = 1 - 4(2) = -7$$

and $y = (2) - 2(2)^2 + 3$
 $= -3$

$$\frac{dy}{dx} = -7 \quad \text{Point } (2, -3)$$

$$y - (-3) = -7(x - 2)$$

$$y + 3 = -7x + 14$$

$$y = -7x + 11$$

d) $y = \frac{4}{\sqrt{x}}$ at $(1, 4)$

$$y = 4x^{-\frac{1}{2}}$$

$$\frac{dy}{dx} = 4 \cdot -\frac{1}{2}x^{-\frac{3}{2}}$$

$$= -2x^{-\frac{3}{2}} = \frac{-2}{x^{\frac{3}{2}}}$$

$x = 1$

$$\frac{dy}{dx} = \frac{-2}{1} = -2$$

Point $(1, 4)$

$$y - 4 = -2(x - 1)$$

$$y - 4 = -2x + 2$$

$$y = -2x + 6$$

(Horizontal Tangents)

3. c)

Find the equation of horizontal tangents to

$$y = \sqrt{x} + \frac{1}{\sqrt{x}}$$

$$\Rightarrow y = x^{\frac{1}{2}} + x^{-\frac{1}{2}}$$

$$\frac{dy}{dx} = \frac{1}{2}x^{-\frac{1}{2}} + -\frac{1}{2}x^{-\frac{3}{2}}$$

(factorise using lowest power)

$$\frac{dy}{dx} = \frac{1}{2}x^{-\frac{3}{2}}(x^{\frac{1}{2}} - 1)$$

Set $\frac{dy}{dx} = 0$

$$\frac{1}{2}x^{-\frac{3}{2}}(x^{\frac{1}{2}} - 1) = 0$$

$$x^{-\frac{3}{2}} \neq 0 \quad x^{\frac{1}{2}} - 1 = 0$$

Find y

$$y = \sqrt{1} + \frac{1}{\sqrt{1}} = 1 + 1$$

$$= 2$$

Horizontal tangent at $(1, 2)$

Equation: $y = 2$

$$\frac{dy}{dx} = \frac{1}{2x^{\frac{1}{2}}} - \frac{1}{2x^{\frac{3}{2}}}$$

$$= \frac{x - 1}{2x^{\frac{3}{2}}}$$

$$\frac{x - 1}{2x^{\frac{3}{2}}} = 0 \Rightarrow x - 1 = 0$$

5. Find another tangent to

$$y = 1 - 3x + 12x^2 - 8x^3$$

which is parallel to the tangent at $(1, 2)$

$$\frac{dy}{dx} = -3 + 24x - 24x^2$$

when $x=1$

$$\begin{aligned}\frac{dy}{dx} &= -3 + 24(1) - 24(1)^2 \\ &= -3\end{aligned}$$

so gradient is -3

Find all x values where gradient $= -3$

$$-3 + 24x - 24x^2 = -3$$

$$24x - 24x^2 = 0$$

$$x - x^2 = 0$$

$$x(1-x) = 0$$

$$x=0 \quad \underbrace{x=1}_{a} \quad) \text{ already have}$$

Find equation of tangent at $x=0$


$$y = 1$$

$$y - 1 = -3(x - 0)$$

$$y - 1 = -3x, \quad y = -3x + 1$$

$$\text{other tangent } y - 2 = -3(x - 1)$$

$$y = -3x + 5$$



6. $y = x^2 + ax + b$

when $x=1$ tangent

$$is 2x+y=6$$

$$y = -2x + 6$$

$$\frac{dy}{dx} = -2 \quad y = 4$$

$(1, 4)$ point on curve.

-2 gradient at that point.

$$\frac{dy}{dx} = 2x + a$$

$$-2 = 2(1) + a$$

$$a = -4$$

$$y = x^2 - 4x + b$$


$$4 = (1)^2 - 4(1) + b$$

$$4 = -3 + b$$

$$b = 7.$$

$$\underline{y = x^2 - 4x + 7}$$

7. Find a and b



What do we know?

- when $x=4 \frac{dy}{dx} = 1$
- $(4, 4+2) = (4, 6)$ is on $y = a\sqrt{x} + b$

① Using $(4, 6)$

$$\begin{aligned} 6 &= a\sqrt{4} + b(4) \\ 6 &= 2a + 4b \\ \Rightarrow 3 &= a + 2b \end{aligned}$$

$$② \frac{dy}{dx} = \frac{a}{2\sqrt{x}} + b$$

$$1 = \frac{a}{2\sqrt{4}} + b$$

$$1 = \frac{a}{4} + b$$

$$4 = a + 4b$$

Equation

$$\begin{aligned} 3 &= a + 2b & ① \\ 4 &= a + 4b & ② \end{aligned}$$

$$② - ① \quad 1 = 2b$$

$$b = \frac{1}{2}$$

$$\begin{aligned} \text{using } ① \quad a &= 3 - 2b \\ &= 3 - 2\left(\frac{1}{2}\right) \\ &= 2 \end{aligned}$$

$$y = 2\sqrt{x} + \frac{x}{2}$$

8. Show equation of

tangent to $y = 2x^2 - 1$

at $x=a$ is

$$4ax - y = 2a^2 + 1$$

Find point:

$$x = a$$

$$y = 2a^2 - 1$$

$$(a, 2a^2 - 1)$$

$$\frac{dy}{dx} = 4x$$

$$= 4a$$

$$y - (2a^2 - 1) = 4a(x - a)$$

$$y - 2a^2 + 1 = 4ax - 4a^2$$

$$(-y) \quad -2a^2 + 1 = 4ax - 4a^2 - y$$

$$(+4a^2) \quad 2a^2 + 1 = 4ax - y$$

$$10) \frac{dy}{dx} = \ln \sqrt{x}$$

$$y = \ln(x^{\frac{1}{2}})$$

$$y = \frac{1}{2} \ln(x)$$

$$\frac{dy}{dx} = \frac{1}{2x}$$

$$y = -1$$

$$\frac{1}{2} \ln(x) = -1$$

$$\ln(x) = 2$$

$$x = e^2$$

$$(e^{-2}, -1)$$

$$\frac{dy}{dx} = \frac{1}{2e^{-2}}$$

$$y + 1 = \frac{1}{2e^{-2}}(x - e^{-2})$$

$$y = \frac{x}{2e^{-2}} - \frac{1}{2} - 1$$

$$y = \frac{e^2}{2}x - \frac{3}{2}$$

Using $\ln x$ and e^x

Find the equation of the tangent to $y = \ln x$ where $y = -1$

• Find x

$$\ln x = -1 \quad (\log_e x = 1)$$

$$x = e^{-1}$$

$$\frac{dy}{dx} = \frac{1}{x} = x^{-1}$$

$$x = e^{-1}, \quad \frac{dy}{dx} = (e^{-1})^{-1} = e^1 = e$$

Point $(e^{-1}, -1)$


$$\frac{dy}{dx} = e$$

$$y + 1 = e(x - e^{-1})$$

$$y + 1 = ex - e^1 e^{-1}$$

$$y + 1 = ex - 1$$

$$y = ex - 2$$



$$12. f(x) = \ln(x(x-2))$$

Domain of log function > 0

$$\Rightarrow x(x-2) > 0$$

$$\text{solve } x(x-2)=0$$

$$\begin{array}{c} x=0 \\ x=2 \end{array}$$

Domain $x < 0 \cup x > 2$

$$\{x | x < 0 \cup x > 2\}$$

$$f'(x) = \frac{1}{x(x-2)} \cdot \frac{d}{dx}(x^2 - 2x)$$

$$= \frac{1}{x(x-2)} \cdot 2x - 2$$

$$= \frac{2(x-1)}{x(x-2)}$$

$$= \frac{2x-2}{x^2-2x}$$

$$= \frac{2x-2}{x(x-2)}$$

$$= \frac{2x+x-2}{x(x-2)}$$

$$= \frac{1}{x} + \frac{1}{x-2}$$

$$x=3, f'(x) = \frac{1}{3} + \frac{1}{1} = \frac{4}{3}$$

$$x=3, f(3) = \ln(3(3-2)) = \ln 3$$

$$y - \ln 3 = \frac{4}{3}(x-3)$$

$$y = \frac{4}{3}x - 4 + \ln 3$$

$$13. y = x^2 e^x$$

$$y = uv \quad \frac{dy}{dx} = uv' + u'v$$

(product rule)

$$\begin{aligned} \frac{dy}{dx} &= x^2 e^x + 2x e^x \\ &= x e^x (x+2) \end{aligned}$$

$$x=1$$

$$\frac{dy}{dx} = e(3) = 3e$$

$$x=1, y=e$$

Gradient: $3e$

Point: $(1, e)$

$$y - e = 3e(x-1)$$

$$y = 3ex - 2e \quad \text{equation of tangent}$$

Find intercepts.

$$x=0 \quad y = -2e$$

$$y=0 \quad x = \frac{2e}{3e}$$

$$x = \frac{2}{3}$$

Intercepts

$$(0, -2e) \quad y\text{-intercept}$$

$$\left(\frac{2}{3}, 0\right) \quad x\text{-intercept}$$

$$14. \quad y = 3xe^{\frac{x}{2}}$$

$$\begin{aligned} \frac{dy}{dx} &= 3x(e^{\frac{x}{2}})' + (3x)' e^{\frac{x}{2}} \\ &= 3x e^{\frac{x}{2}} \cdot \frac{1}{2} + 3e^{\frac{x}{2}} \\ &= 3e^{\frac{x}{2}} \left(\frac{x}{2} + 1 \right) \end{aligned}$$

When $x = -1$

$$\begin{aligned} \frac{dy}{dx} &= 3e^{-\frac{1}{2}} \left(-\frac{1}{2} + 1 \right) \\ &= \frac{3}{2} e^{-\frac{1}{2}} \end{aligned}$$

$$x = -1, \quad y = -3e^{-\frac{1}{2}}$$

$$y + 3e^{-\frac{1}{2}} = \frac{3e^{-\frac{1}{2}}}{2}(x + 1)$$

$$\begin{aligned} y &= \frac{3e^{-\frac{1}{2}}}{2}x + \frac{3e^{-\frac{1}{2}}}{2} - 3e^{-\frac{1}{2}} \\ &= \frac{3e^{-\frac{1}{2}}}{2}x - \frac{3e^{-\frac{1}{2}}}{2} \end{aligned}$$

$$\text{if } x = 0 \quad y = -\frac{3e^{-\frac{1}{2}}}{2}$$

$$\text{if } y = 0 \quad x = 1$$

$$\frac{3e^{-\frac{1}{2}}}{2} \sqrt{\frac{1}{A}} = \frac{3e^{-\frac{1}{2}}}{4}$$

$$A = \frac{3}{4\sqrt{e}} \text{ units}^2$$

15. Find the equation of the tangent to $y = \frac{1}{\sin 2x}$ at the point where

$$x = \frac{\pi}{4}$$

$$u = 1, u' = 0 \quad v = \sin 2x, v' = 2\cos(2x) \quad \left(\frac{u}{v}\right)' = \frac{u'v - uv'}{v^2}$$

$$\begin{aligned} \frac{dy}{dx} &= -\frac{1 \cdot 2\cos(2x)}{\sin^2(2x)} = -\frac{2\cos(2x)}{\sin^2(2x)} \\ &= -2\cot(2x)\csc(x) \end{aligned}$$


$$\text{when } x = \frac{\pi}{4} \quad \frac{dy}{dx} = -2\cot\left(\frac{\pi}{2}\right) = 0$$

$$\text{when } x = \frac{\pi}{4}, \quad y = \frac{1}{\sin\left(\frac{\pi}{2}\right)} = 1$$

Horizontal tangent

$$y = 1$$

27.



$$\text{Common Points } (b, ab^2)$$

$$\text{a) } (b, \ln b)$$

$$\Rightarrow ab^2 = \ln b$$

Common Tangent

$$\frac{dy}{dx} = 2ax \quad , \quad \frac{dy}{dx} = \frac{1}{x}$$

$$x=b \quad 2ab = \frac{1}{b} \Rightarrow 2ab^2 = 1$$

$$\text{b) } \begin{aligned} ab^2 &= \ln b & \textcircled{1} \\ 2ab^2 &= 1 & \textcircled{2} \end{aligned}$$

$$\begin{aligned} 2ab^2 &= 2\ln b \\ 2ab^2 &= 1 \end{aligned}$$

$$\begin{aligned} 2\ln b &= 1 \\ \ln b &= \frac{1}{2} \end{aligned}$$

$$b = e^{\frac{1}{2}} = \sqrt{e}$$

$$\text{Point } (e^{\frac{1}{2}}, \ln e^{\frac{1}{2}}) = (e^{\frac{1}{2}}, \frac{1}{2})$$

$$\text{c) } a = \frac{1}{2b^2}$$

$$a = \frac{1}{2e}$$

$$\text{d) } a = \frac{1}{2e}, \quad b = \sqrt{e}$$

$$\text{Point } (e^{\frac{1}{2}}, \frac{1}{2})$$

$$\text{Gradient } \frac{dy}{dx} = \frac{1}{b} = \frac{1}{\sqrt{e}} = e^{-\frac{1}{2}}$$

$$y - \frac{1}{2} = e^{-\frac{1}{2}}(x - e^{\frac{1}{2}})$$

$$y - \frac{1}{2} = e^{-\frac{1}{2}}x - 1$$

$$y = 2\sqrt{e}x - \frac{1}{2}$$

$$28. \quad P(x) = ax^2$$

$$P'(x) = 2ax$$

$$x=s \quad (s, as^2) \quad P'(s) = 2as$$

$$y - as^2 = 2as(x - s)$$

$$\begin{aligned} y &= 2asx - 2as^2 + as^2 \\ y &= 2asx - as^2 \end{aligned}$$

$$\begin{aligned} x=t \\ y &= 2atx - at^2 \end{aligned}$$

$$2asx - as^2 = 2atx - at^2$$

$$x(2as - 2at) = as^2 - at^2$$

$$x = \frac{a(s^2 - t^2)}{2a(s-t)}$$

$$= \frac{(s+t)(s-t)}{2(s-t)}$$

$$= \frac{s+t}{2}$$

If perpendicular

$$2as \cdot 2at = -1$$

$$y = 2atx - at^2$$

$$x_c = \frac{s+t}{2}$$

$$y = \frac{2at(s+t)}{2} - at^2$$

$$= ast + at^2 - at^2$$

$$= ast$$

$$t = \frac{-1}{4a^2s}$$

$$y = ast$$


$$= as \cdot \frac{-1}{4a^2s}$$

$$= -\frac{1}{4a}$$

1.3 Normals

The product of the gradients of perpendicular lines = -1

There are various proofs, one example below.




For the function $y = f(x)$, and some $x = a$

- $(a, f(a))$ is on the curve
- $-\frac{1}{f'(a)}$ is the gradient of the **normal to the curve** at $x = a$

$$\frac{y - f(a)}{x - a} = -\frac{1}{f'(a)}$$

$$\Rightarrow y = -\frac{1}{f'(a)}(x - a) + f(a)$$

Is the equation of the tangent line



1.3.1 Ex 13B

13B

1. d)
Find the equation of the normal to:

$$y = 8\sqrt{x} - \frac{1}{x^2} \text{ at } x=1$$

Point $x=1$

$$y = 8\sqrt{1} - \frac{1}{1^2} = 7 \quad (1, 7)$$

Gradient:

$$\begin{aligned} y &= 8x^{\frac{1}{2}} - x^{-2} \\ \frac{dy}{dx} &= 8 \cdot \frac{1}{2} x^{-\frac{1}{2}} - -2x^{-3} \\ &= \frac{4}{x^{\frac{1}{2}}} + \frac{2}{x^3} \\ &= \frac{4x^{\frac{5}{2}} + 2}{x^3} \\ \frac{dy}{dx} \perp &= \frac{-x^3}{4x^{\frac{5}{2}} + 2} \end{aligned}$$

$$x=1$$

$$\frac{dy}{dx} \perp = \frac{-1}{4+2} = -\frac{1}{6}$$

$$(1, 7) \text{ and } -\frac{1}{6}$$

$$y-7 = -\frac{1}{6}(x-1)$$

$$y-7 = -\frac{x}{6} + \frac{1}{6}$$

$$y = -\frac{x}{6} + \frac{43}{6}$$

or

$$6y + x = 43$$

$$\text{or } x + 6y - 43 = 0 \quad \circ$$

3. Find the equation of
c) the normal to
 $y = e^{2x-1}$ at $x=1$

Point $(1, e^1)$

$$\frac{dy}{dx} = e^{2x-1} \cdot 2$$

$$= 2e^{2x-1}$$

$$\frac{dy}{dx} \perp = \frac{-1}{2e^{2x-1}} = -\frac{e^{1-2x}}{2}$$

$$x=1$$

$$\frac{dy}{dx} \perp = -\frac{1}{2e}$$

$$y - e = -\frac{1}{2e}(x-1)$$

$$y = -\frac{x}{2e} + \frac{1}{2e} + e$$

$$y = -\frac{xe^{-1}}{2} + \frac{1+2e^2}{2e}$$

or

$$2ey - 2e^2 = -x + 1$$

$$2ey + x = 1 + 2e^2$$

$$x + 2ey = 1 + 2e^2$$

5. Find the points where the normal to

$$y = x^3 - 2x^2 + 1$$

at $x=1$ meets the curve again.

Find equation of normal at $x=1$ and then set equal to y , find intersection points.

$$\frac{dy}{dx} = 3x^2 - 4x$$

$$\frac{dy}{dx} \perp = \frac{-1}{3x^2 - 4x}$$

$$x=1 \quad \frac{dy}{dx} \perp = \frac{-1}{3-4} = 1$$

$$x=1, \quad y = 1 - 2 + 1 = 0$$

$$y - 0 = 1(x-1)$$

$$y = x-1 \quad (\text{normal})$$

$$\text{set } x^3 - 2x^2 + 1 = x-1$$

$$\Rightarrow x^3 - 2x^2 - x + 2 = 0$$

We already have one solution

$$x=1 \Rightarrow x-1 \text{ is a}$$

factor

$$(x-1)(x^2 - x - 2) = 0$$

From deduce

$$\begin{array}{r} x^2 - x - 2 \\ \hline x-1) x^3 - 2x^2 - x + 2 \\ x^2 - x^2 \\ \hline -x^2 - x \\ -x^2 + x \\ \hline -2x + 2 \\ -2x + 2 \\ \hline 0 \end{array}$$

$$(x-1)(x^2 - x - 2)^0$$

$$\Rightarrow (x-1)(x-2)(x+1) = 0$$

Solution $x=1, x=2, x=-1$

so new points

$$x=2 \quad y = (2)^3 - 2(2)^2 + 1$$


$$= 1 \quad (2, 1)$$

$$x = -1$$


$$y = (-1)^3 - 2(-1)^2 + 1$$

$$= -2 \quad (-1, -2)$$

points are $(2, 1), (-1, -2)$



6. Find the equation of the normal to $f(x) = \cos(x)$ which passes through the origin.



let $x = a$
 $\Rightarrow (a, \cos(a))$ on the curve

$$\frac{dy}{dx} = -\sin(x)$$

$$\frac{dy}{dx} \perp = \frac{1}{\sin(x)}$$

$$\frac{dy}{dx} \perp = \frac{1}{\sin(a)}$$

$$\frac{\cos(a) - 0}{a - 0} = \frac{1}{\sin(a)}$$

$$\begin{aligned}\sin(a)\cos(a) &= a \\ 2\sin(a)\cos(a) &= 2a\end{aligned}$$

$$\sin(2a) = 2a$$

Solve $\sin(x) = x$?

We know $\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$

$$\begin{aligned}\Rightarrow \sin(x) &= x \\ \text{when } x &= 0 \\ a &= 0\end{aligned}$$


Point $(0, 1)$

Normal line is vertical

$$\Rightarrow x = 0$$

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$

Proof:



$$\sin(x) < x < \tan(x)$$


$$\Rightarrow \frac{1}{\sin(x)} > \frac{1}{x} > \frac{\cos(x)}{\sin(x)}$$

$$1 > \frac{\sin(x)}{x} > \cos(x)$$

$$\lim_{x \rightarrow 0} \Rightarrow 1 > \frac{\sin(x)}{x} > 1$$

$$\Rightarrow \lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$

7. Find the equation to $y = \sqrt{x}$
from the external point $(4, 0)$



$$y = \sqrt{x}, \frac{dy}{dx} = \frac{1}{2\sqrt{x}}, \frac{dy}{dx} \perp = -2\sqrt{x}$$

let the co ordinate on the
curve be (a, \sqrt{a})

$$\text{so } \frac{dy}{dx} \perp = -2\sqrt{a}$$

$$\text{and } \frac{\sqrt{a} - 0}{a - 4} = -2\sqrt{a}$$

$$\sqrt{a} = -2\sqrt{a}(a-4)$$

$$\sqrt{a} = -2a\sqrt{a} + 8\sqrt{a}$$

$$2a\sqrt{a} - 7\sqrt{a} = 0$$

$$\sqrt{a}(2a-7) = 0$$

$$\sqrt{a} = 0 \quad 2a-7 = 0$$

$$\xrightarrow{a=0} \quad a = \frac{7}{2}$$

reject, because
derivative is not
defined at the
end point of a
curve

$$a = \frac{7}{2}$$

$$\text{Point } \left(\frac{7}{2}, \sqrt{\frac{7}{2}}\right)$$

$$\frac{dy}{dx} \perp = -2\sqrt{\frac{7}{2}}$$

$$y - \sqrt{\frac{7}{2}} = -2\sqrt{\frac{7}{2}}(x - \frac{7}{2})$$

$$y = -2\sqrt{\frac{7}{2}}x + 7\sqrt{\frac{7}{2}} + \sqrt{\frac{7}{2}}$$

$$y = -2\sqrt{\frac{7}{2}}x + 8\sqrt{\frac{7}{2}}$$

$$y = -\sqrt{\frac{4 \times 7}{2}}x + \sqrt{\frac{7 \times 64}{2}}$$

$$= -\sqrt{14}x + \sqrt{7 \times 32}$$

$$= -\sqrt{14}x + \sqrt{16 \times 7 \times 2}$$

$$= -\sqrt{14}x + 4\sqrt{14}$$

1.4 Increasing and Decreasing

When analysing functions, we are often interested in the intervals across which the function is increasing or decreasing.

- If the gradient is entirely positive on an interval, the function is increasing on that interval.
- If the gradient is entirely negative on an interval, the function is decreasing on that interval.


If a function changes from increasing to decreasing, it must pass through a stationary point (or cross a region where the function is not differentiable)

Hence we need to divide up the domain into intervals bounded by stationary points and points where the function is not differentiable.


Finding a single gradient on each interval will tell us whether the function is increasing or decreasing.

We can communicate all of this using a **sign diagram**.

1.4.1 Basic Sign Diagram example



Sign diagram: $f'(x)$ either calculated or observed from a graph



1.4.2 Sign Diagram example with discontinuities


Below is the graph of

$$y = \frac{x-3}{x^2-4}$$


Because division by 0 is undefined, the graph has vertical asymptotes when $x^2 - 4 = 0$
(i.e when $x = 2$ and $x = -2$)

The derivative is $\frac{dy}{dx} = \frac{-x^2 + 6x - 4}{(x^2 - 4)^2}$ and using a graphics calculator to solve $-x^2 + 6x - 4 = 0$,

we have stationary points at $x \approx 0.764$ and $x \approx 5.236$



Sign diagram: $f'(x)$ either calculated or observed from a graph



1.5 Stationary Points

1.5.1 Turning points (minima, maxima)

1.5.2 Stationary points of inflection

1.6 Shape

1.7 Inflection Points

1.8 Understanding functions and their derivatives

Chapter: 2 Applications of differentiation

$$\text{The equation is: } 9a - 4 = 14 + 3a$$

$$\text{Subtract } 3a: \quad 6a - 4 = 14$$

$$\text{Subtract } 4: \quad 6a = 18$$

$$\text{Divide by } 6: \quad a = 3$$

$$A\widehat{B}C$$

$$A\widehat{B}CC$$

$$A\dot{B}C$$

$$N\tilde{a}$$

$$X \sim \mathcal{N}(\mu, \sigma^2)$$

