

SYNTHETIC DEMO DATA — NOT A REAL PATIENT

COMPREHENSIVE PATIENT PROFILE & WEIGHT MANAGEMENT VISIT NOTE

Lakeside Endocrinology & Medical Weight Management

Date of service: 2026-06-22

Provider: Susan Whitfield, MD — Endocrinology / Obesity Medicine

Provider NPI: 1457823690

Clinic phone: (555) 412-7700 Clinic fax: (555) 412-7701

PATIENT

Name: Robert J. Hayes

DOB: 07/22/1968 Sex: Male

Phone: (555) 318-4471

Address: 1190 Maple Crest Drive, Aurora, IL 60504

INSURANCE

Payer: UnitedHealthcare — Choice Plus PPO

Member ID: 948267715 Group: 0708144

Rx BIN: 610279 Rx PCN: 9999 Rx Group: UHCRX01

VITALS / ANTHROPOMETRICS

Height: 70 in (5'10") Weight: 262 lb BMI: 37.6 kg/m²

Blood pressure: 138/86 mmHg

Most recent HbA1c: 7.8% (2026-06-10) Fasting glucose: 156 mg/dL

ACTIVE PROBLEM LIST

1. Type 2 diabetes mellitus, without complications (ICD-10 E11.9)
2. Morbid (severe) obesity due to excess calories (ICD-10 E66.01)
3. Body mass index 37.0-37.9, adult (ICD-10 Z68.37)
4. Essential hypertension (ICD-10 I10)

WEIGHT MANAGEMENT PROGRAM ENROLLMENT

Patient is actively enrolled in the Lakeside Medically Supervised Weight Management Program (enrolled 2025-11-03). He attends monthly visits and is working with a registered dietitian (Karen Mosley, RD) on a reduced-calorie meal plan (~1,500 kcal/day) and a structured activity plan of 150 min/week of moderate exercise. Documented program participation > 6 months.

PRIOR WEIGHT-LOSS THERAPIES TRIED (STEP THERAPY)

- Intensive lifestyle/diet + exercise program x 6 months — < 4% total body weight loss; plateaued.
- Metformin 1,000 mg BID — ongoing for glycemic control; no meaningful weight reduction.
- Saxenda (liraglutide) 3.0 mg subcutaneous daily x 4 months — inadequate weight response (approx. 3% loss) with nausea limiting adherence; discontinued.
- Ozempic (semaglutide) 1.0 mg subcutaneous weekly x 5 months — modest glycemic benefit but insufficient weight reduction at maximally tolerated dose; patient remains in obese range with persistent comorbidities.

ASSESSMENT

57-year-old male with type 2 diabetes and severe obesity (BMI 37.6) and weight-related comorbidities (T2DM, hypertension), enrolled in a supervised weight management program with documented dietitian involvement. Despite lifestyle modification and prior trials of liraglutide and semaglutide (Ozempic) at the doses noted above, he has not achieved a clinically meaningful, sustained weight reduction.

PLAN / REQUESTED MEDICATION

Requesting prior authorization for Wegovy (semaglutide) 0.25 mg/0.5 mL subcutaneous injection pen. Sig: inject 0.25 mg subcutaneously once weekly x 4 weeks (initiation), then titrate per label toward 2.4 mg weekly maintenance as tolerated.

Dispense: 1 carton (4 single-dose pens) = 28-day supply.

Refills: 5 (with planned dose titration).

Medical necessity: BMI > 30 (37.6) with multiple weight-related comorbidities including type 2 diabetes; documented participation in a comprehensive weight management / lifestyle program with dietitian oversight; and inadequate response to prior anti-obesity / GLP-1 therapy (liraglutide, semaglutide). Wegovy is indicated for chronic weight management in this setting.