

Appendix 1

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
Allophrynidae			
<i>Allophryne ruthveni</i>	20.6	4645	(Lynch & Freeman, 1966; Lescure & Marty, 2000)
Alsodidae			
<i>Alsodes nodosus</i> *	75.85	1140	(Palma, 2013; Penna <i>et al.</i> , 1983)
Alytidae			
<i>Alytes cisternasii</i> *	41	1491	(Márquez, 1995)
Arthroleptidae			
<i>Cardioglossa occidentalis</i> *	27.3	4100	(Blackburn <i>et al.</i> , 2008)
<i>Leptopelis argenteus</i>	45	1750	(Channing, 2001)
<i>Leptopelis bocagii</i>	52	600	(Channing & Howell, 2006)
<i>Leptopelis concolor</i>	34	1750	(Schjötz, 1999)
<i>Leptopelis kivuensis</i>	37	1480	(Roelke <i>et al.</i> , 2011)
Batrachylidae			
<i>Batrachyla antartandica</i>	36.7	1951	(Penna & Solis, 1998; Penna & Meier, 2011)
<i>Hylorina sylvatica</i>	56	1252	(Barrio, 1967; Penna & Veloso, 1990)
Bufonidae			
<i>Ansonia hanitschi</i> *	32	5700	(Malkmus <i>et al.</i> , 2002)
<i>Ansonia leptopus</i> *	40	3933	(Wood Jr. <i>et al.</i> , 2008; Matsui, 1982a)
<i>Ansonia longidigita</i> *	50	3500	(Malkmus <i>et al.</i> , 2002)
<i>Ansonia platysoma</i> *	25	8000	(Malkmus <i>et al.</i> , 2002)
<i>Atelopus chiriquiensis</i> *	34	2025	(Savage, 2002; Lötters <i>et al.</i> , 1999)
<i>Atelopus flavescens</i> *	30	2725	(Lescure & Marty, 2000)
<i>Atelopus franciscus</i> *	21	3320	(Lescure & Marty, 2000)
<i>Atelopus peruensis</i> *	38.5	1950	(Coloma <i>et al.</i> , 2000; Lötters <i>et al.</i> , 1999)
<i>Atelopus pulcher</i> *	29.3	2429	(Lötters <i>et al.</i> , 2002)
<i>Atelopus spumarius</i> *	29	3375	(Cocroft <i>et al.</i> , 1990; Lima <i>et al.</i> , 2006)
<i>Atelopus varius</i> *	41	2350	(Savage, 1972; Cocroft <i>et al.</i> , 1990)
<i>Atelopus zeteki</i> *	40	1820	(Savage, 2002; Cocroft <i>et al.</i> , 1990)
<i>Bufo arenarum</i>	112	1100	(Straneck <i>et al.</i> , 1993)
<i>Bufo asper</i> *	100	1031	(Inger & Bacon Jr, 1968; Preininger <i>et al.</i> , 2007)
<i>Bufo beebei</i>	51	3150	(Gallardo, 1965; Tárano, 2010)
<i>Bufo brauni</i> *	65	1120	(Channing & Howell, 2006)
<i>Bufo calamita</i>	71	1477	(Arak, 1988)
<i>Bufo castaneoticus</i>	37.7	1650	(Köhler & Lötters, 1999b)
<i>Bufo coccifer</i>	62	2512	(Mendelson III <i>et al.</i> , 2005)
<i>Bufo cognatus</i>	103	2307	(Wright & Wright, 1949; Cocroft & Ryan, 1995)
<i>Bufo empusus</i>	76	1549	(Schawrtz, 1972; Alonso & Rodríguez, 2003)
<i>Bufo granulosus</i>	70	2906	(Lima <i>et al.</i> , 2006; São-Pedro <i>et al.</i> , 2011)
<i>Bufo gundlachi</i>	34	2669	(Schawrtz, 1972; Alonso & Rodríguez, 2003)
<i>Bufo gutturalis</i>	90	1150	(Harper <i>et al.</i> , 2010; Channing, 2001)
<i>Bufo houstonensis</i>	77	2151	(Dodd Jr., 2013)
<i>Bufo ibarraí</i>	82	1700	(Mendelson III <i>et al.</i> , 2005; Porter, 1966)

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
<i>Bufo juxtasper</i> *	138	718	(Inger, 1964; Matsui, 1982b)
<i>Bufo longinasus</i> *	29	2329	(Schawrtz, 1972; Alonso & Rodríguez, 2003)
<i>Bufo luetkenii</i>	96	1700	(Savage, 2002; Porter, 1966)
<i>Bufo marmoreus</i> *	83	1825	(Suazo-Ortuño <i>et al.</i> , 2007; Porter, 1966)
<i>Bufo melanochlorus</i> *	65	1700	(Savage, 2002)
<i>Bufo nebulifer</i>	98	1475	(Blair, 1956; Dodd Jr., 2013)
<i>Bufo ocellatus</i>	53	1378	(Caldwell & Shepard, 2007)
<i>Bufo pantherinus</i>	91	2500	(Channing, 2001)
<i>Bufo pardalis</i>	110	650	(Cherry & Francillon-Vieillot, 1992; Cherry & Grant, 1994)
<i>Bufo quercicus</i>	32	4750	(Dodd Jr., 2013)
<i>Bufo raddei</i>	67	1513	(Kuzmin & Ischenko, 1997; Stöck <i>et al.</i> , 2000)
<i>Bufo retiformis</i>	47	3113	(Bogert, 1962; Sullivan <i>et al.</i> , 2000)
<i>Bufo schneideri</i>	180	700	(Norman, 1994; Köhler <i>et al.</i> , 1997)
<i>Bufo siculus</i>	86.6	1600	(Lo Valvo & Giacalone, 2013; Stöck <i>et al.</i> , 2008)
<i>Bufo speciosus</i>	78	2621	(Wright & Wright, 1949; Cocroft & Ryan, 1995)
<i>Bufo steindachneri</i>	48.3	2142	(Largen <i>et al.</i> , 1978)
<i>Bufo taladai</i> *	138	768	(Schawrtz, 1959; Alonso & Rodríguez, 2003)
<i>Bufo terrestris</i>	82	2000	(Wright & Wright, 1949; Leary, 2001)
<i>Bufo valliceps</i>	77.8	1480	(Sullivan & Wagner Jr., 1988; Wagner Jr. & Sullivan, 1995)
<i>Dendrophryniscus minutus</i>	17	3900	(Lima <i>et al.</i> , 2006; Lescure & Marty, 2000)
<i>Pedostibes tuberculosus</i> *	38	3782	(Gururaja & Ramachandra, 2006)
<i>Pelophryne misera</i>	21	4000	(Malkmus <i>et al.</i> , 2002)
<i>Schismaderma carens</i>	88	450	(Channing & Howell, 2006)
Calyptocephalellidae			
<i>Calyptocephallela gayi</i>	120	866	(Penna & Veloso, 1990)
Centrolenidae			
<i>Centrolene buckleyi</i> *	34.7	2500	(Guayasamin <i>et al.</i> , 2006; Bernal <i>et al.</i> , 2004)
<i>Centrolene daidaleum</i> *	24.1	6120	(Ruiz-Carranza & Lynch, 1991; Cardozo-Urdaneta & Señaris, 2012)
<i>Centrolene geckoideum</i> *	77	3828	(Lynch <i>et al.</i> , 1983; Grant <i>et al.</i> , 1998)
<i>Centrolene hesperium</i> *	27.3	3630	(Cadle & McDiarmid, 1990)
<i>Centrolene peristictum</i> *	20.6	6878	(Lynch & Duellman, 1973; Maldonado, 2012)
<i>Centrolene savagei</i> *	24.1	6214	(Díaz-Gutiérrez <i>et al.</i> , 2013)
<i>Centrolene venezuelense</i> *	33.8	4050	(Guayasamin <i>et al.</i> , 2009; Señaris & Ayarzagüena, 2005)
<i>Cochranella euknemos</i> *	25	4200	(Savage, 2002)
<i>Cochranella granulosa</i> *	29	4250	(Savage, 2002)
<i>Cochranella mache</i> *	24	5410	(Guayasamin & Bonaccorso, 2004; Ortega-Andrade <i>et al.</i> , 2013)
<i>Espadarana prosoblepon</i> *	28	5758	(Savage, 2002; Jacobson, 1985)
<i>Hyalinobatrachium carlesvilai</i> *	23.6	4838	(Castroviejo-Fisher <i>et al.</i> , 2009)
<i>Hyalinobatrachium colymbiphyllum</i> *	27	4500	(Savage, 2002; Starrett & Savage, 1973)
<i>Hyalinobatrachium crurifasciatum</i> *	24	4300	(Myers & Donnelly, 1997)
<i>Hyalinobatrachium eccentricum</i> *	23	4160	(Myers & Donnelly, 2001)
<i>Hyalinobatrachium fleischmanni</i> *	25.5	5395	(Savage, 2002; Kime <i>et al.</i> , 2000)

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
<i>Hyalinobatrachium fragile</i> *	24.5	3846	(Guayasamin <i>et al.</i> , 2009; Wen <i>et al.</i> , 2012)
<i>Hyalinobatrachium iaspidiense</i> *	20.1	4940	(Lescure & Marty, 2000)
<i>Hyalinobatrachium igniocus</i> *	23	4920	(Noonan & Bonett, 2003; Barrio-Amorós & Castroviejo-Fisher, 2008)
<i>Hyalinobatrachium mondolfii</i> *	22.5	5106	(Señaris & Ayarzagüena, 2001)
<i>Hyalinobatrachium orientale</i> *	24.5	4819	(Cannatella & Lamar, 1986; Castroviejo-Fisher <i>et al.</i> , 2008)
<i>Hyalinobatrachium orocostale</i> *	25.5	3435	(Guayasamin <i>et al.</i> , 2009; Castroviejo-Fisher <i>et al.</i> , 2008)
<i>Hyalinobatrachium pallidum</i> *	22.4	3259	(Rojas-Runjaic <i>et al.</i> , 2012; Cardozo-Urdaneta & Señaris, 2012)
<i>Hyalinobatrachium pellucidum</i> *	21.4	5039	(Wen <i>et al.</i> , 2012; Castroviejo-Fisher <i>et al.</i> , 2009)
<i>Hyalinobatrachium taylori</i> *	21.6	4495	(Lescure & Marty, 2000)
<i>Hyalinobatrachium valerioi</i> *	24	7250	(Savage, 2002; Starrett & Savage, 1973)
<i>Nymphargus griffithsi</i> *	24.2	4107	(Hutter & Guayasamin, 2012)
<i>Nymphargus pluvialis</i> *	26.5	3941	(Catenazzi <i>et al.</i> , 2009)
<i>Rulyrana spiculata</i> *	27.7	4639	(Catenazzi <i>et al.</i> , 2009)
<i>Teratohyla midas</i> *	26.8	4680	(Guayasamin <i>et al.</i> , 2009; Zimmerman, 1983)
<i>Teratohyla pulverata</i> *	29	5950	(Savage, 2002)
<i>Teratohyla spinosa</i> *	20	7000	(Savage, 2002)
<i>Vitreorana antisthenesi</i> *	27	5432	(Guayasamin <i>et al.</i> , 2009; Wen <i>et al.</i> , 2012)
<i>Vitreorana eurygnatha</i> *	22.5	4850	(Heyer <i>et al.</i> , 1990)
<i>Vitreorana gorzulae</i> *	22.5	4787	(Kok & Castroviejo-Fisher, 2008; Castroviejo-Fisher <i>et al.</i> , 2009)
<i>Vitreorana helenae</i> *	20.4	4495	(Guayasamin <i>et al.</i> , 2009; Señaris & Ayarzagüena, 2005)
Ceratophryidae			
<i>Ceratophrys cornuta</i>	72	2176	(Duellman & Lizana, 1994; Marquez <i>et al.</i> , 1995)
<i>Ceratophrys cranwelli</i>	130	1715	(Valetti <i>et al.</i> , 2013)
<i>Ceratophrys ornata</i>	125.8	2000	(Basso, 1990; Salas <i>et al.</i> , 1998)
<i>Chacophrys pierottii</i>	53.1	4357	(Lascano, 2011)
Cycloramphidae			
<i>Cycloramphus boraceiensis</i> *	51.3	1950	(Heyer <i>et al.</i> , 1990)
<i>Thoropa miliaris</i> *	71.5	600	(Heyer <i>et al.</i> , 1990)
Dicroglossidae			
<i>Euphlyctis cyanophlyctis</i>	45	1651	(Roy & Elepfandt, 1993)
<i>Euphlyctis ehrenbergii</i>	66	937	(Khan, 1997; Roy & Elepfandt, 1993)
<i>Fejervarya limnocharis</i>	43	2119	(Prakash, 1988; Márquez & Eekhout, 2006)
<i>Fejervarya sakishimensis</i>	55.5	1030	(Matsui <i>et al.</i> , 2007)
<i>Fejervarya syhadrensis</i>	29.3	3350	(Kuramoto <i>et al.</i> , 2007)
<i>Hoplobatrachus occipitalis</i>	104	1250	(Rödel, 2000; van den Elzen & Kreulen, 1979)
<i>Hoplobatrachus tigerinus</i>	120	520	(Roy & Elepfandt, 1993; Glaw & Vences, 2007)
<i>Limnonectes arathooni</i> *	42	3380	(Smith, 1927; Brown & Iskandar, 2000)
<i>Limnonectes blythii</i> *	105.4	428	(Matsui, 1995)
<i>Limnonectes gyldestolpei</i> *	53	1085	(Garcia-Rutledge & Narins, 2001)
<i>Limnonectes microdiscus</i> *	48.6	875	(Inger, 1954; Kurniati <i>et al.</i> , 2010)
<i>Paa spinosa</i> *	101	834	(Yu & Zheng, 2009)
<i>Sphaerothera breviceps</i>	48	1950	(Minton, 1966; Kanamadi <i>et al.</i> , 1994)
Heleophrynidae			

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
<i>Hadromophryne natalensis</i> *	45	1500	(Channing, 2001)
<i>Heleophryne purcelli</i> *	47	2250	(Channing, 2001)
<i>Heleophryne regis</i> *	43	1800	(Channing, 2001)
Hylidae			
<i>Agalychnis annae</i> *	73.9	1165	(Duellman, 1970)
<i>Agalychnis callidryas</i>	59	2006	(Savage, 2002; Duellman & Pyles, 1983)
<i>Agalychnis litodryas</i>	72.2	1664	(Duellman, 1970)
<i>Agalychnis saltator</i>	54	1800	(Leenders, 2001; Duellman & Pyles, 1983)
<i>Agalychnis spurrelli</i>	75.6	893	(Ortega-Andrade, 2008)
<i>Aplastodiscus albofrenatus</i> *	40	2500	(Lutz, 1973; Heyer <i>et al.</i> , 1990)
<i>Aplastodiscus albosignatus</i> *	52	2630	(Lutz, 1973; Abrunhosa <i>et al.</i> , 2005)
<i>Aplastodiscus arildae</i> *	41.6	2847	(Heyer <i>et al.</i> , 1990; Carvalho Jr <i>et al.</i> , 2006)
<i>Aplastodiscus callipygius</i> *	50.7	1040	(Cruz & Peixoto, 1985; Abrunhosa <i>et al.</i> , 2005)
<i>Argenteohyla siemersi</i>	70	1882	(Ceil, 1980; Cajade <i>et al.</i> , 2010)
<i>Bokermannohyla circumdata</i> *	70	1460	(Lutz, 1973; Carvalho <i>et al.</i> , 2012)
<i>Bokermannohyla hylax</i> *	61.5	2000	(Heyer <i>et al.</i> , 1990; Carvalho <i>et al.</i> , 2012)
<i>Cyclorana brevipes</i>	45	1930	(Tyler & Martin, 1977)
<i>Cyclorana cryptotis</i>	41	1060	(Tyler & Martin, 1977)
<i>Cyclorana cultripes</i>	52	1879	(Tyler & Martin, 1977)
<i>Cyclorana manya</i>	29.9	2500	(Van Beurden & McDonald, 1980)
<i>Dendropsophus allenorum</i>	21.4	3430	(Duellman, 2005)
<i>Dendropsophus anceps</i>	40	3244	(Lutz, 1973; Gomes & Martins, 2006)
<i>Dendropsophus berthaltutae</i>	21	4041	(Lutz, 1973; Forti <i>et al.</i> , 2012)
<i>Dendropsophus bifurcus</i>	28	2958	(Duellman, 1978; Márquez <i>et al.</i> , 1993)
<i>Dendropsophus bipunctatus</i>	25	5426	(Lutz, 1973; Wogel & Pombal Jr, 2007)
<i>Dendropsophus brevifrons</i>	22	4254	(Duellman, 1978; Duellman & Pyles, 1983)
<i>Dendropsophus carnifex</i>	27.7	2490	(Duellman, 1969; De la Riva <i>et al.</i> , 1997)
<i>Dendropsophus ebraccatus</i>	27.8	2577	(Duellman, 1970; Duellman & Pyles, 1983)
<i>Dendropsophus elegans</i>	29.6	3500	(Lutz, 1973; Bastos & Haddad, 1995)
<i>Dendropsophus giesleri</i>	25	3300	(Weygoldt & Peixoto, 1987; Heyer, 1980)
<i>Dendropsophus koechlini</i>	24	4500	(Duellman, 2005)
<i>Dendropsophus labialis</i>	43	1750	(Amézquita, 1999; Guarnizo, 2011)
<i>Dendropsophus leali</i>	23	6200	(Duellman, 2005)
<i>Dendropsophus leucophyllatus</i>	36	2482	(Rodríguez & Duellman, 1994; Márquez <i>et al.</i> , 1993)
<i>Dendropsophus marmoratus</i>	44	1514	(Rodríguez & Duellman, 1994; Duellman & Pyles, 1983)
<i>Dendropsophus microcephalus</i>	24.5	6210	(Duellman, 1970; Táran, 2010)
<i>Dendropsophus minusculus</i>	20.6	9186	(Duellman, 1997; Duellman & Pyles, 1983)
<i>Dendropsophus nanus</i>	22	4200	(Lutz, 1973; Nunes <i>et al.</i> , 2007)
<i>Dendropsophus rhodopeplus</i>	24.2	5670	(Duellman, 2005)
<i>Dendropsophus riveroi</i>	20	5314	(Rodríguez & Duellman, 1994; Márquez <i>et al.</i> , 1993)
<i>Dendropsophus robertmertensi</i>	26.4	5388	(Duellman & Fouquette, 1968; Duellman, 1970)
<i>Dendropsophus rubicundulus</i>	28.3	3500	(Napoli & Caramaschi, 1999; Cardoso & Vielliard, 1985)
<i>Dendropsophus sanborni</i>	17	5100	(Lutz, 1973; Martins & Jim, 2003)

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
<i>Dendropsophus sarayacuensis</i>	29	2990	(Rodríguez & Duellman, 1994; Duellman & Pyles, 1983)
<i>Dendropsophus sartori</i>	26	3217	(Duellman & Fouquette, 1968; Duellman, 1970)
<i>Dendropsophus seniculus</i>	37.7	3969	(Heyer <i>et al.</i> , 1990; Hepp <i>et al.</i> , 2012)
<i>Dendropsophus triangulum</i>	28	2298	(Rodríguez & Duellman, 1994; Duellman & Pyles, 1983)
<i>Duellmanohyla rufiocularis</i> *	30	2320	(Duellman, 1970)
<i>Exerodonta melanomma</i> *	29.9	2336	(Duellman, 1970)
<i>Exerodonta sumichrasti</i> *	27.7	2510	(Duellman, 1970)
<i>Hyla andersonii</i>	51	1100	(Conant & Collins, 1998; Gerhardt, 1974)
<i>Hyla arborea</i>	50	2122	(Arnold, 2003; Castellano <i>et al.</i> , 2002)
<i>Hyla chinensis</i>	32	4000	(Fei <i>et al.</i> , 1999; Kuramoto, 1980)
<i>Hyla chrysoscelis</i>	51	2400	(Wright & Wright, 1949; Gerhardt, 2005)
<i>Hyla femoralis</i>	37	2600	(Wright & Wright, 1949; Blair, 1958b)
<i>Hyla gratiosa</i>	68	1840	(Wright & Wright, 1949; Oldham & Gerhardt, 1975)
<i>Hyla intermedia</i>	50	2388	(Arnold, 2003; Castellano <i>et al.</i> , 2002)
<i>Hyla meridionalis</i>	50	2750	(Arnold, 2003; Oliveira <i>et al.</i> , 1991)
<i>Hyla molleri</i>	45	2750	(Schneider, 1974)
<i>Hyla squirella</i>	36	3344	(Wright & Wright, 1949; Taylor <i>et al.</i> , 2007)
<i>Hyla versicolor</i>	51	2200	(Wright & Wright, 1949; Gerhardt, 2005)
<i>Hylomantis granulosa</i> *	45.8	1828	(Pimenta <i>et al.</i> , 2007)
<i>Hylomantis lemur</i>	40.8	2396	(Duellman, 1970)
<i>Hyloscirtus armatus</i> *	68.5	3600	(Duellman, 1997)
<i>Hyloscirtus colymba</i> *	37	3520	(Duellman, 1970; 1972b)
<i>Hyloscirtus lindae</i> *	68.1	1553	(Duellman & Altig, 1978; Coloma <i>et al.</i> , 2012)
<i>Hyloscirtus pacha</i> *	60.8	1854	(Duellman & Hillis, 1990; Coloma <i>et al.</i> , 2012)
<i>Hyloscirtus palmeri</i> *	45	2500	(Duellman, 1970; Ibáñez <i>et al.</i> , 1999)
<i>Hyloscirtus pantostictus</i> *	63	1570	(Duellman & Berger, 1982; Coloma <i>et al.</i> , 2012)
<i>Hyloscirtus phyllonathus</i> *	34	2588	(Duellman, 1972b)
<i>Hyloscirtus tapichalaca</i> *	63.8	1469	(Kizirian <i>et al.</i> , 2003; Coloma <i>et al.</i> , 2012)
<i>Hypsiboas albopunctatus</i>	60	2168	(Lutz, 1973; Kwet <i>et al.</i> , 2002)
<i>Hypsiboas bischoffi</i>	46.1	1750	(Marcelino <i>et al.</i> , 2009; Pombal Jr., 2010)
<i>Hypsiboas cinerascens</i>	44	1390	(Rodríguez & Duellman, 1994; Duellman & Pyles, 1983)
<i>Hypsiboas ericae</i> *	34	2700	(Caramaschi & Cruz, 2000; Garcia & Haddad, 2008)
<i>Hypsiboas fasciatus</i>	40.3	2530	(Duellman, 2005; Lescure & Marty, 2000)
<i>Hypsiboas joaquina</i> *	56.4	1700	(Garcia <i>et al.</i> , 2003)
<i>Hypsiboas microderma</i> *	34	3300	(Rodríguez & Duellman, 1994; Pyburn, 1977)
<i>Hypsiboas ornatissimus</i>	31	2855	(Lutz, 1973; Lescure & Marty, 2000)
<i>Hypsiboas pardalis</i>	69	1133	(Lutz, 1973; Rodrigues, 2009)
<i>Hypsiboas pellucens</i>	61.6	800	(Cochran & Goin, 1970; Duellman, 1971)
<i>Hypsiboas roraima</i> *	44.9	2616	(Barrio-Amorós <i>et al.</i> , 2011)
<i>Hypsiboas rufitelus</i>	49.2	1600	(Duellman, 1970)
<i>Hypsiboas semiguttatus</i> *	46.1	2450	(Garcia <i>et al.</i> , 2007)
<i>Hypsiboas semilineatus</i>	51	1014	(Lisboa <i>et al.</i> , 2011; Lingnau & Bastos, 2003)
<i>Isthmohyla pseudopuma</i>	41.4	956	(Duellman, 1970)

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
<i>Isthmohyla rivularis</i> *	34	2420	(Duellman, 1970)
<i>Isthmohyla tica</i> *	34.1	2228	(Duellman, 1970)
<i>Litoria gracilentia</i>	38	2750	(Peters, 1869; Günther & Richards, 2000)
<i>Litoria impura</i>	42	1240	(Peters & Doria, 1878; Kraus & Allison, 2004)
<i>Litoria inermis</i>	33	3800	(Davies <i>et al.</i> , 1983)
<i>Litoria latopalmata</i>	39	2900	(Davies <i>et al.</i> , 1983)
<i>Litoria leucova</i> *	35.4	4500	(Johnston & Richards, 1994)
<i>Litoria pallida</i>	34	1963	(Davies <i>et al.</i> , 1983)
<i>Litoria raniformis</i>	79.9	900	(Hamer & Organ, 2008; Ford, 1989)
<i>Litoria rheocola</i> *	35.5	2500	(Hoskin & Goosem, 2010)
<i>Litoria spartacus</i> *	37.5	2919	(Richards & Oliver, 2006)
<i>Litoria thesaurensis</i>	65	1774	(Tyler & Davies, 1978; Kraus & Allison, 2004)
<i>Litoria tornieri</i>	33	2725	(Doughty, 2011)
<i>Litoria wollastoni</i> *	42.7	2950	(Menzies & Zweifel, 1974)
<i>Myersiohyla kanaima</i> *	37.8	3600	(Goin & Woodley, 1969)
<i>Nyctimystes cheesmani</i> *	40	1900	(Kraus, 2012)
<i>Osteocephalus taurinus</i>	85	1501	(Trueb & Duellman, 1971; De la Riva <i>et al.</i> , 1995)
<i>Osteopilus septentrionalis</i>	89	2300	(Savage, 2002; Blair, 1958a)
<i>Pachymedusa dacnicolor</i>	82.6	1727	(Duellman, 1970)
<i>Phyllomedusa atelopoides</i>	37.4	1150	(Duellman <i>et al.</i> , 1988)
<i>Phyllomedusa azurea</i>	43.46	2333	(Álvares, 2008; Guimarães <i>et al.</i> , 2001)
<i>Phyllomedusa bahiana</i>	74.5	960	(Pombal Jr. & Haddad, 1992; Silva & Juncá, 2006)
<i>Phyllomedusa bicolor</i>	103	940	(Lima <i>et al.</i> , 2006; Zimmerman, 1983)
<i>Phyllomedusa burmeisteri</i>	75.8	1500	(Pombal Jr. & Haddad, 1992; Abrunhosa & Wogel, 2004)
<i>Phyllomedusa camba</i>	74	860	(Rodrigues <i>et al.</i> , 2011; Duellman, 2005)
<i>Phyllomedusa centralis</i> *	42	1451	(Bokermann, 1965; Brandão <i>et al.</i> , 2009)
<i>Phyllomedusa distincta</i>	66	1250	(Haddad <i>et al.</i> , 1994)
<i>Phyllomedusa duellmani</i>	54.2	950	(Cannatella, 1982)
<i>Phyllomedusa hypochondrialis</i>	43.73	2047	(Álvares, 2008; De la Riva <i>et al.</i> , 1995)
<i>Phyllomedusa megacephala</i> *	43.2	1722	(Caramaschi, 2006; Giarretta <i>et al.</i> , 2007)
<i>Phyllomedusa neildi</i>	63.8	762	(Barrio-Amoros, 2006)
<i>Phyllomedusa nordestina</i>	42.1	2076	(Caramaschi, 2006; Vilaça <i>et al.</i> , 2011)
<i>Phyllomedusa palliata</i>	43.8	1500	(Duellman, 1974; Zimmerman, 1983)
<i>Phyllomedusa sauvagii</i>	89.2	2000	(Rodrigues <i>et al.</i> , 2007; Halloy & Espinosa, 2000)
<i>Phyllomedusa tarsius</i>	90	870	(Lima <i>et al.</i> , 2006; Zimmerman, 1983)
<i>Phyllomedusa tetraploidea</i>	63.8	1700	(Pombal Jr. & Haddad, 1992)
<i>Phyllomedusa tomopterna</i>	54	1790	(Lima <i>et al.</i> , 2006; Zimmerman, 1983)
<i>Phyllomedusa trinitatis</i>	81	800	(Barrio-Amoros, 2006)
<i>Plectrohyla cyclada</i> *	39.5	2072	(Duellman, 1970)
<i>Pseudacris brachyphona</i>	32	2394	(Wright & Wright, 1949; Cocroft & Ryan, 1995)
<i>Pseudacris clarkii</i>	29	3072	(Duellman, 1970; Cocroft & Ryan, 1995)
<i>Pseudacris feriarum</i>	27.3	2977	(Schawrtz, 1957; Cocroft & Ryan, 1995)
<i>Pseudacris fouquettei</i>	29.8	3138	(Lemmon <i>et al.</i> , 2008)

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
<i>Pseudacris illinoensis</i>	43	2168	(Tucker, 1998; Owen & Tucker, 2006)
<i>Pseudacris kalmi</i>	38.1	2972	(Harper, 1955; Cocroft & Ryan, 1995)
<i>Pseudacris nigrita</i>	28	3149	(Wright & Wright, 1949; Cocroft & Ryan, 1995)
<i>Pseudacris ornata</i>	39	2651	(Lannoo, 2005; Cocroft & Ryan, 1995)
<i>Pseudacris triseriata</i>	32	2776	(Wright & Wright, 1949; Cocroft & Ryan, 1995)
<i>Pseudis bolbodactyla</i>	45	1937	(Caramaschi & Cruz, 1998; Vaz-Silva <i>et al.</i> , 2007)
<i>Ptychohyla euthysanota</i> *	37.3	3100	(Duellman, 1970)
<i>Ptychohyla leonhardschultzei</i> *	35.6	2750	(Duellman, 1970)
<i>Ptychohyla spinipollex</i> *	41.2	4300	(Duellman, 1970)
<i>Scinax berthae</i>	22.2	4886	(Faivovich, 2005; Pereyra <i>et al.</i> , 2012)
<i>Scinax boesemani</i>	31.1	1420	(Duellman, 1997; Duellman & Pyles, 1983)
<i>Scinax boulengeri</i>	48.7	1615	(Duellman, 1970; Duellman & Pyles, 1983)
<i>Scinax crospedospilus</i>	33.3	1350	(Heyer <i>et al.</i> , 1990)
<i>Scinax cruentommus</i>	28	3300	(Duellman, 1972a; 1978)
<i>Scinax fuscovarius</i>	47.1	855	(De la Riva, 1993; Fonte, 2010)
<i>Scinax nasicus</i>	37	1064	(Lutz, 1973; Fonte, 2010)
<i>Scinax proboscideus</i>	39.8	2012	(Duellman, 1972b; Lima <i>et al.</i> , 2004)
<i>Scinax rostratus</i>	45.7	3050	(Duellman, 1970; 1972b)
<i>Scinax ruber</i>	41.2	700	(Duellman & Wiens, 1993; Bernal <i>et al.</i> , 2004)
<i>Scinax staufferi</i>	29	3056	(León, 1969; Duellman, 1970)
<i>Scinax sugillatus</i>	42	2023	(Duellman, 1973)
<i>Scinax uruguayus</i>	25.8	4350	(Langone, 1990; Lingnau, 2009)
<i>Scinax x-signatus</i>	42.5	1110	(Heyer <i>et al.</i> , 1990; Tárano, 2010)
<i>Smilisca baudinii</i>	76	350	(Duellman, 1970)
<i>Smilisca fodiens</i>	62.6	2230	(Duellman, 1970)
<i>Smilisca phaeota</i>	65.5	372	(Duellman, 1970)
<i>Smilisca puma</i>	38.1	740	(Duellman, 1970)
<i>Smilisca sordida</i> *	45	2695	(Duellman, 1970)
<i>Sphaenorhynchus lacteus</i>	41.5	2240	(Duellman, 2005)
<i>Sphaenorhynchus orophilus</i>	32	2350	(Heyer <i>et al.</i> , 1990)
<i>Tlalocohyla loquax</i>	44.7	2323	(Duellman, 1970)
<i>Tlalocohyla picta</i>	21.4	2661	(Duellman, 1970)
<i>Trachycephalus coriaceus</i>	63	2090	(Rodríguez & Duellman, 1994; De la Riva <i>et al.</i> , 1995)
<i>Trachycephalus mesophaeus</i>	85	1014	(Lutz, 1973; Prado <i>et al.</i> , 2003)
<i>Trachycephalus nigromaculatus</i>	86	1990	(Cochran, 1955; Abrunhosa <i>et al.</i> , 2001)
<i>Triprion petasatus</i>	60.8	2096	(Duellman, 1970)
Hylodidae			
<i>Crossodactylus caramaschii</i> *	25.8	5000	(Bastos & Pombal, 1995)
<i>Crossodactylus schmidtii</i> *	30	3306	(Caldart <i>et al.</i> , 2011)
<i>Hylodes dactylocinus</i> *	27	2973	(Pavan <i>et al.</i> , 2001)
<i>Hylodes meridionalis</i> *	37.5	4400	(Pombal Jr. <i>et al.</i> , 2002; Lingnau <i>et al.</i> , 2013)
<i>Hylodes perplicatus</i> *	39.8	3400	(Haddad <i>et al.</i> , 2003)
<i>Hylodes phyllodes</i> *	31.4	5000	(Heyer & Cocroft, 1986)

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
<i>Hylodes sazimai</i> *	28.5	4750	(Heyer, 1982; Haddad & Pombal Jr, 1995)
Hyperoliidae			
<i>Afrixalus delicatus</i>	22.2	4525	(Pickersgill, 2005)
<i>Afrixalus dorsalis</i>	28	3587	(Schiotz, 1999; Köhler <i>et al.</i> , 2005)
<i>Afrixalus fornasini</i>	38	2450	(Channing, 2001)
<i>Afrixalus knysnae</i>	25	3800	(Schiotz, 1999; Channing, 2001)
<i>Afrixalus laevis</i> *	23	3000	(Schiotz, 1999)
<i>Afrixalus paradorsalis</i>	34	2864	(Schiotz, 1999; Köhler <i>et al.</i> , 2005)
<i>Afrixalus stuhlmanni</i>	21.1	4650	(Pickersgill, 2005)
<i>Heterixalus andrakata</i>	29	3200	(Glaw & Vences, 1991; 2007)
<i>Heterixalus betsileo</i>	28	3000	(Glaw & Vences, 2007; 1993)
<i>Heterixalus boettgeri</i>	25	2850	(Glaw & Vences, 2007; 1993)
<i>Heterixalus carbonei</i>	26.4	3200	(Vences <i>et al.</i> , 2000)
<i>Heterixalus luteostriatus</i>	28	3500	(Glaw & Vences, 2007; Andreone <i>et al.</i> , 1994)
<i>Heterixalus madagascariensis</i>	35	3375	(Glaw & Vences, 2007; 1993)
<i>Heterixalus tricolor</i>	26	3000	(Glaw & Vences, 2007; 1993)
<i>Heterixalus variabilis</i>	31	3500	(Glaw & Vences, 2007; 1993)
<i>Hyperolius alticola</i> *	33	2250	(Schiotz, 1999)
<i>Hyperolius argus</i>	34	2000	(Schiotz, 1999)
<i>Hyperolius baumanni</i>	30	3000	(Schiotz, 1999)
<i>Hyperolius cinnamomeoventris</i>	28	3246	(Schiotz, 1999; Sinsch <i>et al.</i> , 2012)
<i>Hyperolius cystocandicans</i>	28	2813	(Schiotz, 1999; Köhler <i>et al.</i> , 2005)
<i>Hyperolius guttulatus</i> *	35	2000	(Schiotz, 1999)
<i>Hyperolius kivuensis</i>	33	2649	(Schiotz, 1999; Sinsch <i>et al.</i> , 2012)
<i>Hyperolius montanus</i>	28	2750	(Schiotz, 1999)
<i>Hyperolius phantasticus</i>	37	2529	(Schiotz, 1999; Köhler <i>et al.</i> , 2005)
<i>Hyperolius picturatus</i> *	31	3850	(Schiotz, 1999)
<i>Hyperolius puncticulatus</i>	33	2000	(Channing & Howell, 2006)
<i>Hyperolius pusillus</i>	21	4950	(Channing, 2001)
<i>Hyperolius torrentis</i> *	38	3500	(Schiotz, 1999)
<i>Hyperolius viridiflavus</i> *	30	4250	(Schiotz, 1999; van den Elzen & Kreulen, 1979)
<i>Kassina maculata</i>	65	1700	(Schiotz, 1999; Channing, 1976)
<i>Kassina senegalensis</i>	49	1000	(Channing & Howell, 2006)
<i>Phlyctimantis verrucosus</i>	52	1600	(Channing & Howell, 2006; Schiotz, 1999)
<i>Semnodactylus wealii</i>	44	2000	(Schiotz, 1999)
Leptodactylidae			
<i>Edalorhina perezi</i>	30.2	2700	(Duellman, 2005)
<i>Engystomops coloradorum</i>	24.1	1060	(Cannatella & Duellman, 1984; Ryan & Drewes, 1990)
<i>Engystomops guayaco</i>	19.4	2114	(Ron <i>et al.</i> , 2005)
<i>Engystomops pustulatus</i>	29.9	947	(Ron <i>et al.</i> , 2004; Ryan & Drewes, 1990)
<i>Engystomops pustulosus</i>	34	546	(Ryan, 1983; Kime <i>et al.</i> , 2000)
<i>Engystomops randi</i>	18.56	1116	(Ron <i>et al.</i> , 2004)
<i>Eupemphix nattereri</i>	42.7	710	(Silva <i>et al.</i> , 2008)

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
<i>Leptodactylus bufonius</i>	53.6	1500	(Heyer, 1978)
<i>Leptodactylus chaquensis</i>	99.7	365	(Heyer & Giaretta, 2009)
<i>Leptodactylus didymus</i>	46.7	780	(Heyer <i>et al.</i> , 1996; Köhler & Lötters, 1999b)
<i>Leptodactylus diedrus</i>	40.4	820	(Heyer, 1994; 1998)
<i>Leptodactylus elenae</i>	45.2	1160	(Heyer, 1978; Heyer <i>et al.</i> , 1996)
<i>Leptodactylus fuscus</i>	52.1	560	(Bernal <i>et al.</i> , 2004)
<i>Leptodactylus gracilis</i>	47.8	1970	(Heyer, 1978; Köhler & Lötters, 1999a)
<i>Leptodactylus griseigularis</i>	51	2770	(Heyer, 1994; Heyer & de Carvalho, 2000)
<i>Leptodactylus knudseni</i>	159	500	(Heyer, 2005)
<i>Leptodactylus labyrinthicus</i>	170	391	(Zina & Haddad, 2005)
<i>Leptodactylus leptodactyloides</i>	48	1200	(Heyer, 1994; Duellman, 2005)
<i>Leptodactylus longirostris</i>	40	2315	(Heyer, 1978; Lescure & Marty, 2000)
<i>Leptodactylus melanonotus</i>	46	1840	(Heyer, 1970; Heyer & de Carvalho, 2000)
<i>Leptodactylus mystaceus</i>	52.2	1056	(Heyer <i>et al.</i> , 1996)
<i>Leptodactylus mystacinus</i>	65	2275	(Heyer <i>et al.</i> , 2003)
<i>Leptodactylus notoaktites</i>	50.8	1230	(Heyer, 1978; Heyer <i>et al.</i> , 1996)
<i>Leptodactylus ocellatus</i>	120	175	(Heyer <i>et al.</i> , 1990; Salas <i>et al.</i> , 1998)
<i>Leptodactylus pentadactylus</i>	177	545	(Savage, 2002; Kime <i>et al.</i> , 2000)
<i>Leptodactylus podicipinus</i>	44.7	1970	(Silva <i>et al.</i> , 2008)
<i>Leptodactylus rhodonotus</i>	79	2000	(Heyer, 1979; Duellman, 2005)
<i>Leptodactylus spixi</i>	43	1634	(Heyer, 1983; Bilate <i>et al.</i> , 2006)
<i>Leptodactylus vastus</i>	180.3	430	(Heyer, 2005)
<i>Lithodytes lineatus</i>	47	2600	(Duellman, 2005)
<i>Paratelmatobius cardosoi</i>	17.9	2550	(Pombal & Haddad, 1999)
<i>Physalaemus albonotatus</i>	32.4	2399	(Rodrigues <i>et al.</i> , 2004; Marquez <i>et al.</i> , 1995)
<i>Physalaemus barrioi</i>	29.45	2265	(Provete <i>et al.</i> , 2012)
<i>Physalaemus biligonigerus</i>	45	568	(Lynch, 1970; Marquez <i>et al.</i> , 1995)
<i>Physalaemus cuvieri</i>	25.9	690	(Heyer <i>et al.</i> , 1990)
<i>Physalaemus ephippifer</i>	28	840	(Martins, 1998; Kaefer <i>et al.</i> , 2011)
<i>Physalaemus signifer</i>	25.9	1210	(Wogel <i>et al.</i> , 2002)
<i>Pleurodema bibroni</i>	38	1960	(Duellman & Veloso, 1977; Kolenc <i>et al.</i> , 2009)
<i>Pleurodema brachyops</i>	49	732	(Duellman & Veloso, 1977; Kime <i>et al.</i> , 2000)
Limnodynastidae			
<i>Neobatrachus pelobatoides</i>	39.6	816	(Roberts <i>et al.</i> , 1991)
<i>Neobatrachus pictus</i>	55	1300	(Roberts, 1997b)
<i>Neobatrachus sudelli</i>	40	1509	(Roberts, 1997a)
Mantellidae			
<i>Aglyptodactylus laticeps</i>	45.4	2350	(Glaw <i>et al.</i> , 1998)
<i>Blommersia blommersae</i>	21.3	6000	(Glaw & Vences, 2002)
<i>Blommersia domerguei</i>	17	4350	(Glaw & Vences, 2007; Vences <i>et al.</i> , 2002)
<i>Blommersia kely</i>	15.9	4700	(Vences <i>et al.</i> , 2002)
<i>Blommersia sarotra</i>	16.8	4592	(Vences <i>et al.</i> , 2002)
<i>Boophis boehmei</i> *	30	2500	(Glaw & Vences, 2007)

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
<i>Boophis goudotii</i>	70	750	(Glaw & Vences, 2007)
<i>Boophis luteus</i> *	40	3250	(Glaw & Vences, 2007; 1992)
<i>Boophis marojezensis</i> *	27	5583	(Glaw <i>et al.</i> , 2001)
<i>Boophis rappiodes</i> *	25.1	3000	(Vences & Glaw, 2002)
<i>Boophis sibilans</i> *	32.5	3300	(Glaw & Thiesmeier, 1993)
<i>Boophis vittatus</i> *	25.5	7250	(Glaw <i>et al.</i> , 2001)
<i>Boophis xerophilus</i>	38.7	2700	(Glaw & Vences, 1997)
<i>Guibemantis tornieri</i>	49	1850	(Glaw & Vences, 2007; Glaw <i>et al.</i> , 2000)
<i>Mantella viridis</i> *	25	4350	(Glaw & Vences, 2007)
<i>Mantidactylus argenteus</i> *	27	4250	(Glaw & Vences, 2007; Vejarano <i>et al.</i> , 2006)
<i>Mantidactylus charlotteae</i> *	26.2	2900	(Vences & Glaw, 2004)
<i>Mantidactylus opiparis</i> *	26.1	3150	(Vences & Glaw, 2004)
Megophyiidae			
<i>Leptobrachium gunungense</i> *	65	1200	(Malkmus, 2006; Malkmus <i>et al.</i> , 2002)
<i>Leptobrachium montanum</i> *	62.7	800	(Malkmus <i>et al.</i> , 2002)
<i>Leptolalax arayai</i> *	29.6	5588	(Matsui, 1997)
<i>Leptolalax oshanensis</i> *	30.7	4512	(Ohler <i>et al.</i> , 2011; Jiang <i>et al.</i> , 2002)
<i>Leptolalax pictus</i> *	36	7000	(Malkmus <i>et al.</i> , 2002)
<i>Megophrys nasuta</i> *	105	2250	(Malkmus <i>et al.</i> , 2002)
<i>Oreolalax omeimontis</i> *	58.4	1071	(Nguyen <i>et al.</i> , 2013; Jiang <i>et al.</i> , 2002)
<i>Xenophrys minor</i> *	32	3456	(Stejneger, 1926; Jiang <i>et al.</i> , 2002)
Microhylidae			
<i>Calluella guttulata</i>	49	1950	(Parker, 1934; Heyer, 1971)
<i>Chiasmocleis hudsoni</i>	17	3795	(Parker, 1940; Rodrigues <i>et al.</i> , 2008)
<i>Chiasmocleis shudikarensis</i>	24.5	3565	(Peloso & Sturaro, 2008; Lescure & Marty, 2000)
<i>Dermatonotus muelleri</i>	62.4	1758	(Giaretta <i>et al.</i> , 2013)
<i>Elachistocleis ovalis</i>	36	3838	(Thomé & Brasileiro, 2007; Lavilla <i>et al.</i> , 2003)
<i>Gastrophryne elegans</i>	25.8	3100	(Nelson, 1963; Nelson, 1973)
<i>Gastrophryne olivacea</i>	30.8	4420	(Nelson, 1973; Loftus-Hills & Littlejohn, 1992)
<i>Glyphoglossus molossus</i>	77	700	(Parker, 1934; Heyer, 1971)
<i>Hamptophryne boliviana</i>	35	2208	(Hödl, 1990; De la Riva <i>et al.</i> , 1996)
<i>Hypopachus variolosus</i>	48	2750	(Parker, 1934; Lee, 2000)
<i>Kaloula pulchra</i>	70	2500	(Malkmus <i>et al.</i> , 2002)
<i>Microhyla butleri</i>	26	2850	(Parker, 1934; Heyer, 1971)
<i>Microhyla pulchra</i>	35	1750	(Parker, 1934; Heyer, 1971)
<i>Micryletta inornata</i>	30	5450	(Parker, 1934; Heyer, 1971)
<i>Phrynomantis bifasciatus</i>	53	1800	(Channing & Howell, 2006)
<i>Phrynomantis microps</i>	47.3	1260	(Grafe, 1999; Rödel, 2000)
<i>Scaphiophryne boribory</i>	60	700	(Glaw & Vences, 2007; Vences <i>et al.</i> , 2003)
<i>Scaphiophryne gottlebei</i>	30	950	(Glaw & Vences, 2007; Andreone <i>et al.</i> , 2005)
<i>Scaphiophryne madagascariensis</i>	49	1000	(Vences <i>et al.</i> , 2003; Vences <i>et al.</i> , 2002)
<i>Scaphiophryne menabensis</i>	42	850	(Glos <i>et al.</i> , 2005)
<i>Scaphiophryne spinosa</i>	48	800	(Glaw & Vences, 2007; Vences <i>et al.</i> , 2003)

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
Myobatrachidae			
<i>Crinia riparia</i> *	25.2	2735	(Littlejohn & Martin, 1965; Odendaal <i>et al.</i> , 1983)
<i>Geocrinia victoriana</i>	27	2676	(Littlejohn & Harrison, 1985)
<i>Pseudophryne bibronii</i>	30	2640	(Chambers <i>et al.</i> , 2006; Byrne, 2008)
<i>Uperoleia laevigata</i>	35	2260	(Robertson, 1986)
Nyctibatrachidae			
<i>Nyctibatrachus major</i> *	55	1250	(Boulenger, 1882; Kuramoto & Joshy, 2001)
Odontophrynidae			
<i>Macrogenioglottus alipioi</i>	100	515	(Dixo & Verdade, 2006; Abravaya & Jackson, 1978)
<i>Odontophrynus achalensis</i> *	49.4	933	(Salas & Di Tada, 1988; Rosset <i>et al.</i> , 2007)
<i>Odontophrynus americanus</i>	58	1035	(Martino & Sinsch, 2002)
<i>Odontophrynus cultripes</i>	60	660	(Caramaschi & Napoli, 2012)
<i>Odontophrynus occidentalis</i> *	60	820	(Martino & Sinsch, 2002)
<i>Proceratophrys avelinoi</i> *	29.2	1600	(Kwet & Faivovich, 2001; Kwet & Baldo, 2003)
<i>Proceratophrys bigibbosa</i> *	43.8	1050	(Kwet & Faivovich, 2001)
<i>Proceratophrys concavitympanum</i> *	57.1	818	(Santana <i>et al.</i> , 2010)
<i>Proceratophrys cristiceps</i> *	50.2	940	(Cruz <i>et al.</i> , 2012; Nunes & Juncá, 2006)
<i>Proceratophrys cururu</i> *	43.1	800	(Eterovick & Sazima, 1998)
<i>Proceratophrys goyana</i> *	46.5	1006	(Martins & Giarretta, 2013)
<i>Proceratophrys melanopogon</i> *	49.9	1179	(Prado & Pombal Jr, 2008; Mângia <i>et al.</i> , 2010)
Pelobatidae			
<i>Pelobates cultripes</i>	85	1075	(Lizana <i>et al.</i> , 1994)
Petropedetidae			
<i>Petropedetes yakusini</i> *	73	1300	(Channing & Howell, 2006)
Phrynobatrachidae			
<i>Phrynobatrachus acridoides</i>	28	2000	(Channing, 2001)
<i>Phrynobatrachus dendrobates</i> *	31	3800	(Channing & Howell, 2006)
Pipidae			
<i>Xenopus ruwenzoriensis</i>	44	1500	(Channing & Howell, 2006)
Ptychadenidae			
<i>Hildebrandtia ornata</i>	65	1400	(Channing, 2001)
<i>Ptychadena anchietae</i>	51	1750	(Harper <i>et al.</i> , 2010; van den Elzen & Kreulen, 1979)
<i>Ptychadena mahnerti</i>	42	3000	(Perret, 1996)
<i>Ptychadena oxyrhynchus</i>	53	1715	(Rödel, 2000; Bwong <i>et al.</i> , 2009)
<i>Ptychadena porosissima</i>	44.1	4091	(Dehling & Sinsch, 2013; Sinsch <i>et al.</i> , 2012)
<i>Ptychadena taenioscelis</i>	35	2450	(Harper <i>et al.</i> , 2010; Bwong <i>et al.</i> , 2009)
Pyxicephalidae			
<i>Cacosternum boettgeri</i>	19	4500	(Boulenger, 1882; Channing <i>et al.</i> , 2013)
<i>Cacosternum capense</i>	31.5	2300	(Channing, 2001; Channing <i>et al.</i> , 2013)
<i>Cacosternum nanum</i>	15	3600	(Channing <i>et al.</i> , 2013)
<i>Microbatrachella capensis</i>	15	4900	(Boulenger, 1910; Channing, 2001)
<i>Poyntonia paludicola</i> *	30	2200	(Channing & Boycott, 1989; Channing, 2001)
<i>Pyxicephalus adspersus</i>	245	225	(Du Preez, 1996; Boeckle <i>et al.</i> , 2009)

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
<i>Pyxicephalus edulis</i>	120	525	(Rödel, 2000; Channing <i>et al.</i> , 1994)
<i>Strongylopus grayii</i>	35	2300	(Channing, 2001)
<i>Tomopterna cryptotis</i>	45	3450	(Channing, 2001)
<i>Tomopterna delalandii</i>	44.8	2500	(Passmore & Carruthers, 1975)
<i>Tomopterna krugerensis</i>	44.8	2500	(Passmore & Carruthers, 1975)
<i>Tomopterna tuberculosa</i>	40	2600	(Channing, 2001)
Ranidae			
<i>Amnirana galamensis</i>	78	2500	(Channing, 2001)
<i>Amolops chunganensis</i> *	39	3425	(Liu <i>et al.</i> , 2000; Matsui <i>et al.</i> , 1993)
<i>Amolops larutensis</i> *	30	5033	(Van Kampen, 1923; Matsui <i>et al.</i> , 1993)
<i>Amolops panhai</i> *	34	6500	(Matsui & Nabhitabhata, 2006)
<i>Amolops wuyiensis</i> *	47	2232	(Zhang <i>et al.</i> , 2013)
<i>Huia cavitimpanum</i> *	52	15966	(Inger & Stuebing, 2005; Arch <i>et al.</i> , 2008)
<i>Huia masonii</i> *	37.8	12000	(Stuart & Chan-Ard, 2005; Boonman & Kurniati, 2011)
<i>Hylarana nicobariensis</i>	46.2	1825	(Inger, 1954; Jehle & Arak, 1998)
<i>Meristogenys jerboa</i> *	44	5650	(Inger & Gritis, 1983; Matsui <i>et al.</i> , 1993)
<i>Meristogenys orphnocnemis</i> *	37.3	7205	(Matsui, 1986; Preininger <i>et al.</i> , 2007)
<i>Odorrana schmackeri</i> *	46	3479	(Liu, 1950; Zhou <i>et al.</i> , 2014)
<i>Odorrana tormota</i> *	33	7000	(Feng <i>et al.</i> , 2002)
<i>Rana adenopleura</i>	51	650	(Matsui & Utsunomiya, 1983)
<i>Rana amamiensis</i> *	69	2850	(Matsui, 1994)
<i>Rana baramica</i>	55.6	2000	(Leong <i>et al.</i> , 2003; Zainudin <i>et al.</i> , 2010)
<i>Rana curtipes</i>	51.4	1222	(Krishna & Krishna, 2005)
<i>Rana dalmatina</i>	59	682	(Hettyey <i>et al.</i> , 2005; Lesbarrères & Lodé, 2002)
<i>Rana erythraea</i>	48	2459	(Brown & Alcala, 1970; Roy <i>et al.</i> , 1995)
<i>Rana forreri</i>	90	850	(Frost, 1982; Savage, 2002)
<i>Rana glandulosa</i>	55	1380	(Brown, 1902; Grafe <i>et al.</i> , 2008)
<i>Rana hosii</i> *	68	4950	(Manthey & Grossmann, 1997; Kurniati <i>et al.</i> , 2010)
<i>Rana laterimaculata</i> *	39	3245	(Leong <i>et al.</i> , 2003)
<i>Rana lessonae</i>	57	1831	(Wycheley <i>et al.</i> , 2002)
<i>Rana livida</i> *	54	5250	(Ao <i>et al.</i> , 2003; Shen <i>et al.</i> , 2011)
<i>Rana luctuosa</i>	68	1500	(Kueh <i>et al.</i> , 2010; Zainudin <i>et al.</i> , 2010)
<i>Rana nigromaculata</i>	81	2043	(Khonsue <i>et al.</i> , 2001; Mu & Zhao, 1990)
<i>Rana picturata</i> *	54	2500	(Zainudin & Sazali, 2012; Zainudin <i>et al.</i> , 2010)
<i>Rana plancyi</i>	50	1840	(Lue, 1990; Ueda, 1994)
<i>Rana pretiosa</i>	75	1000	(Cushman & Pearl, 2007; Briggs, 1987)
<i>Rana septentrionalis</i>	63.1	605	(Leclair & Laurin, 1996; Bevier <i>et al.</i> , 2004)
<i>Rana supranarina</i> *	76.8	1840	(Matsui, 1994)
<i>Rana taylori</i>	78	1350	(Savage, 2002)
<i>Rana temporalis</i> *	55.3	3100	(Hampson & Bennett, 2002; Kadadevaru <i>et al.</i> , 2000)
<i>Rana utsunomiyaorum</i> *	48.1	2326	(Matsui, 1994)
<i>Staurois latopalatus</i> *	47.7	5149	(Boeckle <i>et al.</i> , 2009)
<i>Staurois natator</i> *	35.5	4746	(Emerson, 1997; Preininger <i>et al.</i> , 2007)

Taxa	SVL (mm)	DF (Hz)	Source (first reference always SVL)
<i>Staurois parvus</i> *	23.6	5578	(Matsui <i>et al.</i> , 2007; Grafe <i>et al.</i> , 2012)
<i>Staurois tuberilinguis</i> *	30.2	5240	(Matsui <i>et al.</i> , 2007; Boeckle <i>et al.</i> , 2009)
Rhacophoridae			
<i>Buergeria robusta</i> *	52	1850	(Huang <i>et al.</i> , 2001; Kuramoto, 1986)
<i>Chiromantis doriae</i>	26.5	4300	(Aowphol <i>et al.</i> , 2013)
<i>Chiromantis rufescens</i>	49	1650	(Channing & Howell, 2006)
<i>Chiromantis vittatus</i>	25.2	3900	(Aowphol <i>et al.</i> , 2013)
<i>Chiromantis xerampelina</i>	75	1700	(Channing & Howell, 2006)
<i>Ghatixalus variabilis</i> *	48	3135	(Bossuyt & Dubois, 2001; Kanamadi <i>et al.</i> , 2001)
<i>Kurixalus idiootocus</i>	24.9	2300	(Kuramoto & Wang, 1987)
<i>Polypedates leucomystax</i>	63	1528	(Sheridan <i>et al.</i> , 2010)
<i>Polypedates maculatus</i>	57	1550	(Daniel, 2002; Kanamadi <i>et al.</i> , 1993)
<i>Rhacophorus arboreus</i>	60	1400	(Wilkinson, 2003; Kasuya <i>et al.</i> , 1992)
<i>Rhacophorus chenfui</i>	41	2329	(Wang <i>et al.</i> , 2012; Matsui & Wu, 1994)
<i>Rhacophorus dennysi</i>	96	1321	(Wang <i>et al.</i> , 2012)
<i>Rhacophorus dugritei</i>	40.8	1600	(Ohler <i>et al.</i> , 2000; Matsui & Wu, 1994)
<i>Rhacophorus kio</i>	79.1	890	(Wildenhues <i>et al.</i> , 2011; Ohler & Delorme, 2006)
<i>Rhacophorus malabaricus</i>	48.3	1260	(Das, 2000; Hampson & Bennett, 2002)
<i>Rhacophorus omeimontis</i>	70	972	(Liao & Lu, 2011; Matsui & Wu, 1994)
<i>Rhacophorus reinwardtii</i>	52.5	1300	(Ohler & Delorme, 2006; Kurniati <i>et al.</i> , 2010)
<i>Rhacophorus schlegelii</i>	43	1900	(Maeda & Matsui, 1990; Matsui & Wu, 1994)
Rhinophrynidae			
<i>Rhinophrynus dorsalis</i>	75	1525	(Savage, 2002)
Scaphiopodidae			
<i>Scaphiopus couchii</i>	72	1650	(Vásquez & Pfennig, 2007; Capranica & Moffat, 1975)
<i>Scaphiopus holbrookii</i>	78	1425	(Dodd Jr., 2013; Blair, 1958b)
<i>Scaphiopus hurterii</i>	67	1500	(Strecker, 1910; Awbrey, 1968)
<i>Spea multiplicata</i>	49	1300	(Pfennig & Pfennig, 2005; Pfennig, 2000)

- Abravaya, J.P. & Jackson, J.F. 1978. Reproduction in *Macrogenioglottus alipioi* Carvalho (Anura, Leptodactylidae). *Natural History Museum of Los Angeles County, Contributions to Science* **298**: 1-9.
- Abrunhosa, P.A., Wogel, H. & Pombal Jr., J.P. 2001. Vocalização de quatro espécies de anuros do estado do Rio de Janeiro, sudeste do Brasil (Amphibia, Hylidae, Leptodactylidae). *Boletim do Museu Nacional do Rio de Janeiro* **472**: 1-12.
- Abrunhosa, P.A. & Wogel, H. 2004. Breeding behavior of the leaf-frog *Phyllomedusa burmeisteri* (Anura : Hylidae). *Amphibia-Reptilia* **25**: 125-135.
- Abrunhosa, P.A., Pimenta, B.V.S., Cruz, C.A.G. & Haddad, C.F.B. 2005. Advertisement calls of species of the *Hyla albosignata* group (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional* **63**: 275-282.
- Alonso, R. & Rodríguez, A. 2003. Advertisement calls of Cuban toads of the genus *Bufo* (Anura, Bufonidae). *Phyllomedusa* **2**: 75-82.
- Álvares, G.F.R. 2008. Taxonomia, distribuição geográfica potencial e conservação das espécies de *Phyllomedusa* do grupo *hypochondrialis*, Universidade de Brasília. **Master**: 95.
- Amézquita, A. 1999. Color pattern, elevation and body size in the high Andean frog *Hyla labialis*. *Revista de la Academia Colombiana de Ciencias Exactas Físicas y Naturales* **23**: 231-238.
- Andreone, F., Piazza, R. & Vallan, D. 1994. Evidence for the specific validity and resurrection of *Heterixalus luteostriatus* (Andersson, 1910) (Anura: Hyperoliidae). *Bollettino del Museo regionale di Scienze Naturali Torino* **12**: 391-400.

- Andreone, F., Mattioli, F. & Mercurio, V. 2005. The call of *Scaphiophryne gottlebei*, a microhylid frog from the Isalo Massif, southcentral Madagascar. *Current Herpetology* **24**: 33-35.
- Ao, J.M., Bordoloi, S. & Ohler, A. 2003. Amphibian fauna of Nagaland with nineteen new records from the state including five new records for India. *Zoos' Print Journal* **18**: e1117-1125.
- Aowphol, A., Rujirawan, A., Taksintum, W., Arsirapot, S. & McLeod, D.S. 2013. Re-evaluating the taxonomic status of *Chiromantis* in Thailand using multiple lines of evidence (Amphibia: Anura: Rhacophoridae). *Zootaxa* **3702**: 101-123.
- Arak, A. 1988. Female mate selection in the natterjack toad: active choice or passive attraction? *Behavioral Ecology and Sociobiology* **22**: 317-327.
- Arch, V.S., Grafe, T.U. & Narins, P.M. 2008. Ultrasonic signalling by a Bornean frog. *Biology Letters* **4**: 19-22.
- Arnold, E.N. 2003. *Reptiles and amphibians of Europe*. Princeton University Press, Princeton.
- Awbrey, F.T. 1968. Call discrimination in female *Scaphiopus couchii* and *Scaphiopus hurterii*. *Copeia* **1968**: 420-423.
- Barrio-Amoros, C.L. 2006. A new species of *Phyllomedusa* (Anura : Hylidae : Phyllomedusinae) from northwestern Venezuela. *Zootaxa* **1309**: 55-68.
- Barrio-Amorós, C.L. & Castroviejo-Fisher, S. 2008. Comments on the distribution, taxonomy and advertisement call of the Guyanan glass frog *Hyalinobatrachium igniocolus* (Anura: Centrolenidae). *Salamandra* **44**: 235-240.
- Barrio-Amorós, C.L., Señaris, J.C., Macculloch, R.D., Lathrop, A., Guayasamin, J.M. & Duellman, W.E. 2011. Distribution, vocalization and taxonomic status of *Hypsiboas*

- roraïma* and *H. angelicus* (Amphibia: Anura: Hylidae). *Papéis Avulsos de Zoologia (São Paulo)* **51**: 21-28.
- Barrio, A. 1967. Observaciones etoecologicas sobre *Hylorina sylvatica* Bell (Anura, Leptodactylidae). *Physis* **27**: 153-157.
- Basso, N.G. 1990. Estrategias adaptivas en una comunidad subtropical de anuros. *Cuadernos de Herpetologia* **1**: 1-70.
- Bastos, R.P. & Haddad, C.F.B. 1995. Vocalizações e interações acústicas de *Hyla elegans* (Anura: Hylidae) durante a atividade reprodutiva. *Naturalia* **20**: 165-176.
- Bastos, R.P. & Pombal, J.P. 1995. New species of *Crossodactylus* (Anura: Leptodactylidae) from the Atlantic rain forest of southeastern Brazil. *Copeia* **1995**: 436-439.
- Bernal, M.H., Montealegre, D.P. & Páez, C.A. 2004. Estudio de la vocalización de trece especies de anuros del municipio de Ibagué, Colombia. *Revista de la Academia Colombiana de Ciencias Exactas Físicas y Naturales* **28**: 385-390.
- Bevier, C.R., Larson, K., Reilly, K. & Tat, S. 2004. Vocal repertoire and calling activity of the mink frog, *Rana septentrionalis*. *Amphibia-Reptilia* **25**: 255-264.
- Bilate, M., Wogel, H., Weber, L.N. & Abrunhosa, P.A. 2006. Vocalizações e girino de *Leptodactylus spixi* Heyer, 1983 (Amphibia, Anura, Leptodactylidae). *Arquivos do Museu Nacional* **63**: 235-245.
- Blackburn, D.C., Kosuch, J., Schmitz, A., Burger, M., Wagners, P., Gonwouo, L.N., Hillers, A. & Roedel, M.-O. 2008. A new species of *Cardioglossa* (Anura : Arthroleptidae) from the Upper Guinean forests of West Africa. *Copeia* **2008**: 603-612.
- Blair, W.F. 1956. Call difference as an isolation mechanism in south-western toads (genus *Bufo*). *Texas Journal of Science* **8**: 87-106.

- Blair, W.F. 1958a. Call structure and species groups in US treefrogs (*Hyla*). *The Southwestern Naturalist* **3**: 77-89.
- Blair, W.F. 1958b. Mating call in the speciation of anuran amphibians. *American Naturalist* **92**: 27-51.
- Boeckle, M., Preininger, D. & Hödl, W. 2009. Communication in noisy environments I: acoustic signals of *Staurois latopalmatus* Boulenger 1887. *Herpetologica* **65**: 154-165.
- Bogert, C.M. 1962. Isolation mechanisms in toads of the *Bufo debilis* group in Arizona and western Mexico. *American Museum Novitates* **2100**: 1-37.
- Bokermann, W.C.A. 1965. Três novos batráquios da região central de Mato Grosso, Brasil (Amphibia, Salientia). *Revista Brasileira de Biologia* **25**: 257-264.
- Boonman, A. & Kurniati, H. 2011. Evolution of high-frequency communication in frogs. *Evolutionary Ecology Research* **13**: 197-207.
- Bossuyt, F. & Dubois, A. 2001. A review of the frog genus *Philautus* Gistel, 1848 (Amphibia, Anura, Ranidae, Rhacophorinae). *Zeylanica* **6**: 1-112.
- Boulenger, G.A. 1882. *Catalogue of the Batrachia Salientia s. Ecaudata in the collection of the British Museum*. Collection of the British Museum (Natural History), London.
- Boulenger, G.A. 1910. A revised list of the South African reptiles and batrachians, with synoptic tables, special reference to the specimens in the South African Museum, and descriptions of new species. *South African Museum* **5**: 455-538.
- Brandão, R.A., Álvares, G.F., Crema, A. & Zerbini, G.J. 2009. Natural history of *Phyllomedusa centralis* Bokermann 1965 (Anura: Hylidae: Phyllomedusinae): tadpole and calls. *South American Journal of Herpetology* **4**: 61-68.
- Briggs, J.L. 1987. Breeding biology of the Cascade frog, *Rana cascadae*, with comparisons to *R. aurora* and *R. pretiosa*. *Copeia* **1987**: 241-245.

- Brown, A.E. 1902. A collection of reptiles and batrachians from Borneo and the Loo Choo Islands. *Proceedings of the Academy of Natural Sciences of Philadelphia* **54**: 175-186.
- Brown, R.M. & Iskandar, D.T. 2000. Nest site selection, larval hatching, and advertisement calls, of *Rana arathooni* from southwestern Sulawesi (Celebes) Island, Indonesia. *Journal of Herpetology* **34**: 404-413.
- Brown, W.C. & Alcala, A.C. 1970. Population ecology of the frog *Rana erythraea* in Southern Negros, Philippines. *Copeia* **1970**: 611-622.
- Bwong, B.A., Chira, R., Schick, S., Veith, M. & Lötters, S. 2009. Diversity of ridged frogs (Ptychadenidae: *Ptychadena*) in the easternmost remnant of the Guineo-Congolian rain forest: an analysis using morphology, bioacoustics and molecular genetics. *Salamandra* **45**: 129-146.
- Byrne, P.G. 2008. Strategic male calling behavior in an Australian Terrestrial Toadlet (*Pseudophryne bibronii*). *Copeia* **1**: 57-63.
- Cadle, J.E. & McDiarmid, R.W. 1990. Two new species of *Centrolenella* (Anura: Centrolenidae) from northwestern Peru. *Proceedings of the Biological Society of Washington* **103**: 746-768.
- Cajade, R., Federico Schaefer, E., Ines Dure, M., Ignacio Kehr, A. & Marangoni, F. 2010. Reproductive biology of *Argenteohyla siemersi pedersenii* Williams and Bosso, 1994 (Anura: Hylidae) in northeastern Argentina. *Journal of Natural History* **44**: 1953-1978.
- Caldart, V.M., Iop, S. & Cechin, S.Z. 2011. Vocalizations of *Crossodactylus schmidtii* Gallardo, 1961 (Anura, Hylodidae): advertisement call and aggressive call. *North-Western Journal of Zoology* **7**: 118-124.
- Caldwell, J.P. & Shepard, D.B. 2007. Calling site fidelity and call structure of a neotropical toad, *Rhinella ocellata* (Anura : Bufonidae). *Journal of Herpetology* **41**: 611-621.

- Cannatella, D.C. 1982. Leaf-frogs of the *Phyllomedusa perinesos* group (Anura: Hylidae). *Copeia* **1982**: 501-513.
- Cannatella, D.C. & Duellman, W.E. 1984. Leptodactylid frogs of the *Physalaemus pustulosus* group. *Copeia* **1984**: 902-921.
- Cannatella, D.C. & Lamar, W.W. 1986. Synonymy and distribution of *Centrolenella orientalis* with notes on its life history (Anura: Centrolenidae). *Journal of Herpetology* **20**: 307-317.
- Capranica, R.R. & Moffat, A.J. 1975. Selectivity of the peripheral auditory system of spadefoot toads (*Scaphiopus couchi*) for sounds of biological significance. *Journal of Comparative Physiology* **100**: 231-249.
- Caramaschi, U. & Cruz, C.A.G. 1998. Notas taxonômicas sobre *Pseudis fusca* Garman e *P. bolboactyla* A. Lutz, com a descrição de uma nova espécie correlata (Anura, Pseudidae). *Revista Brasileira de Biologia* **15**: 929-944.
- Caramaschi, U. & Cruz, C.A.G. 2000. Duas espécies novas de *Hyla* Laurenti, 1768 do estado de Goiás, Brasil (Amphibia, Anura, Hylidae). *Boletim do Museu Nacional do Rio de Janeiro, Nova Série, Zoologia* **422**: 1-12.
- Caramaschi, U. 2006. Redefinição do grupo de *Phyllomedusa hypochondrialis*, com redescritção de *P. megacephala* (Miranda-Ribeiro, 1926), revalidação de *P. azurea* Cope, 1862 e descrição de uma nova espécie (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional* **64**: 159-179.
- Caramaschi, U. & Napoli, M.F. 2012. Taxonomic revision of the *Odontophrynus cultripes* species group, with description of a new related species (Anura, Cycloramphidae). *Zootaxa* **3155**: 1-20.

- Cardoso, A.J. & Vielliard, J.M. 1985. Caracterização bio-acústica da população topotípica de *Hyla rubicundula* (Amphibia, Anura). *Revista Brasileira de Zoologia* **2**: 423-426.
- Cardozo-Urdaneta, A. & Señaris, J.C. 2012. Vocalización y biología reproductiva de las ranas de cristal *Hyalinobatrachium pallidum* y *Centrolene daidaleum* (Anura, Centrolenidae) en la sierra de Perijá, Venezuela. *Memoria de la Fundación La Salle de Ciencias Naturales* **70**: 87-105.
- Carvalho Jr, R.R., Galdino, C.A. & Nascimento, L.B. 2006. Notes on the courtship behavior of *Aplastodiscus arildae* (Cruz and Peixoto, 1985) at an urban forest fragment in Southeastern Brazil (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional* **64**: 247-254.
- Carvalho, T.R., Giaretta, A.A. & Magrini, L. 2012. A new species of the *Bokermannohyla circumdata* group (Anura: Hylidae) from southeastern Brazil, with bioacoustic data on seven species of the genus. *Zootaxa* **3321**: 37-55.
- Castellano, S., Cuatto, B., Rinella, R., Rosso, A. & Giacoma, C. 2002. The advertisement call of the European treefrogs (*Hyla arborea*): A multilevel study of variation. *Ethology* **108**: 75-89.
- Castroviejo-Fisher, S., Señaris, J.C., Ayarzagüena, J. & Vilà, C. 2008. Resurrection of *Hyalinobatrachium orcostale* and notes on the *Hyalinobatrachium orientale* species complex (Anura: Centrolenidae). *Herpetologica* **64**: 472-484.
- Castroviejo-Fisher, S., Padial, J.M., Chaparro, J.C., Aguayo, R. & De la Riva, I. 2009. A new species of *Hyalinobatrachium* (Anura: Centrolenidae) from the Amazonian slopes of the central Andes, with comments on the diversity of the genus in the area. *Zootaxa* **2143**: 24-44.

- Catenazzi, A., Rodríguez, L.O. & Donnelly, M.A. 2009. The advertisement calls of four species of glassfrogs (Centrolenidae) from southeastern Peru. *Studies on Neotropical Fauna and Environment* **44**: 83-91.
- Cei, J.M. 1980. Amphibians of Argentina. *Monitore Zoologico Italiano* **2**: 1-609.
- Chambers, J., Wilson, J.C. & Williamson, I. 2006. Soil pH influences embryonic survival in *Pseudophryne bibronii* (Anura : Myobatrachidae). *Austral Ecology* **31**: 68-75.
- Channing, A. 1976. Pre-mating isolation in the genus *Kassina* (Amphibia, Anura, Rhacophoridae) in southern Africa. *Journal of Herpetology* **10**: 19-23.
- Channing, A. & Boycott, R.C. 1989. A new frog genus and species from the mountains of the southwestern Cape, South Africa (Anura: Ranidae). *Copeia* **1989**: 467-471.
- Channing, A., Dupree, L. & Passmore, N. 1994. Status, vocalization and breeding biology of two species of African bullfrogs (Ranidae: *Pyxicephalus*). *Journal of Zoology* **234**: 141-148.
- Channing, A. 2001. *Amphibians of Central and Southern Africa*. Cornell University Press., Ithaca, NY.
- Channing, A. & Howell, K. 2006. *Amphibians of East Africa*. Cornell University Press Ithaca, NY.
- Channing, A., Schmitz, A., Burger, M. & Kielgast, J. 2013. A molecular phylogeny of African Dainty Frogs, with the description of four new species (Anura: Pyxicephalidae: *Cacosternum*). *Zootaxa* **3701**: 518-550.
- Cherry, M.I. & Francillon-Vieillot, H. 1992. Body size, age and reproduction in the leopard toad, *Bufo pardalis*. *Journal of Zoology* **228**: 41-50.
- Cherry, M.I. & Grant, W.S. 1994. Phylogenetic relationships and call structure in four African bufonid species. *South African Journal of Zoology* **29**: 1-10.

Cochran, D.M. 1955. Frogs of Southeastern Brazil. *United States National Museum Bulletin* **206**: 1-423.

Cochran, D.M. & Goin, C.J. 1970. *Frogs of Colombia*. Smithsonian Institution Press, Washington.

Cocroft, R.B., McDiarmid, R.W., Jaslow, A.P. & Ruiz-Carranza, P.M. 1990. Vocalizations of eight species of *Atelopus* (Anura: Bufonidae) with comments on communication in the genus. *Copeia* **1990**: 631-643.

Cocroft, R.B. & Ryan, M.J. 1995. Patterns of advertisement call evolution in toads and chorus frogs. *Animal Behaviour* **49**: 283-303.

Coloma, L.A., Lötters, S. & Salas, A.W. 2000. Taxonomy of the *Atelopus ignescens* complex (Anura : Bufonidae): Designation of a neotype of *Atelopus ignescens* and recognition of *Atelopus exiguus*. *Herpetologica* **56**: 303-324.

Coloma, L.A., Carvajal-Endara, S., Duenas, J.F., Paredes-Recalde, A., Morales-Mite, M., Almeida-Reinoso, D., Tapia, E.E., Hutter, C.R., Toral, E. & Guayasamin, J.M. 2012. Molecular phylogenetics of stream treefrogs of the *Hyloscirtus larinopygion* group (Anura: Hylidae), and description of two new species from Ecuador. *Zootaxa* **3364**: 1-78.

Conant, R. & Collins, J.T. 1998. *A field guide to reptiles and amphibians of Eastern and Central North America*. Horton Mifflin Company, Boston.

Cruz, C.A.G. & Peixoto, O.L. 1985. Espécies verdes de *Hyla*: o complexo “*albosignata*” (Amphibia, Anura, Hylidae). *Arquivos da Universidade Federal Rural do Rio de Janeiro* **7**: 31-47.

Cruz, C.A.G., Nunes, I. & Juncá, F.A. 2012. Redescription of *Proceratophrys cristiceps* (Müller, 1883) (Amphibia, Anura, Odontophrynidae), with description of two new species

- without eyelid appendages from Northeastern Brazil. *South American Journal of Herpetology* **7**: 110-122.
- Cushman, K.A. & Pearl, C.A. 2007. A conservation assessment for the Oregon spotted frog (*Rana pretiosa*), USDA Forest Service and USDI Bureau of Land Management. Oregon.
- Daniel, J.C. 2002. *The book of Indian reptiles and amphibians* Oxford University Press, Mumbai.
- Das, I. 2000. Nomenclatural history and rediscovery of *Rhacophorus lateralis* Boulenger, 1883 (Amphibia: Rhacophoridae). *Current Herpetology* **19**: 35-40.
- Davies, M., Martin, A.A. & Watson, G.F. 1983. Redefinition of the *Litoria latopalmata* species group (Anura: Hylidae). *Transactions of the Royal Society of South Australia* **107**: 87-108.
- De la Riva, I. 1993. A new species of *Scinax* (Anura, Hylidae) from Argentina and Bolivia. *Journal of Herpetology* **27**: 41-46.
- De la Riva, I., Márquez, R. & Bosch, J. 1995. Advertisement calls of eight Bolivian hylids (Amphibia, Anura). *Journal of Herpetology* **29**: 112-118.
- De la Riva, I., Márquez, R. & Bosch, J. 1996. Advertisement calls of four microhylid frogs from Bolivia (Amphibia, Anura). *American Midland Naturalist* **136**: 418-422.
- De la Riva, I., Márquez, R. & Bosch, J. 1997. Description of the advertisement calls of some South American Hylidae (Amphibia: Anura): taxonomic and methodological consequences. *Bonner Zoologische Beiträge* **47**.
- Dehling, J.M. & Sinsch, U. 2013. Diversity of Ridged Frogs (Anura: Ptychadenidae: *Ptychadena* spp.) in wetlands of the upper Nile in Rwanda: Morphological, bioacoustic, and molecular evidence. *Zoologischer Anzeiger* **253**: 143-157.

- Díaz-Gutiérrez, N., Vargas-Salinas, F., Rivera-Correa, M., Andres Rojas-Morales, J., Escobar-Lasso, S., Velasco, J.A., Gutiérrez-Cárdenas, P.D.A. & Amézquita, A. 2013. Description of the previously unknown advertisement call and tadpole of the Colombian endemic glassfrog *Centrolene savagei* (Anura: Centrolenidae). *Zootaxa* **3686**: 289-296.
- Dixo, M. & Verdade, V.K. 2006. Herpetofauna de serrapilheira da Reserva Florestal de Morro Grande, Cotia (SP). *Biota Neotropica* **6**: 1-20.
- Dodd Jr., C.K. 2013. *Frogs of the United States and Canada*. JHU Press, Baltimore.
- Doughty, P. 2011. An emerging frog diversity hotspot in the northwest Kimberley of Western Australia: another new frog species from the high rainfall zone. *Records of the Western Australian Museum* **26**: 209-216.
- Du Preez, L.H. 1996. *Field guide and key to the frogs and toads of the Free State*. University of the Orange Free State, Bloemfontein, South Africa.
- Duellman, W.E. & Fouquette, M.J. 1968. Middle American frogs of the *Hyla microcephala* group. *University of Kansas Publications of the Museum of Natural History* **17**: 517-557.
- Duellman, W.E. 1969. A new species of frog in the *Hyla parviceps* group from Ecuador. *Herpetologica* **25**: 241-247.
- Duellman, W.E. 1970. *The Hylid Frogs of Middle America*. Monograph of the Museum of Natural History, Kansas.
- Duellman, W.E. 1971. The identities of some Ecuadorian hylid frogs. *Herpetologica* **27**: 212-227.
- Duellman, W.E. 1972a. A review of the Neotropical frogs of the *Hyla bogotensis* group. *Occasional Papers of the Museum of Natural History, The University of Kansas* **11**: 1-31.

- Duellman, W.E. 1972b. South American frogs of the *Hyla rostrata* group (Amphibia, Anura, Hylidae). *Zoologische Mededelingen* **47**: 177-192.
- Duellman, W.E. 1973. Descriptions of new hylid frogs from Colombia and Ecuador. *Herpetologica* **29**: 219-227.
- Duellman, W.E. 1974. Taxonomic notes on *Phyllomedusa* (Anura: Hylidae) from the upper Amazon basin. *Herpetologica* **30**: 105-112.
- Duellman, W.E. & Veloso, A. 1977. Phylogeny of *Pleurodema* (Anura, Leptodactylidae): a biogeographic model. *Occasional Papers of the Museum of Natural History, The University of Kansas* **64**: 1-46.
- Duellman, W.E. 1978. The biology of an equatorial herpetofauna in Amazonian Ecuador. *Miscellaneous Publications of the University of Kansas Museum of Natural History* **65**: 1-352.
- Duellman, W.E. & Altig, R. 1978. New species of tree frogs (Family Hylidae) from the Andes of Colombia and Ecuador. *Herpetologica* **34**: 177-185.
- Duellman, W.E. & Berger, T.J. 1982. A new species of Andean treefrog (Hylidae). *Herpetologica* **38**: 456-460.
- Duellman, W.E. & Pyles, R.A. 1983. Acoustic resource partitioning in anuran communities. *Copeia* **1983**: 639-649.
- Duellman, W.E., Cadle, J.E. & Cannatella, D.C. 1988. A new species of terrestrial *Phyllomedusa* (Anura: Hylidae) from southern Peru. *Herpetologica* **44**: 91-95.
- Duellman, W.E. & Hillis, D.M. 1990. Systematics of the *Hyla larinopygion* group *Occasional Papers of the Museum of Natural History, The University of Kansas* **134**: 1-23.

- Duellman, W.E. & Wiens, J.J. 1993. Hylid frogs of the genus *Scinax* Wagler, 1830, in Amazonian Ecuador and Peru. *Occasional Papers of the Museum of Natural History, The University of Kansas* **153**: 1-57.
- Duellman, W.E. & Lizana, M. 1994. Biology of a sit-and-wait predator, the leptodactylid frog *Ceratophrys cornuta*. *Herpetologica* **50**: 51-64.
- Duellman, W.E. 1997. Amphibians of La Escalera Region, Southeastern Venezuela: taxonomy, ecology, and biogeography. *Scientific Papers Natural History Museum, The University of Kansas* **2**: 1-52.
- Duellman, W.E. 2005. *Cusco Amazónico*. Comstock Pub. Associates, Ithaca.
- Emerson, S.B. 1997. Testis size variation in frogs: testing the alternatives. *Behavioral Ecology and Sociobiology* **41**: 227-235.
- Eterovick, P.C. & Sazima, I. 1998. New species of *Proceratophrys* (Anura : Leptodactylidae) from Southeastern Brazil. *Copeia* **1998**: 159-164.
- Faivovich, J. 2005. A new species of *Scinax* (Anura : Hylidae) from Misiones, Argentina. *Herpetologica* **61**: 69-77.
- Fei, L., Ye, C.Y., Huang, Y.A. & Liu, M.Y. 1999. *Atlas of Amphibians of China*. Henan Science and Technical Press, Zhengzhou, China.
- Feng, A.S., Narins, P.M. & Xu, C.H. 2002. Vocal acrobatics in a Chinese frog, *Amolops tormotus*. *Naturwissenschaften* **89**: 352-356.
- Fonte, L.F.M.D. 2010. Variação morfológica e na estrutura do canto em *Scinax granulatus* (PETERS, 1871) (ANURA, HYLIDAE), Universidade Federal do Rio Grande do Sul. **Master**: 76.
- Ford, M.A. 1989. Vocalisations of the green frogs *Litoria aurea* and *Litoria raniformis*, including species-specific male calls. *New Zealand Journal of Zoology* **16**: 17-23.

- Forti, L.R., Miguel Martins, F.A. & Bertoluci, J. 2012. Advertisement call and geographical variation in call features of *Dendropsophus berthalutzae* (Anura: Hylidae) from the Atlantic Rainforest of southeastern Brazil. *Zootaxa* **3310**: 66-68.
- Frost, J.S. 1982. Functional genetic similarity between geographically separated populations of Mexican leopard frogs (*Rana pipiens* complex). *Systematic Zoology* **31**: 57-67.
- Gallardo, J.M. 1965. The species *Bufo granulosus* Spix (Salientia: Bufonidae) and its geographic variation. *Bulletin of the Museum of Comparative Zoology* **134**: 107-138.
- Garcia-Rutledge, E.J. & Narins, P.M. 2001. Shared acoustic resources in an old world frog community. *Herpetologica* **57**: 104-116.
- Garcia, P.C.A., Vinciprova, G. & Haddad, C.F.B. 2003. The taxonomic status of *Hyla pulchella joaquina* (Anura : Hylidae) with description of its tadpole and vocalization. *Herpetologica* **59**: 350-363.
- Garcia, P.C.A., Faivovich, J. & Haddad, C.F.B. 2007. Redescription of *Hypsiboas semiguttatus*, with the description of a new species of the *Hypsiboas pulchellus* group. *Copeia* **2007**: 933-951.
- Garcia, P.C.A. & Haddad, C.F.B. 2008. Vocalizations and comments on the relationships of *Hypsiboas ericae* (Amphibia, Hylidae). *Iheringia Serie Zoologia* **98**: 161-166.
- Gerhardt, H.C. 1974. Behavioral isolation of the tree frogs, *Hyla cinerea* and *Hyla andersonii*. *American Midland Naturalist* **91**: 424-433.
- Gerhardt, H.C. 2005. Advertisement-call preferences in diploid-tetraploid treefrogs (*Hyla chrysoscelis* and *Hyla versicolor*): Implications for mate choice and the evolution of communication systems. *Evolution* **59**: 395-408.

- Giaretta, A.A., de Oliveira Filho, J.C. & Kokubum, M.N.d.C. 2007. A new *Phyllomedusa* Wagler (Anura, Hylidae) with reticulated pattern on flanks from Southeastern Brazil. *Zootaxa* **1614**: 31-41.
- Giaretta, A.A., De Andrade, F.S., Haga, I.A. & Bernardes, C.D.S. 2013. On the advertisement call of *Dermatonotus muelleri* (Boettger, 1885) (Anura, Microhylidae). *Zootaxa* **3700**: 593-596.
- Glaw, F. & Vences, M. 1991. Ein neuer *Heterixalus* aus Madagaskar (Amphibia, Anura, Hyperoliidae). *Acta Biologica Benrodis* **3**: 197-201.
- Glaw, F. & Vences, M. 1992. Zur Kenntnis der Gattungen *Boophis*, *Aglyptodactylus* und *Mantidactylus* (Amphibia: Anura) aus Madagaskar, mit Beschreibung einer neuen Art. *Bonner Zoologische Beiträge* **43**: 45-77.
- Glaw, F. & Thiesmeier, B. 1993. Bioakustische Differenzierung in der *Boophis-luteus*-Gruppe (Anura: Rhacophoridae), mit Beschreibung einer neuen Art und einer neuen Unterart. *Salamandra* **28**: 258-269.
- Glaw, F. & Vences, M. 1993. Zur Bioakustik, Biologie und Systematik der Gattung *Heterixalus* aus Madagaskar (Anura: Hyperoliidae). *Salamandra* **29**: 212-230.
- Glaw, F. & Vences, M. 1997. New species of the *Boophis tephraeomystax* group (Anura: Ranidae: Rhacophorinae) from arid Western Madagascar. *Copeia* **1997**: 572-578.
- Glaw, F., Vences, M. & Bohme, W. 1998. Systematic revision of the genus *Aglyptodactylus*, Boulenger, 1919 (Amphibia : Ranidae), and analysis of its phylogenetic relationships to other Madagascan ranid genera (*Tomopterna*, *Boophis*, *Mantidactylus*, and *Mantella*). *Journal of Zoological Systematics and Evolutionary Research* **36**: 17-37.
- Glaw, F., Vences, M. & Gossmann, V. 2000. A new species of *Mantidactylus* (subgenus *Guibemantis*) from Madagascar, with a comparative survey of internal femoral gland

- structure in the genus (Amphibia : Ranidae : Mantellinae). *Journal of Natural History* **34**: 1135-1154.
- Glaw, F., Vences, M., Andreone, F. & Vallan, D. 2001. Revision of the *Boophis majori* group (Amphibia : Mantellidae) from Madagascar, with descriptions of five new species. *Zoological Journal of the Linnean Society* **133**: 495-529.
- Glaw, F. & Vences, M. 2002. A new sibling species of the anuran subgenus *Blommersia* from Madagascar (Amphibia : Mantellidae : *Mantidactylus*) and its molecular phylogenetic relationships. *Herpetological Journal* **12**: 11-20.
- Glaw, F. & Vences, M. 2007. *A field guide to the amphibians and reptiles of Madagascar*. Verlag, Köln.
- Glos, J., Glaw, F. & Vences, M. 2005. A new species of *Scaphiophryne* from Western Madagascar. *Copeia* **2005**: 252-261.
- Goin, C.J. & Woodley, J.D. 1969. A new tree frog from Guyana. *Zoological Journal of the Linnean Society* **48**: 135-140.
- Gomes, F.B.R. & Martins, I.A. 2006. Amphibia, Anura, Hylidae, *Dendropsophus anceps* (Lutz, 1929): Filling gap, geographic distribution map and vocalization. *Check List* **2**: 22.
- Grafe, T.U. 1999. A function of synchronous chorusing and a novel female preference shift in an anuran. *Proceedings of the Royal Society B-Biological Sciences* **266**: 2331-2336.
- Grafe, T.U., Saat, H.B.M., Hagen, N., Kaluza, B., Berudin, Z.B.H. & Wahab, M.A.B.A. 2008. Acoustic localisation of frog hosts by blood-sucking flies *Corethrella* Coquillett (Diptera: Corethrellidae) in Borneo. *Australian Journal of Entomology* **47**: 350-354.
- Grafe, T.U., Preininger, D., Sztatecsny, M., Kasah, R., Dehling, J.M., Proksch, S. & Hödl, W. 2012. Multimodal communication in a noisy environment: a case study of the Bornean rock frog *Staurois parvus*. *Plos One* **7**: e37965.

Grant, T., Bolivar, W. & Castro, F. 1998. The advertisement call of *Centrolene geckoideum*.
Journal of Herpetology **32**: 452-455.

Guarnizo, C.E. 2011. Effect of topography on genetic divergence and phenotypic traits in tropical frogs, The University of Texas at Austin. **PhD**: 98.

Guayasamin, J.M. & Bonaccorso, E. 2004. A new species of glass frog (Centrolenidae : *Cochranella*) from the lowlands of northwestern Ecuador, with comments on the *Cochranella granulosa* group. *Herpetologica* **60**: 485-494.

Guayasamin, J.M., Bustamante, M.R., Almeida-Reinoso, D. & Funk, W.C. 2006. Glass frogs (Centrolenidae) of Yanayacu Biological Station, Ecuador, with the description of a new species and comments on centrolenid systematics. *Zoological Journal of the Linnean Society* **147**: 489-513.

Guayasamin, J.M., Castroviejo-Fisher, S., Trueb, L., Ayarzagüena, J., Rada, M. & Vilà, C. 2009. Phylogenetic systematics of Glassfrogs (Amphibia: Centrolenidae) and their sister taxon *Allophryne ruthveni*. *Zootaxa* **2100**: 1-97.

Guimarães, L.D., Lima, L.P., Juliano, R.F. & Bastos, R.P. 2001. Vocalizações de espécies de anuros (Amphibia) no Brasil Central. *Boletim do Museu Nacional, Nova Série, Zoologia* **474**: 1-14.

Günther, R. & Richards, S.J. 2000. A new species of the *Litoria gracilentia* group from Irian Jaya (Anura: Hylidae). *Herpetozoa* **13**: 27-43.

Gururaja, K.V. & Ramachandra, T.V. 2006. *Pedostibes tuberculosus* (Malabar tree toad) Advertisement call and distribution. *Herpetological Review* **37**: 75-76.

Haddad, C.F.B., Pombal, J.P. & Batistic, R.F. 1994. Natural hybridization between diploid and tetraploid species of leaf-frogs, genus *Phyllomedusa* (Amphibia). *Journal of Herpetology* **28**: 425-430.

- Haddad, C.F.B. & Pombal Jr, J.P. 1995. A new species of *Hylodes* from Southeastern Brazil (Amphibia: Leptodactylidae). *Herpetologica* **51**: 279-286.
- Haddad, C.F.B., Garcia, P.C.A. & Pombal Jr, J.P. 2003. Redescricao de *Hylodes perplicatus* (Miranda-Ribeiro, 1926) (Amphibia, Anura, Leptodactylidae). *Arquivos do Museu Nacional* **61**: 245-254.
- Halloy, M. & Espinosa, R.E. 2000. Territorial encounters and threat displays in the neotropical frog *Phyllomedusa sauvagii* (Anura: Hylidae). *Herpetological Natural History* **7**: 175-180.
- Hamer, A.J. & Organ, A.K. 2008. Aspects of the ecology and conservation of the growling grass frog *Litoria raniformis* in an urban-fringe environment, Southern Victoria. *Australian Zoologist* **34**: 393-407.
- Hampson, K. & Bennett, D. 2002. Advertisement Calls of Amphibians at Lackunda Estate, Coorg, Karnataka. In: *Frogs of Coorg, Karnataka, India*, Vol. 11 (D. Bennett, ed., pp. 121-135. Viper Press.
- Harper, E.B., Measey, G.J., Patrick, D.A., Menegon, M. & Vonesh, J.R. 2010. *Field guide to the amphibians of the Eastern Arc Mountains and Coastal Forests of Tanzania and Kenya*. Camerapix Publishers International, Nairobi, Kenya.
- Harper, F.A. 1955. A new chorus frog (*Pseudacris*) from the eastern United States. *Natural History Miscellanea Chicago Academic Sciences* **150**: 1-6.
- Hepp, F.S., Luna-Dias, C.D., Gonzaga, L.P. & Carvalho-E-Silva, S.P.D. 2012. Redescription of the advertisement call of *Dendropsophus seniculus* (Cope, 1868) and the consequences for the acoustic traits of the *Dendropsophus marmoratus* species group (Amphibia: Anura: Dendropsophini). *South American Journal of Herpetology* **7**: 165-171.

- Hettzey, A., Török, J. & Hévizi, G. 2005. Male mate choice lacking in the agile frog, *Rana dalmatina*. *Copeia* **2005**: 403-408.
- Heyer, M.M., Heyer, W.R., Spear, S. & De Sá, R.O. 2003. *Leptodactylus mystacinus* (Burmeister) mustached frog. *Catalogue of American Amphibians and Reptiles* **767**: 1-11.
- Heyer, W.R. 1970. Studies on the frogs of the genus *Leptodactylus* (Amphibia: Leptodactylidae). 6. Biosystematics of the *melanonotus* group. *Contributions in Science, Los Angeles County Museum* **191**: 1-48.
- Heyer, W.R. 1971. Mating calls of some frogs from Thailand. *Fieldiana Zoology* **58**: 61-82.
- Heyer, W.R. 1978. Systematics of the *fuscus* group of the frog genus *Leptodactylus* (Amphibia, Leptodactylidae). *Natural History Museum of Los Angeles County* **29**: 1-85.
- Heyer, W.R. 1979. *Systematics of the pentadactylus species group of the frog genus Leptodactylus (Amphibia, Leptodactylidae)*. Smithsonian Institution Press, Washington, D.C.
- Heyer, W.R. 1980. The calls and taxonomic positions of *Hyla giesleri* and *Ololygon opalina* (Amphibia: Anura: Hylidae). *Proceedings of the Biological Society of Washington* **93**: 655-661.
- Heyer, W.R. 1982. Two new species of the frog genus *Hylodes* from Caparaó, Minas Gerais, Brasil (Amphibia: Leptodactylidae). *Proceedings of the Biological Society of Washington* **95**: 377-385.
- Heyer, W.R. 1983. Clarification of the names *Rana mystacea* Spix, 1824, *Leptodactylus amazonicus* Heyer, 1978 and a description of a new species, *Leptodactylus spixi*

- (Amphibia: Leptodactylidae). *Proceedings of the Biological Society of Washington* **96**: 270-272.
- Heyer, W.R. & Cocroft, R.B. 1986. Descriptions of two new species of *Hylodes* from the Atlantic Forests of Brazil (Amphibia: Leptodactylidae). *Proceedings of the Biological Society of Washington* **99**: 100-109.
- Heyer, W.R., Rand, A.S., Cruz, C.A.G., Peixoto, O.L. & Nelson, C.E. 1990. Frogs of Boracéia. *Arquivos de Zoologia* **31**: 231-410.
- Heyer, W.R. 1994. *Variation within the Leptodactylus podicipinus-wagneri complex of frogs (Amphibia: Leptodactylidae)*. Smithsonian Institution Press, Washington, D.C.
- Heyer, W.R., García-Lopez, J.M. & Cardoso, A.J. 1996. Advertisement call variation in the *Leptodactylus mystaceus* species complex (Amphibia: Leptodactylidae) with a description of a new sibling species. *Amphibia-Reptilia* **17**: 7-31.
- Heyer, W.R. 1998. The relationships of *Leptodactylus diedrus* (Anura, Leptodactylidae). *Alytes* **16**: 1-24.
- Heyer, W.R. & de Carvalho, C.M. 2000. Calls and calling behavior of the frog *Leptodactylus natalensis* (Amphibia : Anura : Leptodactylidae). *Proceedings of the Biological Society of Washington* **113**: 284-290.
- Heyer, W.R. 2005. Variation and taxonomic clarification of the large species of the *Leptodactylus pentadactylus* species group (Amphibia: Leptodactylidae) from Middle America, Northern South America, and Amazonia. *Arquivos de Zoologia* **37**: 269-348.
- Heyer, W.R. & Giaretta, A.A. 2009. Advertisement calls, notes on natural history, and distribution of *Leptodactylus chaquensis* (Amphibia: Anura: Leptodactylidae) in Brasil. *Proceedings of the Biological Society of Washington* **122**: 292-305.

- Hödl, W. 1990. Reproductive diversity in Amazonian lowland frogs. *Fortschritte der Zoologie* **38**: 41-60.
- Hoskin, C.J. & Goosem, M.W. 2010. Road impacts on abundance, call traits, and body size of rainforest frogs in Northeast Australia. *Ecology and Society* **15**: 63-70.
- Huang, W.S., Lee, J.K. & Ho, C.H. 2001. Reproductive patterns of two sympatric rhacophorid frogs, *Buergeria japonica* and *B. robusta*, with comments on anuran breeding seasons in Taiwan. *Zoological Science* **18**: 63-70.
- Hutter, C.R. & Guayasamin, J.M. 2012. A new cryptic species of glassfrog (Centrolenidae: *Nymphargus*) from Reserva Las Gralarias, Ecuador. *Zootaxa* **3257**: 1-21.
- Ibáñez, R., Rand, A.S. & Jaramillo, C.A. 1999. *Los anfibios del Monumento Natural Barro Colorado, Parque Nacional Soberanía y áreas adyacentes*. Editorial Mizrachi & Pujol, Panama.
- Inger, R.F. 1954. Systematics and zoogeography of Philippine Amphibia. *Fieldiana Zoology* **39**: 183-531.
- Inger, R.F. 1964. Two new species of frogs from Borneo. *Fieldiana Zoology* **44**: 151-159.
- Inger, R.F. & Bacon Jr, J.P. 1968. Annual reproduction and clutch size in rain forest frogs from Sarawak. *Copeia* **1968**: 602-606.
- Inger, R.F. & Gritis, P.A. 1983. Variation in Bornean frogs of the *Amolops jerboa* species group, with description of two new species. *Fieldiana Zoology* **19**: 1-13.
- Inger, R.F. & Stuebing, R.B. 2005. *A Field Guide to the Frogs of Borneo*, Kota Kinabalu, Borneo.
- Jacobson, S.K. 1985. Reproductive behavior and male mating success in two species of glass frogs (Centrolenidae). *Herpetologica* **41**: 396-404.

- Jehle, R. & Arak, A. 1998. Graded call variation in the Asian cricket frog *Rana nicobariensis*. *Bioacoustics* **9**: 35-48.
- Jiang, J., Xie, F., Fei, L., Ye, C. & Zheng, M. 2002. Mating calls of six forms of pelobatid in Wawu Mountain national forest park, Sichuan, China (Anura: Pelobatidae). *Zoological Research / " Dong wu xue yan jiu" bian ji wei yuan hui bian ji* **23**: 89-94.
- Johnston, G.R. & Richards, S.J. 1994. A new species of *Litoria* (Anura: Hylidae) from New Guinea and a redefinition of *Litoria leucova* (Tyler, 1968). *Memoirs of the Queensland Museum* **37**: 273-279.
- Kadadevaru, G.G., Kanamadi, R.D. & Schneider, H. 2000. Advertisement call of the Indian Bronzed Frog, *Rana temporalis* (Gunther, 1864). *Herpetological Bulletin* **73**: 8-9.
- Kaefer, I.L., Erdtmann, L.K. & Lima, A.P. 2011. The advertisement call of *Physalaemus ephippifer* (Anura: Leiuperidae) from Brazilian Amazonia. *Zootaxa* **2929**: 57-58.
- Kanamadi, R.D., Schneider, H., Hiremath, C.R. & Jirankali, C.S. 1993. Vocalization of the tree frog *Polypedates maculatus* (Rhacophoridae). *Journal of Biosciences* **18**: 239-245.
- Kanamadi, R.D., Hiremath, C.R. & Schneider, H. 1994. Advertisement calls of two anuran amphibians, *Rana tigrina* and *Tomopterna breviceps* *Journal of Biosciences* **10**: 75-80.
- Kanamadi, R.D., Kadadevaru, G.G. & Schneider, H. 2001. Calling behaviour, bioacoustics and distribution of a rhacophorid frog, *Philautus variabilis* (Gunther, 1858). *Amphibia-Reptilia* **22**: 365-372.
- Kasuya, E., Kumaki, T. & Saito, T. 1992. Vocal Repertoire of the Japanese Treefrog, *Rhacophorus arboreus* (Anura: Rhacophoridae). *Zoological Science* **9**: 469-473.
- Khan, M.S. 1997. A new subspecies of common skittering frog *Euphlyctis cyanophlyctis* (Schneider, 1799) from Balochistan, Pakistan. *Pakistan Journal of Zoology* **29**: 107-112.

- Khonsue, W., Matsui, M., Hirai, T. & Misawa, Y. 2001. A comparison of age structures in two populations of a pond frog *Rana nigromaculata* (Amphibia : Anura). *Zoological Science* **18**: 597-603.
- Kime, N.M., Turner, W.R. & Ryan, M.J. 2000. The transmission of advertisement calls in Central American frogs. *Behavioral Ecology* **11**: 71-83.
- Kizirian, D., Coloma, L.A. & Paredes-Recalde, A. 2003. A new treefrog (Hylidae : *Hyla*) from southern Ecuador and a description of its antipredator behavior. *Herpetologica* **59**: 339-349.
- Köhler, J., Reichle, S. & Peters, G. 1997. Advertisement calls of three species of *Bufo* (Amphibia: Anura: Bufonidae) from lowland Bolivia. *Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie)* **562**: 1-8.
- Köhler, J. & Lötters, S. 1999a. Advertisement calls of two Bolivian *Leptodactylus* (Amphibia : Anura : Leptodactylidae). *Amphibia-Reptilia* **20**: 215-219.
- Köhler, J. & Lötters, S. 1999b. Annotated list of amphibian records from the Departamento Pando, Bolivia, with description of some advertisement calls. *Bonner Zoologische Beiträge* **48**: 259-273.
- Köhler, J., Scheelke, K., Schick, S., Veith, M. & Lötters, S. 2005. Contribution to the taxonomy of hyperoliid frogs (Amphibia : Anura : Hyperoliidae): advertisement calls of twelve species from East and Central Africa. *African Zoology* **40**: 127-142.
- Kok, P.J.R. & Castroviejo-Fisher, S. 2008. Glassfrogs (Anura : Centrolenidae) of Kaieteur National Park, Guyana, with notes on the distribution and taxonomy of some species of the family in the Guiana Shield. *Zootaxa* **1680**: 25-53.
- Kolenc, F., Borteiro, C., Baldo, D., Ferraro, D.P. & Prigioni, C. 2009. The tadpoles and advertisement calls of *Pleurodema bibroni* Tschudi and *Pleurodema kriegi* (Muller),

- with notes on their geographic distribution and conservation status (Amphibia, Anura, Leiuperidae). *Zootaxa* **1969**: 1-35.
- Kraus, F. & Allison, A. 2004. Two new treefrogs from Normanby Island, Papua New Guinea. *Journal of Herpetology* **38**: 197-207.
- Kraus, F. 2012. Identity of *Nyctimystes cheesmani* (Anura: Hylidae), with description of two new related species. *Zootaxa* **3493**: 1-26.
- Krishna, S.N. & Krishna, S.B. 2005. Female courtship calls of the litter frog (*Rana curtipes*) in the tropical forests of Western Ghats, South India. *Amphibia-Reptilia* **26**: 431-435.
- Kueh, B.H., Ismail, N., Lau, C.E.S., Albert, J., Siwan, E.S. & Ngidang, V.B.A. 2010. *Rana Luctuosa* (Mahogany Frog). Altitude and maximum size. *Herpetological Review* **41**: 341-342.
- Kuramoto, M. 1980. Mating calls of treefrogs (genus *Hyla*) in the far east, with description of a new species from Korea. *Copeia* **1980**: 100-108.
- Kuramoto, M. 1986. Call structures of the rhacophorid frogs from Taiwan. *Scientific Report of the Laboratory for Amphibian Biology, Hiroshima University* **8**: 45-68.
- Kuramoto, M. & Wang, C.S. 1987. A new rhacophorid treefrog from Taiwan, with comparisons to *Chirixalus eiffingeri* (Anura, Rhacophoridae). *Copeia* **1987**: 931-942.
- Kuramoto, M. & Joshy, S.H. 2001. Advertisement call structures of frogs from Southwestern India, with some ecological and taxonomic notes. *Current Herpetology* **20**: 85-95.
- Kuramoto, M., Joshy, S.H., Kurabayashi, A. & Sumida, M. 2007. The genus *Fejervarya* (Anura: Ranidae) in central Western Ghats, India, with descriptions of four new cryptic species. *Current Herpetology* **26**: 81-105.

- Kurniati, H., Sumadijaya, A., Boonman, A. & Laksono, W.T. 2010. Ecology, distribution and bio-acoustic of amphibians in degraded habitat, Indonesian Institute of Science. Cibinong.
- Kuzmin, S.L. & Ischenko, V.G. 1997. Skeletochronology of *Bufo raddei* from the Gobi Desert. *Journal of Herpetology* **31**: 306-309.
- Kwet, A. & Faivovich, J. 2001. *Proceratophrys bigibbosa* species group (Anura : Leptodactylidae), with description of a new species. *Copeia* **2001**: 203-215.
- Kwet, A., Solé, M., Miranda, T., Melchior, J., Naya, D. & Maneyro, R. 2002. First record of *Hyla albopunctata* Spix, 1824 (Anura: Hylidae) in Uruguay, with comments on the advertisement call. *Boletín de la Asociación Herpetológica Española* **13**: 15-19.
- Kwet, A. & Baldo, D. 2003. Advertisement call of the leptodactylid frog *Proceratophrys avelinoi*. *Amphibia-Reptilia* **24**: 104-107.
- Langone, J.A. 1990. Revalidación de *Hyla uruguayana* Schmidt, 1944 (Amphibia, Anura, Hylidae). *Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo* **12**: 1-9.
- Lannoo, M. 2005. *Amphibian declines: the conservation status of United States species*. University California Press, Berkeley.
- Largen, M.J., Tandy, M. & Tandy, J. 1978. A new species of toad from the Rift Valley of Ethiopia, with observations on the other species of *Bufo* (Amphibia Anura Bufonidae) recorded from this country. *Italian Journal of Zoology* **1**: 1-41.
- Lascano, J.N. 2011. Description of the advertisement and distress call of *Chacophrys pierottii* and comments on the advertisement call of *Lepidobatrachus llanensis* (Anura: Ceratophryidae). *Journal of Natural History* **45**: 2929–2938.

- Lavilla, E.O., Vaira, M. & Ferrari, L. 2003. A new species of *Elachistocleis* (Anura : Microhylidae) from the Andean Yungas of Argentina, with comments on the *Elachistocleis ovalis* - *E. bicolor* controversy. *Amphibia-Reptilia* **24**: 269-284.
- Leary, C.J. 2001. Evidence of convergent character displacement in release vocalizations of *Bufo fowleri* and *Bufo terrestris* (Anura; Bufonidae). *Animal Behaviour* **61**: 431-438.
- Leclair, R. & Laurin, G. 1996. Growth and body size in populations of mink frogs *Rana septentrionalis* from two latitudes. *Ecography* **19**: 296-304.
- Lee, J.C. 2000. *A field guide to the amphibians and reptiles of the Maya world: the lowlands of Mexico, Northern Guatemala, and Belize*. Cornell University Press, Ithaca.
- Leenders, T. 2001. *A guide to amphibians and reptiles of Costa Rica* Zona Tropical, Miami.
- Lemmon, E.M., Lemmon, A.R., Collins, J.T. & Cannatella, D.C. 2008. A new North American chorus frog species (Amphibia : Hylidae : *Pseudacris*) from the South-central United States. *Zootaxa* **1675**: 1-30.
- León, J.R. 1969. The systematics of the frogs of the *Hyla rubra* group in Middle America. *University of Kansas Publications of the Museum of Natural History* **18**: 505-545.
- Leong, T.M., Matsui, M., Yong, H.S. & Hamid, A.A. 2003. Revalidation of *Rana laterimaculata* Barbour et Noble, 1916 from the synonymy of *Rana baramica* Boettger, 1901. *Current Herpetology* **22**: 17-27.
- Lesbarrères, D. & Lodé, T. 2002. Variations in male calls and responses to an unfamiliar advertisement call in a territorial breeding anuran, *Rana dalmatina*: evidence for a "dear enemy" effect. *Ethology Ecology & Evolution* **14**: 287-295.
- Lescure, J. & Marty, C. 2000. *Atlas des amphibiens de Guyane*. Publications Scientifiques du M.N.H.N., Paris.

- Liao, W.B. & Lu, X. 2011. Male mating success in the Omei treefrog (*Rhacophorus omeimontis*): the influence of body size and age. *Belgian Journal of Zoology* **141**: 3-10.
- Lima, A.P., Magnusson, W.E., Menin, M., Erdtmann, L.K., Rodrigues, D.J., Keller, C. & Hödl, W. 2006. *Guia de Sapos da Reserva Adolpho Ducke: Amazônia Central* Áttema Design Editorial, Manaus.
- Lima, L.P., Bastos, R.P. & Giaretta, A.A. 2004. A new *Scinax* Wagler, 1830 of the *S. rostratus* group from Central Brazil (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional* **62**: 505-512.
- Lingnau, R. & Bastos, R.P. 2003. Vocalizações de duas espécies de anuros do sul do Brasil (AMPHIBIA: HYLIDAE). *Arquivos do Museu Nacional* **61**: 203-207.
- Lingnau, R. 2009. Distribuição temporal, atividade reprodutiva e vocalizações em uma assembleia de anfíbios anuros de uma Floresta Ombrófila Mista em Santa Catarina, sul do Brasil, Pontifícia Universidade Católica do Rio Grande do Sul. **PhD**: 94.
- Lingnau, R., Zank, C., Colombo, P. & Kwet, A. 2013. Vocalization of *Hylodes meridionalis* (Mertens 1927) (Anura, Hylodidae) in Rio Grande do Sul, Brazil, with comments on nocturnal calling in the family Hylodidae. *Studies on Neotropical Fauna and Environment* **48**: 76-80.
- Lisboa, E.B.F., Moura, G.J.B., Melo, I.V.C., Andrade, E.V.E. & Figuerêdo Jr., J.M. 2011. Aspectos ecológicos de *Hypsiboas semilineatus* (SPIX, 1824) (AMPHIBIA, ANURA, HYLIDAE) em fragmento de Mata Atlântica, nordeste do Brasil. *Revista Ibero-Americana de Ciências Ambientais, Aquidabã* **2**: 21-30.
- Littlejohn, M.J. & Martin, A.A. 1965. A new species of *Crinia* (Anura: Leptodactylidae) from South Australia. *Copeia* **1965**: 319-324.

- Littlejohn, M.J. & Harrison, P.A. 1985. The functional significance of the diphasic advertisement call of *Geocrinia victoriana* (Anura: Leptodactylidae). *Behavioral Ecology and Sociobiology* **16**: 363-373.
- Liu, C.C. 1950. *Amphibians of Western China*. Chicago Natural History Museum, Chicago.
- Liu, W.Z., Yang, D.T., Ferraris, C. & Matsui, M. 2000. *Amolops bellulus*: A new species of stream-breeding frog from western Yunnan, China (Anura : Ranidae). *Copeia* **2000**: 536-541.
- Lizana, M., Marquez, R. & Martinsanchez, R. 1994. Reproductive biology of *Pelobates cultripes* (Anura: Pelobatidae) in central Spain. *Journal of Herpetology* **28**: 19-27.
- Lo Valvo, M. & Giacalone, G. 2013. Biometrical analyses of a Sicilian green toad, *Bufo siculus* (Stock et al. 2008), population living in Sicily (Italy). *International Journal of Morphology* **31**: 681-686.
- Loftus-Hills, J.J. & Littlejohn, M.J. 1992. Reinforcement and reproductive character displacement in *Gastrophryne carolinensis* and *G. olivacea* (Anura: Microhylidae): a reexamination. *Evolution* **46**: 896-906.
- Lötters, S., Glaw, F., Reichle, S., Köhler, J. & Meyer, E. 1999. Notes on vocalizations in three species of *Atelopus* from Central and South America (Amphibia: Anura). *Herpetozoa* **12**: 79-83.
- Lötters, S., Haas, W., Schick, S. & Böhme, W. 2002. On the systematics of the harlequin frogs (Amphibia: Bufonidae: *Atelopus*) from Amazonia. II: Redescription of *Atelopus pulcher* (Boulenger, 1882) from the eastern Andean versant in Peru. *Salamandra* **38**: 164-184.
- Lue, K.Y. 1990. *The amphibians and reptiles of Taiwan* Society for Wildlife and Nature, Taipei, Taiwan.

- Lutz, B. 1973. *Brazilian species of Hyla*. University of Texas Press, Austin.
- Lynch, J.D. & Freeman, H.L. 1966. Systematic status of a South American frog, *Allophryne ruthveni* Gaige. *University of Kansas Publications of the Museum of Natural History* **17**: 493-502.
- Lynch, J.D. 1970. Systematic status of the American leptodactylid frog genera *Engystomops*, *Eupemphix*, and *Physalaemus*. *Copeia* **1970**: 488-406.
- Lynch, J.D. & Duellman, W.E. 1973. A review of the Centrolenid frogs of Ecuador, with descriptions of new species. *Occasional Papers of the Museum of Natural History, The University of Kansas* **16**: 1-66.
- Lynch, J.D., Ruiz, P.M. & Rueda, J.V. 1983. Notes on the distribution and reproductive biology of *Centrolene geckoideum* Jimenez de la Espada in Colombia and Ecuador (Amphibia: Centrolenidae). *Studies on Neotropical Fauna and Environment* **18**: 239-243.
- Maeda, N. & Matsui, M. 1990. *Frogs and Toads of Japan*. Bun-Ichi Sogo Shuppan, Tokyo.
- Maldonado, A.L.S. 2012. Cuidado parental y selección sexual en *Centrolene peristictum* (Anura: Centrolenidae) Pontificia Universidad Católica del Ecuador. **Master**.
- Malkmus, R., Manthey, U., Vogel, G., Hoffmann, P. & Kosuch, J. 2002. *Amphibians & reptiles of Mount Kinabalu (North Borneo)*. Koeltz Scientific Books, Berlin.
- Malkmus, R. 2006. *Leptobrachium gunungensis* sp. n. (Anura: Pelobatidae) vom Mount Kinabalu, Nord-Borneo. *Mitteilungen aus dem Museum für Naturkunde in Berlin. Zoologisches Museum und Institut für Spezielle Zoologie (Berlin)* **72**: 271-301.
- Mângia, S., Santana, D.J. & Feio, R.N. 2010. Advertisement call of the cycloramphid toad *Proceratophrys melanopogon* (Miranda-Ribeiro, 1926). *South American Journal of Herpetology* **5**: 127-131.

- Manthey, U. & Grossmann, W. 1997. *Amphibien und Reptilien Südasiens*. Natur und Tier Verlag, Münster, Germany.
- Marcelino, V.R., Haddad, C.F.B. & Alexandrino, J. 2009. Geographic distribution and morphological variation of striped and nonstriped populations of the Brazilian Atlantic Forest treefrog *Hypsiboas bischoffi* (Anura: Hylidae). *Journal of Herpetology* **43**: 351-361.
- Marquez, R., De la Riva, I. & Bosch, J. 1995. Advertisement calls of Bolivian Leptodactylidae (Amphibia, Anura). *Journal of Zoology* **237**: 313-336.
- Márquez, R., Delariva, I. & Bosch, J. 1993. Advertisement calls of Bolivian species of *Hyla* (Amphibia, Anura, Hylidae). *Biotropica* **25**: 426-443.
- Márquez, R. 1995. Female choice in the midwife toads (*Alytes obstetricans* and *A. cisternasii*). *Behaviour* **132**: 151-161.
- Márquez, R. & Eekhout, X.R. 2006. Advertisement calls of six species of anurans from Bali, Republic of Indonesia. *Journal of Natural History* **40**: 571-588.
- Martino, A.L. & Sinsch, U. 2002. Speciation by polyploidy in *Odontophrynus americanus*. *Journal of Zoology* **257**: 67-81.
- Martins, I.A. & Jim, J. 2003. Bioacoustic analysis of advertisement call in *Hyla nana* and *Hyla sanborni* (Anura, Hylidae) in Botucatu, São Paulo, Brazil região de Botucatu, Estado de São Paulo, Brasil. *Brazilian Journal of Biology* **63**: 507-516.
- Martins, L.B. & Giaretta, A.A. 2013. Morphological and acoustic characterization of *Proceratophrys goyana* (Lissamphibia: Anura: Odontophrynidae), with the description of a sympatric and related new species. *Zootaxa* **3750**: 301-320.
- Martins, M. 1998. *The frogs of the Ilha de Maracá. Maraca: The Biodiversity and Environment of an Amazonian Rainforest*. John Wiley & Sons Ltd., New York.

Matsui, M. 1982a. Call characteristics of several anuran species from East Kalimantan.

Contributions from the Biological Laboratory, Kyoto University **26**: 131-139.

Matsui, M. 1982b. Amphibians from Sabah II. Acoustic characteristics of three common

anuran species. *Contributions from the Biological Laboratory, Kyoto University* **26**: 123-129.

Matsui, M. & Utsunomiya, T. 1983. Mating call characteristics of the frogs of the subgenus

Babina with reference to their relationship with *Rana adenopleura*. *Journal of Herpetology* **17**: 32-37.

Matsui, M. 1986. Three new species of *Amolops* from Borneo (Amphibia, Anura, Ranidae).

Copeia **1986**: 623-630.

Matsui, M., Wu, G.F. & Yong, H.S. 1993. Acoustitic characteristics of three species of the

genus *Amolops* (Amphibia, Anura, Ranidae). *Zoological Science* **10**: 691-695.

Matsui, M. 1994. A taxonomic study of the *Rana narina* complex, with description of three

new species (Amphibia: Ranidae). *Zoological Journal of the Linnean Society* **111**: 385-415.

Matsui, M. & Wu, G.F. 1994. Acoustic characteristics of treefrogs from Sichuan, China, with

comments on systematic relationship of *Polypedates* and *Rhacophorus* (Anura, Rhacophoridae). *Zoological Science* **11**: 485-490.

Matsui, M. 1995. Calls produced by a "voiceless" frog, *Rana blythi* Boulenger 1920, from

Peninsular Malaysia (Amphibia Anura). *Tropical Zoology* **8**: 325-331.

Matsui, M. 1997. Call characteristics of Malaysian *Leptolalax* with a description of two new

species (Anura: Pelobatidae). *Copeia* **1997**: 158-165.

Matsui, M. & Nabhitabhata, J. 2006. A new species of *Amolops* from Thailand (Amphibia,

Anura, Ranidae). *Zoological Science* **23**: 727-732.

- Matsui, M., Mohamed, M., Shimada, T. & Sudin, A. 2007. Resurrection of *Staurois parvus* from *S. tuberilinguis* from Borneo (Amphibia, Ranidae). *Zoological Science* **24**: 101-106.
- Mendelson III, J.R., Williams, B.L., Sheil, C.A. & Mulcahy, D.G. 2005. Systematics of the *Bufo coccifer* complex (Anura: Bufonidae) of Mesoamerica. *Scientific Papers Natural History Museum, The University of Kansas* **38**: 1-27.
- Menzies, J.I. & Zweifel, R.G. 1974. Systematics of *Litoria arfakiana* of New Guinea and sibling species (Salientia, Hylidae). *American Museum Novitates* **2558**: 1-16.
- Minton, S.A. 1966. A contribution to the herpetology of West Pakistan. *Bulletin of the American Museum of Natural History* **134**: 27-184.
- Mu, Y. & Zhao, E. 1990. Mating Call Structures of the Chinese Frog, *Rana nigromacula* (Amphibia, Anura, Ranidae). *Asiatic Herpetological Research* **3**: 60-63.
- Myers, C.W. & Donnelly, M.A. 1997. A tepui herpetofauna on a granitic mountain (Tamacuari) in the borderland between Venezuela and Brazil: report from the Phipps Tapirapécó Expedition. *American Museum Novitates* **3213**: 1-71.
- Myers, C.W. & Donnelly, M.A. 2001. Herpetofauna of the Yutaje-Corocoro massif, Venezuela: Second report from The Robert G. Goelet American Museum-terramar expedition to the Northwestern Tepuis. *Bulletin of the American Museum of Natural History* **261**: 1-85.
- Napoli, M.F. & Caramaschi, U. 1999. Geographic variation of *Hyla rubicundula* and *Hyla anataliasiasi*, with the description of a new species (Anura, Hylidae). *Alytes* **16**: 165-189.

Nelson, C. 1963. *Gastrophyrne elegans* (Boulenger). Elegant Narrow-mouthed Toad.

Catalogue of American Amphibians and Reptiles, American Society of Ichthyologists and Herpetologists.

Nelson, C.E. 1973. Mating calls of the Microhylinae: descriptions and phylogenetic and ecological considerations. *Herpetologica* **29**: 163-176.

Nguyen, T.Q., Phung, T.M., Le, M.D., Ziegler, T. & Böhme, W. 2013. First record of the genus *Oreolalax* (Anura: Megophryidae) from Vietnam with description of a new species. *Copeia* **2013**: 213-222.

Noonan, B.P. & Bonett, R.M. 2003. A new species of *Hyalinobatrachium* (Anura : Centrolenidae) from the highlands of Guyana. *Journal of Herpetology* **37**: 91-97.

Norman, D.R. 1994. *Anfibios y reptiles del Chaco Paraguayo*. Private printing, San José, Costa Rica.

Nunes, I. & Juncá, F.A. 2006. Advertisement calls of three Leptodactylid frogs in the state of Bahia, northeastern Brazil (Amphibia, Anura, Leptodactylidae), with considerations on their taxonomic status. *Arquivos do Museu Nacional* **64**: 151-157.

Nunes, I., Santiago, R.S. & Juncá, F.A. 2007. Advertisement calls of four Hylid frogs from the state of Bahia, Northeastern Brazil (Amphibia, Anura, Hylidae). *South American Journal of Herpetology* **2**: 89-96.

Odendaal, F.J., Bull, C.M. & Telford, S.R. 1983. The vocabulary of calls of *Ranidella riparia* (Anura: Leptodactylidae). *Copeia* **1983**: 534-537.

Ohler, A., Marquis, O., Swan, S. & Grosjean, S. 2000. Amphibian biodiversity of Hoang Lien Nature Reserve (Lao Cai Province, northern Vietnam) with description of two new species. *Herpetozoa* **13**: 71-87.

- Ohler, A. & Delorme, M. 2006. Well known does not mean well studied: Morphological and molecular support for existence of sibling species in the Javanese gliding frog *Rhacophorus reinwardtii* (Amphibia, Anura). *Comptes Rendus Biologies* **329**: 86-97.
- Ohler, A., Wollenberg, K.C., Grosjean, S., Hendrix, R., Vences, M., Ziegler, T. & Dubois, A. 2011. Sorting out Lalos: description of new species and additional taxonomic data on megophryid frogs from Northern Indochina (genus *Leptolalax*, Megophryidae, Anura). *Zootaxa* **3147**: 1-83.
- Oldham, R.S. & Gerhardt, H.C. 1975. Behavioral isolating mechanisms of the treefrogs *Hyla cinerea* and *H. gratiosa*. *Copeia* **1975**: 223-231.
- Oliveira, M.E., Paillette, M., Rosa, H.D. & Crespo, E.G. 1991. A natural hybrid between *Hyla arborea* and *Hyla meridionalis* detected by mating calls. *Amphibia-Reptilia* **12**: 15-20.
- Ortega-Andrade, H.M. 2008. *Agalychnis spurrelli* Boulenger (Anura, Hylidae): variación, distribución y sinonimia. *Papeís Avulsos de Zoologia (São Paulo)* **48**: 103-117.
- Ortega-Andrade, H.M., Rojas-Soto, O. & Paucar, C. 2013. Novel data on the ecology of *Cochranella mache* (Anura: Centrolenidae) and the importance of protected areas for this critically endangered glassfrog in the Neotropics. *Plos One* **8**: e8137.
- Owen, P.C. & Tucker, J.K. 2006. Courtship calls and behavior in two species of chorus frogs, genus *Pseudacris* (Anura : Hylidae). *Copeia* **2006**: 137-144.
- Palma, C.C.C. 2013. Analisis taxonomico de *Alsodes nodosus* (Dumeril & Bibron, 1841) (amphibia, neobatrachia): antecedentes morfologicos y moleculares Universidad Austral do Chile. Chile. **PhD**: 126.
- Parker, H.W. 1934. *A monograph of the frogs of the family Microhylidae*. Trustees of the British Museum, London.

- Parker, H.W. 1940. XXII.—Undescribed anatomical structures and new species of reptiles and amphibians. *The Annals and Magazine of Natural History* **5**: 257-274.
- Passmore, N.I. & Carruthers, V.C. 1975. A new species of *Tomopterna* (Anura: Ranidae) from the Kruger National Park, with notes on related species. *Koedoe-African Protected Area Conservation and Science* **18**: 31-50.
- Pavan, D., Narvaes, P. & Rodrigues, M.T. 2001. A new species of leptodactylid frog from the Atlantic Forests of South Eastern Brazil with notes on the status and on the speciation of the *Hylodes* species groups. *Papéis Avulsos de Zoologia (São Paulo)* **41**: 407-425.
- Peloso, P.L.V. & Sturaro, M.J. 2008. A new species of narrow-mouthed frog of the genus *Chiasmocleis* Mehely 1904 (Anura, Microhylidae) from the Amazonian rainforest of Brazil. *Zootaxa* **1947**: 39-52.
- Penna, M., Contreras, S. & Veloso, A. 1983. Acoustical repertoires and morphological differences in the ear of two *Alsodes* species (Amphibia: Leptodactylidae). *Canadian Journal of Zoology* **61**: 2369-2376.
- Penna, M. & Veloso, A. 1990. Vocal diversity in frogs of the South American temperate forest. *Journal of Herpetology* **24**: 23-33.
- Penna, M. & Solis, R. 1998. Frog call intensities and sound propagation in the South American temperate forest region. *Behavioral Ecology and Sociobiology* **42**: 371-381.
- Penna, M. & Meier, A. 2011. Vocal strategies in confronting interfering sounds by a frog from the Southern Temperate Forest, *Batrachyla antartandica*. *Ethology* **117**: 1147-1157.
- Pereyra, M.O., Borteiro, C., Baldo, D., Kolenc, F. & Conte, C.E. 2012. Advertisement call of the closely related species *Scinax aromothyella* Faivovich 2005 and *S. berthae* (Barrio

- 1962), with comments on the complex calls in the *S. catharinae* group. *Herpetological Journal* **22**: 133-137.
- Perret, J.L. 1996. Une nouvelle espèce du genre *Ptychadena* (Anura, Ranidae) du Kenya. *Revue Suisse de Zoologie* **103**: 757-766.
- Peters, W.C.H. 1869. Über neue Saurier (*Chaunolaemus multicaudatus*, *Tropidolepisma Richardi* und *Gymnodactylus steudneri*) und Batrachier (*Cyclorhynchus fasciatus* und *Hyla gracilentia*). *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* **1869**: 786-790.
- Peters, W.C.H. & Doria, G. 1878. Catalogo dei rettili e dei batraci raccolti da O. Beccari, L.M. D'Albertis e A.A. Brujin nella sotto-regione Austro-Malese. *Annali del Museo Civico di Storia Naturale di Genova* **13**: 323-450.
- Pfennig, K.S. 2000. Female spadefoot toads compromise on mate quality to ensure conspecific matings. *Behavioral Ecology* **11**: 220-227.
- Pfennig, K.S. & Pfennig, D.W. 2005. Character displacement as the "best of a bad situation": Fitness trade-offs resulting from selection to minimize resource and mate competition. *Evolution* **59**: 2200-2208.
- Pickersgill, M. 2005. The taxonomy and ethology of the *Afraxalus stuhlmanni* complex (Anura: Hyperoliidae). *Steenstrupia* **29**: 1-38.
- Pimenta, B.V.S., Nunes, I. & Cruz, C.A.G. 2007. Notes on the poorly known phyllomedusine frog *Hylomantis aspera* Peters, 1872 (Anura, Hylidae). *South American Journal of Herpetology* **2**: 206-214.
- Pombal, J.P. & Haddad, C.F.B. 1999. Frogs of the genus *Paratelmatobius* (Anura : Leptodactylidae) with descriptions of two new species. *Copeia* **1999**: 1014-1026.

- Pombal Jr., J.P. & Haddad, C.F.B. 1992. Espécies de *Phyllomedusa* do grupo *burmeisteri* do Brasil oriental, com descrição de uma espécie nova (Amphibia, Hylidae). *Revista Brasileira de Biologia* **52**: 217-229.
- Pombal Jr., J.P. 2010. O espaço acústico em uma taxocenose de anuros (Amphibia) do sudoeste do Brasil. *Arquivos do Museu Nacional* **68**: 135-144.
- Pombal Jr., P., Feio, R.N. & Haddad, C.F.B. 2002. A new species of torrent frog genus *Hylodes* (Anura : Leptodactylidae) from Southeastern Brazil. *Herpetologica* **58**: 462-471.
- Porter, K.R. 1966. Mating calls of six Mexican and Central American toads (genus *Bufo*). *Herpetologica* **22**: 60-67.
- Prado, G.M., Borgo, J.H., Abrunhosa, P.A. & Wogel, H. 2003. Comportamento reprodutivo, vocalização e redescrição do girino de *Phrynohyas mesophaea* (Hensel, 1867) do sudeste do Brasil (Amphibia, Anura, Hylidae). *Boletim do Museu Nacional do Rio de Janeiro* **510**: 1-11.
- Prado, G.M. & Pombal Jr, J.P. 2008. Espécies de *Proceratophrys* Miranda-Ribeiro, 1920 com apêndices palpebrais (Anura; Cycloramphidae). *Arquivos de Zoologia* **39**: 1-85.
- Prakash, S. 1988. Genetic studies on *Rana limnocharis*, North Eastern Hill University. Shillong, India.
- Preininger, D., Böckle, M. & Hödl, W. 2007. Comparison of anuran acoustic communities of two habitat types in the Danum Valley Conservation Area, Sabah, Malaysia. *Salamandra* **32**: 129-138.
- Provete, D.B., Garey, M.V., Toledo, L.F., Nascimento, J., Lourenco, L.B., Rossa-Feres, D.d.C. & Haddad, C.F.B. 2012. Redescription of *Physalaemus barrioi* (Anura: Leiuperidae). *Copeia* **2012**: 507-518.

- Pyburn, W.F. 1977. A new hylid frog (Amphibia, Anura, Hylidae) from the Vaupes River of Colombia with comments on related species. *Journal of Herpetology* **11**: 405-410.
- Richards, S.J. & Oliver, P.M. 2006. A new species of torrent-dwelling *Litoria* (Anura: Hylidae) from the Kikori Integrated Conservation and Development Project area, Papua New Guinea. *Salamandra* **42**: 231-238.
- Roberts, J.D., Mahony, M., Kendrick, P. & Majors, C.M. 1991. A new species of burrowing frog, *Neobatrachus* (Anura: Myobatrachidae), from the Eastern wheatbelt of Western Australia. *Records of the Western Australian Museum* **15**: 23-32.
- Roberts, J.D. 1997a. Geographic variation in calls of males and determination of species boundaries in tetraploid frogs of the Australian genus *Neobatrachus* (Myobatrachidae). *Australian Journal of Zoology* **45**: 95-112.
- Roberts, J.D. 1997b. Call evolution in *Neobatrachus* (Anura : Myobatrachidae): Speculations on tetraploid origins. *Copeia* **1997**: 791-801.
- Robertson, J.G.M. 1986. Female choice, male strategies and the role of vocalizations in the Australian frog *Uperoleia rugosa*. *Animal Behaviour* **34**: 773-784.
- Rödel, M.O. 2000. *Herpetofauna of West Africa, Vol. I. Amphibians of the West African Savanna*. Edition Chimaira, Frankfurt, Germany.
- Rodrigues, D.D.J., Uetanabaro, M. & Lopes, F.S. 2004. Reproductive strategies of *Physalaemus nattereri* (Steindachner, 1863) and *P. albonotatus* (Steindachner, 1864) at Serra da Bodoquena, State of Mato Grosso do Sul, Brazil. *Revista Española de Herpetología* **18**: 63-73.
- Rodrigues, D.J., Uetanabaro, M. & Lopes, F.S. 2007. Breeding biology of *Phyllomedusa azurea* Cope, 1862 and *P. sauvagii* Boulenger, 1882 (Anura) from the Cerrado, Central Brazil. *Journal of Natural History* **41**: 1841-1851.

- Rodrigues, D.J., Menin, M., Lima, A.P. & Mokross, K.S. 2008. Tadpole and vocalizations of *Chiasmocleis hudsoni* (Anura, Microhylidae) in Central Amazonia, Brazil. *Zootaxa* **1680**: 55-58.
- Rodrigues, D.J., Noronha, J.D.C., Lima, M.M. & Rosa, A.C. 2011. Amphibia, Anura, Hylidae, *Phyllomedusa boliviana* Boulenger, 1902 and *Phyllomedusa camba* De la Riva, 2000: Distribution extension in central Brazil. *Check List* **7**: 397-399.
- Rodrigues, V.H. 2009. Aspectos da transmissão e processamento de informação em canto de anuros, Universidade de São Paulo. **Master**: 100.
- Rodríguez, L.O. & Duellman, W.E. 1994. *Guide to the frogs of the Iquitos region, Amazonian Peru*. The University of Kansas Natural History Museum Special Publication.
- Roelke, C.E., Greenbaum, E., Kusamba, C., Aristote, M.M. & Smith, E.N. 2011. Systematics and conservation status of two distinct albertine Rift Treefrogs, *Leptopelis karissimbensis* and *Leptopelis kivuensis* (Anura: Arthroleptidae). *Journal of Herpetology* **45**: 343-351.
- Rojas-Runjaic, F.J., Infante-Rivero, E.E. & Cabello, P. 2012. New records and distribution extensions of centrolenid frogs for Venezuela. *Check List* **8**: 819-825.
- Ron, S.R., Cannatella, D.C. & Coloma, L.A. 2004. Two new species of *Physalaemus* (Anura : Leptodactylidae) from Western Ecuador. *Herpetologica* **60**: 261-275.
- Ron, S.R., Coloma, L.A. & Cannatella, D.C. 2005. A new, cryptic species of *Physalaemus* (Anura : Leptodactylidae) from western Ecuador with comments on the call structure of the *P. pustulosus* species group. *Herpetologica* **61**: 178-198.
- Rosset, S.D., Ferraro, D.P., Alcalde, L. & Basso, N.G. 2007. A revision of *Odontophrynus barrioi* (Anura: Neobatrachia): morphology, osteology, vocalizations, and geographic distribution. *South American Journal of Herpetology* **2**: 97-106.

- Roy, D. & Elepfandt, A. 1993. Bioacoustic analysis of frog calls from northeast India. *Journal of Biosciences* **18**: 381-393.
- Roy, D., Borah, B. & Sarma, A. 1995. Analysis and significance of female reciprocal call in frogs. *Current Science* **69**: 265-270.
- Ruiz-Carranza, P.M. & Lynch, J.D. 1991. Ranas Centrolenidae de Colombia III: Nuevas especies de *Cochranella* del grupo *granulosa*. *Lozania* **59**: 1-18.
- Ryan, M.J. 1983. Sexual selection and communication in a Neotropical frog, *Physalaemus pustulosus*. *Evolution* **37**: 261-272.
- Ryan, M.J. & Drewes, R.C. 1990. Vocal morphology of the *Physalaemus pustulosus* species group (Leptodactylidae): morphological response to sexual selection for complex calls. *Biological Journal of the Linnean Society* **40**: 37-52.
- Salas, N.E. & Di Tada, I.E. 1988. Análisis bioacústico del canto nupcial de poblaciones de *Odontophrynus achalensis* y *O. occidentalis* (Anura: Leptodactylidae) en la Provincia de Córdoba. *Boletín de La Asociación Herpetológica Argentina* **4**: 1.
- Salas, N.E., Zavattieri, M.V., Di Tada, I.E., Martino, A.L. & Bridarolli, M.E. 1998. Bioacoustical and etho-ecological features in amphibian communities of southern Cordoba Province (Argentina). *Cuadernos de Herpetologia* **12**: 37-46.
- Santana, D.J., São-Pedro, V.d.A., Bernarde, P.S. & Feio, R.N. 2010. Descrição do canto de anúncio e dimorfismo sexual em *Proceratophrys concavitympanum* Giaretta, Bernarde & Kokubum, 2000. *Papéis Avulsos de Zoologia (São Paulo)* **50**: 167-174.
- São-Pedro, V.A., Medeiros, P.H. & Garda, A.A. 2011. The advertisement call of *Rhinella granulosa* (Anura, Bufonidae). *Zootaxa* **3092**: 60-62.
- Savage, J.M. 1972. The harlequin frogs, genus *Atelopus*, of Costa Rica and Western Panama. *Herpetologica* **28**: 77-94.

- Savage, J.M. 2002. *The amphibians and reptiles of Costa Rica: a herpetofauna between two continents, between two seas*. University of Chicago Press.
- Schawrtz, A. 1957. Chorus frogs (*Pseudacris nigrita* LeConte) in South Carolina. *American Museum Novitates* **1838**: 1-12.
- Schawrtz, A. 1959. A new species of toad, *Bufo cataulaciceps*, from the Isla de Pinos and western Cuba. *Proceedings of the Biological Society of Washington* **72**: 109-120.
- Schawrtz, A. 1972. The native toads (Anura, Bufonidae) of Hispaniola. *Journal of Herpetology* **6**: 217-231.
- Schiøtz, A. 1999. Treefrogs of Africa *Ed. Chimaira*.
- Schneider, H. 1974. Structure of the mating calls and relationships of the European tree frogs (Hylidae, Anura). *Oecologia* **14**: 99-110.
- Señaris, J.C. & Ayarzagüena, J. 2001. Una nueva especie de rana de cristal del género *Hyalinobatrachium* (Anura: Centrolenidae) del Delta del Río Orinoco, Venezuela. *Revista de Biología Tropical* **49**: 1083-1093.
- Señaris, J.C. & Ayarzagüena, J. 2005. *Revisión taxonómica de la Familia Centrolenidae (Amphibia; Anura) de Venezuela*. Publicaciones del Comité Español del Programa Hombre y Biosfera – Red IberoMaB de la UNESCO, Sevilla.
- Shen, J.-X., Xu, Z.-M., Feng, A.S. & Narins, P.M. 2011. Large odorous frogs (*Odorrana graminea*) produce ultrasonic calls. *Journal of Comparative Physiology A-Neuroethology Sensory Neural and Behavioral Physiology* **197**: 1027-1030.
- Sheridan, J.A., Bickford, D. & Su, K.F.-Y. 2010. An examination of call and genetic variation in three wide-ranging Southeast Asian anuran species. *Raffles Bulletin of Zoology* **58**: 369-379.

- Silva, I.S.N. & Juncá, F.A. 2006. Evidence of full species status of the neotropical leaf-frog *Phyllomedusa burmeisteri bahiana* (A. Lutz, 1925) (Amphibia, Anura, Hylidae). *Zootaxa* **1113**: 51-64.
- Silva, R.A., Martins, I.A. & Rossa-Feres, D.d.C. 2008. Bioacústica e sítio de vocalização em taxocenoses de anuros de área aberta no noroeste paulista. *Biota Neotropica* **8**: 123-134.
- Sinsch, U., Luemkemann, K., Rosar, K., Schwarz, C. & Dehling, J.M. 2012. Acoustic niche partitioning in an anuran community inhabiting an Afromontane wetland (Butare, Rwanda). *African Zoology* **47**: 60-73.
- Smith, M. 1927. Contributions to the Herpetology of the Indo-Australian Region. *Proceedings of the Zoological Society of London* **97**: 199-226.
- Starrett, P.H. & Savage, J.M. 1973. The systematic status and distribution of Costa Rican glass-frogs, genus *Centrolenella* (Family Centrolenidae), with description of a new species. *Bulletin of the Southern California Academy of Sciences* **72**: 57-78.
- Stejneger, L. 1926. Two new tailless amphibians from western China. *Proceedings of the Biological Society of Washington* **39**.
- Stöck, M., Bretschneider, P. & Grosse, W.R. 2000. The mating call and male release call of *Bufo raddei* Strauch, 1876 with some phylogenetic implications. *Russian Journal of Herpetology* **7**: 215-226.
- Stöck, M., Sicilia, A., Belfiore, N.M., Buckley, D., Lo Brutto, S., Lo Valvo, M. & Arculeo, M. 2008. Post-Messinian evolutionary relationships across the Sicilian channel: Mitochondrial and nuclear markers link a new green toad from Sicily to African relatives. *Bmc Evolutionary Biology* **8**: 56.

Straneck, R., Olmedo, E.V. & Carrizo, G.R. 1993. *Catalogo de voces de anfibios Argentinos*.

Parte 1. Ediciones Lola, Buenos Aires.

Strecker, J.K. 1910. Description of a new solitary spadefoot (*Scaphiopus hurterii*) from Texas, with other herpetological notes. *Proceedings of the Biological Society of Washington* **23**.

Stuart, B.L. & Chan-Ard, T. 2005. Two new *Huia* (Amphibia : Ranidae) from Laos and Thailand. *Copeia* **2005**: 279-289.

Suazo-Ortuño, I., Alvarado-Díaz, J., Raya-Lemus, E. & Martinez-Ramos, M. 2007. Diet of the Mexican marbled toad (*Bufo marmoreus*) in conserved and disturbed tropical dry forest. *Southwestern Naturalist* **52**: 305-309.

Sullivan, B.K. & Wagner Jr., W.E. 1988. Variation in advertisement and release calls, and social influences on calling behavior in the Gulf Coast toad (*Bufo valliceps*). *Copeia* **1988**: 1014-1020.

Sullivan, B.K., Malmos, K.B., Gergus, E.W.A. & Bowker, R.W. 2000. Evolutionary implications of advertisement call variation in *Bufo debilis*, *B. punctatus*, and *B. retiformis*. *Journal of Herpetology* **34**: 368-374.

Tárano, Z. 2010. Advertisement calls and calling habits of frogs from a flooded savanna of Venezuela. *South American Journal of Herpetology* **5**: 221-240.

Taylor, R.C., Buchanan, B.W. & Doherty, J.L. 2007. Sexual selection in the squirrel treefrog *Hyla squirella*: the role of multimodal cue assessment in female choice. *Animal Behaviour* **74**: 1753-1763.

Thomé, M.T.C. & Brasileiro, C.A. 2007. Dimorfismo sexual, uso do ambiente e abundância sazonal de *Elachistocleis* cf. *ovalis* (Anura: Microhylidae) em um remanescente de Cerrado no estado de São Paulo, sudeste do Brasil. *Biota Neotropica* **7**: 27-33.

- Trueb, L. & Duellman, W.E. 1971. A synopsis of Neotropical hylid frogs, genus *Osteocephalus*. *Occasional Papers of the Museum of Natural History, The University of Kansas* **1**: 1-47.
- Tucker, J.K. 1998. Population status of the Illinois chorus frog (*Pseudacris streckeri illinoensis*) in Madison County, Illinois: Results of 1998 Surveys, University of Illinois
- Tyler, M.J. & Martin, A.A. 1977. Taxonomic studies of some Australian leptodactylid frogs of the genus *Cyclorana* Steindachner. *South Australian Museum* **17**: 261-276.
- Tyler, M.J. & Davies, M. 1978. Species-groups within the Australopapuan hylid frog genus *Litoria* Tschudi. *Australian Journal of Zoology* **26**: 1-47.
- Ueda, H. 1994. Mating calls of the pond frog species distributed in the Far East and their artificial hybrids. *Scientific Report of the Laboratory for Amphibian Biology, Hiroshima University* **13**: 197-232.
- Valetti, J.A., Salas, N.E. & Martino, A.L. 2013. Bioacoustic of the advertisement call of *Ceratophrys cranwelli* (Anura: Ceratophryidae). *Revista de Biología Tropical* **61**: 273-280.
- Van Beurden, E. & McDonald, K.R. 1980. A new species of *Cyclorana* (Anura: Hylidae) from northern Queensland. *Transactions of the Royal Society of South Australia* **104**: 193-195.
- van den Elzen, P. & Kreulen, D.A. 1979. Notes on the vocalisations of some amphibians from the Serengeti National Park, Tanzania. *Bonner Zoologische Beiträge* **30**: 385-403.
- Van Kampen, P.N. 1923. *The Amphibia of the Indo-Australian Archipelago* E. J. Brill, Leiden.
- Vásquez, T. & Pfennig, K.S. 2007. Looking on the bright side: females prefer coloration indicative of male size and condition in the sexually dichromatic spadefoot toad, *Scaphiopus couchii*. *Behavioral Ecology and Sociobiology* **62**: 127-135.

- Vaz-Silva, W., Di-Bernardo, M., Guimarães, L.D.A. & Bastos, R.P. 2007. Territoriality, agonistic behavior, and vocalization in *Pseudis bolbodactylus* A. Lutz, 1925 (Anura: Hylidae) from Central Brazil. *Salamandra* **43**: 35-42.
- Vejarano, S., Thomas, M., Glaw, F. & Vences, M. 2006. Advertisement call and tadpole morphology of the clutch-guarding frog *Mantidactylus argenteus* from eastern Madagascar. *African Zoology* **41**: 164-169.
- Vences, M., Glaw, F., Jesu, R. & Schimmenti, G. 2000. A new species of *Heterixalus* (Amphibia : Hyperoliidae) from Western Madagascar. *African Zoology* **35**: 269-276.
- Vences, M., Andreone, F., Glaw, F., Raminosoa, N., Randrianirina, J.E. & Vieites, D.R. 2002. Amphibians and reptiles of the Ankaratra Massif: reproductive diversity, biogeography and conservation of a montane fauna in Madagascar. *Italian Journal of Zoology* **69**: 263-284.
- Vences, M. & Glaw, F. 2002. Two new treefrogs of the *Boophis rappiodes* group from eastern Madagascar (Amphibia Mantellidae). *Tropical Zoology* **15**: 141-163.
- Vences, M., Raxworthy, C.J., Nussbaum, R.A. & Glaw, F. 2003. A revision of the *Scaphiophryne marmorata* complex of marbled toads from Madagascar, including the description of a new species. *Herpetological Journal* **13**: 69-79.
- Vences, M. & Glaw, F. 2004. Revision of the subgenus *Chonomantis* (Anura : Mantellidae : *Mantidactylus*) from Madagascar, with description of two new species. *Journal of Natural History* **38**: 77-118.
- Vilaça, T.R.A., Silva, J.R.D.S. & Solé, M. 2011. Vocalization and territorial behaviour of *Phyllomedusa nordestina* Caramaschi, 2006 (Anura: Hylidae) from Southern Bahia, Brazil. *Journal of Natural History* **45**: 1823-1834.

- Wagner Jr., W.E. & Sullivan, B.K. 1995. Sexual selection in the Gulf Coast toad, *Bufo valliceps*: female choice based on variable characters. *Animal Behaviour* **49**: 305-319.
- Wang, J., Cui, J., Shi, H., Brauth, S.E. & Tang, Y. 2012. Effects of body size and environmental factors on the acoustic structure and temporal rhythm of calls in *Rhacophorus dennysi*. *Asian Herpetological Research* **3**: 205-212.
- Wen, A., Vasquez, N. & Castroviejo-Fisher, S. 2012. Description of the previously unknown advertisement calls of *Hyalinobatrachium fragile*, *H. pellucidum*, and *Vitreorana antisthenesi* (Amphibia: Centrolenidae). *Zootaxa* **3480**: 80-87.
- Weygoldt, P. & Peixoto, O.L. 1987. *Hyla ruschii* n. sp., a new frog from the Atlantic Forest domain in the state of Espirito Santo, Brazil (Amphibia, Hylidae). *Studies on Neotropical Fauna and Environment* **22**: 237-247.
- Wildenhues, M.J., Bagaturrov, M.F., Schmitz, A., Tran, D.A., Hendrix, R. & Ziegler, T. 2011. Captive management and reproductive biology of Orlov's Treefrog, *Rhacophorus orlovi* Ziegler & Köhler, 2001 (Amphibia: Anura: Rhacophoridae), including larval description, colour pattern variation and advertisement call. *Der Zoologische Garten* **80**: 287-303.
- Wilkinson, J. 2003. *Kinugasa flying frog, Rhacophorus arboreus*. *Grzimek's Animal Life Encyclopedia*. Gale Group, Farmington Hills, Michigan.
- Wogel, H., Abrunhosa, P.A. & Pombal Jr, J.P. 2002. Atividade reprodutiva de *Physalaemus signifer* (Anura, Leptodactylidae) em ambiente temporário Iheringia. *Série Zoologia* **92**: 57-70.
- Wogel, H. & Pombal Jr, J.P. 2007. Comportamento reprodutivo e seleção sexual em *Dendropsophus bipunctatus* (Spix, 1824) (Anura, Hylidae). *Papéis Avulsos de Zoologia (São Paulo)* **47**: 165-174.

- Wood Jr., P.L., Crismer, L.L., Ahmad, N. & Senawi, J. 2008. Two new species of torrent-dwelling toads *Ansonia* Stoliczka, 1870 (Anura: Bufonidae) from Peninsular Malaysia. *Herpetologica* **64**: 321-340.
- Wright, A.H. & Wright, A.A. 1949. *Handbook of Frogs and Toads of the United States and Canada*. Comstock Publishing Company, Ithaca, NY.
- Wycherley, J., Doran, S. & Beebee, T.J.C. 2002. Frog calls echo microsatellite phylogeography in the European pool frog (*Rana lessonae*). *Journal of Zoology* **258**: 479-484.
- Yu, B.G. & Zheng, R.Q. 2009. The advertisement call of the giant spiny frog *Paa spinosa*. *Current Zoology* **55**: 411-415.
- Zainudin, R., Rahman, M.A., Zain, B.M.M., Shukor, M.N., Inger, R.F. & Norhayati, A. 2010. Mating calls description of five species of frogs from the genus *Hylarana* Tschudi 1838 (Amphibia, Anura, Ranidae) from Sarawak, Malaysia. *Sains Malaysiana* **39**: 363-369.
- Zainudin, R. & Sazali, S.N. 2012. A morphometric analysis of *Hylarana signata* group (previously known as *Rana signata* and *Rana picturata*) of Malaysia. *International Journal of Modern Physics: Conference Series* **9**: 199-208.
- Zhang, F., Chen, P. & Zhao, S.Y. 2013. Comparison of mating calls and adaptive strategies of *Amolops wuyiensis* and *Odorrana tormotus* (Anura) in noise-controlled environments. *Zoological Research / "Dong wu xue yan jiu" bian ji wei yuan hui bian ji* **34**: 196-203.
- Zhou, Y.L., Qiu, X., Fang, X.B., Yang, L.Y., Zhao, Y., Fang, T., Zheng, W.H. & Liu, J.S. 2014. Acoustic characteristics of eight common Chinese anurans during the breeding season. *Zoological Research / "Dong wu xue yan jiu" bian ji wei yuan hui bian ji* **35**: 42-50.

Zimmerman, B.L. 1983. A comparison of structural features of calls of open and forest habitat frog species in the Central Amazon. *Herpetologica* **39**: 235-246.

Zina, J. & Haddad, C.F.B. 2005. Reproductive activity and vocalizations of *Leptodactylus labyrinthicus* (Anura: Leptodactylidae) in southeastern Brazil. *Biota Neotropica* **5**: 119-129.