

Department of Physics, Chemistry and Biology

Master's Thesis

# **Search for Dark Matter in the Upgraded High Luminosity LHC at CERN**

## **Impact of ATLAS phase II performance on a mono-jet analysis**

**Sven-Patrik Hallsjö**

Thesis work performed at Stockholms universitet

Linköping, May 27, 2014

LiTH-IFM-EX--YY/NNNN--SE



**Linköpings universitet**  
**TEKNISKA HÖGSKOLAN**

Department of Physics, Chemistry and Biology  
Linköping University  
SE-581 83 Linköping, Sweden



Department of Physics, Chemistry and Biology

# **Search for Dark Matter in the Upgraded High Luminosity LHC at CERN**

## **Impact of ATLAS phase II performance on a mono-jet analysis**

**Sven-Patrik Hallsjö**


Thesis work performed at Stockholms universitet

Linköping, May 27, 2014

Supervisor: **Docent Christophe Clément**  
FYSIKUM Stockholms universitet  
**Professor Magnus Johansson**  
IFM, Linköpings universitet

Examiner: **Professor Magnus Johansson**  
IFM, Linköpings universitet



	<b>Avdelning, Institution</b> Division, Department  Theoretical physics group Department of Physics, Chemistry and Biology SE-581 83 Linköping	<b>Datum</b> Date  2014-05-27										
<b>Språk</b> Language  <input type="checkbox"/> Svenska/Swedish <input checked="" type="checkbox"/> Engelska/English  <input type="checkbox"/> _____	<b>Rapporttyp</b> Report category  <input type="checkbox"/> Licentiatavhandling <input checked="" type="checkbox"/> Examensarbete <input type="checkbox"/> C-uppsats <input type="checkbox"/> D-uppsats <input type="checkbox"/> Övrig rapport <input type="checkbox"/> _____	<b>ISBN</b> _____  <b>ISRN</b> LiTH-IFM-EX-YY/NNNN--SE  <table border="0"> <tr> <td><b>Serietitel och serienummer</b></td> <td><b>ISSN</b></td> </tr> <tr> <td>Title of series, numbering</td> <td>_____</td> </tr> </table>	<b>Serietitel och serienummer</b>	<b>ISSN</b>	Title of series, numbering	_____						
<b>Serietitel och serienummer</b>	<b>ISSN</b>											
Title of series, numbering	_____											
<b>URL för elektronisk version</b>  <a href="http://urn.kb.se/resolve?urn=urn:nbn:se:liu:diva-XXXXX">http://urn.kb.se/resolve?urn=urn:nbn:se:liu:diva-XXXXX</a>												
<table border="0"> <tr> <td><b>Titel</b></td> <td>Sökandet efter mörk materia i den uppgraderade hög luminositets LHC i CERN</td> </tr> <tr> <td><b>Title</b></td> <td>Search for Dark Matter in the Upgraded High Luminosity LHC at CERN</td> </tr> <tr> <td colspan="2"> </td> </tr> <tr> <td><b>Författare</b></td> <td>Sven-Patrik Hallsjö</td> </tr> <tr> <td><b>Author</b></td> <td></td> </tr> </table>			<b>Titel</b>	Sökandet efter mörk materia i den uppgraderade hög luminositets LHC i CERN	<b>Title</b>	Search for Dark Matter in the Upgraded High Luminosity LHC at CERN			<b>Författare</b>	Sven-Patrik Hallsjö	<b>Author</b>	
<b>Titel</b>	Sökandet efter mörk materia i den uppgraderade hög luminositets LHC i CERN											
<b>Title</b>	Search for Dark Matter in the Upgraded High Luminosity LHC at CERN											
<b>Författare</b>	Sven-Patrik Hallsjö											
<b>Author</b>												
<b>Sammanfattning</b> Abstract  <p>If your thesis is written in English, the primary abstract would go here while the Swedish abstract would be optional.</p>												
<table border="0"> <tr> <td><b>Nyckelord</b></td> <td></td> </tr> <tr> <td><b>Keywords</b></td> <td>ATLAS, Beyond standard model physics, CERN, Dark matter, Elementary particle physics, High energy physics, SUSY, something, this is in mythesis.sty</td> </tr> </table>			<b>Nyckelord</b>		<b>Keywords</b>	ATLAS, Beyond standard model physics, CERN, Dark matter, Elementary particle physics, High energy physics, SUSY, something, this is in mythesis.sty						
<b>Nyckelord</b>												
<b>Keywords</b>	ATLAS, Beyond standard model physics, CERN, Dark matter, Elementary particle physics, High energy physics, SUSY, something, this is in mythesis.sty											



## **Abstract**

If your thesis is written in English, the primary abstract would go here while the Swedish abstract would be optional.





## Acknowledgments

I wish to dedicate this thesis to my mathematics teacher Ulf Rydmark without whom I would not have studied physics.

A big thank you to my family, fiancée and friends who have supported me throughout my education.

I want to thank my supervisor Christophe Clément and all those who helped me at Stockholms university.

I also want to thank my examiner Magnus Johansson, who always took time to answer any question from and support his students.

A special thank you to Professor Irina Yakimenko without whom my master years would have been duller and probably impossible.

Finally I want to thank: Someone someone for proofreading my

*Linköping, May 2014*  
*Sven-Patrik Hallsjö*



---

# Contents

<b>Notation</b>	<b>ix</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Research goals . . . . .	2
1.2 Theoretical Background . . . . .	3
1.2.1 Quantum mechanics and quantum field theory . . . . .	3
1.2.2 Effective field theory . . . . .	3
1.2.3 Nuclear, particle and subatomic particle physics . . . . .	3
1.2.4 The standard model of particle physics . . . . .	4
1.2.5 Beyond the standard model: Supersymmetry . . . . .	4
1.2.6 Dark matter . . . . .	4
1.2.7 Search for WIMPS . . . . .	5
1.3 Experimental overview . . . . .	6
1.3.1 LHC . . . . .	6
1.3.2 ATLAS . . . . .	7
1.3.3 Coordinate system . . . . .	7
1.3.4 Calorimeter . . . . .	8
1.3.5 Truth and reconstructed data . . . . .	8
1.3.6 Pile-up . . . . .	8
1.3.7 Jet and missing energy . . . . .	8
1.3.8 Phase II high luminosity upgrade . . . . .	8
<b>2 Method</b>	<b>11</b>
2.1 Monte Carlo simulation, truth data . . . . .	12
2.2 ROOT . . . . .	12
2.3 Validation of smearing functions . . . . .	12
2.3.1 Smearing . . . . .	12
2.3.2 Validation . . . . .	12
2.4 Signal to background ratio . . . . .	13
2.4.1 Selection criteria . . . . .	13
2.4.2 Comparing with published papers . . . . .	13
2.4.3 Figures of merit . . . . .	13
2.5 Other selection criteria and observables . . . . .	13

2.6	Mitigating the effect . . . . .	13
<b>3</b>	<b>Results</b>	<b>15</b>
3.1	Validation of smearing functions . . . . .	15
3.2	Signal to background ratio . . . . .	15
3.3	Mitigating the effect . . . . .	15
<b>4</b>	<b>Discussion</b>	<b>17</b>
<b>5</b>	<b>Conclusions</b>	<b>19</b>
	<b>Bibliography</b>	<b>23</b>

---

# Notation

## ABBREVIATIONS

Abbreviation	Meaning
ATLAS	A large Toroidal LHC ApparatuS
CERN	Organisation européenne pour la recherche nucléaire <sup>1</sup>
LHC	Large Hadron Collider
SUSY	SUperSYmmetry
WIMP	Weakly Interacting Massive Particle
WIMPS	Weakly Interacting Massive ParticleS
QED	Quantum ElectroDynamics
QM	Quantum Mechanics

---

<sup>1</sup>Originally, Conseil Européen pour la Recherche Nucléaire



# 1

---

## Introduction

Discrepancies in measurements of the rotations of galaxies indicate the presence of a large amount of matter which interacts through gravity, though not electromagnetically making it invisible to our telescopes. This matter is commonly referred to as dark matter. Since no known or hypothesised particle in the standard model of particle physics can be used as a candidate for dark matter, this has opened the door for new physics. Aside from dark matter there are other phenomena, such as the neutrino mass and the hierarchy problem, that can not be explained today. One of the proposed models to correct these discrepancies is known as Supersymmetry (SUSY).

At the Organisation européenne pour la recherche nucléaire (CERN) the interest now lies to discover any evidence of SUSY. Among other searches one is fixed at looking for so called weakly interacting massive particles (WIMPS) which may be a candidate for dark matter. It is usually impossible to detect any interaction of dark matter candidates on the subatomic scale, however through looking at proposed interactions, searching for assumed decay channels and inconsistencies in momentum conservation it is hoped that signs will be found. Though as of March 7, 2014 none have been found.

Both these experiments and current theories now show that higher energies are required at the LHC to be able to see any signs. This is why the LHC and all detectors are undergoing a vast upgrading program **reference!**. In this thesis focus will be on the last part of the upgrade due for completion in 2023, known as the high luminosity-LHC phase II upgrade; and also on the ATLAS detector.

In this chapter an introduction to both the theoretical and experimental details required to understand the method and results is given.

## 1.1 Research goals

This research took place at Stockholm University from January 7th until **when?** During the research period the following tasks were set up and performed/answered:

- Implement a C++ programme that loops over the collisions inside the signal and background datasets.
- For each collision retrieve the relevant observables (variables used to extract the signal over the background) and apply "smearing functions" to emulate the effect of the high luminosity on the observables.
- For both signal and background datasets, compare observables before and after smearing. What observables are the least/most affected?
- Implement selection criteria that selects the signal collisions efficiently while reduces significantly the background. In a first step the selection criteria should be taken from existing studies.
- Selection criteria can be evaluated and compared with each other using a figure of merit  $Z$ , that measures the sensitivity of the experiment to the dark matter signal. Calculate  $Z$  for the given selection criteria before and after smearing.
- What is the effect of the high luminosity (smearing) on the value of  $Z$ ?
- Investigate other selection criteria and observables, to mitigate the effect of high luminosity. Use  $Z$  to rank different criteria after smearing.
- Conclude on the effect of the high luminosity on the sensitivity for dark matter and possible ways to mitigate its effects using alternative observables and selection criteria.



## 1.2 Theoretical Background

The following is a short description of the theory which is required to understand this thesis.

### 1.2.1 Quantum mechanics and quantum field theory

In the beginning of the 20th century, some physical phenomena could not be explained by classical physics, for example the ultra-violet disaster of any classical model of black-body radiation, and the photoelectric effect.[1] It was these phenomena that led to the formulation of quantum mechanics (QM), where energy transfer is quantized and particles can act as both waves and particles at the same time. [2]

Combining QM with classical electromagnetism proved harder than expected, colliding a photon(em-field) and an electron (particle/wave) is quite tricky. (Refer to scattering) One idea that came from this was to explain them both in the same framework, field theory. Also, trying to incorporate special relativity into QM suggested a field description. (Metric description and bending of gravitational fields). The culmination of both of these problems is Quantum electrodynamics (QED) which with incredible precision explains electromagnetic phenomena including effects from special relativity.[3]

Something something Lagrangian of these processes something something can be simply viewed through Feynman diagrams. Explain four-vectors quickly. Somehow end up with Feynman diagrams.

Also mention special rel + qm expects and now proven antimatter.

Speak about: Why QM, Lagrangian refer to classical mechanics, end with hand off to particle physics. need to explain observables. Theoretical Particle physics comes from Feynman diagrams!

Dont forget to explain cross-sections from qft.

Find more information in [1, 3, 4].

### 1.2.2 Effective field theory

In general: Feynman diagram with a big circle.

From article:

Which one do we use? What parameters have we joined?

How does this pertain to the rest? info in [5]

### 1.2.3 Nuclear, particle and subatomic particle physics

Can be seen as the experimental counterpart to quantum mechanics. Many could argue that these branches started after Ernest Rutherford famous gold foil experiment (reference), where he discovered that matter is composed of matter with a

nucleus, a lot of empty space and electrons. This and more sparked the curiosity to see what the nucleus was made of and so on...

The discovery of the quark diving of bosons/fermions different generations. Fundamental particles. Basically all of 20th century physics.

Something so that a description of particles are in here, end with standard model. Content should be enough for the rest of the thesis regarding collisions etc. Luminosity!

Explain hadrons...

### 1.2.4 The standard model of particle physics

How and why is there a standard model? give the Lagrangian and refer to all the different interactions that are included. Which then combines QM with subatomic particle physics.

Mention Antimatter! Is it proved? What problems exist?

- No QFT for general relativity! Thus there is no link between gravity and the SM.
- Experimentally it has been shown that neutrinos have mass, though in SM they do not!
- Asymmetry between matter and antimatter can not fully be described.
- No dark matter candidate!
- Dark energy?

### 1.2.5 Beyond the standard model: Supersymmetry

In the early 1970:s similar as QED expansion with antimatter due to (integral which one diverged?). Similarly to this, an expansion with a similar symmetry having bosons instead of fermions and the reverse. These symmetrical particles are known as supersymmetrical partners. The SUSY partner of a boson is denoted as sfermion (squarks and sleptons) whereas the SUSY partner of a fermion is denoted as bosinos (gauginos)

Different problems, hierarchy, etc

Bring up different expansions. Here we will talk about supersymmetry (SUSY) end with neutrino/WIMPS. Minimal Supersymmetric Standard Model

Supersymmetry: Every boson has a supersymmetrical fermion, and the reverse.

Some from, else from licentiates [6]

### 1.2.6 Dark matter

A very quick introduction was given in the beginning of this chapter. Dark matter is the name given to the solution to the discrepancies of galactic rotations.

After this the big question arises, what could this dark matter consist of?

Weakly interacting massive particles (WIMPS) are a candidate to explain Dark matter, it is this candidate which is considered in this thesis. In addition the condition that the WIMPS are Dirac fermions (fermions which are not their own antiparticle) is implied.

In table 1.1 the operators which are integrated out via the effective field theory and are of interest in this thesis are given.

Name	Initial state	Type	Operator
D1	qq	scalar	$\frac{m_q}{M_*^3} \bar{\chi} \chi \bar{q} q$
D5	qq	vector	$\frac{1}{M_*^2} \bar{\chi} \gamma^\mu \chi \bar{q} \gamma_\mu q$
D8	qq	axial-vector	$\frac{1}{M_*^2} \bar{\chi} \gamma^\mu \gamma^5 \chi \bar{q} \gamma_\mu \gamma^5 q$
D9	qq	tensor	$\frac{1}{M_*^2} \bar{\chi} \sigma^{\mu\nu} \chi \bar{q} \sigma_{\mu\nu} q$
D11	gg	scalar	$\frac{1}{4M_*^3} \bar{\chi} \chi \alpha_s (G_{\mu\nu}^a)^2$

**Table 1.1:** From [7]

Where D denotes that the WIMPS are assumed to be Dirac fermions.

Alot here taken from [5]

### 1.2.7 Search for WIMPS

Since the search for WIMPS at the LHC is based on looking at  $E_T^{Miss}$  it will be canonical though the experiment can no establish if a WIMP is stable on a cosmological time scale and thus if it is a Dark matter candidate [7]

Include SUSY-feynman?


What is it? Why at CERN/ATLAS? Candidates? Dark matter is something which does not interact electromagnetically however it does have a gravitational effect on nearby bodies. Cold dark matter? Non-barionic dark matter. Why not bari-  
onic? WIMPS, wimps as candidates. How is this detectable at ATLAS? Finish with this. Refer next chapter and that neutralinos are a candidate.

## 1.3 Experimental overview

What was used in this research and what needs to be explained? Upgrade, pileup etc. Somewhere here explain how the radial coordinate system is defined.

### 1.3.1 LHC

The Large hadron collider (LHC) is a particle accelerator located at CERN near Geneva in Switzerland, see figure 1.1. The accelerator was built to explore physics beyond the standard model and to make more accurate measurements of standard model physics. Before it was shut down for an upgrade in 2012 it was able to accelerate two proton beams to such a velocity that they had an energy of 4 TeV which gives a center of mass energy,  $\sqrt{s} = 8$  TeV. It should be noted that the proton beam is not homogeneous, it is comprised of bunches of protons with enough spacing that bunch collisions can happen independent of each other. Apart from the energy, the ability for an accelerator to produce interactions can be calculated through the instantaneous luminosity of the LHC was  $10^{34} \text{ cm}^{-2}\text{s}^{-1}$  or  $10\text{nb}^{-1}\text{s}^{-1}$  where  $1 \text{ barn(b)} = 10^{-24} \text{ cm}^2$ . All values taken from [8].



*Figure 1.1: Figure showing the LHC and the different detectors[9]*

The instantaneous luminosity can be defined in different ways depending on how the collision takes place. For two collinear intersecting particle beams it is defined as:

$$\mathcal{L} = \frac{fkN_1N_2}{4\pi\sigma_x\sigma_y} \quad (1.1)$$

where  $N_i$  are the number of particles in each of the bunches,  $f$  is the frequency at which the bunches collide,  $k$  the number of colliding bunches in each beam, and  $\sigma_x$  ( $\sigma_y$ ) is the horizontal (vertical) beam size at the interaction point. Since the instantaneous luminosity increases quadratically with more particles in each bunch this would be a good strategy. However aside from the difficulties to create and maintain a beam with more particles, a large  $N_i$  increases the probability for multiple collisions per bunch crossing, referred to as pile-up. Pile up will be a key aspect which is described more in subsection 1.3.6.

The expected number of events can be calculated by using the instantaneous luminosity through the following:

$$N = \sigma \int \mathcal{L} dt := \sigma \mathcal{L} \quad (1.2)$$

where  $\mathcal{L}$  is the luminosity and  $\sigma$  is the cross section which is often measured in barn. The luminosity is a measurement of total number of interactions that have occurred over time. Before the LHC was shut down this values was  $20.8 \text{ fb}^{-1}$ .

The cross section is defined through the integral of the differential cross section, as explained in subsection 1.2.1, over the whole solid angle:

$$\sigma = \oint d\Omega \frac{d\sigma}{d\Omega} \quad (1.3)$$

The cross section is therefore a measure of the effective surface area seen by the impinging particles, and as such is expressed in units of area. The cross section is proportional to the probability that an interaction will occur. It also provides a measure of the strength of the interaction between the scattered particle and the scattering center. Further details can be found in reference [10]

### 1.3.2 ATLAS

As seen in figure 1.1, there are several detectors at CERN. One of these is a large toroidal LHC apparatus (ATLAS) which is a general purpose detector that uses a toroid magnet. Its goal is to observe several different production and decay channels. The detector is composed of three concentric subdetectors, the Inner detector, the Calorimeters and the Muon spectrometer.

Most and more in [11]


### 1.3.3 Coordinate system

The coordinate system of ATLAS, seen in figure 1.2 is a right-handed coordinate system with the x-axis pointing towards the centre of the LHC tunnel, and the z-axis along the tunnel/beam (counter clockwise) seen from above. The y-axis points upward. The origin is define as the interaction point. A cylindrical coordinate system is also used for the transversal plane.  $(R, \phi, Z)$ . For simplicity the pseudorapidity of particles from the primary vertex is defined as:

$$\eta = -\ln\left(\tan \frac{\theta}{2}\right) \quad (1.4)$$

where  $\theta$  is the polar angle (xz-plane) of the particle direction measured from the positive z-axis.  $\eta$  is through this definition invariant under boosts in the z-direction.

It is quite common to calculate the distance between particles and jets in the  $(\eta, \phi)$  plane,  $d = \sqrt{\Delta\eta + \Delta\phi}$



**Figure 1.2:** Figure showing the ATLAS detector and the definition of the coordinate system. Image altered from[12]

### 1.3.4 Calorimeter

### 1.3.5 Truth and reconstructed data

Give a general description.

In this thesis, truth = Montecarlo. Reconstructed data = that which is seen in the detectors, which is the smeared data.

### 1.3.6 Pile-up

### 1.3.7 Jet and missing energy

What is a jet? why are we only looking at transverse missing energy?

### 1.3.8 Phase II high luminosity upgrade

Talk about the upgrade schedule. I am looking at the upgrade which will be done at CERN and will be completed around 2022-2023 and is denoted High Luminosity-LHC Phase 2 upgrade. When this is running the following is expected:

Entity	Expected	Last run (2012)
Luminosity	1000-3000 fb <sup>-1</sup>	20.8 fb <sup>-1</sup>
Pile-up	$\langle \mu \rangle = 200$	$\langle \mu \rangle = 20.7$
Center of mass energy	$\sqrt{s} = 14$ TeV	$\sqrt{s} = 8$ TeV

**Table 1.2:** Expected running values for the Phase II HL-upgraded LHC with older values for comparison. REFERENCE?

Taken from "a short explanation of different terminology by me" Find a cern source. Assumed effects, timespan when will it be done?





# 2

---

## Method

In this chapter the methodology used to perform the tasks given above in section 1.1.

## 2.1 Monte Carlo simulation, truth data

What it is in short detail. How it is set-up and what it produces for us in this context. Where does it come from? Perhaps quick explanation of different programs?

## 2.2 ROOT

A wonderful tool for processing data by programming in C++ and so on... Reference to root homepage?

## 2.3 Validation of smearing functions

Find more information in my presentation. also mention no pile-up dependence of leptons. For the proposed upgrade of the LHC.

### 2.3.1 Smearing

By using the simulation given above one might assume that it would be easy to model and simulate the whole process, from collision to detection in the upgraded LHC. It is possible, but it requires a lot of computing power. Instead one can use one simulation and a mathematical model to calculate the estimated response in the detector. This was validated and used in this thesis to be able to create the data needed for further analysis. The programming code used was the **official something that can be found somewhere**. Several histogram are given regarding this here. Some minor detail about them though most of it should be given in results and discussion also refer perhaps to my presentation? Does one self refer?

Why did we use this, what does it produce for us? A "simple" simulated effect of what can be detected in the detectors without having to run a full simulation.

The functions smear both the energy and the momenta of the four vector. However not the angles. Explain four vector somewhere?

Explain why leptons were not affected by pile-up. Presentation 2?

### 2.3.2 Validation

Parametrization used according to the paper [13]. What results and what did I get/say in my presentation? Use that in results Perhaps even write something better than the original that can be used to explain this again.

Remember for the discussion to mention different types of rms, relative or absolute. and the problem which occurred with this and the papers faults.

## **2.4 Signal to background ratio**

What I am doing now, looking at what signal? What are the different background processes? What and why was the weight used?

Signals should be explained somewhat in the introduction.

### **2.4.1 Selection criteria**

What criteria were used and more importantly why? It is quite important that you can explain why this was used.

### **2.4.2 Comparing with published papers**

10fb-1 paper, also can perhaps use CERN-PH-EP-2012-210

### **2.4.3 Figures of merit**

what is it? how is it calculated?

## **2.5 Other selection criteria and observables**

## **2.6 Mitigating the effect**



# 3

---

## Results

Saving as reference. test citing as: Here we cite Duck [14] [14]. If the above works, remember to edit myreferences.

### **3.1 Validation of smearing functions**

Figures from validation:

### **3.2 Signal to background ratio**

### **3.3 Mitigating the effect**



# 4

---

## Discussion





# 5

---

## Conclusions

Sätt av ett kort kapitel sist i rapporten till att avrunda och föreslå rikningar för framtida utveckling av arbetet.



# Appendix



---

## Bibliography

- [1] B.H. Bransden and C.J. Joachain. *Quantum mechanics*. Pearson Education, second edition, 2000. Cited on page 3.
- [2] Sven-Patrik Hallsjö. Covering the sphere with noncontextuality inequalities. Master's thesis, Linköping University, The Institute of Technology, 2013. URL <http://urn.kb.se/resolve?urn=urn:nbn:se:liu:diva-103663>. Cited on page 3.
- [3] A. Zee. *Quantum Field Theory in a Nutshell*. Princeton University Press, illustrated edition edition, March 2003. ISBN 0691010196. Cited on page 3.
- [4] Herbert Goldstein, Charles P. Poole, and John L. Safko. *Classical Mechanics (3rd Edition)*. Addison-Wesley, 3 edition, June 2001. ISBN 0201657023. Cited on page 3.
- [5] J. Goodman, M. Ibe, A. Rajaraman, W. Shepherd, T. M. P. Tait, and H.-B. Yu. Constraints on dark matter from colliders. *Phys.Rev.D*82:116010,2010, 82 (11):116010, December 2010. doi: 10.1103/PhysRevD.82.116010. Cited on pages 3 and 5.
- [6] S. P. Martin. A Supersymmetry Primer. In G. L. Kane, editor, *Perspectives on Supersymmetry*, number arXiv:hep-ph/9709356v6, page 1, 1998. Cited on page 4.
- [7] ATLAS Collaboration. Search for dark matter candidates and large extra dimensions in events with a jet and missing transverse momentum with the atlas detector. *J. High Energy Phys.*, 04(arXiv:1210.4491. CERN-PH-EP-2012-210):075. 58 p, October 2012. URL <http://cds.cern.ch/record/1485031>. Cited on page 5.
- [8] ATLAS. Atlas luminosity public results. <https://twiki.cern.ch/twiki/bin/view/AtlasPublic/LuminosityPublicResults>, 2013. Accessed: 2014-03-06. Cited on page 6.
- [9] AC Team. The four main LHC experiments, Jun 1999. URL <http://cds.cern.ch/record/40525>. Cited on page 6.

- [10] Werner Herr and B Muratori. Concept of luminosity. 2006. URL <http://cds.cern.ch/record/941318/>. Cited on page 7.
- [11] The ATLAS Collaboration. The atlas experiment at the cern large hadron collider. *Journal of Instrumentation*, 3(08):S08003. 437 p, 2008. URL <https://cdsweb.cern.ch/record/1129811/>. Cited on page 7.
- [12] Joao Pequenaio. Computer generated image of the whole ATLAS detector, Mar 2008. URL <http://cds.cern.ch/record/1095924>. Cited on page 8.
- [13] ATLAS Collaboration. Performance assumptions for an upgraded ATLAS detector at a High-Luminosity LHC. Mar 2013. URL <https://cds.cern.ch/record/1527529/>. Cited on page 12.
- [14] Donald Duck. The history of automatic control. *Duckburg Journal of Science*, 106(3):345–401, 2005. Cited on page 15.

## Upphovsrätt

Detta dokument hålls tillgängligt på Internet — eller dess framtida ersättare — under 25 år från publiceringsdatum under förutsättning att inga extraordinära omständigheter uppstår.

Tillgång till dokumentet innebär tillstånd för var och en att läsa, ladda ner, skriva ut enstaka kopior för enskilt bruk och att använda det oförändrat för icke-kommersiell forskning och för undervisning. Överföring av upphovsrätten vid en senare tidpunkt kan inte upphäva detta tillstånd. All annan användning av dokumentet kräver upphovsmannens medgivande. För att garantera äktheten, säkerheten och tillgängligheten finns det lösningar av teknisk och administrativ art.

Upphovsmannens ideella rätt innefattar rätt att bli nämnd som upphovsman i den omfattning som god sed kräver vid användning av dokumentet på ovan beskrivna sätt samt skydd mot att dokumentet ändras eller presenteras i sådan form eller i sådant sammanhang som är kränkande för upphovsmannens litterära eller konstnärliga anseende eller egenart.

För ytterligare information om Linköping University Electronic Press se förlagets hemsida <http://www.ep.liu.se/>

## Copyright

The publishers will keep this document online on the Internet — or its possible replacement — for a period of 25 years from the date of publication barring exceptional circumstances.

The online availability of the document implies a permanent permission for anyone to read, to download, to print out single copies for his/her own use and to use it unchanged for any non-commercial research and educational purpose. Subsequent transfers of copyright cannot revoke this permission. All other uses of the document are conditional on the consent of the copyright owner. The publisher has taken technical and administrative measures to assure authenticity, security and accessibility.

According to intellectual property law the author has the right to be mentioned when his/her work is accessed as described above and to be protected against infringement.

For additional information about the Linköping University Electronic Press and its procedures for publication and for assurance of document integrity, please refer to its www home page: <http://www.ep.liu.se/>