

TFYA17 Project

Transmission properties in a short biased quantum wire

Patrik Hallsjö, Felix Faber

1 Introduction

2 Theory

3 Results

Table of Contents

1 Introduction

2 Theory

3 Results

Introduction

- This project was dedicated to study an electron transport in a biased quantum wire.
- The goals were to construct a solver for the potential and to calculate the transmission and reflection for an electron sent towards the potential.

$$V(x) = \beta * V_{sd} + V_g * \tanh(s * (x - dx_1)) \\ - (V_{sd} + V_g) * \tanh(s * (x - dx_2))$$

Goals

- To set and to solve a problem with boundary conditions when an electron is injected into a wire.
- To calculate the transmission and reflection coefficients as functions of energy of injected electron and bias.
- To represent solutions in graphical form.
- Optional: to calculate a conductance.

Goals

- To set and to solve a problem with boundary conditions when an electron is injected into a wire.
- To calculate the transmission and reflection coefficients as functions of energy of injected electron and bias.
- To represent solutions in graphical form.
- Optional: to calculate a conductance.

Remark

Matlab was used to achieve the goals.

Table of Contents

1 Introduction

2 Theory

3 Results

Finite derivatives

- $f'(x) = \frac{f(x+h)-f(x+k)}{h-k}$

Finite derivatives

- $f'(x) = \frac{f(x+h)-f(x+k)}{h-k}$
- Forward: $\frac{f(x+h)-f(x)}{h}$

Finite derivatives

- $f'(x) = \frac{f(x+h)-f(x+k)}{h-k}$
- Forward: $\frac{f(x+h)-f(x)}{h}$
- Backward: $\frac{f(x)-f(x-h)}{h}$

Finite derivatives

- $f'(x) = \frac{f(x+h)-f(x+k)}{h-k}$
- Forward: $\frac{f(x+h)-f(x)}{h}$
- Backward: $\frac{f(x)-f(x-h)}{h}$
- Central: $\frac{f(x-h/2)-f(x+h/2)}{h}$

Finite derivatives

- $f'(x) = \frac{f(x+h)-f(x+k)}{h-k}$
- Forward: $\frac{f(x+h)-f(x)}{h}$
- Backward: $\frac{f(x)-f(x-h)}{h}$
- Central: $\frac{f(x-h/2)-f(x+h/2)}{h}$
- Second order: $\frac{f(x-h)-2f(x)+f(x+h)}{h^2}$

Finite steps in differential equations

- Second order differential equation without any first order terms $y''(x) = g(y(x), x)$
- Boundary conditions: $y(x_0) = y_0$ and $y(x_N) = y_N$, on the interval $[x_0, x_N]$.
- $h = d$ and $x = n * d$ yields
$$d * g(y(n * d), n * d) + 2y(n * d) - (y((n + 1) * d) + y((n - 1) * d)) = 0.$$
- $N + 1$ equations with $N + 1$ unknowns.

Quantum barrier problem

(1).

We need to solve the stationary Schrödinger equation with the boundary conditions $y_0 = (I + R) \exp(-ikx_0)$ and $y_N = T \exp(-ikx_N)$. We have put $I = 1$, which gives $y_0 = (1 + R) \exp(-ikx_0)$ to make the equations easier to solve.

Quantum barrier problem

(1).

We need to solve the stationary Schrödinger equation with the boundary conditions $y_0 = (I + R) \exp(-ikx_0)$ and $y_N = T \exp(-ikx_N)$. We have put $I = 1$, which gives $y_0 = (1 + R) \exp(-ikx_0)$ to make the equations easier to solve. We can use the finite difference approximation to solve the stationary Schrödinger equation as a second order ordinary differential equation (ODE). However, with R and T we have $N + 3$ unknowns but only $N + 1$ equations. Two more equations will be needed in order to fully solve the problem.

Quantum barrier problem

(1).

We need to solve the stationary Schrödinger equation with the boundary conditions $y_0 = (I + R) \exp(-ikx_0)$ and $y_N = T \exp(-ikx_N)$. We have put $I = 1$, which gives $y_0 = (1 + R) \exp(-ikx_0)$ to make the equations easier to solve. We can use the finite difference approximation to solve the stationary Schrödinger equation as a second order ordinary differential equation (ODE). However, with R and T we have $N + 3$ unknowns but only $N + 1$ equations. Two more equations will be needed in order to fully solve the problem. The solution to Schrödinger equation is continuous everywhere in the interval because it is a second order ODE, however it also needs to be continuous at the two boundaries. The finite difference approximation of the derivative yields: $y'_0 = y'_1$ and $y'_{N-1} = y'_N$. There are now $N + 3$ unknowns and $N + 3$ equations which will have a unique solution

Table of Contents

1 Introduction

2 Theory

3 Results

Numerical values

The program produces the following data, as well as images of the wave-functions and the potential barriers.

$$R = |R|^2 / (|R|^2 + |T|^2)$$

$$T = |T|^2 / (|R|^2 + |T|^2)$$

$$C = T / (R + T)$$





