






planar graph G























and so on...




separator "tree"





separator "tree"




separator "tree"












G has a **star partition** of width $O(n^{2/3})$




$G \subseteq S \boxtimes K_{O(n^{2/3})}$ where S is a star




$G \subseteq S \boxtimes K_{O(n^{2/3})}$ where S is a star




$G \subseteq S \boxtimes K_{O(n^{2/3})}$ where S is a star




What can we do with parts of size $\tilde{O}(\sqrt{n})$?



Bad News (Linial-Matousek-Sheffet-Tardos 2008): There exists planar G such that every 2-coloring of G has a monochromatic component of size $\Omega(n^{2/3})$




Bad News (Linial-Matousek-Sheffet-Tardos 2008): There exists planar G such that every 2-coloring of G has a monochromatic component of size $\Omega(n^{2/3})$



Bad News (Linial-Matousek-Sheffet-Tardos 2008): There exists planar G such that every 2-coloring of G has a monochromatic component of size $\Omega(n^{2/3})$

$G \subseteq H \boxtimes K_{o(n^{2/3})} \Rightarrow H$ is not 2-colorable.



Bad News (Linial-Matousek-Sheffet-Tardos 2008): There exists planar G such that every 2-coloring of G has a monochromatic component of size $\Omega(n^{2/3})$

$G \subseteq H \boxtimes K_{o(n^{2/3})} \Rightarrow H$ is not 2-colorable $\Rightarrow H$ is not a tree.

