



Journal of Graph Algorithms and Applications
<http://jgaa.info/> vol. 0, no. 0, pp. 0–0 (2005)

Three-Dimensional 1-Bend Graph Drawings

Pat Morin

School of Computer Science
Carleton University
Ottawa, Canada
morin@scs.carleton.ca

David R. Wood

Departament de Matemàtica Aplicada II
Universitat Politècnica de Catalunya
Barcelona, Spain
david.wood@upc.edu

Abstract

We consider three-dimensional grid-drawings of graphs with at most one bend per edge. Under the additional requirement that the vertices be collinear, we prove that the minimum volume of such a drawing is $\Theta(cn)$, where n is the number of vertices and c is the cutwidth of the graph. We then prove that every graph has a three-dimensional grid-drawing with $\mathcal{O}(n^3/\log^2 n)$ volume and one bend per edge. The best previous bound was $\mathcal{O}(n^3)$.

Article Type	Communicated by	Submitted	Revised
concise paper	G. Liotta	April 2004	March 2005

Presented at the 16th Canadian Conference on Computational Geometry (CCCG '04), Concordia University, Montréal, Canada, August 9–11, 2004.

Research of Pat Morin supported by NSERC.

Research of David Wood supported by the Government of Spain grant MEC SB2003-0270, and partially completed at Carleton University, Canada.

1 Introduction

We consider undirected, finite, and simple graphs G with vertex set $V(G)$ and edge set $E(G)$. The number of vertices and edges of G are respectively denoted by $n = |V(G)|$ and $m = |E(G)|$.

Graph drawing is concerned with the automatic generation of aesthetically pleasing geometric representations of graphs. Graph drawing in the plane is well-studied (see [4, 17]). Motivated by experimental evidence suggesting that displaying a graph in three dimensions is better than in two [21, 22], and applications including information visualisation [21], VLSI circuit design [18], and software engineering [23], there is a growing body of research in three-dimensional graph drawing.

A *three-dimensional polyline grid-drawing* of a graph, henceforth called a *polyline drawing*, represents the vertices by distinct points in \mathbb{Z}^3 (called *grid-points*), and represents each edge as a polyline between its endpoints with bends (if any) also at gridpoints, such that distinct edges only intersect at common endpoints, and each edge only intersects a vertex that is an endpoint of that edge. A polyline drawing with at most b bends per edge is called a *b-bend drawing*. A 0-bend drawing is called a *straight-line drawing*.

A folklore result states that every graph has a straight-line drawing. Thus we are interested in optimising certain measures of the aesthetic quality of such drawings. The *bounding box* of a polyline drawing is the minimum axis-aligned box containing the drawing. If the bounding box has side lengths $X-1$, $Y-1$ and $Z-1$, then we speak of an $X \times Y \times Z$ polyline drawing with *volume* $X \cdot Y \cdot Z$. That is, the volume of a polyline drawing is the number of gridpoints in the bounding box. This definition is formulated so that two-dimensional drawings have positive volume. This paper continues the study of upper bounds on the volume and number of bends per edge in polyline drawings. The volume of straight-line drawings has been widely studied [1–3, 5, 6, 9, 11, 12, 14, 16, 20]. Only recently have (non-orthogonal) polyline drawings been considered [11, 13]. Table 1 summarises the best known upper bounds on the volume and bends per edge in polyline drawings.

Cohen *et al.* [3] proved that the complete graph K_n (and hence every n -vertex graph) has a straight-line drawing with $\mathcal{O}(n^3)$ volume, and that $\Omega(n^3)$ volume is necessary. Dyck *et al.* [13] recently proved that K_n has a 2-bend drawing with $\mathcal{O}(n^2)$ volume. The same conclusion can be reached from the $\mathcal{O}(qn)$ volume bound of Dujmović and Wood [11], since trivially every graph has a $(n-1)$ -queue layout. Dyck *et al.* [13] asked the interesting question: what is the minimum volume in a 1-bend drawing of K_n ? The best known upper bound at the time was $\mathcal{O}(n^3)$, while $\Omega(n^2)$ is the best known lower bound. (Bose *et al.* [1] proved that all polyline drawings have $\Omega(n+m)$ volume.)

In this paper we prove two results. The first concerns *collinear* polyline drawings in which all the vertices are in a single line. Let G be a graph, and let σ be a linear order of $V(G)$. Let $L_\sigma(e)$ and $R_\sigma(e)$ denote the endpoints of each edge e such that $L_\sigma(e) <_\sigma R_\sigma(e)$. For each vertex $v \in V(G)$, the set $\{e \in E(G) : L_\sigma(e) \leq_\sigma v <_\sigma R_\sigma(e)\}$ is called the *cut* in σ at v . The *cutwidth* of σ is the maximum size of a cut in σ . The *cutwidth* of G is the minimum cutwidth of a linear order of $V(G)$. Cutwidth is a widely studied graph parameter (see [7]).

Table 1: Volume of 3D polyline drawings of n -vertex graphs with $m \geq n$ edges.

graph family	bends per edge	volume	reference
arbitrary	0	$\mathcal{O}(n^3)$	Cohen <i>et al.</i> [3]
arbitrary	0	$\mathcal{O}(m^{4/3}n)$	Dujmović and Wood [12]
maximum degree Δ	0	$\mathcal{O}(\Delta mn)$	Dujmović and Wood [12]
bounded chromatic number	0	$\mathcal{O}(n^2)$	Pach <i>et al.</i> [20]
bounded chromatic number	0	$\mathcal{O}(m^{2/3}n)$	Dujmović and Wood [12]
bounded maximum degree	0	$\mathcal{O}(n^{3/2})$	Dujmović and Wood [12]
H -minor free (H fixed)	0	$\mathcal{O}(n^{3/2})$	Dujmović and Wood [12]
bounded treewidth	0	$\mathcal{O}(n)$	Dujmović <i>et al.</i> [9]
k -colourable q -queue	1	$\mathcal{O}(kqm)$	Dujmović and Wood [11]
arbitrary	1	$\mathcal{O}(nm)$	Dujmović and Wood [11]
cutwidth c	1	$\mathcal{O}(cn)$	Theorem 1
arbitrary	1	$\mathcal{O}(n^3 / \log^2 n)$	Theorem 2
q -queue	2	$\mathcal{O}(qn)$	Dujmović and Wood [11]
q -queue (constant $\epsilon > 0$)	$\mathcal{O}(1)$	$\mathcal{O}(mq^\epsilon)$	Dujmović and Wood [11]
q -queue	$\mathcal{O}(\log q)$	$\mathcal{O}(m \log q)$	Dujmović and Wood [11]

Theorem 1. *Let G be a graph with n vertices and cutwidth c . The minimum volume for a 1-bend collinear drawing of G is $\Theta(cn)$.*

Theorem 1 represents a qualitative improvement over the $\mathcal{O}(nm)$ volume bound for 1-bend drawings by Dujmović and Wood [11]. Our second result improves the best known upper bound for 1-bend drawings of K_n .

Theorem 2. *Every complete graph K_n , and hence every n -vertex graph, has a 1-bend $\mathcal{O}(\log n) \times \mathcal{O}(n) \times \mathcal{O}(n^2 / \log^3 n)$ drawing with $\mathcal{O}(n^3 / \log^2 n)$ volume.*

It is not straightforward to compare the volume bound in Theorem 2 with the $\mathcal{O}(kqm)$ bound by Dujmović and Wood [11] for k -colourable q -queue graphs (see Table 1). However, since $k \leq 4q$ and $m \leq 2qn$ (see [10]), we have that $\mathcal{O}(kqm) \subseteq \mathcal{O}(q^3n)$, and thus the $\mathcal{O}(kqm)$ bound by Dujmović and Wood [11] is no more than the bound in Theorem 2 whenever the graph has a $\mathcal{O}((n/\log n)^{2/3})$ -queue layout. On the other hand, $kqm \geq m^2/n$. So for dense graphs with $\Omega(n^2)$ edges the $\mathcal{O}(kqm)$ bound by Dujmović and Wood [11] is cubic (in n), and the bound in Theorem 2 is necessarily smaller. In particular, Theorem 2 provides a partial solution to the above-mentioned open problem of Dyck *et al.* [13] regarding the minimum volume of a 1-bend drawing of K_n .

2 Proof of Theorem 1

First we prove the lower bound in Theorem 1.

Lemma 1. *Let G be a graph with n vertices and cutwidth c . Then every 1-bend collinear drawing of G has at least $cn/2$ volume.*

Proof. Consider a 1-bend collinear drawing of G in an $X \times Y \times Z$ bounding box. Let L be the line containing the vertices. If L is not contained in a grid-plane, then $X, Y, Z \geq n$, and the volume is at least $n^3 \geq cn$.

Now assume, without loss of generality, that L is contained in the $Z = 0$ plane. Let σ be a linear order of the vertices determined by L . Let B be the set of bends corresponding to the edges in the largest cut in σ . Then $|B| \geq c$. For every line L' parallel to L , there is at most one bend in B on L' , as otherwise there is a crossing.

First suppose that L is axis-parallel. Without loss of generality, L is the X -axis. Then $X \geq n$. The gridpoints in the bounding box can be covered by YZ lines parallel to L . Thus $YZ \geq |B| \geq c$, and the volume $XYZ \geq cn$.

Now suppose that L is not axis-parallel. Thus $X \geq n$ and $Y \geq n$. The gridpoints in the bounding box can be covered by $Z(X+Y)$ lines parallel to L . Thus $Z(X+Y) \geq |B| \geq c$, and the volume $XYZ \geq XYc/(X+Y) \geq cn/2$. \square

To prove the upper bound in Theorem 1 we will need the following lemma, which is a slight generalisation of a well known result. (For example, Pach *et al.* [20] proved the case $X = Y$). We say two gridpoints v and w in the plane are *visible* if the segment vw contains no other gridpoint.

Lemma 2. *The number of gridpoints $\{(x, y) : 1 \leq x \leq X, 1 \leq y \leq Y\}$ that are visible from the origin is at least $3XY/2\pi^2$.*

Proof. Without loss of generality $X \leq Y$. Let N be the desired number of gridpoints. For each $1 \leq x \leq X$, let N_x be the number of gridpoints (x, y) that are visible from the origin, such that $1 \leq y \leq Y$. A gridpoint (x, y) is visible from the origin if and only if x and y are coprime. Let $\phi(x)$ be the number of positive integers less than x that are coprime with x (Euler's ϕ function). Thus $N_x \geq \phi(x)$, and

$$N = \sum_{x=1}^X N_x \geq \sum_{x=1}^X \phi(x) \approx \frac{3X^2}{\pi^2}.$$

(See [15] for a proof that $\sum_{x=1}^X \phi(x) \approx 3X^2/\pi^2$.) If $X \geq Y/2$, then $N \geq 3XY/2\pi^2$, and we are done. Now assume that $Y \geq 2X$. If x and y are coprime, then x and $y+x$ are coprime. Thus $N_x \geq \lfloor Y/x \rfloor \cdot \phi(x)$. Thus,

$$N \geq \sum_{x=1}^X \left\lfloor \frac{Y}{x} \right\rfloor \cdot \phi(x) \geq \left(\frac{Y-X}{X} \right) \sum_{x=1}^X \phi(x) \approx \frac{3(Y-X)X}{\pi^2} \geq \frac{3XY}{2\pi^2}$$

\square

Now we prove the following strengthening of the upper bound in Theorem 1.

Lemma 3. *Let G be a graph with n vertices and cutwidth c . For all integers $X \geq 1$, G has a 1-bend collinear $X \times \mathcal{O}(c/X) \times n$ drawing with the vertices on the Z -axis. The volume is $\mathcal{O}(cn)$.*

Proof. Let σ be a vertex ordering of G with cutwidth c . For all pairs of distinct edges e and f , say $e \prec f$ whenever $R_\sigma(e) \leq_\sigma L_\sigma(f)$. Then \preceq is a partial order on $E(G)$. A *chain* (respectively, *antichain*) in a partial order is a set of pairwise comparable (incomparable) elements. Thus an antichain in \preceq is exactly a cut in σ . Dilworth's Theorem [8] states that every partial order with no $(k+1)$ -element antichain can be partitioned into k chains. Thus there is a partition of $E(G)$ into chains E_1, E_2, \dots, E_c , such that each $E_i = (e_{i,1}, e_{i,2}, \dots, e_{i,k_i})$ and $R_\sigma(e_{i,j}) \leq_\sigma L_\sigma(e_{i,j+1})$ for all $1 \leq j \leq k_i - 1$.

By Lemma 2 with $Y = \lceil 4\pi^2 c/3X \rceil$, there is a set $S = \{(x_i, y_i) : 1 \leq i \leq c, 1 \leq x_i \leq X, 1 \leq y_i \leq Y\}$ of gridpoints that are visible from the origin. Position the i th vertex in σ at $(0, 0, i)$ on the Z -axis, and position the bend for each edge $e_{i,j}$ at (x_i, y_i, j) . Edges in distinct chains are contained in distinct planes that only intersect in the Z -axis. Thus such edges do not cross. Edges within each chain E_i do not cross since no two edges in E_i are nested or crossing in σ , and the Z -coordinates of the bends of the edges in E_i agrees with the order of their endpoints on the Z -axis, as (imprecisely) illustrated in Figure 1. The bounding box is $X \times \lceil 4\pi^2 c/3X \rceil \times n$, since each chain has at most $n-1$ edges. \square




Figure 1: Construction of collinear 1-bend drawing in Lemma 3.

The constants in Lemma 3 can be tweaked as follows.

Lemma 4. *Let G be a graph with n vertices and cutwidth c . Then G has a 1-bend collinear $3 \times \lceil (c-2)/2 \rceil \times n$ drawing. The volume is at most $3(c-1)n/2$.*

Proof. Let $S = \{(-1, 0), (1, 0)\} \cup \{(x, y) : y \in \{-1, 1\}, -1 \leq x \leq \lceil (c-6)/2 \rceil\}$. Then S consists of at least c gridpoints that are visible from the origin. The result follows from the proof of Lemma 3. \square

Since the cutwidth of K_n is $n^2/4$ we have:

Corollary 1. *The minimum volume for a 1-bend collinear drawing of the complete graph K_n is $\Theta(n^3)$. For all $X \geq 1$, K_n has a 1-bend collinear $X \times \mathcal{O}(n^2/X) \times n$ drawing with the vertices on the Z -axis. Furthermore, K_n has a 1-bend collinear $3 \times \lceil n^2/8 \rceil \times n$ drawing with volume at most $3n^3/8$. \square*

3 Proof of Theorem 2

Let $P = \lceil \frac{1}{2} \log_4 n \rceil$ and $Q = \lceil n/P \rceil$. Let $V(K_n) = \{v_{a,i} : 1 \leq a \leq P, 1 \leq i \leq Q\}$. Position each vertex $v_{a,i}$ at

$$(2a, aQ + i, 0) .$$

For each $1 \leq a \leq P$, the set of vertices $\{v_{a,i} : 1 \leq i \leq Q\}$ induces a complete graph K_Q , which is drawn using Corollary 1 (with the dimensions permuted) in the box

$$[2a, 2a + P] \times [aQ + 1, (a + 1)Q] \times [0, -cQ^2/P] ,$$

for some constant c . For all $1 \leq a < b \leq P$, orient each edge $e = (v_{a,i}, v_{b,j})$, and position the bend for e at

$$r_e = (2a + 1, bQ + j, 4^{P-a}Q - i) ,$$

as (imprecisely) illustrated in Figure 2. We say $v_{a,i}r_e$ is an *outgoing* segment at $v_{a,i}$, and $r_e v_{b,j}$ is an *incoming* segment at $v_{b,j}$.




Figure 2: Construction of 1-bend drawing of K_n in Theorem 2.

Thus the bounding box is $\mathcal{O}(P) \times \mathcal{O}(n) \times \mathcal{O}(4^P Q + Q^2/P)$, which is $\mathcal{O}(\log n) \times \mathcal{O}(n) \times \mathcal{O}(n^{3/2}/\log n + n^2/\log^3 n)$, which is $\mathcal{O}(\log n) \times \mathcal{O}(n) \times \mathcal{O}(n^2/\log^3 n)$. Hence the volume is $\mathcal{O}(n^3/\log^2 n)$. It remains to prove that there are no edge crossings. By Corollary 1 all edges below the $Z = 0$ plane do not cross. We now only consider edges above the $Z = 0$ plane.

Each point in an outgoing segment at $v_{a,i}$ has an X -coordinate in $[2a, 2a+1]$. Thus an outgoing segment at some vertex v_{a_1,i_1} does not intersect an outgoing segment at some vertex v_{a_2,i_2} whenever $a_1 \neq a_2$. Clearly an outgoing segment at v_{a,i_1} is not coplanar with an outgoing segment at v_{a,i_2} whenever $i_1 \neq i_2$, and thus these segments do not cross. Since each bend is assigned a unique gridpoint, any two outgoing segments at the same vertex $v_{a,i}$ do not cross. Thus no two outgoing segments cross.

Each point in an incoming segment at $v_{b,j}$ has a Y -coordinate of $bQ + j$. Thus incoming segments at distinct vertices do not cross. Since each bend is assigned a unique gridpoint, any two incoming segments at the same vertex do not cross. Thus no two incoming segments cross.

To prove that an incoming segment does not cross an outgoing segment, we claim that in the projection of the edges on the $Y = 0$ plane, an incoming segment does not cross an outgoing segment. In the remainder of the proof we work solely in the $Y = 0$ plane, and use (X, Z) coordinates.

The projection in the $Y = 0$ plane of an outgoing segment at a vertex $v_{a,i}$ is the segment

$$s_1 = (2a, 0) \rightarrow (2a + 1, 4^{P-a} Q - i) .$$

The projection in the $Y = 0$ plane of the incoming segment of an edge $(v_{c,k}, v_{d,\ell})$ is the segment

$$s_2 = (2c + 1, 4^{P-c} Q - k) \rightarrow (2d, 0) .$$

For there to be a crossing clearly we must have $c < a < d$. To prove that there is no crossing it suffices to show that the Z -coordinate of s_2 is greater than the Z -coordinate of s_1 when $X = 2a + 1$. Now s_2 is contained in the line

$$Z = \frac{4^{P-c} Q - k}{2c + 1 - 2d} (X - 2d) .$$

Thus the Z -coordinate of s_2 at $X = 2a + 1$ is at least

$$\frac{4^{P-c} Q - Q}{2c + 1 - 2d} (2a + 1 - 2d) .$$

Thus it suffices to prove that

$$\frac{4^{P-c} Q - Q}{2c + 1 - 2d} (2a + 1 - 2d) > 4^{P-a} Q . \quad (1)$$

Clearly (1) is implied if it is proved with $a = c + 1$ and $d = c + 2$. In this case, (1) reduces to

$$\frac{4^{P-c} - 1}{3} > 4^{P-c-1} .$$

That is, $4^{P-c-1} > 1$, which is true since $c \leq P - 2$. This completes the proof.

Acknowledgements

Thanks to Stephen Wismath for suggesting the problem.

References

- [1] PROSENJIT BOSE, JUREK CZYZOWICZ, PAT MORIN, AND DAVID R. WOOD. The maximum number of edges in a three-dimensional grid-drawing. *J. Graph Algorithms Appl.*, 8(1):21–26, 2004.
- [2] TIZIANA CALAMONERI AND ANDREA STERBINI. 3D straight-line grid drawing of 4-colorable graphs. *Inform. Process. Lett.*, 63(2):97–102, 1997.
- [3] ROBERT F. COHEN, PETER EADES, TAO LIN, AND FRANK RUSKEY. Three-dimensional graph drawing. *Algorithmica*, 17(2):199–208, 1996.
- [4] GUISEPPE DI BATTISTA, PETER EADES, ROBERTO TAMASSIA, AND IOANNIS G. TOLLIS. *Graph Drawing: Algorithms for the Visualization of Graphs*. Prentice-Hall, 1999.
- [5] EMILIO DI GIACOMO. Drawing series-parallel graphs on restricted integer 3D grids. In [19], pp. 238–246.
- [6] EMILIO DI GIACOMO AND HENK MEIJER. Track drawings of graphs with constant queue number. In [19], pp. 214–225.
- [7] JOSEP DÍAZ, JORDI PETIT, AND MARÍA SERNA. A survey of graph layout problems. *ACM Comput. Surveys*, 34(3):313–356, 2002.
- [8] ROBERT P. DILWORTH. A decomposition theorem for partially ordered sets. *Ann. of Math.* (2), 51:161–166, 1950.
- [9] VIDA DUJMOVIĆ, PAT MORIN, AND DAVID R. WOOD. Layout of graphs with bounded tree-width. *SIAM J. Comput.*, 34(3):553–579, 2005.
- [10] VIDA DUJMOVIĆ AND DAVID R. WOOD. On linear layouts of graphs. *Discrete Math. Theor. Comput. Sci.*, 6(2):339–358, 2004.
- [11] VIDA DUJMOVIĆ AND DAVID R. WOOD. Stacks, queues and tracks: Layouts of graph subdivisions. *Discrete Math. Theor. Comput. Sci.*, to appear. Layouts of graph subdivisions. In JÁNOS PACH, ed., *Proc. 12th International Symp. on Graph Drawing (GD '04)*, vol. 3383 of *Lecture Notes in Comput. Sci.*, pp. 133–143. Springer, 2004.
- [12] VIDA DUJMOVIĆ AND DAVID R. WOOD. Three-dimensional grid drawings with sub-quadratic volume. In JÁNOS PACH, ed., *Towards a Theory of Geometric Graphs*, vol. 342 of *Contemporary Mathematics*, pp. 55–66. Amer. Math. Soc., 2004.
- [13] B. DYCK, J. JOEVENAZZO, E. NICKLE, J. WILSDON, AND STEPHEN K. WISMATH. Drawing K_n in three dimensions with two bends per edge. Tech. Rep. TR-CS-01-04, Department of Mathematics and Computer Science, University of Lethbridge, 2004.
- [14] STEFAN FELSNER, GIUSEPPE LIOTTA, AND STEPHEN K. WISMATH. Straight-line drawings on restricted integer grids in two and three dimensions. *J. Graph Algorithms Appl.*, 7(4):363–398, 2003.

- [15] GODFREY H. HARDY AND EDWARD M. WRIGHT. *An introduction to the theory of numbers*. Clarendon, 5th edn., 1979.
- [16] TORU HASUNUMA. Laying out iterated line digraphs using queues. In [19], pp. 202–213.
- [17] MICHAEL KAUFMANN AND DOROTHEA WAGNER, eds. *Drawing Graphs: Methods and Models*, vol. 2025 of *Lecture Notes in Comput. Sci.* Springer, 2001.
- [18] F. THOMSON LEIGHTON AND ARNOLD L. ROSENBERG. Three-dimensional circuit layouts. *SIAM J. Comput.*, 15(3):793–813, 1986.
- [19] GUISEPPE LIOTTA, ed. *Proc. 11th International Symp. on Graph Drawing (GD '03)*, vol. 2912 of *Lecture Notes in Comput. Sci.* Springer, 2004.
- [20] JÁNOS PACH, TORSTEN THIELE, AND GÉZA TÓTH. Three-dimensional grid drawings of graphs. In BERNARD CHAZELLE, JACOB E. GOODMAN, AND RICHARD POLLACK, eds., *Advances in discrete and computational geometry*, vol. 223 of *Contemporary Mathematics*, pp. 251–255. Amer. Math. Soc., 1999.
- [21] COLIN WARE AND GLENN FRANCK. Viewing a graph in a virtual reality display is three times as good as a 2D diagram. In ALLEN L. AMBLER AND TAKYUKI D. KIMURA, eds., *Proc. IEEE Symp. Visual Languages (VL '94)*, pp. 182–183. IEEE, 1994.
- [22] COLIN WARE AND GLENN FRANCK. Evaluating stereo and motion cues for visualizing information nets in three dimensions. *ACM Trans. Graphics*, 15(2):121–140, 1996.
- [23] COLIN WARE, DAVID HUI, AND GLENN FRANCK. Visualizing object oriented software in three dimensions. In *Proc. IBM Centre for Advanced Studies Conf. (CASCON '93)*, pp. 1–11. 1993.