Reducción de Problema

Autor

1 Hamiltonian Cycle

We will consider de problem of the Hamiltonian cycle of undirected graphs. Therefore an edge would have two 'sources' instead of a source and a target as it is regarded on directed graphs. Prior to the reduction a little lemma will be prove.

Lemma 1.1. Being $G = (V = \{v_1, ..., v_n\}, E = \{e_1, ..., e_m\})$ a graph. The set $\{e_{i_1}, ..., e_{i_n}\} \subset E$ is a hamiltonian cycle if, and only if, each vertex is the source to exactly two edges and the path $\{e_{i_1}, ..., e_{i_n}\} \subset E$ is connected.

Proof. If each vertex is the source of an edge, then every vertex is accesible by an edge. Also, as every vertex has exactly two edges, each connected component of the graph would be a clycle. As the graph is connected there is only one of such components.

In order to make the reducction we will represent with boolean clauses these two condition:

• We will start defining the variables $e_1, ..., e_n$ that will represent if the edge e_i is choose for the path. Also, if a vertex e_i has as sources v_j, v_k then the variables e_{i,v_j} and e_{i,v_k} will be also defined. The first set of formulas to consider will be:

$$e_i \iff e_{i,v_j} \iff e_{i,v_k} \qquad \forall i \in 1,...,m, \forall j,k \in 1,...,n$$

Note than if e_j does not have as source v_j then $e_{j,v_j} \iff 0$. To ensure that each vertex is the source of exactly two edges we will define these clauses:

$$\bigwedge_{k=1}^{m} \left(\bigwedge_{i=1}^{m} \bigvee_{\substack{j=1\\j\neq i}}^{m} e_{j,v_k} \right)$$

In order two ensure that each vertes is source to at least two edges. Then to ensure that there would not be more than two:

$$\bigwedge_{h=1}^{m} \bigwedge_{\substack{i=1\\j=1\\k=1}}^{n} \neg e_{i,v_h} \lor \neg e_{j,v_h} \lor \neg e_{k,v_h}$$

• To prove the connectivity we will use the connectivity matrix. Henceforth all matrix will be consider as $n \times n$ -sized matrix. Given $A = (a_{i,j})$ such that $a_{i,j} = 1$ if, and olny if, there is a edge between v_i and v_j , otherwise $a_{i,j} = 0$. Then consider $A^k = (a_{i,j}^*)$, it happens that if $(a_{i,j}^*) = 1$ then there is a path of exactly length k. Then to check the connectivity

we will define $A' = \sum_{i=0}^n A^i$ and defining the formula:

$$(a'_{1,1} \wedge \dots \wedge a'_{1,n})$$
 (1)

Matriz product could be seen as a boolean operation (for the purpose that we reach): Given $A = (a_{i,j}), B = (b_i, j)$ and $C = A \cdot B$ then

$$c_{i,j} = (a_{i,1} \wedge b_{1,j}) \vee \dots \vee (a_{i,n} \wedge b_{n,j})$$

As we do not care about the exact value of the sum in A' but only whether $a'_{i,j}$ is greater than 0 we could condider as sum the or operation element-wise. This prove that the expresion (1) is a formula, a bit laborious to do by hand but quite computable.