Statistical Inference and Two-Phase Audit Design for Contaminated Measurements in Clinical Research: The catmech R Package

Patrick Iben
Independent Biostatistician
patrickiben@gmail.com | ORCID: 0009-0005-2443-7973

November 6, 2026

Abstract

Continuous measurements in multicenter clinical trials, electronic health record (EHR) disease registries, and observational epidemiological cohorts are frequently compromised by transcription mistakes, unit conversion errors, assay calibration shifts, and missing values. Traditional statistical packages address missing covariates under Missing at Random (MAR) assumptions, discarding untrusted observations and imputing from scratch. In source-data verification (SDV) and clinical audit designs, however, recorded values are present but unreliable, and a gold-standard reference measurement is audited on only a fraction of records. Naive analysis of recorded values suffers attenuation bias, complete-case analysis of audited records inflates standard errors, and standard multiple imputation needlessly discards the precision of valid recordings. This paper presents `catmech`, an R package providing a unified, auditable framework for clinical measurement verification, two-phase prospective audit power planning, predictive calibration checking, downstream multi-endpoint regression pooling, and regulatory audit trail generation. We derive the analytical conditional posteriors for reference-calibrated contamination mixture models, formalize prospective power calculation using two-phase influence function expansions, and establish cluster-robust covariance estimators with Barnard-Rubin finite-sample degrees of freedom for linear models (CV3 jackknife), generalized linear models (CR1S Huber-White-Liang-Zeger sandwich), and Cox proportional hazards survival models (event-based model covariance and clustered Lin-Wei sandwich). We evaluate the package across 10 frozen validation archives (over 500,000 requests), including the synthetic contamination study (52,950 requests), two public paired-discrepancy benchmarks on ClinicalTrials.gov ($N = 27,208$) and NHANES 2017–2018 ($N = 3,908$; 33,800 requests), clustered variance evaluations (148,500 requests), and facility-level recounts from the Rwanda Health Management Information System and psychology effect-size recounts (45,600 requests). Retaining recorded measurements reduces root mean squared error by 43% to 48% over discarding unaudited values on ClinicalTrials.gov and NHANES. We articulate an evidence-based decision rule and report all pre-specified negative and inconclusive findings: the contamination mixture model avoids severe attenuation bias and narrows confidence intervals by 44% relative to complete-case analysis under simulated gross, separable errors, whereas simple linear imputation is preferred under subtle, diffuse real-world discrepancies. Finally, `catmech_audit_trail()` records SHA-256 hashes of the input data, assumptions, and completed datasets to support documentation under FDA 21 CFR Part 11 and ICH E9; it does not by itself make an analysis system compliant.

1 Introduction

In multicenter randomized clinical trials, real-world electronic health record (EHR) registries, and epidemiological cohorts, continuous biological and physiological measurements are central to primary and secondary study endpoints. Circulating biomarkers (e.g., cardiac troponin, serum creatinine, viral load), hemodynamic measurements (e.g., systolic blood pressure), and radiographic dimensions (e.g., tumor response diameters) guide clinical decisions and estimate treatment effects. However, continuous recordings in multi-site clinical registries are inevitably subject to data contamination, including assay lot calibration shifts, clerical transcription errors, decimal slips, and inconsistent unit conversions [Carroll et al., 2006, Angelopoulos et al., 2023].

To safeguard data integrity, clinical trial sponsors and regulatory authorities mandate source-data verification (SDV) and targeted clinical audits, wherein trial monitors review primary source documents (e.g., clinical chart re-abstractions, central laboratory re-assays) for a subsample of patients [U.S. Food and Drug Administration, 2003, International Conference on Harmonisation, 1998]. When discrepancies between the originally recorded measurement and the audited reference standard are uncovered, biostatisticians and trialists confront an inferential dilemma:

1. **Naively trusting recorded measurements**: Treating recorded numbers as exact introduces classic errors-in-variables. For continuous exposures, non-differential measurement error attenuates regression coefficients toward the null, obscuring true treatment effects and biological gradients [Carroll et al., 2006]. For non-linear models (e.g., logistic regression or Cox proportional hazards), measurement contamination causes complex residual confounding and severe coverage failure.
2. **Restricting analysis to audited cases (complete-case analysis)**: Analyzing only the audited subset ($\mathcal{A} \subset \{1, \ldots, N\}$, typically 10% to 25% of the cohort) protects against contamination bias but drastically reduces effective sample size, inflating standard errors, widening confidence intervals, and sacrificing statistical power [Neyman, 1938, Breslow et al., 2003].
3. **Discarding recorded values and applying standard multiple imputation**: Imputing unaudited values under standard Missing at Random (MAR) assumptions using packages such as `mice` [van Buuren and Groothuis-Oudshoorn, 2011, White et al., 2011] or `Amelia` [Honaker et al., 2011] avoids bias under correct specification. However, treating recorded values as completely unobserved discards the valuable information present in records that were recorded accurately [Rubin, 1996].

To overcome these limitations, we introduce `catmech`, an R package for auditable measurement verification, two-phase clinical audit power planning, predictive calibration diagnostics, and multi-endpoint model pooling. The package unites two-phase sampling theory [Cochran, 1977, Breslow et al., 2003, Robins et al., 1994] with reference-calibrated Bayesian mixture modeling [Huang and Wand, 2013], Barnard-Rubin finite-sample inference [Barnard and Rubin, 1999], and cryptographic data provenance [U.S. Food and Drug Administration, 2003].

2 Statistical theory and methodology

2.1 The data generating and contamination framework

Let $i \in \{1, \ldots, N\}$ index patients enrolled across C clinical sites. Let $T_i \in \mathbb{R}$ denote the latent true biological measurement. Conditional on clinical covariates $X_i \in \mathbb{R}^p$ (including the analysis outcome and predictors in the imputation model), the latent measurement follows:

$$T_i \mid X_i \sim \mathcal{N}(X_i\beta, \sigma_T^2)$$

Let $\delta_i \in \{0, 1\}$ represent an unobserved recording contamination indicator, where $\delta_i = 1$ indicates that the recorded value was corrupted during observation or transcription. The contamination probability $\pi_{s_i} = \mathbb{P}(\delta_i = 1 \mid S_i = s_i)$ may depend on facility strata or clinical monitoring quality flags S_i (`corruption_strata`), and the additive error distribution may vary across error strata g_i (`error_strata`). Writing (π, μ_e, σ_e^2) for a single stratum to simplify notation, the observed recorded measurement W_i is generated as:

$$W_i = (1 - \delta_i)T_i + \delta_i(T_i + \epsilon_i), \quad \epsilon_i \sim \mathcal{N}(\mu_e, \sigma_e^2)$$

where ϵ_i represents an additive error mechanism with mean recording bias μ_e and error variance σ_e^2.

In Phase II of the audit design, an independent gold-standard reference measurement R_i is audited on a subset $\mathcal{A} \subset \{1, \ldots, N\}$ with sample size $n_{\text{audit}} = |\mathcal{A}|$:

$$R_i = T_i + \eta_i, \quad \eta_i \sim \mathcal{N}(0, \sigma_{\text{ref}}^2)$$

where σ_{ref} (`reference_sd`, defaulting to 0) is a user-declared sensitivity constant representing known reference assay noise, with $\sigma_{\text{ref}}^2 = 0$ corresponding to an exact recount.

2.2 Collapsed slice-within-Gibbs sampling posteriors and linear reference imputation

In `catmech_contamination()`, the estimated parameter vector is $\Theta = (\beta, \sigma_T^2, \pi, \mu_e, \sigma_e^2)$. For exact recounts ($\sigma_{\text{ref}} = 0$), the target regression parameters receive the standard noninformative Jeffreys prior $p(\beta, \sigma_T^2) \propto 1/\sigma_T^2$ (or a flat prior on β with a scaled inverse-χ^2 prior on σ_T^2 when $\sigma_{\text{ref}} > 0$). Contamination probabilities receive Beta priors $\pi \sim \text{Beta}(a_\pi, b_\pi)$, and recording error parameters receive a normal-scaled-inverse-χ^2 prior $\mu_e \mid \sigma_e^2 \sim \mathcal{N}(\mu_0, \sigma_e^2/\kappa_0)$ and $\sigma_e^2 \sim \text{Scaled-Inv-}\chi^2(\nu_0, \tau_0^2)$. During Markov Chain Monte Carlo (MCMC) estimation in R, the latent indicators and true values (δ_i, T_i) on unaudited rows are analytically integrated out when updating $\text{logit}(\pi)$, μ_e, and $\frac{1}{2}\log(\sigma_e^2)$ via univariate stepping-out slice sampling (supplemented by a scale-ridge slice step when fewer than 5 discrepancies are observed), followed by QR-based conditional draws of (β, σ_T^2) and exact conditional draws of (δ_i, T_i).

Specifically, for unaudited patients ($i \notin \mathcal{A}$), integrating out the unobserved latent value $T_i \mid X_i \sim \mathcal{N}(X_i\beta, \sigma_T^2)$ under $\delta_i = 1$ yields the convolved marginal predictive distribution $W_i \mid \delta_i = 1, X_i, \Theta \sim \mathcal{N}(X_i\beta + \mu_e,\, \sigma_T^2 + \sigma_e^2)$. The conditional posterior probability of recording corruption $p_i = \mathbb{P}(\delta_i = 1 \mid W_i, X_i, \Theta)$ is therefore evaluated analytically at each Gibbs iteration as:

$$p_i = \frac{\pi\phi(W_i;\, X_i\beta + \mu_e,\, \sigma_T^2 + \sigma_e^2)}{\pi\phi(W_i;\, X_i\beta + \mu_e,\, \sigma_T^2 + \sigma_e^2) + (1-\pi)\phi(W_i;\, X_i\beta,\, \sigma_T^2)}$$

where $\phi(w;\, m, v)$ denotes the Gaussian probability density function evaluated at w with mean m and variance v.

Conditional on drawing $\delta_i = 0$ for an unaudited record, $T_i = W_i$ deterministically. Conditional on drawing $\delta_i = 1$, T_i is sampled from the conjugate Gaussian conditional posterior:

$$T_i \mid \delta_i = 1, W_i, X_i, \Theta \sim \mathcal{N}(\mu_{T|W}, \sigma_{T|W}^2)$$

$$\mu_{T|W} = \sigma_{T|W}^2 \left(\frac{X_i\beta}{\sigma_T^2} + \frac{W_i - \mu_e}{\sigma_e^2} \right), \quad \sigma_{T|W}^2 = \left(\frac{1}{\sigma_T^2} + \frac{1}{\sigma_e^2} \right)^{-1}$$

For audited patients ($i \in \mathcal{A}$), R_i directly anchors T_i (Setting $T_i = R_i$ when $\sigma_{\text{ref}} = 0$, or drawing from the Gaussian precision-weighted combination of R_i, W_i, and $X_i\beta$ when $\sigma_{\text{ref}} > 0$). Multiple imputations $T^{(1)}, \ldots, T^{(m)}$ are drawn from post-burn-in iterations after verifying Gelman-Rubin $\hat{R}$ and effective sample size convergence thresholds [Gelman and Rubin, 1992].

As a parsimonious alternative when discrepancies are diffuse rather than discrete, `catmech_reference()` implements Gaussian reference imputation accelerated in C++17 via `RcppArmadillo`. When called with `use_recorded = TRUE` (referred to in our benchmarks as Simple Linear MI, or `cole_mi`), `catmech_reference()` includes the recorded value W_i directly as a continuous linear predictor alongside X_i in the audited calibration regression $T_i \mid W_i, X_i \sim \mathcal{N}(\gamma_0 + \gamma_W W_i + X_i\gamma_X, \sigma_{T|W,X}^2)$ without positing a discrete contamination indicator δ_i. When called with `use_recorded = FALSE` (the v0.3 default, `reference_normal`), `catmech_reference()` discards W_i on unaudited rows and imputes T_i solely from X_i.

2.3 Two-phase audit sample size and power planning

Prior to re-abstracting medical charts or commissioning secondary lab assays, investigators must budget the audit sample size n_{audit}. Using semiparametric influence function theory for two-phase estimators [Neyman, 1938, Cochran, 1977, Breslow et al., 2003, Angelopoulos et al., 2023], the asymptotic variance of an efficient parameter estimator at audit sampling fraction $f = n_{\text{audit}}/N$ satisfies:

$$\text{Var}(\hat{\theta}_f) = \frac{1}{N}\text{Var}(\psi_i) + \frac{1-f}{fN}\text{Var}(\psi_i - \mathbb{E}[\psi_i \mid G_i])$$

where ψ_i is the efficient influence function of θ and $G_i = (W_i, Y_i, X_i)$ denotes the Phase I auxiliary variables observed on the full cohort (or $G_{0,i} = (Y_i, X_i)$ when discarding W_i).

The precision ratio $\rho(f)$, defined as the ratio of confidence interval width at audit fraction f relative to an ideal census audit ($f = 1$), is:

$$\rho(f) = \frac{\text{Width}(f)}{\text{Width}(1)} = \sqrt{1 + k\frac{1-f}{f}}, \quad k = \frac{\text{Var}(\psi_i - \mathbb{E}[\psi_i \mid G_i])}{\text{Var}(\psi_i)}$$

Because the noise parameter k is estimated with uncertainty, the realized precision ratio $\hat{\rho}(f)$ is stochastic. `catmech_audit_power()` evaluates the statistical power to achieve a target precision ratio $\rho \leq \rho_{\text{target}}$:

$$\text{Power}(n_{\text{audit}}) = \mathbb{P}\left(\sqrt{1 + \hat{k}\frac{1-f}{f}} \leq \rho_{\text{target}}\right)$$

by simulating log-normal Monte Carlo draws (`n_sim = 2000`) of $\hat{k}$, either from prospective error-rate and correlation parameters or calibrated to the bias-corrected and accelerated (BCa) bootstrap interval bounds produced by `catmech_audit_plan()`. In pre-specified validation testing, `catmech_audit_plan()` returned a `NO-GO` verdict because its median error in predicting interval width at a 5% audit fraction was 11.3% to 16.8% (exceeding the pre-specified 10% tolerance); accordingly, `catmech_audit_plan()` is documented as an exploratory calculation rather than a validated sample-size guarantee.

2.4 Multi-endpoint model pooling with Barnard-Rubin degrees of freedom

Parameter estimates $\hat{\theta}_j$ and variance-covariance matrices $\hat{U}_j$ across $j \in \{1, \ldots, m\}$ completed datasets are combined using Rubin's rules [Rubin, 1987]:

$$\bar{\theta} = \frac{1}{m}\sum_{j=1}^{m}\hat{\theta}_j, \quad \bar{U} = \frac{1}{m}\sum_{j=1}^{m}\hat{U}_j, \quad B = \frac{1}{m-1}\sum_{j=1}^{m}(\hat{\theta}_j - \bar{\theta})(\hat{\theta}_j - \bar{\theta})^\top$$

$$T = \bar{U} + \left(1 + \frac{1}{m}\right)B, \quad \hat{\lambda} = \frac{(1 + 1/m)\text{diag}(B)}{\text{diag}(T)}$$

Standard Rubin degrees of freedom $\nu_m = (m-1)\hat{\lambda}^{-2}$ assume infinite complete-data sample size. In clinical audits where sample sizes, cluster counts, or event counts are constrained, this assumption leads to anti-conservative confidence intervals. `catmech` implements the Barnard & Rubin (1999) small-sample adjustment [Barnard and Rubin, 1999]:

$$\nu_{obs} = \left(\frac{\nu_{com} + 1}{\nu_{com} + 3}\right)\nu_{com}(1 - \hat{\lambda}), \quad \nu_{BR} = \frac{\nu_m \nu_{obs}}{\nu_m + \nu_{obs}}$$

where ν_{com} is the complete-data degrees of freedom. Downstream pooling covers three core clinical endpoint families:

1. **Continuous endpoints (`catmech_pool_lm`)**: When `cluster = NULL` (unclustered), fits ordinary least squares models with standard covariance and $\nu_{com} = N - p$. When `cluster` is specified across C clinical sites, computes the closed-form delete-one-cluster jackknife (`cv3`) covariance $\frac{C-1}{C}\sum_{c=1}^{C}(\hat{\beta}_{(-c)} - \hat{\beta})(\hat{\beta}_{(-c)} - \hat{\beta})^\top$ on each completed dataset and pools with $\nu_{com} = C - 1$.
2. **Binary and count endpoints (`catmech_pool_glm`)**: Fits generalized linear models (logistic and Poisson). When `cluster = NULL`, uses standard inverse-information covariance with $\nu_{com} = N - p$. When `cluster` is specified across C clinical sites, sets $\nu_{com} = C - 1$ and computes the CR1S finite-sample-corrected Huber–White–Liang–Zeger cluster-robust sandwich covariance [Liang and Zeger, 1986]:

   $$\hat{V}_{\text{sandwich}} = \frac{C}{C-1}\frac{N-1}{N-p}\hat{J}^{-1}\left(\sum_{c=1}^{C}s_c s_c^\top\right)\hat{J}^{-1}$$

   where $s_c = \sum_{i \in \text{cluster } c}\frac{\partial \ell_i}{\partial \theta}$ are cluster score residuals and $\hat{J}^{-1}$ is the unclustered GLM covariance matrix. Point estimates and intervals are optionally exponentiated to yield odds ratios or incidence rate ratios.

3. **Time-to-event survival endpoints (`catmech_pool_coxph`)**: Fits Cox proportional hazards models via `survival::coxph()`. When `cluster = NULL`, uses the model-based inverse partial Fisher information matrix and defines complete-data degrees of freedom strictly as $\nu_{com} = d - p$, where d is the number of observed failure events and p is the parameter count. When `cluster` is specified across C clinical sites, computes the cluster-summed Lin & Wei (1989) score-residual sandwich covariance [Lin and Wei, 1989] and sets $\nu_{com} = C - 1$. Hazard ratios $\mathrm{HR} = \exp(\bar{\theta})$ and confidence intervals are computed using the Barnard-Rubin ν_{BR} t-distribution.

2.5 Out-of-sample predictive calibration checking

To screen for predictive miscalibration, `catmech_holdout_check()` partitions audited records into K stratified (or cluster-grouped) cross-validation folds and refits the imputation workflow on the remaining $K - 1$ folds $\mathcal{D}_{-k}$. Because the mixture predictive distribution has a discrete point mass at W_i (retained with probability $1 - p_i$) and uses m finite completions, `catmech_holdout_check()` computes the tie-randomized empirical rank Probability Integral Transform (PIT) [Dawid, 1984] for each held-out reference record $i \in \mathcal{A}_k$:

$$u_i = \frac{\sum_{j=1}^{m} \mathbb{I}(T_{i,-k}^{(j)} < R_i^*) + U_i \left(\sum_{j=1}^{m} \mathbb{I}(T_{i,-k}^{(j)} = R_i^*) + 1 \right)}{m + 1}, \quad U_i \sim \mathrm{Uniform}(0, 1)$$

where $R_i^* = W_i$ when $|W_i - R_i|$ lies within `agreement_tol` under exact recounts ($\sigma_{\mathrm{ref}} = 0$) and $R_i^* = R_i$ otherwise. The diagnostic flags `miscalibration_detected` if a two-sided Kolmogorov–Smirnov test of u_i against Uniform$(0, 1)$ yields $p < 0.01$, if the 95% Wilson upper bound on empirical predictive interval coverage falls below the nominal level, or if the corruption probability calibration z-statistic exceeds $z_{0.995}$. As shown in our empirical validation below, held-out row checks have low statistical power to detect regression slope bias, so `catmech_holdout_check()` is a standalone diagnostic screen rather than a proof of model validity.

3 Software architecture and comparison with existing packages

The `catmech` package is structured into modular S3 components designed for computational efficiency, numerical stability, and regulatory auditability:

Module	Core functions	Mathematical and computational role
Audit planning	`catmech_audit_plan()`, `catmech_audit_power()`	Two-phase influence function ratio estimation (exploratory) and Monte Carlo power curves.
Data diagnostics	`catmech_recording_audit()`, `catmech_holdout_check()`	Stratum-specific discrepancy auditing (Jeffreys beta intervals) and out-of-sample rank-PIT checks.
Imputation engines	`catmech_contamination()`, `catmech_reference()`	Reference-calibrated collapsed slice-within-Gibbs mixture model and Gaussian reference MI (`RcppArmadillo`).
Model pooling	`catmech_pool_lm()`, `catmech_pool_glm()`, `catmech_pool_coxph()`, `catmech_compare_strategies()`	Multi-endpoint pooling with Rubin rules, Barnard-Rubin df, cluster-robust covariance, and strategy comparison.
Provenance	`catmech_sensitivity()`, `catmech_audit_trail()`	Multi-scenario sensitivity envelopes and SHA-256 provenance reports for regulatory documentation.

While several mature R packages address missing data or measurement error, none combine two-phase audit power planning, reference-calibrated contamination mixture modeling, multi-endpoint pooling with cluster-robust covariance and Barnard-Rubin degrees of freedom, and cryptographic provenance tracking:

Capability	`catmech`	`mice` [van Buuren and Groothuis-Oudshoorn, 2011]	`simex` [Lederer and Küchenhoff, 2006]	`mecor` [Nab et al., 2021]	`Amelia` [Honaker et al., 2011]
Contamination Mixture	**Yes** (Gibbs)	No	No	No	No
Gold-Standard Audits	**Yes** (Two-phase)	Partial (MAR)	No (Replicates)	Yes (Reg. Cal.)	Partial (Priors)
Audit Power Planning	**Yes** (MC)	No	No	No	No
Calibration Screen	**Yes** (Rank PIT)	No	No	No	Partial (`overimpute`)
Linear Model Pooling	**Yes** (BR + CV3)	Yes (BR df)	Yes (SIMEX)	Yes (Reg. Cal.)	Yes (Rubin df)
GLM (Logit/Poisson)	**Yes** (Sandwich)	Yes (BR df)	Yes (SIMEX)	No	Yes (Rubin df)
Cox PH Pooling	**Yes** (Lin-Wei)	Partial (`coxph`)	No	No	No
SHA-256 Audit Trail	**Yes**	No	No	No	No

In confirmatory clinical trials, primary estimands are frequently time-to-event survival endpoints or binary response endpoints rather than continuous linear outcomes. All `catmech` pooling functions export `broom`-style S3 methods (`tidy()`, `glance()`) and S3 forest-plot visualization (`plot()`), while `catmech_compare_strategies()` compares four benchmark strategies (`validation_subsample` complete-case, `naive_trust_recorded` naive analysis, `linear_mi_recorded` using `catmech_reference(use_recorded = TRUE)`, and `contamination_model` using `catmech_contamination()`) side by side for continuous (`lm`) and generalized linear (`glm`) endpoints.

4 Illustrative clinical trial workflow

The following reproducible R workflow illustrates the `catmech` pipeline across all eight analytical stages on a simulated 10-site clinical cohort ($N = 1,000$, 25% reference audit $n_{\text{audit}} = 250$):

```r
library(catmech)
library(survival)

# 1. Simulate a 10-site cohort with recording contamination and a 25% reference audit
set.seed(42)
cohort <- simulate_catmech_lab(n = 1000, seed = 42)
cohort$site <- rep(seq_len(10), length.out = nrow(cohort))
audited <- sample.int(nrow(cohort), size = 250)
cohort$lab_ref <- ifelse(seq_len(nrow(cohort)) %in% audited, cohort$lab_true, NA_real_)
cohort$adverse_event <- as.integer(cohort$outcome > median(cohort$outcome))
cohort$time <- rexp(nrow(cohort), rate = exp(0.25 * (cohort$lab_true - 6)))
cohort$event <- rbinom(nrow(cohort), 1, 0.75)

# 2. Prospective audit sample size and statistical power planning
power_plan <- catmech_audit_power(
  N = 1000, target_ratio = 1.25, power_target = 0.80,
  error_rate = 0.20, correlation = 0.85, fractions = seq(0.10, 0.50, by = 0.05)
```

```
)

# 3. Discrepancy screening across clinical sites (exact 95% Jeffreys beta intervals)
audit_screen <- catmech_recording_audit(
  data = cohort, value = "lab_value", reference = "lab_ref",
  by = "site", agreement_tol = 0.5
)

# 4. Reference-calibrated contamination mixture model (or catmech_reference(use_recorded = TRUE))
fit <- catmech_contamination(
  data = cohort, value = "lab_value", reference = "lab_ref",
  predictors = c("age", "sex", "inflammation", "outcome"),
  corruption_strata = "outlier_flag", m = 10, seed = 2026
)

# 5. Out-of-sample predictive calibration check (5-fold cross-validated rank PIT)
calib_check <- catmech_holdout_check(fit = fit, folds = 5L, m = 20L, seed = 123L)

# 6. Multi-endpoint downstream regression pooling (LM with CV3, GLM with CR1S, Cox PH with Lin-Wei)
pool_continuous <- catmech_pool_lm(
  result = fit, formula = outcome ~ lab_value + age + sex, cluster = "site"
)
pool_binary <- catmech_pool_glm(
  result = fit, formula = adverse_event ~ lab_value + age + sex,
  family = binomial(link = "logit"), cluster = "site", exponentiate = TRUE
)
pool_survival <- catmech_pool_coxph(
  result = fit, formula = Surv(time, event) ~ lab_value + age + sex,
  cluster = "site", exponentiate = TRUE
)

# 7. Side-by-side comparison of all four audit strategies for continuous or GLM endpoints
comp <- catmech_compare_strategies(
  data = cohort, value = "lab_value", reference = "lab_ref",
  formula = outcome ~ lab_value + age + sex, m = 10, seed = 2026
)

# 8. SHA-256 provenance report (supports FDA 21 CFR Part 11 / ICH E9 documentation)
report <- catmech_audit_trail(
  object = fit, pooled = pool_survival, format = "markdown", path = "audit_trail.md"
)
```

Printing `pool_survival` reports pooled log-hazard point estimates, clustered Lin-Wei sandwich standard errors, Barnard-Rubin degrees of freedom (with $\nu_{com} = C - 1 = 9$ for $C = 10$ sites), exponentiated hazard ratios (`estimate_exp`), and 95% confidence intervals:

```
-- Catmech Pooled Cox PH Model --
Completions (m): 10 | Observations (n): 1000 | Events: 735

      term estimate std.error     df   lower  upper estimate_exp lower_exp upper_exp
 lab_value   0.1632    0.0314 6.5933  0.0881 0.2384       1.1773    1.0921    1.2692
       age   0.0056    0.0041 7.4010 -0.0038 0.0151       1.0056    0.9962    1.0152
      sexM  -0.0541    0.0433 7.3074 -0.1555 0.0474       0.9474    0.8560    1.0485
```

5 Synthetic simulation studies and benchmarks

To evaluate the statistical properties of `catmech`, we conducted pre-specified Monte Carlo simulation studies across 10 validation archives (over 500,000 total requests). The primary reference-calibrated contamination study (`validation/contamination-study`) executed 52,950 planned requests across 183 method-scenario cells spanning 4 primary cells (`clean`, `flagged_errors`, `missed_flags`, and `site_shift_declared`, 1,000 cohorts each) and 23 stress cells (150 cohorts each), with cohort size $N = 600$, linear regression target slope $\beta = 0.800$ on the latent covariate T_i, a 25% reference audit ($n_{\text{audit}} = 150$), and $m = 20$ imputations. Seven core strategies were evaluated across the 27 scenarios (with 3 noisy-reference/assay-noise stress scenarios running only `contamination`, and 6 scenarios additionally evaluating standard and bootstrap-donor predictive mean matching `reference_pmm` and `reference_pmm_abb` to total 52,950 requests), yielding 52,900 returned intervals, 16 expected design refusals (14 in `reference_05pct` and 2 in `reference_10pct` due to sparse strata), 33 MCMC convergence refusals, and 1 singular raking fit.

5.1 Scenario 1: Unflagged, gross, and misspecified recording contamination

The table below compares the seven core strategies across the primary unflagged contamination scenario (`missed_flags`, 1,000 cohorts, where recording errors occur in unflagged records) and the mixed gross-and-small error stress scenario (`error_gross_and_small`, 150 cohorts, combining 10× gross transcription outliers with mild recording noise). The seven strategies are: **Oracle** (regressing on true T_i for all $N = 600$ rows), **Flag Masking** (`v02_mask_normal`, setting flagged observations $S_i = 1$ to `NA` for normal MI while trusting unflagged observations $S_i = 0$ as exact truth), **Complete-Case** (`gold_only`, analyzing only the $n_{\text{audit}} = 150$ audited rows), **Discard Recorded MI** (`reference_normal`, `catmech_reference(use_recorded = FALSE)`), **Simple Linear MI** (`cole_mi`, `catmech_reference(use_recorded = TRUE)`), **Generalized Raking** (`raking`, two-phase calibration estimator), and **Contamination Mixture** (`catmech_contamination()`):

Scenario & Strategy	$\hat{\beta}$	Bias (MCSE)	Emp SD	Mean SE	Cov.	Width	RMSE
Unflagged Errors ($N_{\text{sim}} = 1,000$)							
Oracle (Latent T Known)	0.7978	-0.0022 (0.0012)	0.0394	0.0411	95.7%	0.1614	0.0394
Flag Masking (`v02_mask_normal`)	0.7185	-0.0815 (0.0015)	0.0472	0.0444	55.9%	0.1750	0.0942
Complete-Case (`gold_only`)	0.7988	-0.0012 (0.0026)	0.0815	0.0837	95.7%	0.3309	0.0814
Discard Recorded MI (`reference_normal`)	0.7896	-0.0104 (0.0021)	0.0676	0.0684	95.4%	0.2769	0.0684
Simple Linear MI (`cole_mi`)	0.7806	-0.0194 (0.0018)	0.0557	0.0528	92.4%	0.2101	0.0590
Generalized Raking (`raking`)	0.7853	-0.0147 (0.0020)	0.0631	0.0589	92.9%	0.2331	0.0648
Contamination Mixture	**0.7952**	**-0.0048 (0.0014)**	**0.0446**	**0.0467**	**94.9%**	**0.1842**	**0.0449**
Gross & Small ($N_{\text{sim}} = 150$)							
Oracle (Latent T Known)	0.7970	-0.0030 (0.0034)	0.0411	0.0412	94.7%	0.1617	0.0411
Flag Masking (`v02_mask_normal`)	0.0609	-0.7391 (0.0055)	0.0674	0.0155	0.7%	0.0614	0.7422
Complete-Case (`gold_only`)	0.7962	-0.0038 (0.0064)	0.0787	0.0845	95.3%	0.3339	0.0785
Discard Recorded MI (`reference_normal`)	0.7890	-0.0110 (0.0055)	0.0671	0.0695	95.3%	0.2816	0.0678

Scenario & Strategy	$\hat{\beta}$	Bias (MCSE)	Emp SD	Mean SE	Cov.	Width	RMSE
Simple Linear MI (`cole_mi`)	0.5646	-0.2354 (0.0261)	0.3200	0.0617	64.7%	0.2562	0.3964
Generalized Raking (`raking`)	0.5156	-0.2844 (0.0281)	0.3436	0.0693	53.7%	0.2740	0.4451
Contamination Mixture	**0.7883**	**-0.0117 (0.0035)**	**0.0432**	**0.0436**	**94.0%**	**0.1714**	**0.0447**

In the `missed_flags` scenario, trusting unflagged recordings (`v02_mask_normal`) suffers severe attenuation bias (-0.0815), collapsing 95% coverage to 55.9%. Complete-case analysis (`gold_only`) and discarding recorded values (`reference_normal`) remain unbiased but forfeit precision (mean CI widths of 0.3309 and 0.2769). The `catmech` contamination mixture model recovers unbiased estimates (bias: -0.0048), achieves 94.9% coverage, and narrows confidence intervals by 33.5% relative to discarding recorded values (`reference_normal`) and by 44.3% relative to complete-case analysis (`gold_only`). When gross transcription errors are mixed with small discrepancies (`error_gross_and_small`), both simple linear MI (bias: -0.2354, coverage: 64.7%) and generalized raking (bias: -0.2844, coverage: 53.7%) break down because gross outliers corrupt the linear calibration slope, whereas `catmech_contamination()` preserves 94.0% coverage and low RMSE (0.0447).

Across the four primary 1,000-cohort cells (`clean` with no recording errors, `flagged_errors` where errors are confined to flagged rows, `missed_flags` where errors occur in unflagged rows, and `site_shift_declared` where site-level additive shifts are combined with occasional larger errors), `catmech_contamination()` met the pre-specified $[0.925, 0.975]$ Wilson coverage screen in three cells (96.5%, 96.3%, and 94.9% coverage) and narrowly missed the screen in `site_shift_declared` at 93.7% coverage (Wilson interval $[0.9202, 0.9505]$, bias -0.0085, MCSE 0.0016), while reducing mean interval width by 30% to 38% across all four primary cells relative to `reference_normal`.

To provide a balanced counterweight to the primary cells, the 23 stress cells (150 cohorts each) deliberately violated model assumptions under a pre-specified failure threshold of coverage $< 90.0\%$ or $|\text{bias}| > 0.030$. Under this rule, `catmech_contamination()` failed in three stress cells:

1. **Heavy-tailed target distribution (`heavy_tail_target`)**: Bias reached $+0.0330$ with 87.3% coverage because the Gaussian mixture model misclassifies genuine heavy-tailed extremes in T_i as recording errors and shrinks them toward $X_i\beta$; by contrast, `reference_normal` (93.3% coverage, bias $+0.0130$), `cole_mi` (92.0% coverage, bias -0.0003), and `raking` (92.0% coverage, bias -0.0060) held nominal coverage.
2. **Non-ignorable audit selection on latent truth with errors present (`nonignorable_missed`)**: When audit selection depended directly on the unobserved true value T_i in the presence of unflagged recording errors, `catmech_contamination()` suffered bias of -0.0506 and 79.3% coverage (whereas design-consistent `raking` achieved bias $+0.0070$ and 87.3% coverage). When records were clean (`nonignorable_clean`), coverage held at 92.5% but 17 of 150 fits (11.3%) were refused for slow MCMC mixing.
3. **Undeclared reference measurement noise (`noisy_undeclared`)**: When reference measurements R_i contained undeclared assay noise ($\sigma_{\text{ref}} > 0$ left at the default `reference_sd = 0`), bias was -0.0261 and coverage fell to 88.7% (triggering the near-universal-disagreement warning in all 150 cohorts). Declaring `reference_sd` (`noisy_declared`) restored bias to -0.0071 and coverage to 91.8% (147/150 returned, vs. 52.7% coverage for `reference_normal` and 38.0% for `cole_mi`).

Two additional stress regimes were borderline or exhibited elevated refusal rates: bimodal two-sided errors (`error_bimodal`) was right at the boundary with bias -0.0293 and 90.0% coverage; range-check flags computed directly from recorded values (`flag_from_recorded`) had 10 of 150 (6.7%) convergence refusals (94.3% coverage among returned fits); and sparse 5% reference audits (`reference_05pct`) had 17 of 150 (11.3%) refusals (14 unsupported-stratum design refusals and 3 convergence refusals, with 91.7% coverage among returned intervals). Finally, in an 8-scenario evaluation of `catmech_holdout_check()` (40 cohorts per

cell), false alarms were absent under the true model (0/40 in `clean` and `missed_flags`), but alarm power against structural misspecification was low: it flagged 18 of 40 cohorts (45%) in `site_shift_undeclared`, only 1 of 40 in `heavy_tail_target` (with 6 unresolved fold failures), and 0 of 40 in `nonlinear_target`, because held-out rows test observation-level prediction rather than regression slope bias.

5.2 Scenario 2: Finite-sample variance and clustered multilevel imputation

In multicenter trials with C clinical sites, facility-level clustering induces correlation that distorts naive variance estimators. We evaluated 148,500 requests (148,486 returned intervals and 14 refusals: 12 rank-deficient audited designs on `iron` and 2 zero-residual-variance fits on `stillbirths`) across simulated clustered cohorts and facility-level recounts of 11 maternal and child health indicators from the Rwanda Health Management Information System (HMIS) [Nshimyiryo et al., 2020]. Standard small-sample cluster-robust sandwich standard errors (`cr1s`) and Bell-McCaffrey / Pustejovsky-Tipton bias-reduced linearization with Satterthwaite degrees of freedom (`cr2`, which inflated interval widths by up to 95% under random-intercept imputation and passed only 23 of 44 simulated cells) failed the pre-specified eligibility rules. The delete-one-cluster jackknife (`cv3`) restored 100% complete-data recount coverage (`gold_only`) to 92.0%–97.5% across the 11 Rwanda HMIS indicators and narrowly passed pre-specified eligibility under random-intercept outcome imputation (`catmech_reference(cluster = )`, achieving $\geq 92.0\%$ coverage on 9 of 11 Rwanda indicators, with `syphilis` at 90.0% and `iron` at 91.9%, and passing 36 of 44 simulated cells while reducing mean distance from nominal 95% coverage from 3.18 to 1.47 percentage points).

Specifically, this 148,500-request study also established two major operational boundaries:

1. **Imputation shortfall under linear MI on clustered recounts**: Applying `cv3` variance did not rescue unclustered linear MI (`catmech_reference(use_recorded = TRUE)` without `cluster`) on the 11 Rwanda HMIS indicators, where coverage remained below 92% on 7 of 11 indicators (dropping to 81.0% on `iron`, 81.6% on `anc1_early`, 84.2% on `syphilis`, and 86.1% on `pnc`), leaving the overall accuracy verdict as `PARTIAL` (and simulated rare-count recounts averaged 87.5% coverage under `cv3`).
2. **Predictor-slope attenuation under clustered random-intercept MI**: When the audited variable T_i is a **predictor** in the outcome regression (`Y ~ T + x + stratum`) rather than the outcome, clustered random-intercept imputation (`catmech_reference(cluster = )`) attenuated the estimated slope toward zero by up to -16.1% with 84.4% coverage at a 25% audit under `cv3` (and -10.1% bias, 86.0% coverage at a 50% audit), whereas fitting `catmech_reference(use_recorded = TRUE)` without `cluster` and pooling with `catmech_pool_lm(cluster = )` yielded -3.0% bias and 95.4% coverage. Accordingly, `catmech_reference(cluster = )` remains experimental and should only be used when the audited variable is the analysis outcome (`catmech_pool_lm()` issues an explicit warning if `catmech_reference(cluster = )` is used on an imputed predictor).

6 Real-world empirical benchmarks

6.1 Public paired-discrepancy registry and survey benchmarks

Because no public record-level clinical trial chart SDV dataset exists, we first evaluated `catmech` on two large-scale public datasets containing real paired recording discrepancies on every row without synthetic injection (`validation/REAL_DATA_SYNTHESIS.md`): ClinicalTrials.gov ($N = 27,208$ studies, comparing sponsor-reported log enrollment against participant-flow log Started counts in a regression of log study duration in months on log enrollment, phase, and sponsor class) and the National Health and Nutrition Examination Survey (NHANES) 2017–2018 [Centers for Disease Control and Prevention, 2020] ($N = 3,908$ adults, comparing self-reported weight against examination-measured weight in kg in a regression of systolic blood pressure in mmHg on weight, age, sex, and race). Across 33,800 planned requests (1,000 resampled cohorts of size $N = 600$ at the primary 25% audit fraction $n_{\text{audit}} = 150$, plus 200 cohorts per cell at 5%, 10%, 50%, and targeted audit fractions), 33,613 intervals were returned alongside 76 unsupported-design refusals, 83 MCMC convergence refusals, and 28 singular raking errors:

Dataset	Metric	Naive (Trust Rec.)	Discard Rec.	Contamination	Linear MI	Full Ref. (Oracle)
ClinicalTrials.gov ($N = 27,208$)	Bias	+0.0043	-0.0003	+0.0068	+0.0001	-0.0003
Discrepancy rate: 14.2%	Coverage	91.4%	92.3%	91.2%	91.8%	91.2%
Target: 0.0357 log mo / log N	Mean Width	0.097	0.205	0.106	0.102	0.096
	RMSE	0.0282	0.0539	0.0308	0.0286	0.0275
NHANES 2017–2018 ($N = 3,908$)	Bias	-0.0180	-0.0063	-0.0066	-0.0025	+0.0001
Discrepancy rate (> 1 kg): 67.3%	Coverage	92.7%	94.2%	94.5%	93.9%	94.9%
Target: 0.127 mmHg / kg	Mean Width	0.149	0.297	0.155	0.154	0.140
	RMSE	0.0422	0.0785	0.0408	0.0409	0.0357

On both datasets, retaining recorded values reduced RMSE by 43% to 48% and narrowed confidence intervals by 48% compared to discarding unaudited values (`Discard Rec.`, `reference_normal`). Note that on ClinicalTrials.gov, the Full-Reference Oracle itself covered 91.2% because homoskedastic OLS standard errors (0.0244) slightly underestimated empirical sampling variability (0.0275) under non-constant residual variance, a gap inherited by all methods. Moreover, on ClinicalTrials.gov at a 25% audit, naively trusting recorded values (`Naive (Trust Rec.)`) achieved an RMSE of 0.0282 and bias of +0.0043, whereas `catmech_contamination()` had slightly higher bias (+0.0068) and RMSE (0.0308), and at a sparse 5% audit fraction `catmech_contamination()` was weakest (returning 164 of 200 intervals [18% refusal rate] with bias +0.0159 and RMSE 0.0423, vs. 200/200, bias +0.0062, and RMSE 0.0356 for `Linear MI`). On NHANES at 25%, where self-reported weight bias was substantial (-0.0180, or 14% of the target slope), both `Linear MI` (bias -0.0025, RMSE 0.0409) and `Contamination` (bias -0.0066, RMSE 0.0408) corrected the naive bias (and at a 10% NHANES audit, `Contamination` achieved lower RMSE [0.0435 vs. 0.0490] than `Linear MI` with 14/200 refusals).

Most importantly, because real-world discrepancies in both datasets were small relative to the conditional spread of $T_i \mid X_i$, **per-record posterior corruption probabilities p_i had near-chance discrimination on real data**: AUC for identifying discrepant unaudited rows at a 25% audit was **0.543** on ClinicalTrials.gov (Brier score 0.127 vs. 0.121 for base-rate prevalence alone; posterior mean RMSE 0.277 vs. 0.252 for raw recorded values) and **0.500** on NHANES (Brier score 0.222 vs. 0.220 for prevalence; posterior mean RMSE 4.81 vs. 4.73 kg for raw recorded values), compared to an AUC of 0.962 in the synthetic `missed_flags` cell. Consequently, individual record-level error probabilities p_i must not be used to adjudicate individual patient records on diffuse real-world data. While the KS alarm in `catmech_holdout_check()` stayed quiet on these cohorts (firing in only 2 of 38 completed checks on ClinicalTrials.gov and 1 of 40 on NHANES), the continuous held-out Brier and RMSE skill scores correctly indicated that the mixture model had no predictive edge over raw recorded values.

6.2 Real-audit regression slope recovery benchmarks

We further conducted a pre-specified 45,600-request real-audit benchmark (`validation/REAL_AUDIT_SYNTHESIS.md`) evaluating regression slope recovery on the Rwanda HMIS facility register recounts [Nshimyiryo et al., 2020] alongside 498 paired effect-size recounts from psychology meta-analyses [Maassen et al., 2020]. Under three pre-specified decision rules at a 25% audit fraction:

1. **Rule 1 (Does a partial 25% audit with `Linear MI` beat raw recorded data?)**: A 25% audit with `catmech_reference(use_recorded = TRUE)` paid clearly on only **1 of 4 Rwanda slopes** (`ANC 4th visits`, where errors scaled with facility size [$r = +0.477$] and recovery RMSE fell to 0.458× that


of raw recorded data), tied on PNC (0.923× in the initial 500 cohorts, 1.009× [0.969, 1.048] on 2,000 fresh seeds, and 1.136× against the full-pool slope), and increased recovery RMSE relative to raw recorded values on Syphilis (2.393×) and Iron (1.125×).

2. **Rule 2 (Does `catmech_contamination()` add value over `Linear MI`?)**: The pre-specified verdict on Rwanda HMIS was `INCONCLUSIVE`. Although `catmech_contamination()` had lower point recovery RMSE than `Linear MI` on PNC (0.053 vs. 0.072, ratio 0.745) and Iron (0.112 vs. 0.156, ratio 0.717), it failed the pre-specified coverage condition on **all three primary slopes** and failed the $\geq 90\%$ interval return-rate condition on **two of three**: clustered 95% coverage was **71.6%** on PNC (468/500 [93.6%] returned, vs. 85.4% coverage for `Linear MI` and 95.2% for the recount oracle), **66.6%** on Syphilis (404/500 [80.8%] returned, RMSE 0.195 vs. 0.182 for `Linear MI` and 0.076 for raw recorded), **35.4%** on Iron (424/500 [84.8%] returned, vs. 79.5% coverage for `Linear MI`), and **49.7%** on the size-dependent `ANC 4th visit` stress slope (RMSE 0.124 vs. 0.085 for `Linear MI`, retaining 63% of reported bias). Across all 20 Rwanda cells, `catmech_contamination()` was closest to 95% coverage in only 1 of 20 cells (worst case 35.4%), compared to 10 of 20 cells for `Linear MI` (worst case 79.5%), and per-record error AUCs were 0.501 to 0.582.
3. **Rule 3 (Psychology effect-size recount controls where reported bias is near zero)**: On the 498 psychology effect-size recounts [Maassen et al., 2020], where raw reported slopes already closely matched recounted truth, `Linear MI` at a 25% audit inflated recovery RMSE relative to raw recorded values by 2.705× (standardized mean difference) and 1.578× (correlation r), whereas `catmech_contamination()` stayed close to the recorded values (1.306× and 0.950×).

Together, these real-audit results demonstrate that the advantage of `catmech_contamination()` under gross, separable errors remains **synthetic-only** (because Rwanda's recount discrepancies are moderate and non-separable), and that on moderate real-world audit discrepancies `Linear MI` provides substantially more reliable interval coverage than the contamination mixture model.

7 Methodological decision framework

Based on our synthetic and empirical findings, we recommend the following evidence-based protocol for clinical data audits:



1. **Gross, separable contamination**: When recording errors represent large, discrete corruption events (e.g., decimal transcription slips or 10× unit errors) and target residuals are approximately Gaussian under ignorable audit selection, `catmech_contamination()` prevents severe attenuation bias while maintaining near-oracle precision.
2. **Subtle, diffuse, or moderate real-world discrepancies**: When errors represent continuous rounding, mild reporting bias, or moderate non-separable recount differences (as in ClinicalTrials.gov, NHANES, and Rwanda HMIS), simple linear imputation (`catmech_reference(use_recorded = TRUE)`) is the recommended baseline, avoiding MCMC convergence refusals and mixture interval undercoverage.

When the audited variable is a predictor in a clustered study, fit `catmech_reference(use_recorded = TRUE)` without `cluster` and pool with `catmech_pool_lm(cluster = )`.

8 Discussion: simulation-to-practice divergence and methodological implications

A central contribution of this work is documenting the divergence between synthetic Monte Carlo simulations and empirical real-world audits, including full disclosure of pre-specified negative and inconclusive findings.

In the statistical methodology literature, novel measurement-error and imputation models are predominantly benchmarked on synthetic datasets where contamination is gross, bimodal, and generated from the exact parametric family assumed by the estimator. Under those synthetic conditions (`missed_flags` and `error_gross_and_small`), `catmech_contamination()` eliminates the severe attenuation bias that compromises simple linear imputation (bias of -0.0117 vs. -0.2354 in `error_gross_and_small`) while narrowing confidence intervals by 44.3% relative to complete-case analysis in `missed_flags` (0.1842 vs. 0.3309). Even in synthetic stress testing, however, our pre-specified counterfactual scenarios expose clear failure boundaries: heavy-tailed target distributions (`heavy_tail_target`, 87.3% coverage), audit selection on the latent true value (`nonignorable_missed`, 79.3% coverage), and undeclared reference noise (`noisy_undeclared`, 88.7% coverage).

Our empirical investigations across ClinicalTrials.gov, NHANES 2017–2018, and the Rwanda HMIS facility recount audit reinforce a sobering practical lesson: real-world recording discrepancies are rarely gross or separable into discrete clean and contaminated components. On ClinicalTrials.gov and NHANES, simple linear MI (`catmech_reference(use_recorded = TRUE)`) matched or slightly beat the contamination mixture model in RMSE (0.0286 vs. 0.0308 on ClinicalTrials.gov; 0.0409 vs. 0.0408 on NHANES) with zero MCMC overhead, while on Rwanda HMIS the contamination mixture model suffered severe clustered interval undercoverage (35.4% to 71.6%) and 6% to 19% convergence refusal rates. Across all three real datasets, per-record corruption probabilities p_i performed near chance (AUC 0.500 to 0.582).

This simulation-to-practice divergence demonstrates why `catmech` is designed around an empirical diagnostic workflow rather than a single flagship estimator:

1. Exploratory discrepancy screening via `catmech_recording_audit()` to quantify error magnitudes and between-stratum heterogeneity before choosing an imputation strategy;
2. Out-of-sample predictive cross-validation via `catmech_holdout_check()`, comparing held-out Brier and RMSE skill scores against raw recorded values rather than relying on the low-power KS uniformity alarm alone;
3. Direct side-by-side strategy comparison via `catmech_compare_strategies()` so study teams can verify whether the data support mixture modeling, linear MI, complete-case analysis, or unadjusted recorded values.

9 Regulatory documentation and data governance

Clinical trial analyses are expected to be documented and reproducible: FDA 21 CFR Part 11 [U.S. Food and Drug Administration, 2003] governs electronic records and audit trails, and ICH E9 [International Conference on Harmonisation, 1998] asks that data handling and statistical methods be prespecified and documented. `catmech_audit_trail()` supports that documentation. It does not certify compliance, which depends on the validated system, access controls, and procedures around the analysis. It generates provenance records in Markdown, JSON, or standalone HTML, capturing:

1. Full execution environment metadata (R version, operating system platform, package version, UTC timestamp).
2. Cryptographic SHA-256 checksums of the raw input dataset, model specification, and each of the m completed datasets.

3. MCMC convergence diagnostics recorded on the audit trail (`convergence_passed` pass/fail status and `mcmc_iterations` count, with full Gelman-Rubin $\hat{R}$ and bulk/tail effective sample size vectors stored in `result$model_diagnostics`).
4. Pooled parameter estimates and multi-scenario sensitivity envelopes.

10 Quality assurance and verification

`catmech` has been developed under an automated software verification harness:

- **Unit tests**: 1,754 unit tests executed via `testthat`, achieving 0 failures, 0 warnings, and 0 skips.
- **CRAN conformance**: `R CMD check --as-cran` reports 0 errors and 0 warnings.
- **Automated artifact verification**: Verified against an independent Python verification suite (`verify-artifacts.py`, `verify-release-archive.py`, and `verify_paper_numbers.py`) confirming exact correspondence across 10 frozen simulation and empirical benchmark archives and over 500,000 replicate requests.

11 Availability and software details

`catmech` is open-source software distributed under the MIT license. It requires R ($\geq$ 4.1.0) and a C++17 compiler, utilizing the non-base CRAN packages `Rcpp`, `RcppArmadillo`, `digest`, `jsonlite`, `mice`, `pan`, `posterior`, `survival`, `truncnorm`, and `yaml` (alongside base R packages `graphics`, `grDevices`, `parallel`, `stats`, and `utils`). Source code, documentation, and vignettes are available at https://github.com/patrickiben/catmech.

References

Anastasios N. Angelopoulos, Stephen Bates, Emmanuel Candès, Michael I. Jordan, and Tijana Zrnic. Prediction-powered inference. *Science*, 382(6671):669–674, 2023. doi: 10.1126/science.adi6000.

John Barnard and Donald B. Rubin. Small-sample degrees of freedom with multiple imputation. *Biometrika*, 86(4):948–955, 1999. doi: 10.1093/biomet/86.4.948.

Norman E. Breslow, Brad McNeney, and Jon A. Wellner. Large sample theory for semiparametric regression models with two-phase, outcome dependent sampling. *The Annals of Statistics*, 31(4):1110–1139, 2003. doi: 10.1214/aos/1059655907.

Raymond J. Carroll, David Ruppert, Leonard A. Stefanski, and Ciprian M. Crainiceanu. *Measurement Error in Nonlinear Models: A Modern Perspective*. Chapman & Hall/CRC, Boca Raton, FL, 2nd edition, 2006. doi: 10.1201/9781420010138.

Centers for Disease Control and Prevention. National Health and Nutrition Examination Survey (NHANES) 2017–2018, 2020. URL https://www.cdc.gov/nchs/nhanes/.

William G. Cochran. *Sampling Techniques*. John Wiley & Sons, New York, 3rd edition, 1977.

A. Philip Dawid. Present position and potential developments: Some personal views: Statistical theory: The prequential approach. *Journal of the Royal Statistical Society: Series A (General)*, 147(2):278–292, 1984. doi: 10.2307/2981683.

Andrew Gelman and Donald B. Rubin. Inference from iterative simulation using multiple sequences. *Statistical Science*, 7(4):457–472, 1992. doi: 10.1214/ss/1177011136.

James Honaker, Gary King, and Matthew Blackwell. Amelia II: A program for missing data. *Journal of Statistical Software*, 45(7):1–47, 2011. doi: 10.18637/jss.v045.i07.

Alan Huang and M. P. Wand. Simple marginally noninformative prior distributions for covariance matrices. *Bayesian Analysis*, 8(2):439–452, 2013. doi: 10.1214/13-BA815.

International Conference on Harmonisation. *ICH Harmonised Tripartite Guideline: Statistical Principles for Clinical Trials (E9)*. International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use, 1998. URL https://database.ich.org/sites/default/files/E9_Guideline.pdf.

Wolfgang Lederer and Helmut Küchenhoff. A short introduction to the SIMEX and MCSIMEX. *R News*, 6 (4):26–31, 2006. URL https://CRAN.R-project.org/package=simex.

Kung-Yee Liang and Scott L. Zeger. Longitudinal data analysis using generalized linear models. *Biometrika*, 73(1):13–22, 1986. doi: 10.1093/biomet/73.1.13.

D. Y. Lin and L. J. Wei. The robust inference for the Cox proportional hazards model. *Journal of the American Statistical Association*, 84(408):1074–1078, 1989. doi: 10.1080/01621459.1989.10478874.

Esther Maassen, Marcel A. L. M. van Assen, Michèle B. Nuijten, Anton Olsson-Collentine, and Jelte M. Wicherts. Reproducibility of individual effect sizes in meta-analyses in psychology. *PLOS ONE*, 15(5): e0233107, 2020. doi: 10.1371/journal.pone.0233107.

Linda Nab, Maarten van Smeden, Ruth H. Keogh, and Rolf H. H. Groenwold. mecor: An R package for measurement error correction in linear regression models with a continuous outcome. *Computer Methods and Programs in Biomedicine*, 208:106238, 2021. doi: 10.1016/j.cmpb.2021.106238.

Jerzy Neyman. Contribution to the theory of sampling human populations. *Journal of the American Statistical Association*, 33(201):101–116, 1938. doi: 10.1080/01621459.1938.10503378.

Alphonse Nshimyiryo, Catherine M. Kirk, Stephen M. Sauer, Emmanuel Ntawuyirusha, Andrew Muhire, Felix Sayinzoga, and Bethany Hedt-Gauthier. Health management information system (HMIS) data verification: A case study in four districts in Rwanda. *PLOS ONE*, 15(7):e0235823, 2020. doi: 10.1371/journal.pone.0235823.

James M. Robins, Andrea Rotnitzky, and Lue Ping Zhao. Estimation of regression coefficients when some regressors are not always observed. *Journal of the American Statistical Association*, 89(427):846–866, 1994. doi: 10.1080/01621459.1994.10476818.

Donald B. Rubin. *Multiple Imputation for Nonresponse in Surveys*. John Wiley & Sons, New York, 1987. doi: 10.1002/9780470316696.

Donald B. Rubin. Multiple imputation after 18+ years. *Journal of the American Statistical Association*, 91 (434):473–489, 1996. doi: 10.1080/01621459.1996.10476908.

U.S. Food and Drug Administration. *Guidance for Industry: Part 11, Electronic Records; Electronic Signatures: Scope and Application*. U.S. Food and Drug Administration, Rockville, MD, 2003. URL https://www.fda.gov/regulatory-information/search-fda-guidance-documents/part-11-electronic-records-electronic-signatures-scope-and-application.

Stef van Buuren and Karin Groothuis-Oudshoorn. mice: Multivariate imputation by chained equations in R. *Journal of Statistical Software*, 45(3):1–67, 2011. doi: 10.18637/jss.v045.i03.

Ian R. White, Patrick Royston, and Angela M. Wood. Multiple imputation using chained equations: Issues and guidance for practice. *Statistics in Medicine*, 30(4):377–399, 2011. doi: 10.1002/sim.4067.