A False Floor: How a Weak Baseline and a Curated Benchmark Produced an "Information Limit" on a Lethal Recognition Task that a Stronger Baseline then Refuted

Anonymous authors
Paper under double-blind review

Abstract

Claims that a model *cannot* do something (impossibility, information limits, irreducible error) are treated as durable conclusions, yet they are as prone to artifact as claims that a model *can*. We document a worked example end to end. Investigating the canonical "should I eat this mushroom?" decision, we built a careful, reproducible, adversarially-verified study and concluded that the task is **information-limited**: on real image embeddings a strong aggregate separability (AUROC 0.94) coexisted with an apparently irreducible ~5% confidently-wrong "false-eat" residual that *no method in a panel of ~18* could remove, with formal divergence-floor and data-processing arguments to explain why, and decision-curve analysis showing abstention dominates. The conclusion survived an adversarial-verification process that had already caught two of our own *positive* results as rigged. It was wrong anyway. Within the same pipeline and the same pairs, a stronger frozen encoder (DINOv2) cut the false-eat to near zero: $\leq$**0.7%** per cell un-cropped ($\leq$1.0% under a substrate crop; pooled 95% upper bound **0.4%**), where the weak encoder left **0.1–21.3%**, lower within every pair — including the freshly-scraped textbook "you-need-a-microscope" confusions (honey-mushroom vs deadly *Galerina*, where it cleared 0 of 150 held-out poisons). What matters is the encoder-dependence, not the absolute rate. (The strong-encoder cells are near-zero rather than exact zeros: at the original $\approx$60-specimen sampling they read a bare "0.0%" whose $\approx$4.9% upper bound overlapped the original residual; at full sampling they read 0.1–0.7%, with a single-cell bound $\approx$2% and a pooled bound $\approx$0.4%.) An information limit is a property of the *photograph* and would bind every encoder equally, so a stronger backbone that beats a weaker one within every pair is incompatible with one: the "information limit" was a **weak-baseline artifact**. We show this across **six lethal look-alike decisions spanning five biological groups** (fungi, plants, reptiles, arachnids, amphibians), **three anchored to externally-published indistinguishability claims** (coral-snake mimicry, (Pfennig et al., 2001); brown-recluse misidentification, (Vetter, 2000); poison-frog mimicry, (Symula et al., 2001)). In every group the weak encoder leaves a large false-safe rate (often near-chance, clearing up to half of the dangerous specimens) that a stronger frozen encoder beats within every pair: to near zero in five of six groups, and always far below the weak baseline. One small-n poison-frog pair is the honest exception, at ~10% against the weak encoder's 33%. We offer the representation-invariance test as a portable protocol for frozen-representation limit claims, not a general theorem about all safety-negatives, and we distill the post-mortem (a weak baseline, a curated benchmark mistaken for the field, and a true-but-empirically-vacuous theory whose antecedent we never checked) into a checklist. The lesson generalizes beyond mushrooms: *benchmark separability is not field safety, and a limitation claim is only as strong as the baseline that fails to beat it.*

1. Introduction

A large fraction of machine-learning writing argues that a method *helps*. A smaller, influential fraction argues that something *cannot be done* — an information limit, an irreducible error, an impossibility. Such negative claims are valuable precisely because they redirect effort: if a task is information-limited, the rational response is to stop building models and go collect a different signal. They are also unusually *sticky*, because they feel like theorems and are rarely re-litigated.

This paper is a fully-worked cautionary study of how a negative claim can be confidently produced, defended against adversarial scrutiny, and still be false — anchored on one mushroom self-refutation and replicated across five further look-alike decisions in other kingdoms (§3.1–§3.2), three of them re-examinations of externally-published indistinguishability claims, to show the failure is not domain-specific. We did not set out to write it. We set out to answer whether "should I eat this mushroom?" is answerable from a photograph, concluded after months of work that it is not — that the limit is *information*, not method — and wrote that paper. Then, preparing it for review, we did the one thing the original investigation never did: we replaced a deliberately sovereignty-preserving but weak frozen encoder with a stronger one. The central result did not survive.

We report the full sequence (the wrong conclusion, the refutation, and the anatomy of the error) because the failure mode appears general: safety-negative claims inherit the weakness of their strongest-failing baseline, and curated benchmarks systematically understate field difficulty in one direction while overstating model limits in the other.

Related work, and our delta. That weak or poorly-tuned baselines inflate apparent findings is well documented on the *positive* side: metric-learning gains that evaporate under fair comparison (Musgrave et al., *A Metric Learning Reality Check*, 2020), recommender progress that does not survive tuned baselines (Ferrari Dacrema et al., 2019), and rank instability across configurations (Dehghani et al., 2021; Bouthillier et al., 2021). That curated benchmarks diverge from deployment is the dataset-bias and distribution-shift program (Torralba & Efros, 2011; Koh et al., 2021; Hendrycks et al., 2021). Our contribution is the **negative-side analogue and its asymmetry.** A limitation/impossibility claim inherits the weakness of its strongest-*failing* baseline exactly as a positive claim inherits its weakest-*tuned* one — but the community has standard instrumentation for the positive case (strong baselines, ablations) and almost none for the negative, and, more sharply, a *self-verification discipline calibrated to catch rigged wins is structurally blind to over-claimed losses* (§4d). Two nearest neighbors deserve explicit distinction. **McGreivy & Hakim (2024, *Nat. Mach. Intell.* 6:1256–1269)** find that 79% of ML-for-PDE papers claiming to beat standard solvers compared against weak baselines, with widespread under-reporting of negatives — the closest positive-side analogue; our delta is the *negative-claim* direction and the representation-invariance instrument that operationalizes it. And on *this exact task*, Kuznetsov et al. (2026) reach the *opposite* verdict — that AI tools cannot safely identify wild mushrooms (the best they tested still failed ≈15% of real-world cases, none consistently correct) — but from deployed, often weak or black-box systems; our results re-examine that conclusion and locate the failure in the deployed model, not an information limit, which is precisely the distinction this paper exists to draw. We make this concrete with **six case studies across five biological groups** (§3–§3.2), three of them re-examinations of externally-published "look-alike" claims, a checklist (§5), and a symmetric "negative-claim red-team" protocol (§5.1). We do not claim a measured base rate of false limits in the literature; establishing one is future work.

---

2. The investigation, as we built and believed it

The task. A frozen image encoder over real GBIF photographs of edible species (chanterelle, field mushroom *Agaricus campestris*, morel) and poisonous ones (jack-o'-lantern, the Destroying Angel *Amanita bisporigera*, the open-set Death Cap *A. phalloides*), reduced to a one-step decision in {eat, abstain} with a deliberately asymmetric cost: eating an edible is a meal, eating a poison is death.

The headline result *as we found it.* On real I-JEPA (ViT-H/14, mean-pooled) latents, the lethal pair was "separable" (AUROC 0.94) — and it did not matter: the actual eat decision cleared **7 of 81** held-out

Destroying Angels (8.6% on the split, ~4.9% ± 2.5% over 20 splits), and *no abstention tuning removed them* (7 eaten at every confidence threshold). The cleared specimens were confidently-wrong consensus — low predicted risk, low ensemble disagreement, on-manifold. A 0.94 aggregate AUROC coexisting with a deadly residual the safety machinery could not catch: the "94% → liver failure" phenomenon on a real mushroom.

The sweep. We then pushed ~18 methods at the residual — more data, fine-grained metric learning, resampling, the full f-divergence/optimal-transport family, generative and diffusion augmentation, metadata fusion, conformal prediction, hierarchical planning, active-inference exploration, persistent homology, 3D reconstruction, Morse-theoretic shape analysis, and macro↔micro generative modeling. Each was run with fair baselines; the most important claims also with adversarial verification. None removed the residual. Decision-curve analysis showed that at the field-standard lethal cost ratio (FungiCLEF `c_PSC = 100`) no operating point beat "eat nothing." A frontier VLM refused every deadly specimen — but, we argued, from world knowledge of famous species, not from overcoming the underlying limit.

The theory. We formalized the negative (Appendix A): a *coincidence/divergence floor* (where two class-conditional feature distributions coincide, every f-divergence and Wasserstein vanish and Bayes error is the prior), a *data-processing bound* (any function of the photo — a sharper representation, a generated image, an upscale, a topological summary — is bounded by the photo's Bayes floor), and an *independence condition* (only a channel with information about edibility *beyond* the photo can help). These say: if the discriminating feature is not in the image, no method recovers it.

The discipline. Twice during this work we produced a result that *looked* like a method win and was not — a hierarchical-planning improvement that adversarial-verification panels decomposed as a rigged baseline and an objective swap. We caught both, reported them, and adopted adversarial verification as the project's spine. We were, we believed, being unusually careful.

This is the paper we were ready to submit. It is reproduced, essentially intact, as the companion document; we preserve it deliberately as the specimen.

---

3. The refutation

Preparing for review, a referee-style objection was obvious: *the backbone is a frozen, mean-pooled I-JEPA; did you try a stronger representation?* We had, in fact, shown that fine-tuning a domain encoder helped *in-category* — but we had filed that under "ceiling up, floor stays" and not pursued frozen alternatives. So we ran the sweep we should have run first.

Five real pairs × three strong frozen encoders. We assembled five genuine field-confusion edible/deadly pairs (field-mushroom/Destroying-Angel, chanterelle/jack-o'-lantern, morel/false-morel, oyster/jack-o'-lantern, puffball/death-cap-button) and encoded each with three paradigmatically different strong frozen encoders: masked-latent **I-JEPA** (Assran et al., 2023) (ViT-H/14), self-distilled **DINOv2** (Oquab et al., 2024) (ViT-L/14), and contrastive **CLIP** (Radford et al., 2021) (ViT-L/14). Per cell, the actual false-eat at 80% edible-yield, 10 seeds.

mean over 5 pairs	I-JEPA-H14	DINOv2-L	CLIP-L
false-eat at 80%	1.3% (95% UCB 2.2%, n=720)	0.1% (95% UCB 0.4%, n=720)	0.0% (95% UCB 0.4%, n=720)
open-set eaten	27%	0%	3%

Bounds are one-sided 95% Clopper-Pearson on the false-eat rate. ***n=720*** *is the five pairs* ***pooled*** *— ~150 unique held-out poison specimens per pair (the per-seed 30% test split over* all available images*,* ***not*** *seed-multiplied; the 10 seeds reshuffle the same pool). The strong encoders are* ***near-zero but not exactly zero****: DINOv2's pooled estimate is 0.1% (its Agaricus/Destroying-Angel cell alone reads 0.3%), CLIP's 0.0%, both with pooled upper bound ≈0.4%. At the original ≈60-specimen-per-cell sampling these same cells read a bare "0.0%" with*

a ≈4.9% per-cell upper bound that overlapped* the original ~4.9% ± 2.5% residual (§2) — i.e. the exact zero was itself an undersampling artifact, the very effect this paper warns against (§4). The encoder gap is unchanged: weak I-JEPA's pooled false-eat is 1.3% (UCB 2.2%), an order of magnitude above the strong encoders.*

The residual did not persist across encoders. **DINOv2 and CLIP drove the in-distribution false-eat to near zero — DINOv2 ≤0.3% on every one of the five pairs (pooled 0.1%, UCB 0.4%), CLIP ≤0.1%.** On the *identical* Agaricus-vs-Destroying-Angel task where I-JEPA leaves 3.5%, DINOv2 leaves 0.3% (UCB 2.0%). A center-crop confound test (background removed) left DINOv2 essentially unchanged (≤0.2% per cell, pooled 95% UCB 0.4%), so it was reading the fungus, not the habitat.

Field degradation. We tested whether the curated success survives the regime where foraging deaths occur: downsample, blur, **occlude the bottom third** (the un-excavated base — the single most common field failure, which removes the volva), and color-jitter the genuine lookalike pairs. DINOv2 held at **0.0–0.7%** (95% UCB ≤3.1%, n=150/cell); a lookalike open-set of novel white toxins (n=1151) was eaten 0% (95% UCB 0.5%).

The decisive test (the hardest pairs we could find). The above pairs are separable when photographed well; a fair test of an *information* limit needs the cases mycologists call genuinely undecidable from a photo. We freshly scraped the textbook "requires-a-microscope" confusions and tested them with the easy pair as a positive control:

pair (10-seed AUROC / false-eat at 80)	I-JEPA	DINOv2	CLIP
[control] Agaricus / Destroying-Angel	0.97 / 3.5%	1.00 / 0.3% (95% UCB 2.0%, n=150)	1.00 / 0.1% (95% UCB 2.0%, n=150)
honey-mushroom / deadly *Galerina* (most fatal real confusion)	0.97 / 3.5%	**1.00 / 0.0% (95% UCB 2.0%, n=150)**	1.00 / 0.1% (95% UCB 2.0%, n=150)
woodtuft / deadly *Galerina* ("separable only by spore ornamentation")	0.95 / 4.8%	**0.98 / 0.7% (95% UCB 3.1%, n=150)**	0.98 / 1.9% (95% UCB 5.1%, n=150)
honey-mushroom / sulphur-tuft	0.88 / 21.3%	1.00 / 0.4% (95% UCB 2.5%, n=120)	0.98 / 1.4% (95% UCB 5.2%, n=120)
puffball / earthball	0.98 / 2.1%	1.00 / 0.0% (95% UCB 2.5%, n=120)	1.00 / 0.4% (95% UCB 2.5%, n=120)

Bounds are one-sided 95% Clopper-Pearson on the false-eat rate; ***n*** *is the number of* ***unique held-out poison specimens*** *(the per-seed 30% test split over* all available images*,* ***not*** *seed-multiplied).* ***n*** *differs by row because the species differ in image count: the control and both* Galerina* rows have `n=150`, the sulphur-tuft and earthball rows `n=120`. Two cells (honey/*Galerina* and puffball/earthball, DINOv2) hold at an exact 0 of 150/120; the rest are **small but nonzero** (DINOv2 0.3–0.7%, CLIP 0.1–1.9%) — at the original ≈60–90-specimen sampling several of these read a bare "0.0%", so the higher-powered re-run shows those zeros were undersampling artifacts. The weak I-JEPA baseline's one-sided 95% upper bounds run 6.3–28.8%. Point estimates are the full-image re-run; the encoder-dependence conclusion is unchanged.*

DINOv2 performed nearly perfectly on the pairs we selected *because* they are supposed to be undecidable from a photo. A substrate-stripping crop (the one plausible reprieve — *Armillaria* on living hardwood vs *Galerina* on conifer logs is a habitat cue) left DINOv2 low (**0.0–1.0%**, 95% UCB ≤4.1%), while the *weak* I-JEPA stayed high and degraded further on the canonical pair (honey/*Galerina* 3.5→4.1%) — confirming that the weak encoder relied on contextual cues that the strong encoder did not require. *(All sub-tests above — field degradation, ±background crop, and substrate-strip crop — were re-run on all available images, matching the two main pair tables; DINOv2's false-eat stays ≤1.0% per cell across every regime, CLIP ≤3.8% (only under the substrate-strip crop).)*

Across three regimes — curated pairs (± background crop), field degradation with base occlusion, and open-set — spanning nine lethal pairs, all re-run on every available image, the strong encoder held the false-eat to ≤1.0% (DINOv2; ≤0.7% un-cropped, curated-pooled 95% UCB 0.4%) and ≤3.8% (CLIP, under substrate-strip crop) where the weak encoder left 0.1–21.3% — lower within every pair. The curated GBIF pairs are the benchmark regime; the field-degraded, base-occluded, and open-set results are the harder regime where deaths occur, and the worst-case individual specimen (§6) remains untested. The decisive logic is not the absolute rate (the strong-encoder cells are near-zero, not exact zeros) but its *encoder-dependence*: an information limit binds the photograph, hence every encoder equally, so DINOv2 beating I-JEPA *within every pair* is incompatible with one. The ~5% "information-limited residual" the original paper was built on tracked the backbone, not the photo.

3.1 A second case, to test whether the failure is mushroom-specific

A single self-refutation invites the obvious objection — perhaps mushrooms are special, or our weak encoder was uniquely ill-suited to them. So we ran the identical protocol on a *second, independent lethal decision in a different kingdom*: the deadliest look-alike confusions in the carrot family (Apiaceae) — **poison hemlock** (*Conium maculatum*, the plant that killed Socrates) and **water hemlock** (*Cicuta maculata*, the most violently toxic plant in North America) against edible **wild carrot / Queen Anne's lace** (*Daucus carota*) and **cow parsley** (*Anthriscus sylvestris*) — a confusion that kills foragers every year. Fresh GBIF scrape, same three encoders, same eval, same Clopper-Pearson bounds (n = 90 held-out poison specimens per pair):

pair (10-seed AUROC / false-eat at 80, 95% UCB)	I-JEPA	DINOv2	CLIP
poison-hemlock / wild-carrot	0.93 / **9.0%** (≤15.5%)	1.00 / 0.2% (≤3.3%)	0.99 / 0.2% (≤3.3%)
water-hemlock / wild-carrot	0.96 / **5.8%** (≤11.3%)	1.00 / 0.0% (≤3.3%)	1.00 / 0.2% (≤3.3%)
poison-hemlock / cow-parsley	0.90 / **16.2%** (≤24.5%)	0.99 / 0.8% (≤5.2%)	0.98 / 1.4% (≤5.2%)

The pattern reproduces, more clearly than in mushrooms. The weak encoder leaves a **measured 5.8–16.2% false-eat** (a residual that, reported alone, would read as a substantial information limit on a deadly decision), and a stronger frozen encoder reduces it to 0–0.8% within every pair (the I-JEPA intervals, ≤11–24%, do not overlap the DINOv2 point estimates). The representation-invariance test (§5.1, item 2) flags both domains identically: a limit one encoder sees and another does not is a property of the encoder, not the photograph. The failure is **not mushroom-specific**; it is a generic hazard of anchoring a safety-negative on a weak representation. (Both these cases are our own constructions; §3.2 takes the next step and re-examines three *externally published* indistinguishability claims.)

3.2 The same protocol across five biological groups — three external claims

To test the failure mode at scale, we ran the identical three-encoder protocol on a battery of well-known lethal look-alike confusions spanning five biological groups, **three of them anchored to externally-published claims of visual indistinguishability** — so the "limit" is not one we authored:

case (group)	external "look-alike" claim	I-JEPA false-safe at 80%	DINOv2
mushrooms (fungi)	— (this work)	0.1–21.3%	0–0.7%
hemlock / wild carrot (plants)	— (this work)	5.8–16.2%	0–0.8%
coral snake / kingsnake (reptiles)	(Pfennig et al., 2001)	4.2–22.4%	**0.0%**

case (group)	external "look-alike" claim	I-JEPA false-safe at 80%	DINOv2
brown recluse / house spider (arachnids)	(Vetter, 2000)	10.3–**48.9%**	0–2.0%
poison-frog mimic / model (amphibians)	(Symula et al., 2001)	12.7–**43.3%**	0.3–10.0%
deadly berries (plant fruit)	— (foraging literature)	0.2–19.9%	**0.0%**

(Full per-pair tables with Clopper-Pearson bounds in Appendix C. Per our own no-bare-zeros rule (§5.1, item 5), the DINOv2 "0.0%" entries are 0 observed of ≈90 held-out poison specimens per pair — one-sided 95% upper bound ≤3.3%, not exact zeros; the amphibian row is small-n, see below.) The pattern holds across all six groups: the weak encoder leaves a large, measured false-safe rate, frequently **near-chance** (the recluse / house-spider and the *fantastica* / *imitator* pairs reach AUROC 0.73 and 0.72, clearing roughly half of the dangerous specimens), and a stronger frozen encoder lowers it within every pair, to near-zero in every group but the small-sample poison frogs (qualifier iii). A reviewer shown only the I-JEPA column would reasonably conclude that several of these decisions sit at an information limit; for the three external rows the literature says exactly that: coral snakes and their mimics are "near-indistinguishable" (Pfennig et al., 2001), brown recluses are massively over-reported because non-experts cannot tell them from harmless house spiders (Vetter, 2000), poison-frog mimics co-evolved to be visually identical to their models (Symula et al., 2001). None of these external indistinguishability claims binds a strong frozen encoder.

Three honest qualifiers. (i) These are species distributions on (often whole-organism) GBIF photographs, so they test *species separability*, not the isolated diagnostic confusion a forager faces with a single berry cluster or a partial specimen; the field and worst-case caveats of §6 apply unchanged. (ii) The pattern is **not universal** — the bilberry / deadly-nightshade pair is easy even for the weak encoder (0.2%) — and that is the point: the encoder-dependence appears precisely where a weak baseline *would* have been read as a limit, and is absent where it would not. The representation-invariance test (§5.1, item 2) separates the two automatically. (iii) The poison frogs are the one group where the strong-encoder advantage, though present in every pair point-wise, is **not yet statistically decisive**: the mimic/model species are photograph-poor, giving only n=22–30 held-out specimens on three of the four pairs (DINOv2 up to 10.0%, upper bound ≤25.9%), so only the n=90 *Allobates* / *Ameerega bilinguis* pair is conclusive. We report the frog case as suggestive, not confirmatory (per-pair bounds in Appendix C.1).

---

4. Anatomy of the error

How did a careful, reproduced, adversarially-verified investigation produce and defend a false limit? Four compounding causes, each general.

(a) The baseline was weak, and a negative inherits its baseline. A claim that "no method beats error ε" is exactly as strong as the *best* method that fails to. We used frozen mean-pooled I-JEPA for principled reasons (sovereignty, a fixed feature space for the divergence-floor proof) and never asked whether it was *strong enough to anchor a negative*. It was not. Under a stronger frozen encoder the same residual was an order of magnitude smaller (near zero, ≤0.7% per cell): it was a property of the backbone, not the photograph. An ablation (`ablation_agg.py`) isolates the cause, and finds that the baseline was weak in *two* ways at once. First, the pretraining objective: replacing I-JEPA's mean-pool with max-pool does not rescue it (both stay near-chance, 51% and 56% false-safe on the recluse pair), so I-JEPA's masked-latent features are weak for this discrimination regardless of how they are pooled. Second, the aggregation is independently lossy: mean-pooling DINOv2's patch tokens rather than using its CLS embedding degrades that same hard pair from 2.1% to 25.7% (max-pool recovers most, 5.5%). Our original baseline combined both handicaps, and we checked neither.

(b) Curated data was mistaken for the field. GBIF research-grade photographs are *selected to show the diagnostic features*. A model that scores ~0 on them has learned that the species differ — which they do, in photographable ways, even for "microscope" pairs (the microscope is needed for *certain* identification, not for a good *probabilistic* guess). We treated benchmark separability as a proxy for field decidability; it is the opposite — curated benchmarks understate field difficulty and overstate the model's limits at once.

(c) The theory was true but empirically vacuous. Our formal floor and data-processing propositions are correct: *if* the photo lacks the discriminating information, no function of it recovers it. We never verified the antecedent. For every real lethal pair we could obtain, the antecedent is false at the distribution level — the photo *does* carry enough signal — so the propositions, while valid, said nothing about these pairs. A true theorem with an unverified antecedent is not evidence.

(d) Adversarial verification checked the wrong axis. Our verification panels were genuinely effective — they reproduced every number and caught two rigged *positive* results. But they audited *correctness* (do the numbers reproduce?) and *self-deception toward positives* (is this win rigged?). They never asked the question that mattered for a *negative*: *is the baseline strong enough that its failure is informative?* A verification discipline tuned to catch over-claimed wins is blind to over-claimed losses.

The unifying point is that we instrumented heavily against finding a false positive and not at all against asserting a false negative; that asymmetry is the central lesson. We establish it from six case studies across five biological groups (three re-examining externally-published indistinguishability claims). This is an existence proof and a protocol, not a measured base rate over the literature; how often published limit claims are refuted by the strong-baseline test is the question §5.1 operationalizes.

---

5. A checklist for safety-negative claims

From the post-mortem, the stress tests a limitation/impossibility claim should pass before it is trusted:

1. **Strong-baseline floor.** Re-run the negative against the *strongest reasonable* baseline, not the most principled or convenient one. A residual that vanishes under a stronger backbone was never a limit. Report the strongest baseline that *fails*, and treat the claim as bounded by it.
2. **Antecedent verification.** If the negative rests on a theorem ("no function of X helps"), test that the theorem's antecedent ("X lacks the information") actually holds on the data, empirically — do not assume it.
3. **Hard-data, not curated-proxy.** Stress the claim on the genuinely-difficult regime it concerns (here: field degradation, occlusion, the expert-acknowledged-undecidable cases), scraped fresh — not on the curated benchmark, which is selected against difficulty.
4. **Symmetric verification.** If adversarial verification is used, give it a lens aimed at the negative: *is this loss real, or an artifact of a weak setup?* — not only *is this win rigged?*
5. **Benchmark ≠ field.** State explicitly what the benchmark number does and does not bound. Separability on curated data bounds nothing about field safety, in either direction.

5.1 The negative-claim red-team (the lens our verification lacked)

Our adversarial-verification panels were effective against rigged *positives* and useless against an over-claimed *negative*, because they never asked the negative-specific questions. We distill those into a red-team a verification process should run against any limitation/impossibility claim before it is trusted — the protocol that, applied to our own original paper, would have killed it in an afternoon:

1. **Strongest-baseline, not principled-baseline.** Name the single strongest method that could plausibly beat the claim; if the negative's baseline was chosen for principle or convenience (ours: a frozen, mean-pooled encoder for a clean proof), *replace it with the strongest available and re-measure.* Bound the claim by the strongest baseline that still fails.

2. **Representation-invariance test.** Re-run the claim across ≥2 paradigmatically different representations (here: masked-latent, self-distilled, contrastive). An *information* limit must bind all of them; a *representation* artifact will not. A within-instance gap between encoders refutes an information limit *regardless of the absolute error level* — the single most diagnostic check, and the one we omitted.
3. **Antecedent audit.** If the negative invokes a theorem ("no function of X helps"), *measure* its antecedent ("X lacks the discriminating information") on the data; do not assume it. A valid theorem with an unverified antecedent is not evidence.
4. **Fresh hard-data, not curated proxy.** Test the claim on the genuinely-difficult regime it concerns, scraped fresh for the purpose — not the curated benchmark, which is selected against difficulty.
5. **No bare zeros.** Report every "0 / no-method-helps" cell with a one-sided upper confidence bound. A 0-of-`n` is not 0; under an asymmetric cost the difference between "0%" and "≤5% with 95% confidence" can flip the conclusion.
6. **Cost-honest conclusion.** Re-state the conclusion against the *upper* bound under the task's real loss: does the negative still hold if the true rate sits at the top of its confidence interval?

The diagnostic asymmetry is item 2: a representation-invariance test is inexpensive and conclusive, and was the one check the original investigation never performed.

---

6. What remains true (and the real safety point)

The collapse of the information-limit thesis is not a safety endorsement — and the corrected picture is itself a safety argument, just a different one.

- **Strong encoders separate these species' *distributions* very well; this does not make foraging safe.** Three concerns survive untouched: *open-set true novelty* (a toxic species the model never saw), the *cost asymmetry* (under `c_PSC = 100`, even a tiny tail false-eat can fail the decision-curve test), and the *worst-case individual specimen* — the single *Galerina* photographed at the moment it is truly identical to an *Armillaria*, or an isolated button-stage *Amanita* egg — which a random-sample 0% cannot bound and which we could not obtain life-stage-filtered data to test. At full sampling the strong encoders cleared only a small fraction of held-out poisons (DINOv2 ≤0.7% per cell, pooled 95% UCB 0.4%) — low, but no longer a clean zero, and a random-sample rate this low still cannot bound the worst-case specimen above; absence of a measured tail in a curated sample is not a field guarantee.
- **The danger flips from "the model can't" to "the model is trusted because it can."** A model that aces even the microscope pairs invites exactly the over-trust the app-safety literature documents (higher confidence in AI *reduces* correct decisions in high-risk identification). A near-perfect benchmark score is a *reason for more caution*, not less, for a lethal decision.
- **The formal propositions remain valid** as a statement about genuinely information-poor observations — they simply do not apply to the real lethal pairs we tested. They would apply to the worst-case corner above, which is the honest residual of the original thesis: narrow, real, and undemonstrated.

---

7. Conclusion

We built a careful, reproduced, adversarially-verified case that a lethal recognition task is information-limited, and we were wrong: the limit was a weak-baseline artifact — a stronger frozen encoder beat the weak one *within every pair*, which an information limit forbids, across curated, field-degraded, and open-set regimes including the textbook-undecidable pairs, and then again across six lethal look-alike decisions in five biological groups — three of them re-examining externally-published "near-indistinguishable" claims that a strong encoder violates. We report the sequence because the failure is general: a limitation claim is only as strong

as the baseline that fails to beat it, curated benchmarks understate field difficulty while overstating model limits, and a verification discipline built to catch false positives will miss false negatives. The contribution is methodological rather than mycological — a worked demonstration, including its own collapse, that safety-*negative* claims demand the same adversarial stress as safety-positive ones, with a checklist for supplying it. *Research vehicle, never a forager. When in doubt, DON'T EAT.*

Appendix A — Formal statements (true, and empirically vacuous for the real pairs)

These are the propositions the original investigation used to *explain* the limit. They are correct. The finding of §4(c) is that their antecedent — "the photograph lacks the discriminating information" — is **false** for every real lethal pair we could obtain (§3), so the propositions, though valid, bound nothing about those pairs. We state them precisely both because they are the only durable formal content and because seeing the antecedent explicitly makes the vacuity clear.

Setup. Let X be the photograph on space $\mathcal{X}$, $Y \in \{0,1\}$ the edibility label (0 = edible, 1 = deadly), and $\varphi : \mathcal{X} \to \mathbb{R}^d$ a fixed feature map (the encoder). Write the class-conditional feature laws $P_0 = \mathrm{Law}(\varphi(X) \mid Y{=}0)$ and $P_1 = \mathrm{Law}(\varphi(X) \mid Y{=}1)$, the posterior $\eta(z) = P(Y{=}1 \mid \varphi(X){=}z)$, and the prior $\pi = P(Y{=}1)$. A decision rule is a measurable $g : \mathbb{R}^d \to \{\mathrm{eat}, \mathrm{abstain}\}$; the **false-eat** rate is $\mathrm{FE}(g) = P(g{=}\mathrm{eat} \mid Y{=}1)$ and the **yield** is $\mathrm{Yld}(g) = P(g{=}\mathrm{eat} \mid Y{=}0)$. All three statements concern a *fixed* modality X (a single RGB photograph).

Proposition 1 (Coincidence / divergence floor). Fix a target yield α. Among rules with yield $\geq \alpha$, the minimum false-eat is the Bayes value $\mathrm{FE}^*(\alpha)$, attained by thresholding the likelihood ratio dP_1/dP_0 (Neyman–Pearson). On any region R where $P_0 = P_1$: (i) every f-divergence and every Wasserstein distance between $P_0|_R$ and $P_1|_R$ is 0; (ii) $dP_1/dP_0 = 1$, $\eta \equiv \pi$, and $I(\varphi(X); Y \mid \varphi(X) \in R) = 0$; (iii) no rule separates the classes on R, so any rule clears poisons and edibles in equal proportion there. *Proof.* Each f-divergence and each W_p vanishes iff the measures coincide; $dP_1/dP_0 = 1$ gives $\eta = \pi$ by Bayes, and $I = 0$ by conditional independence; with η constant on R the Neyman–Pearson statistic is constant there. ■

Proposition 2 (Data-processing bound on post-hoc methods). Let $Z = g(X)$ be *any* measurable transform of the photograph: a different representation, a generated or upscaled image, a topological summary, a *predicted* microscopy. Then $Y \to X \to Z$ is a Markov chain, so $I(Z;Y) \leq I(X;Y)$, and for every yield α, $\mathrm{FE}^*_Z(\alpha) \geq \mathrm{FE}^*_X(\alpha)$: no function of the photograph beats the photograph's own Bayes floor. *Proof.* The data-processing inequality, plus monotonicity of the constrained Bayes risk in the decision σ-algebra ($\sigma(Z) \subseteq \sigma(X)$). ■

Proposition 3 (When an extra channel helps: the independence condition). For corroboration rules (eat only when both X and an auxiliary W clear), W lowers the false-eat below $\mathrm{FE}^*_X(\alpha)$ **iff** $I(W; Y \mid X) > 0$. (i) A redundant channel $W = h(X, \xi)$ with $\xi \perp\!\!\!\perp Y$ has $I(W; Y \mid X) = 0$ and cannot help. (ii) An independent channel $W \perp\!\!\!\perp X \mid Y$ with per-class reliability ρ multiplies the false-eat by $(1 - \rho)$, at a yield cost. *Proof.* If $W \perp\!\!\!\perp X \mid Y$, then FE factors as $\mathrm{FE}^*_X(\alpha) \cdot (1 - \rho)$; if $I(W; Y \mid X) = 0$, then g_W is conditionally independent of Y given $\{g_X{=}\mathrm{eat}\}$, leaving FE unchanged. ■

The vacuity, stated plainly. Propositions 1–3 are conditional: *if* a coincidence region of nonzero mass exists for a real lethal pair, *then* no method escapes it. Section 3 measured FE^*_X directly with a strong encoder and found it near zero ($\leq 0.7\%$ per cell; pooled 95% UCB 0.4%) across every pair tested, including the textbook-undecidable ones. The coincidence region of Proposition 1 therefore has at most negligible mass for these pairs, and the precondition for Proposition 2 to bind (a large FE^*_X) does not hold. The theory is intact; it does not describe the regime these pairs occupy. It would describe the worst-case corner of Section 6 (a single genuinely-coincident specimen), which remains undemonstrated for lack of life-stage-filtered data.

Appendix B — The original investigation

Preserved verbatim as the companion document (`MANUSCRIPT.md`): the ~18-method sweep, the decision-curve and conformal apparatus, the VLM analysis, and the two self-caught rigged positives. It is the specimen this paper dissects, not an endorsed result.

Appendix C — Reproducibility

All experiments are runnable in `~/Documents/EatThisMushroom/`. The refutation: `pairs_encoders.py` (five pairs × three encoders, ±crop), `hard_regime.py` (field degradation + base occlusion + lookalike open-set), `scrape_hard.py` + `hard_pairs.py` (the needs-a-microscope pairs, ±substrate-strip crop). The two tables in §3 are the **full-image** re-run (`FULL=1`, which uses every cached image and writes a separate `__full` feature cache, preserving the original 200/300-image arrays); both scripts print, per cell, the number of unique held-out poison test specimens and a one-sided 95% Clopper-Pearson upper bound on the false-eat rate. The second case study (§3.1): `scrape_plants.py` (the deadly Apiaceae) + `case2_plants.py`. The §3.2 multi-taxon battery: `scrape_snakes.py` + `case3_snakes.py` (coral-snake mimicry), `scrape_more.py` + `case4_spiders.py` (brown-recluse misID) and `case5_frogs.py` (poison-frog mimicry), `scrape_berries.py` + `case6_berries.py` (deadly berries) — each the same three-encoder protocol with Clopper-Pearson bounds and a corrupt-image-robust loader. The objective-vs-aggregation ablation (§4a): `ablation_agg.py`, which re-encodes three hard pairs keeping patch tokens and compares I-JEPA {mean, max} against DINOv2 {CLS, mean, max}. The original investigation: the method-sweep scripts enumerated in the companion document. All GBIF image scrapes are reproducible via the `species/match` + `occurrence/search` endpoints in the `scrape_*.py` files.

Appendix C.1 — Per-pair tables for the §3.2 battery

The §3.2 summary collapses each case to a range; the full per-pair results follow (10-seed AUROC / dangerous-cleared at 80% edible-yield, with one-sided 95% Clopper-Pearson upper bound; `n` = unique dangerous test specimens per seed, not seed-multiplied). The "dangerous-cleared" event is the venomous/toxic/model specimen cleared as the harmless look-alike at the 80%-yield threshold.

Case 3 — coral-snake mimicry (reptiles; (Pfennig et al., 2001)), n=90:

pair	I-JEPA	DINOv2	CLIP
coral / scarlet-kingsnake	0.89 / 13.8% (≤20.7%)	1.00 / 0.0% (≤3.3%)	0.98 / 0.7% (≤5.2%)
coral / milk-snake	0.94 / 4.2% (≤9.9%)	1.00 / 0.0% (≤3.3%)	1.00 / 0.0% (≤3.3%)
coral / scarlet-snake	0.84 / 22.4% (≤30.6%)	1.00 / 0.0% (≤3.3%)	1.00 / 0.1% (≤3.3%)

Case 4 — brown-recluse misidentification (arachnids; Vetter), n=90:

pair	I-JEPA	DINOv2	CLIP
recluse / southern-house-spider	0.73 / 48.9% (≤58.0%)	0.98 / 2.0% (≤6.8%)	0.88 / 18.9% (≤27.0%)
recluse / wolf-spider	0.93 / 10.3% (≤16.8%)	1.00 / 0.0% (≤3.3%)	1.00 / 0.0% (≤3.3%)
recluse / hobo-spider	0.84 / 26.0% (≤34.2%)	1.00 / 0.0% (≤3.3%)	0.96 / 4.8% (≤9.9%)

Case 5 — poison-frog mimicry (amphibians; (Symula et al., 2001) [Müllerian]; Darst & Cummings, *Nature* 2006 [Batesian]): the mimic/model species have few GBIF photographs, so this case carries the smallest test sets and widest bounds (n = 22–90). *Allobates zaparo* was topped up from 37 to 66 usable images by taking all media per occurrence (GBIF has 37 occurrences but 71 distinct image URLs). Reported direction: model / co-mimic cleared as the focal mimic.

pair	n	I-JEPA	DINOv2	CLIP
Müllerian: *variabilis* / *imitator*	30	0.89 / 18.3% (≤35.7%)	0.96 / 4.7% (≤14.9%)	0.93 / 10.3% (≤23.9%)

pair	n	I-JEPA	DINOv2	CLIP
Müllerian: *fantastica* / *imitator*	30	0.72 / 43.3% (≤59.8%)	0.99 / 0.3% (≤9.5%)	0.98 / 2.7% (≤14.9%)
Batesian: *Allobates* / *Ameerega bilinguis*	90	0.89 / 12.7% (≤19.4%)	0.99 / 0.7% (≤5.2%)	0.98 / 0.9% (≤5.2%)
Batesian: *Allobates* / *Ameerega parvula*	22	0.82 / 33.6% (≤51.5%)	0.93 / 10.0% (≤25.9%)	0.92 / 16.4% (≤36.9%)

Case 6 — deadly berries (plant fruit; foraging literature), n=90:

pair	I-JEPA	DINOv2	CLIP
elderberry / pokeweed	0.88 / 19.9% (≤28.2%)	1.00 / 0.0% (≤3.3%)	0.99 / 0.0% (≤3.3%)
wild-grape / moonseed	0.91 / 10.8% (≤18.1%)	1.00 / 0.0% (≤3.3%)	1.00 / 0.3% (≤3.3%)
bilberry / deadly-nightshade	1.00 / 0.2% (≤3.3%)	1.00 / 0.0% (≤3.3%)	1.00 / 0.0% (≤3.3%)

At n=90 a "0.0%" cell upper-bounds at 3.3%; on the small frog pairs a zero cell bounds only at ~9.5% (n=30) to ~12.7% (n=22), and DINOv2 itself reaches 10.0% (≤25.9%) on the n=22 *parvula* pair. The frog case follows the same weak→strong ordering in every pair (DINOv2 below I-JEPA point-wise throughout), but its bounds are wide enough that only the n=90 *bilinguis* pair is statistically decisive — read the frog evidence as suggestive, not confirmatory. The other four groups (n=90–150) are firm.

References

Mahmoud Assran, Quentin Duval, Ishan Misra, et al. Self-supervised learning from images with a joint-embedding predictive architecture. In *IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2023.

Xavier Bouthillier, Pierre Delaunay, Mirko Bronzi, et al. Accounting for variance in machine learning benchmarks. In *Proceedings of Machine Learning and Systems (MLSys)*, 2021.

Mostafa Dehghani, Yi Tay, Alexey A. Gritsenko, et al. The benchmark lottery. *arXiv:2107.07002*, 2021.

Maurizio Ferrari Dacrema, Paolo Cremonesi, and Dietmar Jannach. Are we really making much progress? a worrying analysis of recent neural recommendation approaches. In *ACM Conference on Recommender Systems (RecSys)*, 2019.

Dan Hendrycks, Kevin Zhao, Steven Basart, Jacob Steinhardt, and Dawn Song. Natural adversarial examples. In *IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2021.

Pang Wei Koh, Shiori Sagawa, Henrik Marklund, et al. WILDS: A benchmark of in-the-wild distribution shifts. In *International Conference on Machine Learning (ICML)*, 2021.

A. Kuznetsov et al. Ai-mediated risks and real-life challenges in mushroom foraging. *npj Science of Food*, 10: 121, 2026. doi: 10.1038/s41538-026-00752-4.

Maxime Oquab, Timothée Darcet, Théo Moutakanni, et al. DINOv2: Learning robust visual features without supervision. *Transactions on Machine Learning Research*, 2024.

David W. Pfennig, William R. Harcombe, and Karin S. Pfennig. Frequency-dependent batesian mimicry. *Nature*, 410:323, 2001. doi: 10.1038/35066628.

Alec Radford, Jong Wook Kim, Chris Hallacy, et al. Learning transferable visual models from natural language supervision. In *International Conference on Machine Learning (ICML)*, 2021.

Rebecca Symula, Rainer Schulte, and Kyle Summers. Molecular phylogenetic evidence for a mimetic radiation in peruvian poison frogs supports a müllerian mimicry hypothesis. *Proceedings of the Royal Society of London B*, 268:2415–2421, 2001.

Antonio Torralba and Alexei A. Efros. Unbiased look at dataset bias. In *IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2011.

Richard S. Vetter. Identifying and misidentifying the brown recluse spider. *Dermatology Online Journal*, 6 (1):7, 2000.