

STABLE LEARNING AND NO-ARBITRAGE PRICING BASED ON SENTIMENTS ¹

BY PATRICK H. O'CALLAGHAN ²

Australian Institute for Business and Economics

University of Queensland

5

A stable learner is a prediction model that generalizes to new samples without re-evaluation of the current kernel. In market settings, a learner is stable if, and only if, its pricing kernel is arbitrage-free. Stable pricing kernels preserve information and support the efficient markets hypothesis. We derive the current kernel on the basis of sentiments (a mapping from samples to rankings) for a broader class of stable learners. This extension of Gilboa and Schmeidler (2003) is justified by US Treasury bond market data for 1961-2023. When the data is insufficiently rich, stability requires out-of-sample validation: sentient agents may simulate or imagine the impact of novel data; artificial ones may engage in leave-one-type-out cross-validation; in market settings, one may bypass sentiments and directly rule out Dutch books. Our framework embeds current sentiments in a system of potential generalizations to enable within-sample testing of stability for any form of external validation.

10

15

20

From the past, the present acts prudently, lest it spoil future action.

Titian: Allegory of Prudence

1. INTRODUCTION

IN GENERAL, PREDICTION REQUIRES IMAGINATION: the capacity to go beyond the current training set of past cases. To paraphrase (Hume, 1739/1896, pages 9

¹I thank Itzhak Gilboa for encouraging this project at an early stage. I thank Khoa Hoang for helpful discussions and pointers to the data.

This version has time stamp 22:13 (AEST), 31st March, 2023. Past versions can be found at <https://arxiv.org/abs/1904.02934> and <https://github.com/patrickocal/stable-learner>.

²Email address p.ocallaghan@uq.edu.au and ORCID iD 0000-0003-0606-5465

and 10): *the memory is tied down, by original impressions* [past cases], *without any power of variation; in contrast, the imagination is free to transpose and change its ideas.* Well-established neuroscientific evidence points to an important role for the imagination and the degrees of freedom or flexibility that it provides in improving prediction and thus the chances of survival (Bartlett, 1932; Mullally & Maguire, 2013; Suddendorf & Corballis, 2007). In machine learning, where learners (prediction models) are artificial, the practice of splitting the data into a training set and a test set is a standard (Hastie et al., 2009). Cases in the training set resemble memories whereas cases in the test set resemble instances of the imagination in the sense that they allow the learner to explore unseen and potentially novel data.

The goal of external validation is generalization: the ability to adapt appropriately to previously unseen data. The key to generalization is stability (Bousquet & Elisseeff, 2002; Mukherjee et al., 2006; Poggio et al., 2004) Loosely speaking, a learner is stable if small changes in the data yield small changes in its predictions: continuity of the learning map from training sets to weighting functions (kernels). McCloskey and Cohen, 1989 provides famous examples of unstable learning where training on new cases causes current kernels to unravel in what is called catastrophic interference.

In this paper, we derive the learner's kernel on basis of sentiments. We formalise *sentiments* as a mapping from samples of past cases to rankings of underlying objects of interest. To account for all kinds of intelligence, we provide a framework that captures the behaviour of learners with the ability to externally validate their model. The axioms we provide are agnostic in relation to the method of external validation. The goal is instead to allow an observer to determine whether or not the learner is able to generalise to richer training sets of past cases in a consistent manner.

Stability is the key axiom in the present derivation of a learner's kernel on the basis of qualitative rankings, given any finite resampling of past cases.³ The learner's kernel is a subjective assessment of how similar past cases are to the current prediction problem. Stable learners generalize to novel case types without

³Here a *kernel* is simply a real-valued weighting function of two variables.

needing to: re-evaluate past rankings; generate intransitive rankings; or suppress novel (transitive) rankings. In essence, a stable learner’s kernel is not only stable in a literal sense, it also allows for meaningful, unbiased generalisations. 55

Stable learners stand in contrast to those that learn by doing. Children are encouraged to learn by making (noncatastrophic) mistakes. They are free to modify their beliefs and behaviour after the event: the intransitive rankings that novel case types may reveal (**weinstein**) are of little consequence. But what of a market maker that exposes their pricing kernel either directly or indirectly (by buying and selling securities) on a financial market? We show that a market maker fails to be stable if, and only if, the returns on investment that they offer provide a free lunch to other agents today (i.e. before any novel case types arrive and before they have a chance to re-evaluate). This equivalence between stability (when novel data arrives tomorrow) and no-arbitrage (today) is at the heart of the present paper and it has a number of interesting implications.⁴ 60 65

First, at the individual level, unstable learners are unlikely to survive.

Second, we can express stable pricing kernels as an empirical (geometric) mean. This means that, in stable markets (i.e. those with only stable learners), prices are efficient stores of information. A stable market price formation process is modular in cases and separable in securities. This implies that, if we are also given the order that past cases arrived, we can identify the stable market pricing kernel (i.e. the weighting function implied by market prices). 70 75

Third, what if, in advance, learners “massage” their kernels to ensure they are arbitrage-free (and thus stable)? This alternative method of external validation is reminiscent of De Finetti’s Dutch books argument for the formula of conditional probabilities: isn’t this what bookies do? But then how then should an observer distinguish the stable learners that are prudent (i.e. ones that actually use their imagination or cross-validates their model)? This observational ambivalence carries over to the aggregate level. The market pricing kernel might be stable even if none of the individual agents in the market are prudent. This is a Deus-ex-machina or invisible-hand property: the market process provides benefits independently of 80

⁴In general, the learner is stable if, and only if, its kernel forms a groupoid over eventualities. This concept corresponds to the Jacobi equations in GS03.

the wisdom of its agents. All other things being equal, markets where arbitrage is easier to exploit should have more stable market pricing kernels.

The second implication speaks directly to the timeless topic of market efficiency Fama (1970) and Malkiel (2003). Stable pricing kernels provide an explanation for why we might expect a passive (i.e. buy-and-hold) strategy to perform at least as well as any other strategy. (The semi-martingale hypothesis of Fama (1970).) This explanation is as follows. If the market pricing kernel is stable, then the success of an active strategy depends solely on its ability to predict the impact of the next case type. Whilst this does simplify the task, it also means that the impact of the next case is unlikely to be large. This is the nature of the empirical mean: the most likely cases are those that have occurred most frequently in the past. The impact of a new case is likely to be crowded out by the frequency of those that it resembles. On the flip side, unstable pricing kernels come with arbitrage opportunities and learners modifying their kernels. Moreover, by virtue of the fact that unstable market pricing kernels cannot be expressed as an empirical mean: price movements are likely to be larger; passive strategies are less attractive; active strategies may indeed be less volatile.

The third implication motivates the present generalization of GS03. Similar to GS03, the learner is endowed with a mapping from finite resamplings of past cases to rankings of eventualities (sentiments). Via the basic axioms, all learners (stable or not) are characterised by a kernel that, for every pair of eventualities, gives rise to a linear function on samples of past cases. With sufficiently rich data, prudence (or the threat of arbitrage) is superfluous. Who needs an imagination when raw memory will do? To be more precise, a translation of the main result of GS03 to the present context tells us that a learner is guaranteed to be stable whenever resampling of past cases generates a \mathcal{A} -diversity of rankings: for every subset of $k \leq 4$ eventualities, all $k!$ total orderings (of those k eventualities) feature in sentiments. The \mathcal{A} -diversity axiom is sufficient but not necessary for a kernel that, for every single eventuality, gives rise to a linear function on samples of past cases. The stability axiom allows us to substantially weaken \mathcal{A} -diversity to partial- β -diversity: for every subset of $k \leq 3$ eventualities, at least k total orderings feature in sentiments. Our main theorem then leads to the following

hypothesis: we expect unstable learning to arise in settings where \mathcal{I} -diversity fails to hold and where prudence is required to achieve stability: i.e. where the learners are inexperienced (short of relevant data) and where arbitrage opportunities are difficult to exploit. 120

Before turning to the model, let us consider some examples. These examples feature two kinds of learner: one who only looks back at past cases (within-sample, Inny), and another who also considers the potential impact of novel types of case (the prudent, Pru).

Example 1 (A setting where \mathcal{I} -diversity holds). *A tali (a primitive four-sided die) is rolled over and over again.⁵ So far, it yields only ones, twos and threes, so that the empirical distribution, conditional upon the current data, assigns zero probability to fours. By resampling (with replacement), we obtain the empirical distribution conditional on potential samples of past cases. This conditional empirical distribution induces a single ranking of the set $X = \{1, 2, 3\}$ of eventualities (outcomes) for each sample. The resulting rankings map generates the arrangement of hyperplanes in fig. 1. To each region of this arrangement, sentiments assigns a distinct total ranking. In dice-like problems, the inherent symmetry between outcomes and past cases ensures all $3! = 6$ total rankings of X arise.* 125
130

Our learners, Inny and Pru, forecast the outcome of the next roll. Both reveal plausibility rankings that agree with fig. 1. By virtue of \mathcal{I} -diversity, their ranking maps generalise to higher dimensions (four case types) without any need for external validation. The data is sufficiently rich for making predictions about X . 135

Example 2 (A non-market setting where \mathcal{I} -diversity fails to hold). *Each month, the federal reserve announces its target overnight rate. Inny and Pru wish to forecast the next announcement. The consensus is that there will be a rate rise of 0, 25 or 50 basis points. Inny and Pru have observed the same past announcements. Current circumstances are such that they both find that past cases generate just three distinct case types t_1 , t_2 and t_3 where the indices capture the (increasing) degree to which the fed is “behind the curve”. At (samples containing only cases of type) t_3 , Inny and Pru agree with the plausibility ranking $(0, 25, 50)$, so that larger* 140
145

⁵Similar to a modern tetrahedral die, the winning number lands face down.




Figure 1: Each rational point in this simplex corresponds to a sample of past case types: t_1, t_2 and t_3 . The total ranking $(1, 2, 3)$ arises at D (one is least likely and three is most likely). This reflects the relative frequencies of past cases in D . The inverse ranking $(3, 2, 1)$ arises at D' . Within each hyperplane $H^{\{x,y\}}$, faces x and y are equally plausible. The three hyperplanes form a centered arrangement.

rate rises are more likely. At t_1 , the opposite ranking holds: both find that smaller rate rises are more likely. At t_2 , they agree ranking is $(\{0, 50\}, 25)$, so that a 25 rise is most likely, and the others are equally plausible.

Figure 2 shows some subtle differences between their current (bold) rankings 150 maps. These subtle differences turn out to be substantial once we extend each learner's rankings map to a hypothetical fourth case type. To test for stability, the observer assigns t_4 the novel ranking $(25, 0, 50)$ and then extends both learners' current rankings maps to samples that include cases of type t_4 . Pru's extended rankings map can be chosen to form a congruent arrangement with six regions. In 155 contrast, Inny's extended rankings map forms an incongruent arrangement with seven regions. Samples in the seventh region give rise to the intransitive ranking $(25, 0, 50, 25)$. We may not be able to observe Pru's imagination (cross-validation method), but since incongruence is a generic property for triples of hyperplanes, we can prove that he is almost certainly using one. 160

Example 3 (a bond-market setting that we develop in section 4). Inny and Pru are now fair market makers of Zeros (zero-coupon treasury bonds).⁶ The

⁶Similar to a fair insurer, a fair market maker sets the market spread to zero.




Figure 2: In contrast with fig. 1, the current (bold) rankings maps features four regions and the current hyperplane arrangement is uncentered.

compound-interest formula for the accumulation process of such bonds is

$$a^{(x,y)} = (1 + r^{\{x,y\}})^{-x+y},$$

where $r^{\{x,y\}}$ is the yield to maturity on a forward contract that accrues interest 165
between dates x and y . If x is later than y , then the contract is to sell, and the market maker pays this yield, so that $r^{\{y,x\}} = r^{\{x,y\}}$. Let $X \subseteq \mathbb{R}_+$ index a suitable sequence of trading dates with $0 \in X$ being the spot date. The log-accumulation process has no arbitrages provided it satisfies

$$\text{for every distinct } x, y, z \in X, \quad \log a^{(x,y)} = \log a^{(x,z)} + \log a^{(z,y)}.^7 \quad (1) \quad 170$$

In eq. (6), reference to the data that drives the market-makers' prices is suppressed. When $\log a^{(x,y)}$ depends on past cases, it forms a vector that is normal (i.e. orthogonal) to a hyperplane such as those that separate regions in ranking maps such as fig. 1 or fig. 2. When the data is rich enough to generate the full diversity rankings (as in example 1), both Inny and Pru set prices that are currently 175
arbitrage-free. Otherwise, only Pru satisfies this property. This is true even

⁷To see this, suppose that, for $x = 0 < y < z$, Inny sets $a^{(x,y)} < a^{(x,z)} \cdot a^{(z,y)}$. Pru would do well to buy the spot contract (x, y) and sell both the spot contract (x, z) and the forward contract (z, y) . This yields a risk-free profit of $a^{(x,z)} \cdot a^{(z,y)} - a^{(x,y)} > 0$.

though they both observe the same collection of past cases. Equivalently, only Pru sets prices with the canonical form of an empirical (geometric) mean

$$B(x, D) = \prod_{c \in D} (1 + r_c^{\{0, x\}})^{-x/n} \quad (2)$$

for every date x and finite sample D that we can generate by resampling (with replacement) from past cases. 180

Example 4 (A multi-sectoral investment setting). Suppose Inny and Pru are potential representative agents in a multi-sectoral macroeconomy (Atalay, 2017; Long & Plosser, 1983). Instead of maturity dates, as in example 3, X is a variety of sectors with x_0 being the household (i.e. consumption sector). The following no-arbitrage equations may be derived from first-order conditions of the social planner's constrained optimisation problem provided we specify an explicit functional form for the instantaneous flow of utility. But here we take a nonparametric approach: we avoid the specification of a utility function and proceed from past cases to beliefs and decisions. 185

For every pair of sectors x and y and nonempty set D of past cases, let $S_D^{(x,y)}$ denote a suitably normalised quantity of sector x inputs that the social planner invests in order to produce sector y outputs, conditional on D ; then, for every sector z , the corresponding no-arbitrage equations are 190

$$\log S_D^{(x,y)} = \log S_D^{(x,z)} + \log S_D^{(z,y)}.^8 \quad (3)$$

Just as in example 3, when the data is sufficiently rich to generate a sufficient diversity of rankings, Inny and Pru are indistinguishable. Otherwise, only the prudent social planner, Pru, has a rankings map that satisfies the above no-arbitrage conditions; only Pru sets the required rate of return on investment (with the gross return $R(x_0, \cdot)$ to consumption normalised to one) according to the canonical form of an empirical mean 195

$$R(x, D) = \prod_{c \in D} (1 + r_c^{\{x_0, x\}})^{1/|D|} \quad (4)$$

⁸We can confirm that, for rather general model specifications, these equations hold in our traditional simulations with explicit utility functions. 200

for every sector x and finite sample D that we can generate by resampling (with replacement) from past cases; finally, only Pru satisfies the inter-sectoral (and inter-temporal) consumption Euler equations

$$\log S_D^{(x,y)} = \log R(y, D) - \log R(x, D). \quad (5)$$

Example 5 (Supervised learning). We are given a training set $D \subset X \times Y$ consisting of n i.i.d cases (i.e. examples) drawn from some unknown probability distribution on $(X \times Y)^n$. That is $D = \{(x_i, y_i) : i = 1, \dots, n\}$. The basic goal of supervised learning is to use the training set D to “learn” a function $f_D : X \rightarrow Y$ such that, for any given new value x_{new} , f_D provides an accurate prediction of the associated value y_{new} . That is, we hope that $f_D(x_{\text{new}})$ is suitably close to y_{new} .

Example 6 (Choosing factors). In examples 1 to 4, X respectively consists of a set of indices for faces of a tali, rate movements, maturity dates and sectors. A different, causal interpretation is possible if we let each $x \in X$ denote distinct combination of risk factors such as the market factor, firm size and book-to-market equity (Fama & French, 2015). Thus each x forms a model with which to forecast stock or bond market returns (Fama & French, 2015; Gu et al., 2020; Harvey & Liu, 2021). Such an interpretation brings our notation closer to the standard form that one finds in the statistical learning literature (Cucker & Smale, 2001; Mukherjee et al., 2006; Poggio & Smale, 2003).

Let (Y, \leq^*) denote a suitably ordered criterion space for evaluating models.⁹ The statement $x \leq_D x'$ is interpreted as “At D , x' is a more plausible model than x ”. The learner subscribes to this view if, and only if, the function $f_D : X \rightarrow Y$, $f_D(x') \leq^* f_D(x)$.

Our goal is to derive a stable learning map $D \mapsto f_D$ on the basis of sentiments $D \mapsto \leq_D$ where D is any finite sample of past cases.¹⁰

The rest of the paper is organised as follows. In section 2 we present the model and key definitions. In section 3 we present the axioms and main representation

⁹We might induce \leq^* via a norm on Y so that $y \leq^* y'$ if, and only if, for the true value y^* , $\|y - y^*\| \leq \|y' - y^*\|$.

¹⁰The term learning map is introduced in Mukherjee et al. (2006). There stability is defined in terms of continuity of the learning map, something that we prove in proposition 1, below.

theorem. In section 4 we develop example 3 by deriving a representation result for a bond-market setting. In section 4 we also discuss the implications for market efficiency identification of pricing kernels in more detail. In section 5 we provide a more detailed comparison of our main theorem with that of GS03 as well as matters such as second-order induction (Argenziano & Gilboa, 2019).

2. MODEL

THE PRIMITIVES OF OUR MODEL consist of the nonempty sets X and \mathbb{C}^f . For the current prediction problem, we interpret members of X as *eventualities* and members of \mathbb{C}^f as *current* cases. Our first departure from GS03 is to allow for two forms of current case. That is we take \mathbb{C}^f to be the union $\mathbb{C} \cup [f]$ of a set \mathbb{C} of constant, *past* cases and a set $[f]$ of copies of the variable, *free* case f .

Remark. *By way of analogy with computer memory, a natural implementation is as follows. Take any case $c \in \mathbb{C}^f$ to consist of a pair $p \times m$: a pointer p that references a memory location and the memory content m . For $c \in \mathbb{C}$, just as in the setting of GS03, content of m is meaningful. For $c \in [f]$, although m is empty (assigned a “garbage” value), the allocation is itself valuable. Pairs $c, d \in [f]$ are indistinguishable in terms of content and are in this sense copies of f .*

The learner is free to assign novel, imagined content to f and before going on to explore the prediction problem with this content in mind. As we will show, an observer (or more sophisticated learner) who observes past cases, but not the imagination, can nonetheless use f to analyse the learners’ behaviour.

With case *resampling* from the literature on bootstrapping in mind, let

$$\mathbb{D} \stackrel{\text{def}}{=} \{D \subseteq \mathbb{C} : \#D < \infty\}$$

denote the set of (finite) *determinate or constant databases* or memories and let \mathbb{D}^f denote the corresponding set of all finite subsets of \mathbb{C}^f . Finally, throughout the sequel, we take \mathfrak{C} to denote a member of $\{\mathbb{C}, \mathbb{C}^f\}$ without reference to the latter set. Similarly,

$$\mathfrak{D} = \begin{cases} \mathbb{D} & \text{if, and only if, } \mathfrak{C} = \mathbb{C}, \text{ and} \\ \mathbb{D}^f & \text{otherwise.} \end{cases}$$

SENTIMENTS. We now take a first step towards formalising the ranking maps of figs. 1 and 2. (For the translation from databases to rational vectors, see section A.) For each D in \mathbb{D} , the learner is endowed with a well-defined plausibility ranking \leq_D in the set $\text{rel}(X)$ of binary relations on X . Denote the symmetric and asymmetric parts of \leq_D by \simeq_D and $<_D$ respectively.

Sentiments $D \mapsto \leq_D$ are thus a vector in $\text{rel}(X)^{\mathbb{D}}$ of the form

$$\leq_{\mathbb{D}} \stackrel{\text{def}}{=} \langle \leq_D : D \in \mathbb{D} \rangle.$$

This subtle departure from the form $\{\leq_D : D \in \mathbb{D}\}$ of GS03 is closer in structure to visual representations (figs. 1 and 2) and the graph $\{D \times \leq_D : D \in \mathbb{D}\}$.¹¹

CASE TYPES. As in GS03, two past cases $c, d \in \mathbb{C}$ are of the same *case type* if, and only if, the marginal information of c is everywhere equal to the marginal information of d . Formally, $c \sim^{\star} d$ if, and only if, for every $D \in \mathbb{D}$ such that $c, d \notin D$, $\leq_{D \cup \{c\}} = \leq_{D \cup \{d\}}$. By observation 1 of GS03, \sim^{\star} is an equivalence relation on \mathbb{C} . The equivalence classes of \sim^{\star} generate a partition \mathbb{T} of case types.

We extend \sim^{\star} to $\mathbb{C}^{\mathfrak{f}}$ by taking $[f]$ to be an equivalence class of its own, so that, for every $c \in \mathbb{C}$, $c \not\sim^{\star} f$. We let $\mathbb{T}^{\mathfrak{f}}$ denote the corresponding partition of $\mathbb{C}^{\mathfrak{f}}$. Like GS03, we also extend \sim^{\star} to $\mathbb{D}^{\mathfrak{f}}$ by treating databases that contain the same number of each case type as equivalent. That is, $C \sim^{\star} D$ if, and only if, for every $t \in \mathbb{T}^{\mathfrak{f}}$, the numbers $\#(C \cap t)$ and $\#(D \cap t)$ of that case type coincide. To enable a surjection $D \mapsto \langle \#(D \cap t) : t \in \mathbb{T}^{\mathfrak{f}} \rangle$ from databases to counting vectors, we impose

Richness Assumption. *For every $t \in \mathbb{T}^{\mathfrak{f}}$, there are infinitely many cases in t .*

GENERALIZATIONS of sentiments $\leq_{\mathbb{D}}$ to $\mathbb{D}^{\mathfrak{f}}$ require a suitable structure. The following definition provides that structure. It also simplifies the exposition by accommodating generalizations that restrict attention to subsets Y of X .

Definition 1. $\mathcal{R} \stackrel{\text{def}}{=} \langle \mathcal{R}_D : D \in \mathfrak{D} \rangle$ is a generalization (or Y -generalization) of $\leq_{\mathbb{D}}$ if, for some nonempty $Y \subseteq X$, the following all hold:

1. for every $D \in \mathbb{D}$ and every $x, y \in Y$, $x \mathcal{R}_D y$ if, and only if, $x \leq_D y$,

¹¹I thank Maxwell B. Stinchcombe for bringing this point to my attention.

2. for every $D \in \mathfrak{D}$, \mathcal{R}_D belongs to $\text{rel}(Y)$,
3. for every $D \in \mathfrak{D}$ and every $c, d \in \mathfrak{C} - D$, if $c \sim^\star d$ then $\mathcal{R}_{D \cup \{c\}} = \mathcal{R}_{D \cup \{d\}}$.

A generalization $\mathcal{R}_{\mathfrak{D}}$ is proper if $\mathfrak{D} = \mathbb{D}^f$ and improper otherwise.

Observe that $\leq_{\mathbb{D}}$ is an (improper) generalization of itself: take $\mathfrak{D} = \mathbb{D}$ and $Y = X$. Indeed part 1 of the definition implies that, for every Y -generalization \mathcal{R} and every $D \in \mathbb{D}$, \mathcal{R} is simply the restriction $\leq_D \cap Y^2$ of \leq_D to Y . We therefore refer to part 1 of the definition of a generalization as the preservation or *nonrevision* condition. 285

Part 2 of the definition of a generalization ensures that, for every proper Y -generalization \mathcal{R} and every $D \in \mathbb{D}^f$, \mathcal{R}_D is a well-defined binary relation on Y . Let \mathcal{I}_D and \mathcal{P}_D respectively denote the symmetric and asymmetric parts of \mathcal{R}_D . 290

Part 3 of the definition ensures that, for proper generalizations \mathcal{R} , the partition \mathbb{T}^f of case types generated by \sim^\star is at least as fine the partition generated by the equivalence relation generated by \mathcal{R} . Two cases $c, d \in \mathbb{C}^f$ are *equivalent with respect to \mathcal{R}* , written $c \sim^{\mathcal{R}} d$, if, for every $D \in \mathbb{D}$ such that $c, d \notin D$, $\mathcal{R}_{D \cup \{c\}} = \mathcal{R}_{D \cup \{d\}}$. This notion allows us to partition the set of proper generalizations as follows. 295

Definition. A proper generalization \mathcal{R} is either regular or novel. It is novel whenever $[\mathfrak{f}]$ is a distinct equivalence class of $\sim^{\mathcal{R}}$, so that, for every $c \in \mathbb{C}$, $c \not\sim^{\mathcal{R}} \mathfrak{f}$.

For any given novel generalization \mathcal{R} , $[\mathfrak{f}]$ is the unique novel case type that $\sim^{\mathcal{R}}$ generates. We impose this restriction, not because we think the imagination of learners is constrained in this way, or because the model would not work that way, but rather because one degree of freedom is sufficient for our purposes. From an evolutionary perspective: Pru represents a minimal deviation from Inny. 300

For every regular generalization \mathcal{R} , there exists $c \in \mathbb{C}$ such that $c \sim^{\mathcal{R}} \mathfrak{f}$. Thus, there are as many regular generalizations as there are past case types (i.e. $\#\mathbb{T}$). Yet every regular Y -generalization \mathcal{R} is *equivalent* to the unique improper Y -generalization $\langle \leq_D \cap Y^2 : D \in \mathbb{D} \rangle$ in the sense of 305

Observation 1. For every regular Y -generalization \mathcal{R} and improper Y -generalization $\hat{\mathcal{R}}$, for every $C \in \mathbb{D}^f$, there exists $D \in \mathbb{D}$ such that $C \sim^{\mathcal{R}} D$ and $\mathcal{R}_C = \hat{\mathcal{R}}_D$.¹²

¹²This means that there is a canonical embedding of $\{C \times \mathcal{R}_C : C \in \mathbb{D}^f\}$ in $\{D \times \hat{\mathcal{R}}_D : D \in \mathbb{D}\}$. The converse embedding follows from the nonrevision condition of definition 1.

See proof on page 57.

310

3. AXIOMS AND MAIN RESULTS

THE BASIC AXIOMS of GS03, which we rewrite in terms of generalizations, are the following. In each of these axioms, \mathcal{R} is an arbitrary Y -generalization of $\leq_{\mathbb{D}}$.

A0 (Transitivity axiom for \mathcal{R}). *For every $D \in \mathfrak{D}$, \mathcal{R}_D is transitive.*

A1 (Completeness axiom for \mathcal{R}). *For every $D \in \mathfrak{D}$, \mathcal{R}_D is complete.*

315

A2 (Combination axiom for \mathcal{R}). *For every disjoint $C, D \in \mathfrak{D}$ and every $x, y \in Y$, if $x \mathcal{R}_C y$ and $x \mathcal{R}_D y$, then $x \mathcal{R}_{C \cup D} y$; and if $x \mathcal{P}_C y$ and $x \mathcal{R}_D y$, then $x \mathcal{P}_{C \cup D} y$.*

A3 (Archimedean axiom for \mathcal{R}). *For every disjoint $C, D \in \mathfrak{D}$ and every $x, y \in Y$, if $x \mathcal{P}_D y$, then there exists $k \in \mathbb{Z}_{++}$ such that, for every pairwise disjoint collection $\{D_j : D_j \sim^{\mathcal{R}} D \text{ and } C \cap D_j = \emptyset\}_1^k$ in \mathfrak{D} , $x \mathcal{P}_{C \cup D_1 \cup \dots \cup D_k} y$.*

320

THE DIVERSITY AXIOMS that now follow require that \mathbb{D} is sufficiently rich to support Y -generalizations \mathcal{R} with a variety of *total* orderings: i.e. complete, transitive and antisymmetric ($x \mathcal{R}_D y$ and $y \mathcal{R}_D x$ implies $x = y$). Let $\text{total}(\mathcal{R})$ denote the set $\{R : \text{for some } C \in \mathfrak{D}, R = \mathcal{R}_C \text{ is total}\}$ of total orders that feature in \mathcal{R} . For $k = 4$, the following axiom is a restatement of the diversity axiom of GS03.

325

Diversity (k -Diversity axiom). *For every $Y \subseteq X$ of cardinality $n = 2, \dots, k$, every regular Y -generalization \mathcal{R} of $\leq_{\mathbb{D}}$ is such that $\#\text{total}(\mathcal{R}) = n!$.*

We say *k -diversity holds on Z* if the axiom holds with Z in the place of X . By lowering the bar for the required number of total orders, the following axiom substantively weakens 3-diversity and *a fortiori* 4-diversity.

330

A4' (Partial 3-diversity). *For every $Y \subseteq X$ with cardinality $n = 2$ or 3 , every regular X -generalization \mathcal{R} of $\leq_{\mathbb{D}}$ is such that $\#\text{total}(\mathcal{R}) \geq n$.*

Example 2 (and example 7 of the appendix) provide examples of $\leq_{\mathbb{D}}$ satisfying A0–A4', but not 3-diversity. For settings where the basic axioms hold, the next observation shows that partial-3-diversity is equivalent to

335

A4 (Conditional-2-diversity). *For every three distinct elements $x, y, z \in X$, one of the sets $\{D' : x <_{D'} y\}$ and $\{D' : y <_{D'} x\}$ contains both C and D such that $z <_C x$ and $x <_D z$. If $\#X = 2$, then 2-diversity holds on X .*

Observation 2. *For $\leq_{\mathbb{D}}$ satisfying A0–A3, conditional-2-diversity and partial-3-diversity are equivalent.*

340

Proof. This follows by virtue of proposition 4 and the translation of section A. \square

Partial-3-diversity is therefore stronger than 2-diversity and, moreover, it plays the dual role of guaranteeing the existence and uniqueness of the representation. Via example 8, partial-3-diversity is weakest axiom with these properties.

THE STABILITY AXIOM is our final requirement. It is distinguished by the fact that it imposes structure on novel generalizations. First a definition that reduces the set of generalizations we test when establishing stability.

345

Definition. *A proper generalization \mathcal{R} of $\leq_{\mathbb{D}}$ is testworthy if it satisfies A1–A3 and, for some $D \in \mathbb{D}$ such that \mathcal{R}_D is total, $\mathcal{R}_{\mathfrak{f}}$ is the inverse of \mathcal{R}_D .¹³*

Testworthy generalizations are distinguished by the ranking at \mathfrak{f} and the fact that they need not satisfy A0. Stability requires that for any testworthy generalization we can find a suitable perturbation that also satisfies A0.

350

Definition. *Let \mathcal{R} and $\acute{\mathcal{R}}$ be generalizations of $\leq_{\mathbb{D}}$. $\acute{\mathcal{R}}$ is a perturbation of \mathcal{R} if $\acute{\mathcal{R}}_{\mathfrak{f}} = \mathcal{R}_{\mathfrak{f}}$ and, moreover, a diverse perturbation if $\#\text{total}(\acute{\mathcal{R}}) \geq \#\text{total}(\mathcal{R})$.*

A diverse perturbation of \mathcal{R} does not suppress the novel, transitive rankings that \mathcal{R} generates. If we allowed for nondiverse (i.e. dogmatic) perturbations, the learner could continue to “hide” intransitive rankings. Our main axiom is then

355

4-Stability. *For every $Y \subseteq X$ with cardinality 3 or 4, every testworthy Y -generalization of $\leq_{\mathbb{D}}$ that is novel has a diverse perturbation that satisfies A0–A3.*

Since every proper generalization is either regular or novel, the next observation confirms that 4-diversity holds whenever 4-stability holds vacuously; that is, whenever there are no novel generalizations for us to test.

360

¹³Recall that the inverse \mathcal{R}_D^{-1} of \mathcal{R}_D satisfies $x \mathcal{R}_D^{-1} y$ if, and only if, $y \mathcal{R}_D x$.

Observation 3. *Let $\leq_{\mathbb{D}}$ satisfy A0–A4. For every $Y \subseteq X$ of cardinality 3 or 4, the set of testworthy Y -generalizations is nonempty. If, for some Y , every testworthy Y -generalization is regular, then 4-diversity holds on Y .* 365

Proof. Respectively, these two statements follow via lemma 3.2 and lemma 3.1. \square

It is natural to ask whether 4-stability is simply requiring that, on Y such that $\leq_{\mathbb{J}}$ fails to satisfy 4-diversity, there exists a testworthy Y -generalization that is novel and satisfies 4-diversity. The proof of theorem 3 (see for instance fig. 5), shows otherwise. When the basic axioms hold, 4-diversity implies that the current ranking map features a centered arrangement of hyperplanes (for every $Y \subseteq X$ of cardinality four). In contrast, 4-stability does not even imply the existence of a centered extended ranking map. 370

OUR MAIN THEOREM involves real-valued function \mathbf{v} on the product $X \times \mathbb{C}$. We view \mathbf{v} as a matrix and $\mathbf{v}(x, \cdot)$ as one of its rows. The matrix \mathbf{v} is a *representation* of $\leq_{\mathbb{D}}$ whenever it satisfies 375

$$\begin{cases} \text{for every } x, y \in X \text{ and every } D \in \mathbb{D}, \\ x \leq_D y \quad \text{if, and only if,} \quad \sum_{c \in D} \mathbf{v}(x, c) \leq \sum_{c \in D} \mathbf{v}(y, c). \end{cases}$$

The matrix \mathbf{v} *respects case equivalence* (with respect to $\leq_{\mathbb{D}}$) if, for every $c, d \in \mathbb{C}$, $c \sim^* d$ if, and only if, the columns $\mathbf{v}(\cdot, c)$ and $\mathbf{v}(\cdot, d)$ are equal.

Theorem 1 (Part I, Existence). *Let there be given X , $\mathbb{C}^{\mathfrak{f}}$, $\leq_{\mathbb{D}}$ and associated generalizations, as above, such that the richness condition holds. Then (1.i) and (1.ii) are equivalent.* 380

(1.i) A0–A4 and 4-stability hold for $\leq_{\mathbb{D}}$.

(1.ii) There exists a matrix $\mathbf{v} : X \times \mathbb{C} \rightarrow \mathbb{R}$ satisfying (1.a) and (1.b):

(1.a) \mathbf{v} is a representation of $\leq_{\mathbb{D}}$ that respects case equivalence; 385

(1.b) no row of \mathbf{v} is dominated by any other row, and for every three distinct elements $x, y, z \in X$ and $\lambda \in \mathbb{R}$, $\mathbf{v}(x, \cdot) \neq \lambda \mathbf{v}(y, \cdot) + (1 - \lambda) \mathbf{v}(z, \cdot)$.¹⁴

¹⁴Observe that $\mathbf{v}(x, \cdot) - \mathbf{v}(z, \cdot)$ and $\mathbf{v}(y, \cdot) - \mathbf{v}(z, \cdot)$ are noncollinear if, and only if, the affine independence condition of (1.b) holds.

Our uniqueness result is identical to that of GS03.

Theorem 1 (Part II, Uniqueness). *If (1.i) [or (1.ii)] holds, then the matrix \mathbf{v} is unique in the following sense: for every other matrix $\mathbf{u} : X \times \mathbb{C} \rightarrow \mathbb{R}$ that represents $\leq_{\mathbb{D}}$, there is a scalar $\lambda > 0$ and a matrix $\beta : X \times \mathbb{C} \rightarrow \mathbb{R}$ with identical rows (i.e. with constant columns) such that $\mathbf{u} = \lambda \mathbf{v} + \beta$.* 390

A VARIETY OF STATISTICAL METHODS that are related to theorem 1 are discussed in GS03. In the next section, we provide a more detailed application to no-arbitrage asset pricing. This application also appeals to intermediate results in the proof of theorem 1, so let us unpack the key steps. 395

The first step in the proof of theorem 1 (see section A is to translate all the above concepts to the setting of rational vectors indexed by case types: as in figs. 1 and 2, i.e. vectors in $\mathbb{R}_+^{\mathbb{T}}$. Theorem 2 is the corresponding theorem for rational vectors. Definitions and intermediate results in the proof of theorem 2 are the subject of section B. For now, we maintain the discussion of these results in terms of samples (i.e. databases). 400

In section B, we show that A1–A4 alone yield a more general matrix representation than that of theorem 1. In particular, a matrix $v^{(\cdot, \cdot)} : X^2 \times \mathbb{C} \rightarrow \mathbb{R}$, $x \times y \times c \mapsto v_c^{(x,y)}$. For each pair $x, y \in X$, the row $v^{(x,y)}$ of this matrix is a pairwise representation provided that, for every $x, y \in X$ and $D \in \mathbb{D}$,

$$x \leq_D y \text{ if, and only if, } \sum_{c \in D} v_c^{(x,y)} \geq 0.$$

(In the context of fig. 1, the projection of each such $v^{(x,y)}$ onto case-type space is orthogonal to the hyperplane $H_+^{\{x,y\}} \subset \mathbb{R}_+^{\mathbb{T}}$.)

A4 is characterised by the property that no row of $v^{(\cdot, \cdot)}$ dominates any another; nor is any pair collinear. The final key to the proof of theorem 1, stability, and the connection with no-arbitrage pricing is 405

Definition. *For $Y \in 2^X$, $v^{(\cdot, \cdot)} : Y^2 \times \mathbb{C} \rightarrow \mathbb{R}$ satisfies the groupoid property (Jacobi identity) whenever, for every $x, y, z \in Y$, $v^{(x,z)} = v^{(x,y)} + v^{(y,z)}$.*

When the Jacobi identity holds on X , instead of the $\binom{\#X}{2}$ rows of $v^{(\cdot, \cdot)}$, we only need $\#X$ vectors in $\mathbb{R}^{\mathbb{C}}$ to summarise $\leq_{\mathbb{D}}$. This is the crux of theorem 1. 410

Corollary 1 (a characterisation of stability). *Let the number of case types be finite and let $\leq_{\mathbb{D}}$ satisfy A4. Then $\leq_{\mathbb{D}}$ satisfies 4-stability if, and only if, $\leq_{\mathbb{D}}$ has a pairwise representation $v^{(\cdot, \cdot)}$ that satisfies the Jacobi identity. Moreover, for every other pairwise representation $u^{(\cdot, \cdot)}$, there exists $\lambda > 0$, such that $u^{(\cdot, \cdot)} = \lambda v^{(\cdot, \cdot)}$.* 415

See proof on page 57.

This suffices for us to formalise the log-accumulation process of example 3.

4. APPLICATION TO NO-ARBITRAGE ASSET PRICING

Fixed income securities (henceforth *bonds*) are commonly traded over the counter with certain parties (typically investment banks) acting as market makers. Similar to bookmakers in betting markets, market makers profit from the bid-ask spread. Unlike bookmakers, however, they do seize on perceived opportunities to make money by trading with other market makers. To simplify the exposition, we assume the market makers are *fair*, so that the bid-ask spread is zero. 420

Recall that a (fixed income) *forward* is a contract between two parties to exchange a bond at a given date, price and amount in the future. Building on example 3, let X denote the set of settlement/maturity dates for forwards associated with given zero-coupon bond (a *Zero*) issued by the same entity. In particular, take $X = \{x_0, x_1, \dots\}$ such that $x_0 = 0$ and $x_k < x_{k+1}$ for all feasible k . 425

Remark (on zero-coupon bonds). *Although coupon-paying bonds are far more common, Zeros provide a natural asset for studying no-arbitrage pricing (Barillas & Nimark, 2019). Zeros are generated by “stripping” coupon-paying bonds of their coupons enroute to deriving benchmark zero-coupon yield curves.¹⁵ Since coupon stripping is reversible, restricting attention to Zeros is without loss of generality. Indeed, modulo surmountable complications relating to the uncertainty of the cashflows, the arguments that follow extend to dividend-paying stocks.* 430 435

Let D^* denote the (unique) current history of time-series data that are relevant to this market for Zeros. We generate the set \mathbb{C} of past cases by taking each case

¹⁵Recall that Zeros are a common way to derive the implied yield curve. They are generated by taking a coupon-paying bond price and subtracting of the present value of each of its coupons. These present values are calculated using price of Zeros with shorter maturities (Brealey et al., 2020).

$c \in \mathbb{C}$ to consist of market-relevant data from a given time interval (a block of time periods) in the past. Block resampling of time series data was originally developed by Kunsch (1989) and Politis and Romano (1994). (For applications to finance problems see Harvey and Liu (2021) and White (2000).) For the present purposes, blocks need to be chosen so that, for any $k > 0$ and resampling $D = \{c_1, \dots, c_k\}$, what matters is the number of repetitions of a given case type, not the order of the cases that form the pseudo-time series. In other words, $D \sim^* D'$ for any permutation D' of the cases in D . It is straightforward to verify that this is indeed the case for the stationary bootstrap of Politis and Romano (1994, section 2).

The free case \mathfrak{f} has no additional structure beyond that of sections 2 and 3.

Before turning to the interpretation of $\leq_{\mathbb{D}}$ in the present setting, we introduce current market prices. From the market maker's perspective, current prices reflect the rest of the market's view on yields to maturity conditional on D^* . Since we take current market prices (and rates) to be fixed, we suppress reference to D^* . The market maker observes past cases, and current prices and forms a view about how prices might change when the information changes. The compound-interest formula for the *market accumulation process* (at current market rates) is then

$$a^{(x,y)} = \left(1 + r^{\{x,y\}}\right)^{-x+y},$$

where $r^{\{x,y\}}$ is the market's implied forward yield for the contract that accrues interest between dates x and y . If x is later than y , then the contract is to sell, and the market maker pays this yield, so that $r^{\{y,x\}} = r^{\{x,y\}}$. For each $x \in X$, the *market price* of a Zero that pays out one-dollar at date x is defined as $b(x) \stackrel{\text{def}}{=} a^{(x,0)}$.

Let $X \subseteq \mathbb{R}_+$ index a suitable sequence of trading dates with $0 \in X$ being the spot date. It is well known that a Zero that pays out one dollar at time $x > 0$ is arbitrage-free if, and only if, the log-accumulation process satisfies

$$\text{for every } x, y, z \in X, \quad \log a^{(x,y)} = \log a^{(x,z)} + \log a^{(z,y)}.^{16} \quad (6)$$

This no-arbitrage condition is a special case of the Jacobi identity.

¹⁶To see this, w.l.o.g. suppose that, for some $x < y < z$, $a^{(x,z)} < a^{(x,y)}a^{(y,z)}$. The market maker would do well to sell the contract (x, z) and buy the contracts (x, y) and (y, z) . In the absence of counterparty risk, the difference between interest paid and received is risk-free.

We now explain how a market maker might infer her own subjective accumulation process from past cases. Our market maker thinks in terms of economic profits (relative to the market price). For every finite resampling D and every date x and y , let $x \leq_D y$ if, and only if, at D , the market maker finds y (weakly) more plausible than x in answer to the question

470

“Hold current market prices fixed and consider a one-dollar investment today. Given D , which maturity will yield a higher return?”

Remark (alternative interpretation). *It is also possible to interpret $\leq_{\mathbb{D}}$ in terms of statements of the market maker’s intention to buy or sell forwards. Let (x, y) be shorthand for the forward contract where the buyer accumulates interest between dates x and y . For $x < y$, accumulating interest over (y, x) simply means selling (x, y) . Holding current prices fixed, for each D in \mathbb{D} , we have $x \leq_D y$ if, and only if, given D , the market maker would buy (x, y) . Under this interpretation, if at D^* the market maker agrees with the market, then $x \simeq_{D^*} y$ for every $x, y \in X$. Thus, if the market maker agrees with the market, then her sentiments are centered as in fig. 1. Our point is that the market maker may not agree with the market and thus the rankings map may be uncentered as in fig. 2. To operationalise this interpretation, simply suppress reference to $a^{(\cdot)}$ in what follows.*

475

480

Next, we introduce the market maker’s (possibly negative and subjective) markup function. This is a markup relative to current rates $r^{\{\cdot\}}$. A *markup function* $\mu : X^2 \times \mathbb{C} \rightarrow \mathbb{R}$, is characterised by three conditions: for time intervals of length zero, the yield is zero; fair pricing; and case equivalence. These are, respectively, formalised as follows: for every $x, y \in X$ and every $c, d \in \mathbb{C}$, $\mu_c^{\{x, x\}} = 0$; $\mu_c^{\{y, x\}} = \mu_c^{\{x, y\}}$; and $c \sim^* d$ if, and only if, $\mu_c^{\{x, y\}} = \mu_d^{\{x, y\}}$. We extend to \mathbb{D} by taking $\mu_D^{\{x, y\}}$ to be the (geometric) *mean markup conditional on D*

$$1 + \mu_D^{\{x, y\}} = \prod_{c \in D} (1 + \mu_c^{\{x, y\}})^{\frac{1}{|D|}}.$$

The market maker’s subjective forward yield conditional on D is then the following modification of the market yield $r^{\{\cdot\}}$: for every $x, y \in X$ and $D \in \mathbb{D}$

$$1 + \rho_D^{\{x, y\}} = (1 + r^{\{x, y\}}) \cdot (1 + \mu_D^{\{x, y\}}).$$

In turn, the market maker's subjective accumulation process $A^{(\cdot, \cdot)} : X^2 \times \mathbb{D} \rightarrow \mathbb{R}$ modifies the market accumulation process $a^{(\cdot, \cdot)}$. For every $x, y \in X$ and $D \in \mathbb{D}$,

$$A_D^{(x, y)} := \left(1 + \rho_D^{\{x, y\}}\right)^{-x+y} = a^{(x, y)} \cdot \left(1 + \mu_D^{\{x, y\}}\right)^{-x+y}.$$

Note that the market maker agrees with the current market accumulation process whenever the cases in D^* are such that the positive markups countervail those that are negative. That is, for every $x, y \in X$, $\mu_{D^*}^{(x, y)} = 0$. 485

The market maker's (subjective) empirical bond price function $B : X \times \mathbb{D} \rightarrow \mathbb{R}$ modifies the market price $b : X \rightarrow \mathbb{R}$. Thus, for a Zero with a one-dollar face value the price at time x , conditional on D is

$$B(x, D) := A_D^{(x, 0)} = b(x) \cdot \left(1 + \mu_D^{\{0, x\}}\right)^{-x}.$$

This reflects the inverse relationship between bond prices and yields.

When D^* belongs to \mathbb{D} , the number of case types is finite, the following corollary of theorem 1 characterises 4-stability in the Zero-market setting.

Corollary 2. *Let $\leq_{\mathbb{D}}$ satisfy A4. Then $\leq_{\mathbb{D}}$ satisfies 4-stability if, and only if, there exists empirical implied yield and empirical bond price functions, such that* 490

$$\begin{cases} \text{for every } x, y \in X \text{ and every } D \in \mathbb{D}, \\ x \leq_D y \quad \text{if, and only if,} \quad B(x, D) \geq B(y, D). \end{cases} \quad (*)$$

Moreover, for every other empirical bond price function \hat{B} that satisfies $(*)$, there exists a scalar $\lambda > 0$ such that $\log \hat{B} = \lambda \log B$; and, for every $D \in \mathbb{D}$, the associated accumulation process $A_D^{(\cdot, \cdot)}$ is arbitrage-free. 495

Proof of corollary 2. Via corollary 1, $\leq_{\mathbb{D}}$ satisfies 4-stability if, and only if, there exists a pairwise Jacobi representation $v^{(\cdot, \cdot)} : X^2 \times \mathbb{C} \rightarrow \mathbb{R}$. Recalling that $0 \in X$, for every $x \in X$ and $D \in \mathbb{D}$, let

$$B(x, D) = \exp \left(-\frac{1}{|D|} \sum_{c \in D} v_c^{(0, x)} \right). \quad (7)$$

For the proof of $(*)$, suppose that w.l.o.g., $x \leq_D y$, so that $\sum_{c \in D} v_c^{(y, x)} \leq 0$. Via the Jacobi identity we have $\sum_{c \in D} \{v^{(y, 0)} + v^{(0, x)}\} \leq 0$. Via the representation property $v^{(y, y)} = 0$. Another application of the Jacobi identity yields $v^{(y, 0)} = -v^{(0, y)}$. Thus, 500

$$-\log B(x, D) + \log B(y, D) = \frac{1}{|D|} \sum_{c \in D} \left\{ v_c^{(0, x)} - v_c^{(0, y)} \right\} \leq 0.$$

We now show that the bond price is a suitable function of the empirical yield function. For every $x, y \in X$ and $D \in \mathbb{D}$, take $\log A_D^{(x,y)} = \frac{1}{|D|} \sum_{c \in D} v_c^{(x,y)}$. That is,

$$\log a^{(x,y)} + \frac{y-x}{|D|} \sum_{c \in D} \log(1 + \mu_c^{\{x,y\}}) = \frac{1}{|D|} \sum_{c \in D} v_c^{(x,y)}.$$

Take $D = \{c\}$ and note that the Jacobi identity implies $v_c^{(x,y)} = -v_c^{(y,x)}$. Thus

$$\log a^{(x,y)} + (y-x) \log(1 + \mu_c^{\{x,y\}}) = -\log a^{(y,x)} - (x-y) \log(1 + \mu_c^{\{y,x\}}).$$

Moreover, note that $\log a^{(x,y)} = -\log a^{(y,x)}$, so that, by cancelling terms, we obtain

$$\log(1 + \mu_c^{\{x,y\}}) = \log(1 + \mu_c^{\{y,x\}}), \quad (8)$$

so that $\mu_c^{\{x,y\}} = \mu_c^{\{y,x\}}$. Then $v_c^{(x,x)} = 0 = \log a_c^{(x,x)}$ implies that $\mu_c^{\{x,x\}} = 0$. Finally, note that for $c \sim^* d$, the property $\mu_c^{\{x,y\}} = \mu_d^{\{x,y\}}$ is inherited from $v_c^{(x,y)} = v_d^{(x,y)}$, so that we have an empirical implied yield function. All of the above arguments are reversible, so that, given the existence of such empirical yield and bond price functions satisfying (*), it follows that $\leq_{\mathbb{D}}$ satisfies \mathcal{I} -stability. 505

If \hat{B} is another empirical bond price function that satisfies (*), then via eq. (7) and corollary 1, for some $\lambda > 0$, $\log \hat{B} = \lambda \log B$. The fact that, for every D , $A_D^{(\cdot, \cdot)}$ is arbitrage-free follows by virtue of the fact that the Jacobi identity holds element-wise for $v^{(\cdot, \cdot)}$. In particular, since, for every $x, y, z \in X$ and $c \in D$, $\log A_c^{(x,y)} = v_c^{(x,y)}$, 510

$$\log A_D^{(x,y)} = \frac{1}{|D|} \sum_{c \in D} \log A_c^{(x,y)} = \frac{1}{|D|} \sum_{c \in D} \{\log A_c^{(x,z)} + \log A_c^{(z,y)}\}.$$

Taking exponents, we obtain the no-arbitrage condition $A_D^{(x,y)} = A_D^{(x,z)} \cdot A_D^{(z,y)}$. \square 515

***prop-fourdiv-empty here ***

The uniqueness result in corollary 2 is stronger than in the general setting of part II of theorem 1. This is a consequence of the fact that our empirical bond yield satisfies the property $r^{(x,x)} = 0 = \mu^{(x,x)}$, for every $x \in X$.

OUR DIVERSITY AXIOM, A4, has a straightforward interpretation in the present setting. Given \mathcal{I} -stability, \mathcal{D} -diversity implies that, for every distinct x and y , there exist c and d such that $v_c^{(x,y)} < 0 < v_d^{(x,y)}$. This, via the arguments that lead 520

to eq. (8), is equivalent to both $\mu_d^{\{x,y\}} < 0 < \mu_c^{\{x,y\}}$ and $\rho_c^{\{x,y\}} < r^{\{x,y\}} < \rho_d^{\{x,y\}}$. In words, 2-diversity requires that the market maker's data is rich enough to contain at least one case where her markup between date x and y is positive, and at least one where it is negative. Conditional-2-diversity generalizes this notion to require that for every distinct x, y, z , $\mu_C^{\{x,y\}} < 0 < \mu_D^{\{x,y\}}$ for some C and D such that $\mu_C^{\{x,z\}} \cdot \mu_D^{\{x,z\}} > 0$. An equivalent characterisation is also available in terms of the affine independence condition (1.b) of theorem 1 for the matrix $\beta : X \times \mathbb{C} \rightarrow \mathbb{R}$, $x \times c \mapsto \beta(x, c) = B(x, \{c\})$.

525

530

The considerably stronger conditional-3-diversity arises (implicitly) in the final step of our proof of theorem 3. When it holds for $\leq_{\mathbb{D}}$, stability is unnecessary (provided A0–A3 hold). In terms of mark ups, we may characterise conditional-3-diversity as, for every distinct x, y, z and w , one of the half spaces $\{D : \mu_D^{\{x,w\}} > 0\}$ or $\{D : \mu_D^{\{x,w\}} < 0\}$ contains D_1, \dots, D_6 and permutations π_1, \dots, π_6 of x, y and z such that for $i = 1, \dots, 6$,

$$\mu_{D_i}^{\{\pi_i(x), 0\}} < \mu_{D_i}^{\{\pi_i(y), 0\}} < \mu_{D_i}^{\{\pi_i(z), 0\}}.$$

Yet 4-diversity is stronger still, requiring the above to hold on both half spaces.

This disparity between conditional-2-diversity and 4-diversity reflects the value of experience or of rich data. It also reflects the value of a prudent imagination when the data fails to be rich. Outside of market settings, this may be as far as we can go, but in the present context we can say more.

535

MARKET EFFICIENCY We begin with a proposition that justifies our claim that when 4-stability holds, so will the efficient markets hypothesis in the usual (Fama, 1970) sense: passive (buy-and-hold) strategies outperform active ones. To this end, let $\beta : X \times \mathbb{D}^f \rightarrow \mathbb{R}_+$ be an empirical bond price function such that its restriction to $X \times \mathbb{D}$ coincides with B of corollary 2 and let

$$g_c(x, D) := \frac{\beta(x, D \cup \{c\}) - \beta(x, D)}{\beta(x, D)}$$

denote the *proportional price increment* for the maturity date x given the data D , following the arrival of a new case c . By new case, we mean that c can be regular or novel. Under stable pricing, the proportional price increment converges to zero exponentially.

Proposition 1. *For every $x \in X$, $D \in \mathbb{D}$ of cardinality n and $c \in \mathbb{C}^f - D$,*

$$1 + g_c(x, D) = \left((1 + \rho_c^{\{0, x\}})^{-x} / B(x, D) \right)^{\frac{1}{n+1}} \leq (1 + \rho)^{\frac{2x}{n+1}}$$

where $\rho = \operatorname{argmax} \left\{ (1 + \rho_d^{\{0, x\}})^{-x} : d \in D \cup \{c\} \right\}$.

540

Proof. Let $1, \dots, n$ be an enumeration of D and identify c with $n+1$. Moreover, let us suppress reference to $\{0, x\}$, so that $\rho_{n+1} = \rho_c^{\{0, x\}}$. We first derive the equality. By corollary 2, and the fact that β agrees with B on $X \times \mathbb{D}$, we obtain

$$\beta(x, D) = B(x, D) = \prod_i^n (1 + \rho_i)^{\frac{-x}{n}}.$$

and, by manipulating exponents, we obtain

$$\beta(x, D \cup \{c\}) = \left(\prod_i^{n+1} (1 + \rho_i)^{\frac{-x}{n}} \right)^{\frac{n}{n+1}}.$$

Substituting for $B(x, D)$ and noting that $\frac{n}{n+1} = 1 - \frac{1}{n+1}$, we arrive at the expression

$$1 + g_c(x, D) = \frac{B(x, D)^{1 - \frac{1}{n+1}} \cdot (1 + \rho_{n+1})^{\frac{-x}{n+1}}}{B(x, D)}.$$

For the inequality, note that $B(x, D)^{-1} = \exp(\frac{1}{n} \sum_1^n \nu_i)$ where $\nu_i = x \log(1 + \rho_i)$. Let $\nu_{n+1} = x \log(1 + \rho_{n+1})$. Then, for $\nu := \max\{|\nu_i| : i = 1, \dots, n+1\}$, we have

$$\frac{1}{n} \sum_1^n \nu_i \leq \left| \frac{1}{n} \sum_1^n \nu_i \right| \leq \frac{1}{n} \sum_1^n |\nu_i| \leq \nu = x \log(1 + \rho).$$

Then, since \exp is an increasing function,

$$B(x, D)^{-1} \leq \exp(\nu) = (1 + \rho)^x$$

Mutatis mutandis, the same argument yields $(1 + \rho_{n+1})^{-x} \leq (1 + \rho)^x$. Extending this pair of bounds extend to the product brings us to the desired inequality. \square

Key to the proof of proposition 1 is that the new pricing kernel β coincides with the old one B on $X \times \mathbb{D}$. Proposition 1 thus demonstrates that \mathcal{L} -stability implies the usual notion of stability of statistical learning (Bousquet & Elisseeff, 2002; Mukherjee et al., 2006; Poggio et al., 2004). That is to say it implies continuity of the learning map (here induced by the ranking map) $L : \bigsqcup_{i \geq 1} Z^n \rightarrow \mathcal{H}$ from the sample space to the hypothesis space (Mukherjee et al., 2006).

545

The following corollary of proposition 1 confirms that, for stable pricing kernels, the impact of the arrival of a given case type is decreasing in its frequency. This is a decreasing marginal impact of information property: the information carried by a common case type is already baked into the price. We discuss the impact of this result on market efficiency following the statement and proof.

Corollary 3. *For $D \in \mathbb{D}$ of cardinality n , let \mathbb{T}_D be the set of case types that feature in D and let $c, d \in \mathbb{C} - D$ be cases of type $s \in \mathbb{T}_D$. Then, for every $x \in X$,*

$$|g_d(x, D \cup \{c\})| < |g_c(x, D)|$$

and, moreover, if n_s is the frequency of s in D , then $g_c(x, D)$ tends to 0 as $n_s \rightarrow n$.

Proof. Take $\gamma_c = g_c(x, D)$ and $\gamma_d = g_d(x, D \cup \{c\})$. For any c' of type t , let $\rho_t = \rho_{c'}^{\{0, x\}}$. We claim that

$$1 + \gamma_c = \left((1 + \rho_s)^{n-n_s} / \prod_{t \neq s} (1 + \rho_t)^{n_t} \right)^{\frac{-x}{n(n+1)}}. \quad (9)$$

First note that, since, for each $t \in \mathbb{T}_D$, D contains $n_t > 0$ cases of type t , we have

$$B(x, D) = \left(\prod_{t \in \mathbb{T}_D} (1 + \rho_t)^{n_t} \right)^{\frac{-x}{n}} = \left(\prod_{t \neq s} (1 + \rho_t)^{n_t} \cdot (1 + \rho_s)^{n_s} \right)^{\frac{-x}{n}}$$

Then a substitution for $B(x, D)$ in the expression for $1 + \gamma_c$ of proposition 1 followed by a straightforward manipulation of exponents yields eq. (9).

Let θ denote the main ratio (inside the brackets) of eq. (9). For convergence of γ_c to 0 note that $n_s \rightarrow n$ implies that, for every $t \neq s$, $n_t \rightarrow 0$. Thus both the numerator and the denominator of θ tend to one and $\gamma_c \rightarrow 0$.

For monotonicity, since d also belongs to s , we extend eq. (9) to obtain

$$1 + \gamma_d = \left((1 + \rho_s)^{(n+1)-(n_s+1)} / \prod_{t \neq s} (1 + \rho_t)^{n_t} \right)^{\frac{-x}{(n+1)(n+2)}}. \quad (10)$$

Note that the exponent of the main numerator in eq. (10) simplifies to $n - n_s$. Thus, the main ratio in eq. (10) is also equal to θ . In both eq. (9) and eq. (10), the exponent involving x is negative, so that, via the connection with θ ,

$$\gamma_c < 0 \quad \text{iff} \quad \theta > 1 \quad \text{iff} \quad \gamma_d < 0. \quad (11)$$

Combining eq. (9) and eq. (10), substituting θ and simplifying the exponent:

$$\frac{1 + \gamma_d}{1 + \gamma_c} = \theta^{2x \cdot \frac{n+1}{k}}, \quad (12)$$

where $k = n(n+1)^2(n+2)$. Since the exponent here is positive, we arrive at

$$\frac{1 + \gamma_d}{1 + \gamma_c} < 1 \quad \text{iff} \quad \theta < 1. \quad (13)$$

Via eq. (11), there are two cases to consider. The most obvious is $\gamma_d, \gamma_c > 0$, for then $\theta < 1$ and $\gamma_d < \gamma_c$ follows from eq. (13). The remaining case is $\gamma_c, \gamma_d < 0$ so that, via eq. (11), $\theta > 1$. Then, since $1 + \gamma_d = 1 - |\gamma_d|$ (and, mutatis mutandis, the same is true of γ_c), via eq. (13), we arrive at

$$\frac{1 - |\gamma_d|}{1 - |\gamma_c|} > 1 \quad \text{iff} \quad \theta > 1.$$

A simple rearrangement then yields the desired inequality. \square

In terms of market efficiency, implies that, if we assume that, more often than not, the future resembles the past in the sense that past cases with a higher frequency are more likely to continue to arrive more frequently, corollary 3 means that the most likely cases are crowded out by the volume other cases of the same type. When the pricing kernel is stable, the only way an active strategy can exploit the information in past cases is to more accurately predict the type of the case. But, since the marginal return to the most frequent cases is diminishing, this will be an uphill struggle.

Yet corollary 3 goes further. It tells us that there is value in diversity, value to the information discovery process, for novel case types are more rewarding. This explains why financial institutions spend vast resources on research. When the price kernel is stable, the largest price movements come out of “left field”. This frequentist idea of market efficiency allows research and active strategies to co-exist with passive buy-and-hold. Most of the time buy-and-hold is likely to be better, but if one can also discover the nature of less frequent, novel cases prior to their arrival, one can improve by actively researching the novel case generation process. In this way, an active imagination and hard work can help an entrepreneurial market maker to go beyond prudence. (And also beyond the formal scope of this paper.)

With unstable pricing kernels, the story is very different, whilst the yield accumulation process $A = \{A_D^{(x,y)} : x, y \in X, D \in \mathbb{D}\}$ is well-defined, since the groupoid property fails to hold, the pricing kernel is not. Whilst it is mathematically possible to derive equivalent statements to proposition 1 and ?? for A , it is somewhat meaningless in the presence of the resulting arbitrage opportunities that are present and likely to cause instability to A . Instead, the value of A is that we can study out-of-equilibrium market activity in the absence of a well-defined pricing kernel (in the sense of ??). To our knowledge, this feature is absent in the literature. 590
595

Malkiel (2003, p.60) points out that “Markets can be efficient even if many market participants are quite irrational.” Here, we can show that markets can be efficient, in the sense that the pricing kernel is stable (and thus acting as if it were guided by a prudent market maker), even if no agent is prudent. Given the above discussion, this possibility should be clear when the data is rich. But what if the data fails to be \mathcal{I} -diverse? 600

Throughout the following, we suppose that conditional- \mathcal{I} -diversity holds. For in its absence, uniqueness and existence are not guaranteed.

Recall that in our derivation above, we only assumed the market’s accumulation and bond price were available conditional on the current set D^* . Suppose the analyst is able to extend the market accumulation process and condition on any finite sample and, moreover, is able to identify an arbitrage-free bond price $b : X \times \mathbb{D} \rightarrow \mathbb{R}$. (Below we provide a rudimentary algorithm for this generalization.) Then our results say that the market bond price induces a rankings map $\leq_{\mathbb{D}}$ that satisfies \mathcal{I} -stability. Prudent pricing emerges from market activity even in the absence of prudent agents. 605
610

This inductive notion of market efficiency differs from the more Bayesian and forward-looking definitions of Fama (1970) and Malkiel (2003). It is not a form of weak efficiency, for we may define cases to include more than historical data on prices. We may include features such as firm size or price-earnings data or indeed any of the many other factors that are extensively discussed in the literature (Fama & French, 2015; Gu et al., 2020; Harvey & Liu, 2021). 615

In our view, the fact that, in market settings, stable pricing can arise even

in the absence of both rich data (4-diversity) and prudent agents is a tribute to the power of markets: it is an instance of Adam Smith's invisible hand at its best. While the threat of arbitrage may supplant the imagination and provide an alternative form of external validation, stability also provides a way to identify or "memorize" past cases. This is a more operational, non-behavioural form of efficiency.

STRUCTURAL BREAKS AND SECOND-ORDER INDUCTION. Of course some new types of cases might for good reason require a re-evaluation of past ones. Such cases lead to the formation of a new rankings map $\preceq_{\mathbb{D}}$. We would expect such cases to be less common than other forms of novel case since they represent structural breaks or regime changes. Agents that are sufficiently imaginative or privately informed about such cases may be able to profit from the associated upheaval once they arrive. But novel cases that are "independent" of past cases (in the sense that they do not cause a re-evaluation) should not generate arbitrage opportunities. This is the essence of stability.

ALGORITHM FOR IDENTIFYING THE PRICING KERNEL FROM PAST CASES Under stable pricing, cases combine in a modular way, markets assimilate the information in new types of cases without re-pricing the information of old case. This is the role of the nonrevision property of generalizations (see definition 1). Price movements reflect the marginal information of new cases.

We now describe a classification method for deciding the nature of a new case and the implications for market maker(s) who themselves act as analysts. Consider the arrival of j new cases c_1, \dots, c_j . Under stable pricing, the analyst takes new price movements to reflect the market view on the information value of new cases. That is, for each $i = 1, \dots, j$, the analyst estimates the markup associated with case c_i to be $\hat{\mu}^{\{x,y\}} = b(x, D^* \cup \{c_i\}) - b(x, D^*)$. This is reasonable provided she already has used the history of prices to estimate $b : X \times \mathbb{D} \rightarrow \mathbb{R}$. Moreover suppose that, for each $i = 1, \dots, k-1$, there exists $d \in D^*$ such that for every $x, y \in X$, $\hat{\mu}_{c_i}^{\{x,y\}} = \mu_d^{\{x,y\}}$. She concludes that each of the cases c_1, \dots, c_{k-1} is a "copy" of some past case type. Then $D^* \cup \{c_1, \dots, c_{k-1}\}$ belongs to \mathbb{D} and the analyst's market model is unchanged: all that has changed is the location of the

current sample D^\star in \mathbb{D} .

Case c_k is different. The analyst finds that price movements are such that, for every $d \in D^\star$ (and hence every case in \mathbb{C}), $\hat{\mu}_{c_k}^{\{\cdot, \cdot\}} \neq \mu_d^{\{\cdot, \cdot\}}$. This reveals c_k to be a new type of case. The analyst then generalizes her model of the market to let $\hat{\mathbb{C}}$ and $\hat{\mathbb{D}}$ to respectively be the new sets of all cases and finite resamplings that she can generate from $\hat{D} = D^\star \cup \{c_k\}$. Assuming no-arbitrage/prudent pricing, there no need for her to update $\mu_d^{\{\cdot, \cdot\}}$ for $d \in D^\star$. That is, the new market pricing function $\hat{b} : X \times \hat{\mathbb{D}} \rightarrow \mathbb{R}$ satisfies $\hat{b}(\cdot, D) = b(\cdot, D)$ for every $D \in \mathbb{D}$. It is in this sense that markets deliver efficiency via prudent pricing.

Now consider two possibilities for c_{k+1}, \dots, c_j . The first is that the price movements associated with all of these cases are similar to cases in $\hat{\mathbb{D}}$. That is, for each $i = k+1, \dots, j$, there exists $d \in \hat{D}$ such that $\hat{\mu}_{c_i}^{\{\cdot, \cdot\}} = \mu_d^{\{\cdot, \cdot\}}$. This is what we would expect if new case types arrive infrequently. It would represent a consolidation of her updated model \hat{b} and $\hat{\leq}_{\hat{\mathbb{D}}}$.

The second possibility is where many of the cases c_k, \dots, c_j turn out to be novel. If new case types do indeed arrive infrequently, then it is likely to reflect a re-evaluation of cases in D^\star and a structural break from the past. In this scenario, it may well be worth checking to see if a re-evaluation or even re-specification of past cases reduces the number of case types: thus generating a more parsimonious model.

5. DISCUSSION OF THE AXIOMS AND THEOREM 1

We begin by restating the existence part of the main theorem of GS03.

Theorem (GS03, existence). *Let there be given X , \mathbb{C} and $\leq_{\mathbb{D}}$, as above, such that the richness condition holds. Then (i) and (ii) are equivalent.*

(i) A0–A3 and 4-diversity hold for $\leq_{\mathbb{D}}$.

(ii) *There exists a matrix $\mathbf{v} : X \times \mathbb{C} \rightarrow \mathbb{R}$ satisfying (a) and (b):*

(a) *\mathbf{v} is a representation of $\leq_{\mathbb{D}}$ that respects case equivalence;*

(b) *For every four distinct elements $x, y, z, w \in X$ and every $\lambda, \mu, \theta \in \mathbb{R}$ such that $\lambda + \mu + \theta = 1$, $\mathbf{v}(x, \cdot) \not\leq \lambda \mathbf{v}(y, \cdot) + \mu \mathbf{v}(z, \cdot) + \theta \mathbf{v}(w, \cdot)$. For $\#X < 4$, no row is dominated by an affine combination of other rows.*

Although diversity axioms play an important technical role, they are not obviously behavioral. Instead, diversity axioms impose restrictions on what is beyond the learner's control and on what is central to inductive inference: experience. We contend that D^* may not be so rich as to support \leq_D satisfying 4-diversity. That is to say, there may exist $Y \subseteq X$ such that $\#Y = 4$, and such that the data is 685 insufficiently rich to support all $4! = 24$ strict rankings.

A casual comparison of condition (1.b) and (b) confirms that the present framework accommodates the less experienced or equivalently those settings where the data is not rich. By doing so, we have identified an important role for the imagination and in particular, a prudent imagination. Our results show how inexperienced 690 learners that are prudent can survive the initial phase before going on to become experienced learners in their own right.

We have shown that, even in the absence of prudent learners, provided the market structure allows agents to exploit arbitrage opportunities, market pricing is prudent. By this we mean that it is as if a prudent market maker were guiding 695 the price formation process. This novel form of efficiency is grounded in inductive inference. Moreover, it implies a modular nature to the way information is built into market prices. It implies stability in the value of information in past cases.

REFERENCES

- Argenziano, R., & Gilboa, I. (2019). Second-order induction in prediction problems. *Proceedings of the National Academy of Sciences*, 116(21), 10323–10328. <https://doi.org/10.1073/pnas.1901597116> 700
- Atalay, E. (2017). How important are sectoral shocks? *American Economic Journal: Macroeconomics*, 9(4), 254–280. <https://doi.org/10.1257/mac.20160353> 705
- Barillas, F., & Nimark, K. (2019). Speculation and the bond market: An empirical no-arbitrage framework. *Management Science*, 65(9), 4179–4203. <https://doi.org/10.1287/mnsc.2018.3027>
- Bartlett, F. C. (1932). *Remembering: A study in experimental and social psychology*. Cambridge University Press. 710

- Bousquet, O., & Elisseeff, A. (2002). Stability and generalization. *The Journal of Machine Learning Research*, 2, 499–526. <https://www.jmlr.org/papers/v2/bousquet02a.html>
- Brealey, R. A., Brealey, R., Brealey, D., Brealey, R., Richard, B., Brealey, R. A., Myers, S. C., Allen, F., BREALEY, Mohanty, P., & Brealey, R. (2020). *Principles of corporate finance 13th edition*. McGraw Hill. 715
- Cucker, F., & Smale, S. (2001). On the mathematical foundations of learning. *Bulletin of the American Mathematical Society*, 39(1), 1–49. <https://doi.org/10.1090/s0273-0979-01-00923-5>
- Dimca, A. (2017). *Hyperplane arrangements*. Springer International Publishing. 720
<https://doi.org/10.1007/978-3-319-56221-6>
- Fama, E. F. (1970). Efficient capital markets: A review of theory and empirical work. *The Journal of Finance*, 25(2), 383. <https://doi.org/10.2307/2325486>
- Fama, E. F., & French, K. R. (2015). A five-factor asset pricing model. *Journal of Financial Economics*, 116(1), 1–22. <https://doi.org/10.1016/j.jfineco.2014.10.010> 725
- Gilboa, I., & Schmeidler, D. (2001). *A theory of case-based decisions*. Cambridge University Press. <https://doi.org/10.1017/cbo9780511493539>
- Gilboa, I., & Schmeidler, D. (2003). Inductive inference: An axiomatic approach. *Econometrica*, 71(1), 1–26. <https://doi.org/10.1111/1468-0262.00388> 730
- Gu, S., Kelly, B., & Xiu, D. (2020). Empirical asset pricing via machine learning (A. Karolyi, Ed.). *The Review of Financial Studies*, 33(5), 2223–2273. <https://doi.org/10.1093/rfs/hhaa009>
- Harvey, C. R., & Liu, Y. (2021). Lucky factors. *Journal of Financial Economics*, 141(2), 413–435. <https://doi.org/10.1016/j.jfineco.2021.04.014> 735
- Hastie, T., Tibshirani, R., & Friedman, J. (2009). *The elements of statistical learning*. Springer New York. <https://doi.org/10.1007/978-0-387-84858-7>
- Hume, D. (1896). *A treatise of human nature*. Clarendon Press. (Original work published 1739) 740

- Kunsch, H. R. (1989). The jackknife and the bootstrap for general stationary observations. *The Annals of Statistics*, 17(3). <https://doi.org/10.1214/aos/1176347265>
- Long, J. B., & Plosser, C. I. (1983). Real business cycles. *Journal of Political Economy*, 91(1), 39–69. <https://doi.org/10.1086/261128> 745
- Malkiel, B. G. (2003). The efficient market hypothesis and its critics. *Journal of Economic Perspectives*, 17(1), 59–82. <https://doi.org/10.1257/089533003321164958>
- McCloskey, M., & Cohen, N. J. (1989). Catastrophic interference in connectionist networks: The sequential learning problem. In *Psychology of learning and motivation* (pp. 109–165). Elsevier. [https://doi.org/10.1016/s0079-7421\(08\)60536-8](https://doi.org/10.1016/s0079-7421(08)60536-8) 750
- Mukherjee, S., Niyogi, P., Poggio, T., & Rifkin, R. (2006). Learning theory: Stability is sufficient for generalization and necessary and sufficient for consistency of empirical risk minimization. *Advances in Computational Mathematics*, 25(1-3), 161–193. <https://doi.org/10.1007/s10444-004-7634-z> 755
- Mullally, S. L., & Maguire, E. A. (2013). Memory, imagination, and predicting the future. *The Neuroscientist*, 20(3), 220–234. <https://doi.org/10.1177/1073858413495091>
- Orlik, P., & Terao, H. (1992). *Arrangements of hyperplanes*. Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-662-02772-1> 760
- Poggio, T., Rifkin, R., Mukherjee, S., & Niyogi, P. (2004). General conditions for predictivity in learning theory. *Nature*, 428(6981), 419–422. <https://doi.org/10.1038/nature02341>
- Poggio, T., & Smale, S. (2003). The mathematics of learning: Dealing with data. *Notices of the AMS*, 50(5), 537–544. <https://www.ams.org/notices/200305/fea-smale.pdf> 765
- Politis, D. N., & Romano, J. P. (1994). The stationary bootstrap. *Journal of the American Statistical Association*, 89(428), 1303–1313. <https://doi.org/10.1080/01621459.1994.10476870>
- Sagan, B. (1999). Why the characteristic polynomial factors. *Bulletin of the American Mathematical Society*, 36(2), 113–133. <https://doi.org/10.1090/s0273-0979-99-00775-2> 770

- Stanley, R. (2007). An introduction to hyperplane arrangements. In *Geometric combinatorics* (pp. 389–496). American Mathematical Society. <https://doi.org/10.1090/pcms/013/08> 775
- Suddendorf, T., & Corballis, M. C. (2007). The evolution of foresight: What is mental time travel, and is it unique to humans? *Behavioral and Brain Sciences*, 30(3), 299–313. <https://doi.org/10.1017/s0140525x07001975>
- White, H. (2000). A reality check for data snooping. *Econometrica*, 68(5), 1097–1126. <https://doi.org/10.1111/1468-0262.00152> 780

APPENDIX A. THE PROOF OF THEOREM 1

Similar to GS03, we translate the model into one where databases are represented by vectors, the dimensions of which are case types. To allow us to focus on aspects of the present model, proceed directly to rational vectors and present the axioms and a corresponding theorem (theorem 2) which, as we confirm, holds if, and only if, theorem 1 does. The proof of theorem 2 can be found in section B. 785

CASE TYPES AS DIMENSIONS. From our definition of case types in section 2, $\mathbb{T} = \mathbb{C}_{/\sim\star}$ and $\mathbb{T}^f \stackrel{\text{def}}{=} \mathbb{T} \cup [\mathfrak{f}]$. Let \mathfrak{T} be a free variable in $\{\mathbb{T}, \mathbb{T}^f\}$. When no possible confusion should arise, we use \mathfrak{f} as shorthand for $[\mathfrak{f}]$. It is straightforward to show that the following construction would work if instead we were to work with any partition \mathbb{T} of \mathbb{C} that is at least as fine as \mathbb{T} . The present construction is the one with the lowest feasible number $\#\mathbb{T}$ of dimensions. 790

TRANSLATION TO COUNTING VECTORS. Let \mathbb{Z}_+ denote the set of nonnegative integers and \mathbb{Z}_{++} those that are (strictly) positive. Let $\mathbb{L} \subseteq \mathbb{Z}_+^{\mathbb{T}}$ denote the set of counting vectors $L : \mathbb{T} \rightarrow \mathbb{Z}_+$ such that $\{t : L(t) \neq 0\}$ is finite and let \mathbb{L}^f denote the corresponding subset of $\mathbb{Z}_+^{\mathbb{T}^f}$. Then let 795

$$\mathbb{L} = \begin{cases} \mathbb{L} & \text{if, and only if, } \mathfrak{T} = \mathbb{T}, \text{ and} \\ \mathbb{L}^f & \text{otherwise.} \end{cases}$$

Modulo notation, the following construction is identical to GS03. For every $D \in \mathbb{D}$, let $L_D : \mathbb{T} \rightarrow \mathbb{Z}_+$ denote the function $t \mapsto L_D(t) = \#(D \cap t)$. For each $D \in \mathbb{D}$, let $\leq_{L_D} \stackrel{\text{def}}{=} \leq_D$. We need to establish that $\leq_{\mathbb{L}} \stackrel{\text{def}}{=} \langle \leq_L : L \in \mathbb{L} \rangle$ is well-defined. For 800

every $L \in \mathbb{L}$, the richness assumption (on \mathbb{T}^f) guarantees the existence of $D \in \mathbb{D}$ such that $L_D = L$. By definition, \sim^* is such that, for every $C, D \in \mathbb{D}$, $C \sim^* D$ if, and only if, $L_C = L_D$. Straightforward mathematical induction on the cardinality of C shows that $C \sim^* D$ implies $\leq_C = \leq_D$. This construction of $\leq_{\mathbb{L}}$ ensures that the same notion of equivalence that we introduced in observation 1 also applies here. Thus, $\leq_{\mathbb{L}} \equiv \leq_{\mathbb{D}}$. 805

TRANSLATION TO RATIONAL VECTORS. Similarly, let \mathbb{Q}_+ denote the nonnegative rationals and \mathbb{Q}_{++} those that are (strictly) positive. Take $\mathbb{J} \subseteq \mathbb{Q}_+^{\mathbb{T}}$ to be the set of rational vectors with $\{t \in \mathbb{T} : J(t) \neq 0\}$ finite and take \mathbb{J}^f to denote the corresponding subset of $\mathbb{Q}_+^{\mathbb{T}^f}$. For each $J \in \mathbb{J}$, by virtue of the fact that \mathbb{Z}_{++} is well-ordered and J has finite support, there exists (unique) minimal $k_J \in \mathbb{Z}_{++}$ such that $L_J \stackrel{\text{def}}{=} k_J J$ belongs to \mathbb{L} . Let $\leq_J \stackrel{\text{def}}{=} \leq_{L_J}$. (This definition acquires meaning below once we translate and apply the combination axiom.) In this way, $\leq_{\mathbb{J}} = \langle \leq_J : J \in \mathbb{J} \rangle$ is well-defined, and we may introduce axioms for $\leq_{\mathbb{J}}$ directly: i.e. without first introducing axioms for $\leq_{\mathbb{L}}$. We first demonstrate that $\leq_{\mathbb{J}}$ and $\leq_{\mathbb{D}}$ are equivalent. First note that, for every $I, J \in \mathbb{J}$ such that $L_I = L_J$, $\leq_I = \leq_J$. Then, let $L' = L_J$ and take any D such that $L_D = L'$. Then $\leq_J = \leq_D$. The reverse embedding follows by virtue of the fact that $\mathbb{L} \subset \mathbb{J}$. Thus, $\leq_{\mathbb{J}} \equiv \leq_{\mathbb{D}}$. 810
815

CONSTRUCTION OF GENERALIZATIONS OF $\leq_{\mathbb{J}}$. We follow common practice by letting 2^X denote the collection of nonempty subsets $Y \subseteq X$. For each $Y \in 2^X$, we will denote the set of regular, novel and testworthy Y -generalizations (of $\leq_{\mathbb{D}}$ or $\leq_{\mathbb{J}}$) by $\text{reg}(Y, \cdot)$, $\text{nov}(Y, \cdot)$ and $\text{test}(Y, \cdot)$ respectively. Recalling that every Y -generalization is either regular or novel, let $\text{ext}(Y, \cdot)$ denote the set of all Y -generalizations. We now clarify what it means to be a generalization of $\leq_{\mathbb{J}}$. 820

For each $t \in \mathfrak{T}$, we take $\delta_t : \mathfrak{T} \rightarrow \mathbb{R}$ to be the function satisfying $\delta_t(s) = 1$ if $s = t$ and $\delta_t(s) = 0$ otherwise. (When \mathfrak{T} is finite, these are simply the basis vectors for $\mathbb{R}^{\mathfrak{T}}$.) When we wish to emphasise that the vectors belong to in $\mathbb{R}^{\mathbb{T}^f}$, then, for each \mathbb{T}^f , we will write δ_t^f . Let 825

$$\mathbb{I} = \begin{cases} \mathbb{J} & \text{if, and only if, } \mathfrak{T} = \mathbb{T}, \text{ and} \\ \mathbb{J}^f & \text{otherwise.} \end{cases}$$

For every $I \in \mathbb{J}$ and $J \in \mathfrak{J}$, we write $I \equiv J$ whenever $I = J$ or $J = I \times 0$. (In the latter case, $J(t) = I(t)$ for every $t \in \mathbb{T}$ and $J(\mathfrak{f}) = 0$.) This notion reflects the fact that, for the purposes of the present model, such I and J are equivalent. 830

Definition 2. $\mathcal{R} = \langle \mathcal{R}_J : J \in \mathfrak{J} \rangle$ is a generalization, and in particular a Y -generalization, of $\leq_{\mathbb{J}}$ if, and only if, for some nonempty $Y \subseteq X$ both the following hold 835

1. for every $J \in \mathfrak{J}$, $\mathcal{R}_J \in \text{rel}(Y)$, $\mathcal{I}_J \stackrel{\text{def}}{=} \mathcal{R}_J \cap \mathcal{R}_J^{-1}$ and $\mathcal{P}_J \stackrel{\text{def}}{=} \mathcal{R}_J - \mathcal{R}_J^{-1}$;
2. for every $J \in \mathbb{J}$ and $L \in \mathfrak{J}$ such that $J \equiv L$, $\mathcal{R}_L = \leq_J \cap (Y^2)$.

A generalization $\mathcal{R}_{\mathfrak{J}}$ (of $\leq_{\mathbb{J}}$) is proper if $\mathfrak{J} = \mathbb{J}^{\mathfrak{f}}$ and improper otherwise. A proper generalization is either regular or novel. \mathcal{R} is novel if, for every $s \in \mathbb{T}$, there exists I in \mathbb{J} such that, for $J = I \times 0$ (in $\mathbb{J}^{\mathfrak{f}}$), we have $\mathcal{R}_{J+\delta_s^{\mathfrak{f}}} \neq \mathcal{R}_{J+\delta_{\mathfrak{f}}^{\mathfrak{f}}}$. 840

For every regular Y -generalization \mathcal{R} of $\leq_{\mathbb{D}}$ such that $Y = X$, observation 1 implies $\mathcal{R} \equiv \leq_{\mathbb{D}}$. And, via $\leq_{\mathbb{J}} \equiv \leq_{\mathbb{D}}$ and transitivity of equivalence, we conclude that \mathcal{R} is equivalent to $\leq_{\mathbb{J}}$. Two sets of generalizations are isomorphic if there exists a canonical isomorphism between equivalent generalizations.

Lemma 1.1 (proof on page 57). For every $Y \in 2^X$, $\text{reg}(Y, \leq_{\mathbb{J}})$ is isomorphic to $\text{reg}(Y, \leq_{\mathbb{D}})$ and $\text{nov}(Y, \leq_{\mathbb{J}})$ is isomorphic to $\text{nov}(Y, \leq_{\mathbb{D}})$. 845

AXIOMS AND THEOREM. We restate the axioms for Y -generalizations \mathcal{R} of $\leq_{\mathbb{J}}$.

A0^b For every $J \in \mathfrak{J}$, \mathcal{R}_J is transitive on Y .

A1^b For every $J \in \mathfrak{J}$, \mathcal{R}_J complete on Y .

A2^b For every $I, J \in \mathfrak{J}$, every $x, y \in Y$ and every $\lambda, \mu \in \mathbb{Q}_{++}$, if $x \mathcal{R}_I y$ and $x \mathcal{R}_J y$, then $x \mathcal{R}_{\lambda I + \mu J} y$; moreover, if $x \mathcal{P}_I y$ and $x \mathcal{R}_J y$, then $x \mathcal{P}_{\lambda I + \mu J} y$. 850

A3^b For every $I, J \in \mathfrak{J}$ and every $x, y \in Y$ if $x \mathcal{P}_J y$, then there exists $0 < \lambda < 1$ such that, for every $\mu \in \mathbb{Q} \cap (\lambda, 1)$, $x \mathcal{P}_{(1-\mu)I + \mu J} y$.

For $k = 2, 3, 4$, k -diversity is defined for generalizations of $\leq_{\mathbb{J}}$ in exactly the same way. We continue to use the term k -diversity in this setting. The following are conditional-2-diversity and partial-3-diversity respectively. 855

A4^b For every three distinct elements $x, y, z \in Y$, one of the two subsets $\{J' : x <_{J'} y\}$ and $\{J' : y <_{J'} x\}$ of \mathbb{J} contains both I and J such that $z <_I x$ and $x <_J z$. If $\#Y = 2$, then 2-diversity holds on Y .

A4'^b For every $Y' \subseteq Y$ with cardinality $n = 2$ or 3 , every Y' -generalization \mathcal{R} of $\leq_{\mathbb{J}}$ is such that $\#\text{total}(\mathcal{R}) \geq n$. 860

A proper generalization \mathcal{R} of $\leq_{\mathbb{J}}$ is *testworthy* if it satisfies A1^b–A3^b and, for some $J \in \mathbb{J}$ such that $\mathcal{R}_{J \times 0}$ is total, $\mathcal{R}_{\dagger} = \mathcal{R}_{J \times 0}^{-1}$. Thus, for each $Y \in 2^X$, $\text{test}(Y, \leq_{\mathbb{D}}) \simeq \text{test}(Y, \leq_{\mathbb{J}})$. For any pair of generalizations \mathcal{R} and $\hat{\mathcal{R}}$, $\hat{\mathcal{R}}$ is a perturbation of \mathcal{R} if $\mathcal{R}_{\dagger} = \hat{\mathcal{R}}_{\dagger}$. Moreover, $\hat{\mathcal{R}}$ is a diverse perturbation if $\#\text{total}(\hat{\mathcal{R}}) \leq \#\text{total}(\mathcal{R})$. 865

4-P^b For every $Y \subseteq X$ of cardinality 3 or 4, every testworthy Y -generalization of $\leq_{\mathbb{J}}$ that is novel has a diverse perturbation that satisfies A0^b–A3^b.

The following result corresponds to claim 2 of GS03. Its proof is a consequence of mathematical induction and the combination axiom. 870

Lemma 1.2. *If $\mathcal{R}_{\mathfrak{J}}$ and $\hat{\mathcal{R}}_{\mathfrak{E}}$ are equivalent and the latter satisfies A2, then for every $J \in \mathfrak{J}$ and every rational number $q > 0$, we have $\mathcal{R}_{qJ} = \mathcal{R}_J$.*

The fact that $\leq_{\mathbb{J}} \equiv \leq_{\mathbb{D}}$ immediately implies that $\leq_{\mathbb{J}}$ satisfies A0^b, A1^b and A4^b if, and only if, the corresponding axiom holds for $\leq_{\mathbb{D}}$. In general, we have the following result, which then also yields the equivalence for the stability axiom. 875

Lemma 1.3 (proof on page 58). *For $\mathcal{R}_{\mathfrak{J}} \equiv \hat{\mathcal{R}}_{\mathfrak{D}}$, $\mathcal{R}_{\mathfrak{J}}$ satisfies A2^b–A3^b if, and only if, $\hat{\mathcal{R}}_{\mathfrak{D}}$ satisfies A2–A3.*

The matrix $\mathbf{v} : X \times \mathbb{T} \rightarrow R$ is a *representation of $\leq_{\mathbb{J}}$* whenever it satisfies

$$\begin{cases} \text{for every } x, y \in X \text{ and every } J \in \mathbb{J}, \\ x \leq_J y \quad \text{if, and only if,} \quad \sum_{t \in \mathbb{T}} \mathbf{v}(x, t)J(t) \leq \sum_{t \in \mathbb{T}} \mathbf{v}(y, t)J(t). \end{cases} \quad (\text{b})$$

We observe that, via the definition of case types, there exists a representation of $\leq_{\mathbb{D}}$ that respects case equivalence if, and only if there exists a representation of $\leq_{\mathbb{J}}$. The above translation and results imply that theorem 1 is equivalent to 880

Theorem 2. *Let there be given X , \mathbb{T}^f , $\leq_{\mathbb{J}}$ and associated generalizations, as above. Then (2.i) and (2.ii) are equivalent.*

(2.i) $A0^b$ – $A4^b$ and 4 - P^b hold for $\leq_{\mathbb{J}}$ on X .

885

(2.ii) *There exists a matrix $\mathbf{v} : X \times \mathbb{T} \rightarrow \mathbb{R}$ that satisfies both:*

(2.a) \mathbf{v} is a representation of $\leq_{\mathbb{J}}$; and

(2.b) *no row of \mathbf{v} is dominated by any other row, and, for every three distinct elements $x, y, z \in X$, $\mathbf{v}(x, \cdot) - \mathbf{v}(z, \cdot)$ and $\mathbf{v}(y, \cdot) - \mathbf{v}(z, \cdot)$ are noncollinear (i.e. linearly independent).*

890

Moreover, \mathbf{v} is unique in the sense of theorem 1 part II, with (2.ii) replacing (1.ii) and \mathbb{T} replacing \mathbb{C} .

APPENDIX B. THE PROOF OF THEOREM 2

For any pair of vectors $\acute{v}, J : \mathfrak{T} \rightarrow \mathbb{R}$ such that $J \in \mathfrak{J}$, the linear operator

$$J \mapsto \langle \acute{v}, J \rangle \stackrel{\text{def}}{=} \sum_{\{t: J(t) > 0\}} v(t) \cdot J(t).$$

895

is well-defined and real-valued by virtue of the fact that J has finite support.

In our proof, we build on GS03 to directly prove all results regardless of the cardinality of X and \mathbb{T} . To facilitate this approach, we first introduce the notion of an essentialization.

Let $\mathbb{R}^{\oplus \mathfrak{T}}$ denote the vectors in $\mathbb{R}^{\mathfrak{T}}$ that have finite support and observe that $\mathfrak{J} \subseteq \mathbb{R}^{\oplus \mathfrak{T}}$ is the dense subset of rational vectors. Given a vector $\acute{v} : \mathfrak{T} \rightarrow \mathbb{R}$, we associate the following subsets of $\mathbb{R}^{\oplus \mathfrak{T}}$: $\acute{H} = \{J : \langle \acute{v}, J \rangle = 0\}$, $\acute{G} = \{J : \langle \acute{v}, J \rangle > 0\}$, and $\acute{F} = \{J : \langle \acute{v}, J \rangle \geq 0\}$. For any finite collection $\acute{\mathcal{V}} = \{\acute{v}_1, \dots, \acute{v}_n\}$ of such vectors, let $\acute{\mathcal{H}}$ denote the associated collection or *arrangement* of hyperplanes in $\mathbb{R}^{\oplus \mathfrak{T}}$. let $\acute{S} = \text{span } \acute{\mathcal{V}}$ denote the \mathbf{r} -dimensional linear span of $\acute{\mathcal{V}}$. Then \acute{S} is a well-defined inner-product space in its own right and let $\langle \cdot, \cdot \rangle_{\acute{S}} : \acute{S}^2 \rightarrow \mathbb{R}$ denote the inner product. Let $\acute{p} : \mathbb{R}^{\oplus \mathfrak{T}} \rightarrow \acute{S}$ denote the orthogonal projection. Then observe that, for every $i = 1, \dots, n$, and $J \in \mathbb{R}^{\oplus \mathfrak{T}}$, $\langle \acute{v}_i, J \rangle = \langle \acute{v}_i, \acute{p}(J) \rangle$. Moreover, note that, since $\acute{v}_i \in \acute{S}$, $\langle \acute{v}_i, J \rangle = \langle \acute{v}_i, \acute{p}(J) \rangle_{\acute{S}}$. The *essentialization* $\mathcal{H}_{\acute{S}}$ of the arrangement $\acute{\mathcal{H}}$

900

905

is the arrangement we obtain by orthogonally projecting $\acute{\mathcal{H}}$ onto \acute{S} . That is, for every $H_{\acute{S}} \in \mathcal{H}_{\acute{S}}$, there exists $\acute{H} \in \acute{\mathcal{H}}$ such that $\acute{H} = \acute{p}^{-1}(H_{\acute{S}})$. In the literature on arrangements of hyperplanes, it is common to work with the essentialization of an arrangement by default. We therefore identify $\acute{\mathcal{H}}$ and $\mathcal{H}_{\acute{S}}$ and suppress reference to the latter subscript whenever no possible confusion may arise. The main benefit of essentializations is that they will allow us to work in finite dimensions whenever we consider a finite subset $Y \subseteq X$: regardless of the cardinality of \mathfrak{T} .

Our domain of interest is $\mathbb{R}_+^{\oplus \mathfrak{T}}$ and not the whole of $\mathbb{R}^{\oplus \mathfrak{T}}$. In order to be able to apply results from the literature on hyperplane arrangements without adjusting for boundaries, we find it useful to work within the relative interior \acute{S}_{++} of $\acute{S}_+ = \acute{p}(\mathbb{R}_+^{\oplus \mathfrak{T}})$. Observe \acute{S}_{++} is an open subset of the \acute{r} -dimensional linear space \acute{S} . For $\acute{H} \in \mathcal{H}$, take $\acute{H}_{++} = \acute{H} \cap \acute{S}_{++}$ to be the (strictly) *positive* null-space of $\langle \acute{v}, \cdot \rangle_{\acute{S}}$ and, for $0 \not\leq \acute{v} \not\leq 0$, \acute{G}_{++} and \acute{F}_{++} are, respectively, the open and closed half-spaces of \acute{S}_{++} associated with \acute{v} . (For such \acute{v} , we also refer to \acute{H}_{++} as a hyperplane in \acute{S}_{++} .) We refer to the non-negative counterpart of these sets as \acute{H}_+ , \acute{G}_+ and \acute{F}_+ . For any finite $Y \in 2^X$ consider the matrix $\acute{v}^{(\cdot, \cdot)} : Y^2 \times \mathfrak{T} \rightarrow \mathbb{R}$. For a given $x, y \in Y$ and row $\acute{v}^{(x, y)} : \mathfrak{T} \rightarrow \mathbb{R}$ of $\acute{v}^{(\cdot, \cdot)}$, the associated sets are $\acute{H}_{++}^{\{x, y\}}$, $\acute{G}_{++}^{(x, y)}$ and $\acute{F}_{++}^{(x, y)}$ respectively. STEP B.1 (characterisation of A1^b–A3^b, 2-diversity and novel generalizations). The following proposition corresponds to lemma 1 of GS03 and gives meaning to the statement “the arrangement generated by a generalization”.

PROPOSITION 2. *For every $Y \subseteq X$, A1^b–A3^b and 2-diversity hold for the Y -generalization $\acute{\mathcal{R}}$ if, and only if, there exists $\acute{v}^{(\cdot, \cdot)} : Y^2 \times \mathbb{T} \rightarrow \mathbb{R}$ such that,*

- (i) *for every $x, y \in Y$ and $J \in \mathfrak{J}$, $x \acute{\mathcal{R}}_J y$ if, and only if, $\langle \acute{v}^{(x, y)}, J \rangle > 0$; and*
- (ii) *for every $x, y \in Y$, there exists $s, t \in \mathbb{T}$ such that $\acute{v}^{(x, y)}(s) < 0 < \acute{v}^{(x, y)}(t)$.*

Moreover, $\acute{v}^{(x, y)}$ is unique upto multiplication by a positive scalar.

See proof on page 59. We refer to a matrix $\acute{v}^{(\cdot, \cdot)}$ that satisfies condition (ii) of proposition 2 as a *2-diverse matrix*. We refer to a matrix $\acute{v}^{(\cdot, \cdot)}$ that satisfies proposition 2 as a 2-diverse (pairwise) matrix representation of \mathcal{R} .

LEMMA 2.1 (two-diverse pairwise representation). *Let $\acute{\mathcal{R}}$ be a Y -generalization of $\leq_{\mathbb{J}}$. $\acute{\mathcal{R}}$ satisfies A1^b–A3^b and 2-diversity holds on Y , if, and only if, there exists a*

matrix $\acute{v}^{(\cdot, \cdot)} : Y^2 \times \mathfrak{T} \rightarrow \mathbb{R}$ such that, for every $x, y \in Y$, row $\acute{v}^{(x, y)} : \mathfrak{T} \rightarrow \mathbb{R}$ and its associated spaces satisfy 940

- (i) $\acute{H}_+^{\{x, y\}} \cap \mathbb{Q}^\mathfrak{T} = \{J : x \acute{\mathcal{I}}_J y\}$ and $\acute{G}_+^{(x, y)} \cap \mathbb{Q}^\mathfrak{T} = \{J : x \acute{\mathcal{P}}_J y\}$,
- (ii) $\acute{G}_{++}^{(x, y)}$ and $\acute{G}_{++}^{(y, x)}$ are both nonempty if $x \neq y$ and both empty otherwise,
- (iii) $\acute{H}_{++}^{\{y, x\}} = \acute{H}_{++}^{\{x, y\}}$ (and in particular $\acute{v}^{(y, x)} = -\acute{v}^{(x, y)}$),
- (iv) $\acute{H}_{++}^{\{x, y\}}$ is the unique hyperplane in $\mathbb{R}_{++}^\mathfrak{T}$ that separates $\{J : x \acute{\mathcal{P}}_J y\}$ and $\{J : y \acute{\mathcal{P}}_J x\}$. 945

Moreover, $\acute{\mathcal{R}}$ is novel if, and only if, for every $t \neq \mathfrak{f}$, $\acute{v}^{(\cdot, \cdot)}(t) \neq \acute{v}^{(\cdot, \cdot)}(\mathfrak{f})$.

See proof on page 59.

AN *arrangement* is a collection of hyperplanes in $\mathbb{R}_{++}^\mathfrak{T}$ or $\mathbb{R}^\mathfrak{T}$. (A *hyperplane* in $\mathbb{R}_{++}^\mathfrak{T}$ is the positive kernel of some nonzero vector.) The chief result, from the mathematics of arrangements, to which we extensively appeal is Zaslavsky's theorem 950
two form of which we state below. For any given generalization \mathcal{R} , Zaslavsky's theorem allows us to use information about the intersections of hyperplanes in the arrangement to identify $\#\text{total}(\mathcal{R})$. It does so by counting the collection \mathcal{G}_{++} of open and connected subsets of $\mathbb{R}^\mathfrak{T} - \bigcup\{H_{++} : H_{++} \in \mathcal{A}\}$ are the *chambers* or *regions* 955
of the arrangement. In the present setting, each chamber corresponds to a *CAR ranking* of the elements of Y : a complete, antisymmetric and reflexive (but possibly intransitive) ranking R . Every CAR ranking can be succinctly represented as a CAR list. For instance, take $Y = \{x, y, z\}$ and $x R y R z$, then the corresponding 960
list

$$l = \begin{cases} (x, y, z) & \text{if } R \text{ is transitive} \\ (x, y, z, x) & \text{if } R \text{ is intransitive.}^{17} \end{cases}$$

The notation generalizes without exception to sets of cardinality 4. E.g., (x, y, z, x, w) represents the CAR ranking that is intransitive over $\{x, y, z\}$ and such that w dominates every other member.

THE INTERSECTION SEMILATTICE of any arrangement \mathcal{A} is the partially ordered 965
(by reverse inclusion) set \mathcal{L} of intersections of members of \mathcal{A} . The unique minimal

element is obtained by taking the intersection A^\emptyset over the empty subarrangement \mathcal{A}^\emptyset of \mathcal{A} to obtain the ambient space itself. That is $A^\emptyset = \mathbb{R}^\mathfrak{T}$ or $\mathbb{R}_{++}^\mathfrak{T}$, depending on whether we are considering the lattice \mathcal{L} or the lattice \mathcal{L}_{++} respectively. In GS03, as a consequence of 4-diversity, \mathcal{H}_{++} is always central. In our setting, it is only \mathcal{H} that is guaranteed to be central. In general an arrangement is central, if, and only if, its intersection semilattice has a unique maximal element (Stanley, 2007, proposition 2.3). Thus, if \mathcal{H}_{++} is *centerless*, then \mathcal{L}_{++} is a meet semilattice with multiple maxima: as in example 7. Extending our notation: if $Y = \{x, y, z, w\}$, then the unique intersection A^Y is the (nonempty) *center* of $\mathcal{A}^Y = \mathcal{H}$. By $A^{\{x,y,z\}}$, we mean the intersection over $\mathcal{A}^{\{x,y,z\}} \stackrel{\text{def}}{=} \{H^{\{i,j\}} : i \neq j \text{ in } \{x, y, z\}\}$. Finally, by $A^{\{x,y\}\{z,w\}}$, we mean the intersection over $\mathcal{A}^{\{x,y\}\{z,w\}} \stackrel{\text{def}}{=} \{H^{\{x,y\}}, H^{\{z,w\}}\}$.

ZASLAVSKI'S THEOREM provides two distinct methods for counting the number of regions in an arrangement. The first states that $\#\mathcal{G}$ is equal to the sum of the absolute values of the Möbius function $\mu : \mathcal{L} \rightarrow \mathbb{Z}$ which is defined recursively via

$$\mu(A) = \begin{cases} 1 & \text{if } A = A^\emptyset \\ -\sum\{\mu(B) : A \subsetneq B\} & \text{otherwise.}^{18} \end{cases} \quad (14)$$

The above definition of Zaslavski's theorem is explicitly provided by Sagan (1999). Specialised to the present setting, the more common (see Dimca, 2017; Orlik & Terao, 1992; Stanley, 2007) “rank” version of Zaslavski's theorem is

$$\#\mathcal{G} = \sum_{\substack{\mathcal{A} \subseteq \mathcal{H} \\ \mathcal{A} \text{ central}}} (-1)^{|\mathcal{A}| - \text{rank}(\mathcal{A})}, \quad (15)$$

where *central* means that $\bigcap\{H : H \in \mathcal{A}\}$ is nonempty, and $\text{rank}(\mathcal{A})$ is the dimension of the space spanned by the normals to the hyperplanes in \mathcal{A} .¹⁹

EXAMPLE 7 (a comparison of \mathcal{L} and \mathcal{L}_{++}). Let $X = \{x, y, z\}$, $\mathbb{T} = \{s, t\}$, and $u^{(x,y)} = 1 \times -1$ and $u^{(y,z)} = 2 \times -1$ denote vectors in $\mathbb{R}^\mathbb{T}$. We now appeal to the Jacobi identity and take $u^{(x,z)} = u^{(x,y)} + u^{(y,z)} = 3 \times -2$ and extend to the remaining pairs in X^2 using lemma 2.1. Since these vectors are pairwise noncollinear and

¹⁹Equivalently, $\text{rank}(\mathcal{A})$ is the dimension of the orthogonal complement in $\mathbb{R}^\mathfrak{T}$ (or $\mathbb{R}_{++}^\mathfrak{T}$) of the intersection over \mathcal{A} . Since the intersection over the empty arrangement \mathcal{A}^\emptyset is the ambient space $\mathbb{R}^\mathfrak{T}$ (or $\mathbb{R}_{++}^\mathfrak{T}$), the only vector in $\mathbb{R}^\mathfrak{T}$ that is orthogonal to the ambient space is 0, $\text{rank}(\mathcal{A}^\emptyset) = 0$.




Figure 3: The intersection semilattice $\mathcal{L}_{++} = \mathcal{L} - A^{\{x,y,z\}}$.

$\#\mathbb{T} = 2$, the associated arrangement $\mathcal{H}_{++} = \{H_{++}^{\{x,y\}}, H_{++}^{\{y,z\}}, H_{++}^{\{x,z\}}\}$ consists of three pairwise disjoint lines that partition $\mathbb{R}_{++}^{\mathbb{T}}$ and \mathcal{G}_{++} has cardinality 4. We now confirm this using Zaslavski's theorem.

In the present setting $A_{++}^{\emptyset} = \mathbb{R}_{++}^{\mathbb{T}}$ and, via eq. (14), $\mu(A_{++}^{\emptyset}) = 1$. Then, since $\mathbb{R}_{++}^{\mathbb{T}}$ is the unique element in \mathcal{L}_{++} that (strictly) contains each hyperplane in \mathcal{H}_{++} , eq. (14) yields $\mu(A) = -\mu(A_{++}^{\emptyset})$ for each $A \in \mathcal{H}_{++}$. Now since the hyperplanes in \mathcal{H}_{++} are disjoint, there are no further elements in \mathcal{L}_{++} and we observe that

$$|\mathcal{G}_{++}| = \sum_{A \in \mathcal{L}_{++}} |\mu(A)| = 4.$$

In contrast, although the structure of \mathcal{L} is otherwise isomorphic to \mathcal{L}_{++} , since $\{0\} \subset \mathbb{R}^{\mathbb{T}}$ is a subset of every hyperplane in \mathcal{H} , $\{0\}$ is the center $A^{\{x,y,z\}}$ of \mathcal{H} and the maximal element of \mathcal{L} . Via eq. (14) and the calculations of the previous paragraph, we obtain $\mu(A^{\{x,y,z\}}) = -(\mu(A^{\emptyset}) - 3\mu(A^{\emptyset})) = 2$. Thus,

$$\#\mathcal{G} = \sum_{A \in \mathcal{L}} |\mu(A)| = 6 = 3!.$$

REMARK 1 (The relationship between \mathcal{L} and \mathcal{L}_{++}). Let $\hat{\mathcal{R}}$ be a Y -generalization with 2-diverse representation $\acute{u}^{(\cdot, \cdot)}$. Since, for every distinct x and y in Y , $\acute{H}^{\{x,y\}}$ contains the origin, $\acute{\mathcal{H}}$ is centered. As we see in example 7, this is not the case for $\acute{\mathcal{H}}_{++}$ where $\acute{\mathcal{H}}_{++}$ is centerless and each of its members is maximal in $\acute{\mathcal{L}}_{++}$.

In GS03, 4-diversity guarantees that, for every $Y \subseteq X$ of cardinality 2, 3 or 4, the improper Y -generalization generates a centered arrangement in $\mathbb{R}_{++}^{\mathbb{T}}$. The fact that $\mathbb{R}_{++}^{\mathbb{T}}$ is open in $\mathbb{R}^{\mathbb{T}}$ ensures that the dimension of any $L \in \acute{\mathcal{L}}$ is equal to its counterpart $L_{++} \in \acute{\mathcal{L}}_{++}$ provided the latter exists. Thus, $\acute{\mathcal{L}}_{++}$ and $\acute{\mathcal{L}}$ are isomorphic if, and only if, $\acute{\mathcal{H}}_{++}$ is centered. For the same reason, $\acute{\mathcal{G}}_{++}$ and $\acute{\mathcal{G}}$ are isomorphic if, and only if, $\acute{\mathcal{H}}_{++}$ is centered.

We now abstract a useful property from example 7.

1015

PROPOSITION 3 (polar opposite rankings). *If $\leq_{\mathbb{J}}$ satisfies $A1^b$ – $A3^b$ and 2-diversity, then, for every $Y \subseteq X$ of cardinality 3 or 4, the improper Y -generalization \mathcal{R} is such that, for some $J, L \in \mathbb{J}$, $\mathcal{R}_J = \mathcal{R}_L^{-1}$ belongs to $\text{total}(\mathcal{R})$.*

See proof on page 60.

EXAMPLE 8 (insufficiency of 2-diversity). *Let $X = [0, 1]^2$ and let \leq^{lex} denote the lexicographic ordering on X . Let $\mathbb{T} = \{s, t\}$, and, for each $J \in \mathbb{J}$, let*

$$\leq_J = \begin{cases} X^2 & \text{if } J(s) = J(t); \\ \leq^{\text{lex}} & \text{if } J(s) < J(t); \\ (\leq^{\text{lex}})^{-1} & \text{otherwise.} \end{cases}$$

Recall that if $\leq_J = X^2$, then \leq_J is symmetric and hence equal to \simeq_J . Thus, for every distinct $x, y \in X$, $H_{++}^{\{x, y\}} = \{J \in \mathbb{R}_{++}^{\mathbb{T}} : J(s) = J(t)\}$. Via lemma 2.1, $\leq_{\mathbb{J}}$ has a two-diverse matrix representation $v^{(\cdot, \cdot)}$. But via lemma 2.2, below, conditional-2-diversity fails to hold. The fact that $\leq_{\mathbb{J}}$ fails to satisfy part (2.a) of theorem 2 follows from the fact that \leq_J is lexicographic for every J outside H .

1025

STEP B.2 (for $\mathbb{T} < \infty$, characterisations of $A4^b$). A matrix $v^{(\cdot, \cdot)}$ that satisfies the conditions of the next lemma is a *conditionally-2-diverse (pairwise) representation*.

LEMMA 2.2 (conditionally-2-diverse representation). *Let \mathcal{R} be a Y -generalization of $\leq_{\mathbb{J}}$ with 2-diverse matrix representation $v^{(\cdot, \cdot)}$. Then $A4^b$ holds on Y if, and only if, for every three distinct elements $x, y, z \in Y$, $v^{(x, z)}$ and $v^{(y, z)}$ are noncollinear.*

1030

See proof on page 61. The following is a translation of observation 2.

PROPOSITION 4 (on $A4^b$ and $A4'^b$). *For $\leq_{\mathbb{J}}$ satisfying $A0^b$ – $A3^b$, conditional-2-diversity and partial-3-diversity are equivalent.*

1035

See proof on page 61.

STEP B.3 (for $\mathbb{T} < \infty$, a characterisation of 4-stability). The following Jacobi identity plays a central role in the proof of GS03.

DEFINITION. *For $Y \in 2^X$, the matrix $v^{(\cdot, \cdot)} : Y^2 \times \mathfrak{T} \rightarrow \mathbb{R}$ satisfies the Jacobi identity whenever, for every $x, y, z \in Y$, $v^{(x, z)} = v^{(x, y)} + v^{(y, z)}$.*

1040

For any given Y -generalization \mathcal{R} , the *Jacobi identity holds for \mathcal{R}* whenever it holds for some pairwise representation $v^{(\cdot, \cdot)}$ of \mathcal{R} . Moreover, in this case, $v^{(\cdot, \cdot)}$ is a *Jacobi representation*. Finally, if the Y -generalization \mathcal{R} is improper and the Jacobi identity holds for \mathcal{R} , we simply say that the *Jacobi identity holds on Y* . Consider

1045

k -JAC. *For every $Y \subseteq X$ with $3 \leq \#Y \leq k$, the Jacobi identity holds on Y .*

We will work with 3-Jac and 4-Jac in particular. The following lemma is the special case of theorem 3 of section C where \mathbb{T} is finite. When \mathbb{T} is finite, for every Y , the set of testworthy Y -generalizations that are novel is nonempty. In this case, 4-stability implies $A0^b$ – $A3^b$ via nonrevision of rankings (part 2 of definition 2).

1050

LEMMA 2.3. *For $\leq_{\mathbb{J}}$ satisfying $A4^b$, 4-stability holds if, and only if, 4-Jac holds.*

STEP B.4 (for $\#\mathbb{T} < \infty$, the concluding arguments in the proof of theorem 2).

In step B.2 we showed that $A1^b$ – $A4^b$ hold if, and only if, $\leq_{\mathbb{J}}$ has a conditionally 2-diverse pairwise representation. In step B.3, we showed that, when $\leq_{\mathbb{J}}$ satisfies $A0^b$ – $A4^b$, a necessary and sufficient condition for 4-Jac is 4- P^b . In step B.5, via (mathematical) induction, we showed that conditionally 2-diverse Jacobi representations of Y -generalizations of $\leq_{\mathbb{J}}$ such that $\#Y = 4$, can be patched together to obtain a conditionally 2-diverse Jacobi representation of $\leq_{\mathbb{J}}$ (on all of X , regardless of cardinality). The fact that $A0^b$ is necessary for a Jacobi representation follows from GS03. As a consequence, $A0^b$ – $A4^b$ and 4- P^b are necessary and sufficient for a conditionally 2-diverse Jacobi representation of $\leq_{\mathbb{J}}$. The following argument then completes the proof of theorem 2.

1055

1060

Let $v^{(\cdot, \cdot)}$ be a (conditionally 2-diverse) Jacobi representation of $\leq_{\mathbb{J}}$ and define $\mathbf{v} : X \times \mathbb{T} \rightarrow \mathbb{R}$ as follows. Fix arbitrary $w \in X$, and let $\mathbf{v}(w, \cdot) = 0$. Then, for every $x \in X$, let $\mathbf{v}(x, \cdot) = v^{(w, x)}$. Recalling that $v^{(w, x)} = -v^{(x, w)}$, note that, since the rows of $v^{(\cdot, \cdot)}$ satisfy the Jacobi identity, for every $x, y \in X$, we have

1065

$$v^{(x, y)} = v^{(x, w)} + v^{(w, y)} = -\mathbf{v}(x, \cdot) + \mathbf{v}(y, \cdot).$$

To see that (2.a) holds note that, for every $J \in \mathbb{J}$, we have $x \leq_J y$, if, and only if, $0 \leq \langle v^{(x, y)}, J \rangle$, if, and only if, $\langle \mathbf{v}(x, \cdot), J \rangle \leq \langle \mathbf{v}(y, \cdot), J \rangle$. For (2.b), note that, since $v^{(\cdot, \cdot)}$ is a conditionally 2-diverse pairwise representation, for every $x, y \in X$,

1070

$$0 \leq v^{(x, y)} = -\mathbf{v}(x, \cdot) + \mathbf{v}(y, \cdot).$$

Finally, for every $z \in X$, we have, for every $\lambda \in \mathbb{R}$, $v^{(z,x)} \neq \lambda v^{(z,y)}$. Equivalently,

$$v(x, \cdot) \neq (1 - \lambda)\mathbf{v}(z, \cdot) + \lambda\mathbf{v}(y, \cdot).$$

Theorem 1 part II, on uniqueness, follows from lemma 2.4 and, without modification, part 3 of the proof of theorem 2 of GS03 (see page 23). 1075

STEP B.5 (for arbitrary X and $\mathbb{T} < \infty$, the induction argument). The present step is closely related to lemma 3 and claim 9 of GS03. There the authors establish that, when 4-diversity holds, 3-Jac is a necessary and sufficient condition for the (global) Jacobi identity to hold on X . GS03 relies on the fact that 4-diversity implies linear independence of $\{v^{(x,y)}, v^{(y,z)}, v^{(z,w)}\}$ for every four distinct elements $x, y, z, w \in X$. In the present setting, where conditional-2-diversity only implies linear independence of pairs $\{v^{(x,y)}, v^{(y,z)}\}$, the Jacobi identity requires 4-Jac. 1080

LEMMA 2.4 (Jacobi representation). *Let $\leq_{\mathbb{J}}$ have a conditionally-2-diverse representation $u^{(\cdot, \cdot)}$. Then 4-Jac holds if, and only if, $\leq_{\mathbb{J}}$ has a Jacobi representation $v^{(\cdot, \cdot)}$. Moreover, for every Jacobi representation $\mathbf{v}^{(\cdot, \cdot)}$ of $\leq_{\mathbb{J}}$ there exists $\lambda > 0$ satisfying $\mathbf{v}^{(\cdot, \cdot)} = \lambda v^{(\cdot, \cdot)}$.* 1085

See proof on page 62.

STEP B.6 (the case of arbitrary X and \mathbb{T}). Note that the inner product we have been working with throughout is well-defined on the infinite-dimensional, linear subspace $\mathbb{R}^{\oplus \mathfrak{I}} \subset \mathbb{R}^{\mathfrak{I}}$ of vectors with finite support. Indeed, for every $v \in \mathbb{R}^{\mathfrak{I}}$ and $J \in \mathbb{R}_+^{\oplus \mathfrak{I}}$, the inner product is a finite sum 1090

$$\langle v, J \rangle = \sum_{\{t: J(t) > 0\}} v(t)J(t).$$

Next note that \mathfrak{I} is just the set of rational-valued vectors in $\mathbb{R}^{\oplus \mathfrak{I}}$. As such, the notions of orthogonality and hyperplanes carry over to the present, infinite-dimensional, setting. W.l.o.g., let $\acute{\mathcal{R}}$ be an improper Y -generalization of $\leq_{\mathbb{J}}$ that satisfies A1^b-A4^b. When $\#\mathbb{T} = \infty$, lemma 2.1, does not hold (since the interior of $\mathbb{R}^{\oplus \mathfrak{I}}$ is empty). Nevertheless, the preceding results extend to the present setting via the notion of an *essentialization*. First note that, via theorem 2 of GS03, there exists a representation $\acute{u}^{(\cdot, \cdot)}$ of $\acute{\mathcal{R}}$. The construction is the same as the one we 1095

present in proof of lemma 2.1. For every $Y \subseteq X$ of cardinality 2, 3 or 4, the representation $\acute{u}^{(\cdot, \cdot)}$ of $\acute{\mathcal{R}}$ has finite rank \acute{r} . The *essentialization* $\text{ess}(\acute{\mathcal{H}})$ of the associated arrangement $\acute{\mathcal{H}}$ in $\mathbb{R}^{\oplus \mathbb{T}}$ is the arrangement we obtain via the by orthogonally projecting $\acute{\mathcal{H}}$ onto the (\acute{r} -dimensional) span of $\acute{u}^{(\cdot, \cdot)}$. Let $p : \mathbb{R}^{\oplus \mathbb{T}} \rightarrow S = \text{span}\{\acute{u}^{(\cdot, \cdot)}\}$ denote this projection. Then, for every $x, y \in Y$, any $J \in \mathfrak{J}$, J is the sum of $p(J)$ and a term that lies in the null space of $\langle \acute{u}^{(x, y)}, \cdot \rangle$. Thus, for every $J \in \mathbb{R}^{\oplus \mathbb{T}}$, $\langle \acute{u}^{(x, y)}, J \rangle = \langle \acute{u}^{(x, y)}, p(J) \rangle$. As such, all the structure of $\acute{\mathcal{H}}$ in $\mathbb{R}^{\oplus \mathbb{T}}$ is preserved by $\text{ess}(\acute{\mathcal{H}})$ in the finite-dimensional subspace S . On this space, define an inner product $\langle \cdot, \cdot \rangle_S$. 1100

For $n = \acute{r}$, take $\mathcal{T} = \{T_i : i = 1, \dots, n\}$ to be an orthonormal basis for S . Then let $T : \mathbb{R}^n \rightarrow S$ to be the matrix with columns \mathcal{T} . Then, for every $J \in \mathbb{R}^n$, $T(J) = J_1 T_1 + \dots + J_n T_n$, where $J_i \in \mathbb{R}$ for each i . Via orthonormality of \mathcal{T} , it then follows that, for every pair of vectors $v, J \in \mathbb{R}^n$, we have 1105

$$\langle T(v), T(J) \rangle_S = \sum_{i,j=1}^n v_i J_j \langle T_i, T_j \rangle_S = \langle v, J \rangle_{\mathbb{R}^n}.$$

Let $\acute{\mathbf{u}}^{(\cdot, \cdot)}$ denote the matrix with rows $\left\{ \acute{\mathbf{u}}^{(x, y)} \stackrel{\text{def}}{=} (T^{-1} \circ P)(\acute{u}^{(x, y)}) : x, y \in Y \right\}$. Then let $\acute{\mathcal{R}} = \langle \acute{\mathcal{R}}_J : J \in \mathbb{Q}_+^n \rangle$ denote the Y -generalization generated by $\acute{\mathbf{u}}^{(\cdot, \cdot)}$ in \mathbb{Q}_+^n , for every $x, y \in Y$ and every $J \in \mathbb{Q}_+^n$, satisfy $x \acute{\mathcal{R}}_J y$ if, and only if, $\langle \acute{\mathbf{u}}^{(x, y)}, J \rangle_{\mathbb{R}^n} \geq 0$. Then, by construction, $\langle \acute{\mathbf{u}}^{(x, y)}, J \rangle_{\mathbb{R}^n} \geq 0$ if, and only if $\langle \acute{u}^{(x, y)}, T(J) \rangle_S \geq 0$. Thus, $x \acute{\mathcal{R}}_J y$ if, and only if, $x \acute{\mathcal{R}}_{T(J)} y$. 1110

Next note that, with the exception of lemma 2.4 of step B.5, all results in the proof of theorem 2 involve Y -generalizations \mathcal{R} of $\leq_{\mathbb{J}}$ for $\#Y = 2, 3$ or 4 . Thus, via the above equivalence, all these results extend to arbitrary \mathbb{T} . And, since step B.5 holds for arbitrary \mathbb{T} and X , the proof of theorem 2 is indeed complete. 1115

APPENDIX C. STATEMENT AND PROOF OF THEOREM 3

Theorem 3. *For $\leq_{\mathbb{J}}$ satisfying $A0^b$ – $A4^b$, 4-stability holds if, and only if, the Jacobi identity holds (for some pairwise representation of $\leq_{\mathbb{J}}$).* 1120

Via lemma 2.4, it suffices to show that 4-stability is equivalent to 4-Jac. Then by construction, 4-stability and 4-Jac are conditions that apply independently to each subset Y of X that has cardinality 3 or 4. Throughout the present section,

$Y \subseteq X$ has cardinality 3 or 4 and \mathcal{R} denotes the improper Y -generalization of $\leq_{\mathbb{J}}$. This will allow us to apply the translation of step B.6 and work in a finite dimensional space. But first we need to rule out the following case. 1130

SUPPOSE THERE IS NO TESTWORTHY Y -GENERALIZATION THAT IS NOVEL. In this case, 4-stability holds vacuously on Y . The following lemma then confirms that 4-diversity holds and theorem 2 of GS03 applies, so that 4-Jac holds on Y .

Lemma 3.1. *If every testworthy Y -generalization of $\leq_{\mathbb{J}}$ is regular, then $|\mathbb{T}| = \infty$ and 4-diversity holds on Y .* 1135

Proof of lemma 3.1. Let $|Y| = 4$. (The proof for case where $|Y| = 3$ is similar and omitted.) Via lemma 3.2, the set of testworthy Y -generalizations is nonempty. Let $\hat{\mathcal{R}}$ be a testworthy Y -generalization, so that there exists J in \mathbb{J} such that $\hat{\mathcal{R}}_{J \times 0}$ is total and equal to the inverse of $\hat{\mathcal{R}}_{\mathfrak{f}}$. Let $P \stackrel{\text{def}}{=} \hat{\mathcal{P}}_{\mathfrak{f}}$. Thus P is the asymmetric part of a total ordering of Y . Via lemma 2.1, let $\hat{v}^{(\cdot, \cdot)}$ be the matrix representation of $\hat{\mathcal{R}}$ and let \hat{v}^P denote the restriction of $\hat{v}^{(\cdot, \cdot)}$ to $P \times \mathbb{T}^{\mathfrak{f}}$. 1140

Claim 3.1.1. *For every vector $\eta^P = \langle \eta^{(x,y)} \in \mathbb{R}_{++} : (x,y) \in P \rangle$, there exist $s, t \in \mathbb{T}$ such that $\hat{v}^P(s) = \eta^P$ and $\hat{v}^P(t) = -\eta^P$.*

Proof of claim 3.1.1. By way of contradiction, suppose there exists $\eta^P \in \mathbb{R}_{++}^P$ such that, for every s in \mathbb{T} , $\hat{v}^P(s) \neq \eta^P$. That is η^P such that, for every s in \mathbb{T} , there exists (x,y) in P such that $\hat{v}^{(x,y)}(s) \neq \eta^{(x,y)}$. This property will suffice for the existence of a testworthy Y -generalization $\acute{\mathcal{R}}$ that is novel. Define $\acute{v}^{(\cdot, \cdot)} : X^2 \times \mathbb{T}^{\mathfrak{f}} \rightarrow \mathbb{R}$ as follows. For each (x,y) in P , let 1145

$$\acute{v}^{(x,y)}(s) \stackrel{\text{def}}{=} \begin{cases} \eta^{(x,y)} & \text{if } s = \mathfrak{f}, \\ \hat{v}^{(x,y)}(s) & \text{otherwise.} \end{cases} \quad \text{1150}$$

For every (x,y) in P^{-1} , since (y,x) in P , take $\acute{v}^{(x,y)} = -\acute{v}^{(y,x)}$. Finally, since P is the asymmetric part of a total ordering, for every remaining (x,y) in Y^2 , $x = y$, so let $\acute{v}^{(x,y)} = 0$. Observe that, by construction, for every s in \mathbb{T} , $\acute{v}^{(\cdot, \cdot)}(s) \neq \acute{v}^{(\cdot, \cdot)}(\mathfrak{f})$. This allows us to appeal to lemma 2.1 and take $\acute{\mathcal{R}}$ to be the novel generalization that $\acute{v}^{(\cdot, \cdot)}$ generates. Moreover, $\acute{\mathcal{R}}$ is also testworthy. For together $P = \hat{\mathcal{P}}_{\mathfrak{f}}$ and $\hat{\mathcal{P}}_{\mathfrak{f}} = \hat{\mathcal{P}}_{J \times 0}^{-1}$ imply that $\acute{\mathcal{R}}_{\mathfrak{f}} = \acute{\mathcal{R}}_{J \times 0}^{-1}$ since, for every (x,y) in P , we have 1155

$$\langle \acute{v}^{(x,y)}, J \times 0 \rangle = \langle \hat{v}^{(x,y)}, J \times 0 \rangle < 0 < \eta^{(x,y)} = \acute{v}^{(x,y)}(\mathfrak{f}).$$

This contradiction implies that, for every $\eta^P \in \mathbb{R}_{++}^P$, there exists s in \mathbb{T} such that $\hat{v}^P(s) = \eta^P$. *Mutatis mutandis*, a repetition of the preceding argument by contradiction confirms that there exists t in \mathbb{T} such that $\hat{v}^P(t) = -\eta^P$. \square 1160

Claim 3.1.1 implies that, when every testworthy Y -generalization is regular, the cardinality of \mathbb{T} is equal to the cardinality of \mathbb{R}^P . We now show that 4-diversity holds on Y . Let R denote an arbitrary total ordering of Y . We show that, for some K in \mathbb{J} , $\langle \hat{v}^{(x,y)}, K \rangle \geq 0$ if, and only if, (x, y) in R . Claim 3.1.1 ensures that we can choose s in \mathbb{T} such that, for some $0 < \epsilon < 1$ 1165

$$\hat{v}^{(x,y)}(s) = \begin{cases} 1 + \epsilon & \text{if } (x, y) \text{ in } R \cap P, \\ 1 - \epsilon & \text{if } (x, y) \text{ in } R^{-1} \cap P. \end{cases}$$

Via claim 3.1.1, take t in \mathbb{T} such that, for every (x, y) in P , $\hat{v}^{(x,y)}(t) = -1$. Let $K := \delta_s + \delta_t$ in \mathbb{J}^f , so that $\langle \hat{v}^{(x,y)}, K \rangle = \hat{v}^{(x,y)}(s) + \hat{v}^{(x,y)}(t)$. By evaluating terms and observing that $\epsilon > 0$ we obtain

$$\langle \hat{v}^{(x,y)}, K \rangle = \begin{cases} (1 + \epsilon) - 1 > 0 & \text{if } (x, y) \text{ in } R \cap P, \\ (1 - \epsilon) - 1 < 0 & \text{if } (x, y) \text{ in } R^{-1} \cap P. \end{cases} \quad 1170$$

Since (x, y) in $R^{-1} \cap P^{-1}$ if, and only if, (y, x) in $R \cap P$ (and, similarly, (x, y) in $R \cap P^{-1}$ if, and only if (y, x) in $R^{-1} \cap P$), we appeal to $\hat{v}^{(x,y)} = -\hat{v}^{(y,x)}$ and obtain

$$\langle \hat{v}^{(x,y)}, K \rangle = \begin{cases} -(1 + \epsilon) + 1 < 0 & \text{if } (x, y) \text{ in } R^{-1} \cap P^{-1}, \\ -(1 - \epsilon) + 1 > 0 & \text{if } (x, y) \text{ in } R \cap P^{-1}. \end{cases}$$

Since P is the asymmetric part of a total ordering we conclude that, for every $x \neq y$, $\langle \hat{v}^{(x,y)}, K \rangle$ has the right sign. Finally, for $x = y$, $\langle \hat{v}^{(x,y)}, K \rangle = 0$. \square 1175

As a consequence of lemma 3.1, for the remainder of the proof of theorem 3 we work under the assumption that the set of testworthy Y -generalizations that are novel is nonempty.

FOR THE REMAINDER OF THIS SECTION, REDEFINE $\mathbb{T} := \mathbb{T}_Y$, where recall \mathbb{T}_Y is defined in step B.6. Thus, in the present section, \mathbb{T} is the finite set that indexes the translation to \mathbb{R}^n of step B.6. Then $\mathbb{T}^f = \mathbb{T} \cup [\mathbf{f}]$ as before and indeed all other notation remains the same. Whereas $\mathbb{T} = \mathbb{T}_Y$ is now finite, the set $\mathbb{C}_{/\sim \kappa}$ of equivalence classes of $\sim^{\mathcal{R}}$ in \mathbb{C} may be infinite as we have seen in lemma 3.1. 1180

Since A1^b–A4^b hold, the pairwise representation $u^{(\cdot,\cdot)} : Y^2 \times \mathbb{T} \rightarrow \mathbb{R}$ of \mathcal{R} is conditionally 2-diverse. Lemma 2.2 implies that $u^{(\cdot,\cdot)}$ has row rank $\mathbf{r} \geq 2$. 1185

Recall the definition of central arrangements. We now show that the set of testworthy generalizations with a central arrangement is always nonempty. (The

Lemma 3.2. *For every $J \in \mathbb{J}$ such that \mathcal{R}_J is total, there exists a Y -generalization $\hat{\mathcal{R}}$ with $\hat{\mathcal{R}}_{\mathfrak{f}} = \mathcal{R}_J^{-1}$. Moreover, its arrangement $\hat{\mathcal{H}}_{++}$ is central; its representation $\hat{u}^{(\cdot,\cdot)}$ has rank $\hat{\mathbf{r}} = \mathbf{r}$; and 4-Jac holds for $\hat{\mathcal{R}}$ if, and only if, it holds for \mathcal{R} .* 1190

Proof. Let $J \in \mathbb{J}$ such that \mathcal{R}_J is total. If, for some $t \in \mathbb{T}$, $J(t) = 0$, then, via $\binom{\#Y}{2}$ applications of A3^b (or the continuity properties of the inner product), there exists there $L \in \mathbb{R}_{++}^{\mathbb{T}}$ such that $\mathcal{R}_L = \mathcal{R}_J$. Thus, w.l.o.g., take $J(t) > 0$ for every $t \in \mathbb{T}$. For some rational $0 < \iota < 1$, let $\hat{J} = (1 - \iota)J \times \iota$, so that \hat{J} lies on the (relative) interior of $\mathbb{J}^{\mathfrak{f}}$. Now, for every x, y in Y , let 1195

$$\hat{\eta}^{(x,y)} := -\frac{1 - \iota}{\iota} \langle u^{(x,y)}, J \rangle,$$

and let $\hat{u}^{(x,y)} \stackrel{\text{def}}{=} u^{(x,y)} \times \hat{\eta}^{(x,y)}$, so that $\langle \hat{u}^{(x,y)}, \hat{J} \rangle = 0$. Let $\hat{\mathcal{R}}$ be the associated Y -generalization, so that by construction $\hat{\mathcal{R}}_{\mathfrak{f}} = \hat{\mathcal{R}}_J^{-1}$. Since $\hat{J} \in \hat{H}^{\{x,y\}}$ for every distinct $x, y \in Y$, $\hat{\mathcal{H}}$ is central. By construction, note that $\mathbf{r} \leq \hat{\mathbf{r}} \leq \mathbf{r} + 1$. Moreover, recalling the rank–nullity theorem, we have $\hat{\mathbf{r}} = \#\mathbb{T}^{\mathfrak{f}} - \dim(\ker(\hat{u}^{(\cdot,\cdot)}))$. Observe that \hat{J} belongs to $\ker(\hat{u}^{(\cdot,\cdot)}) - \ker(u^{(\cdot,\cdot)}) \times 0$. Thus, $\ker(\hat{u}^{(\cdot,\cdot)})$ is of dimension one more than $\ker(u^{(\cdot,\cdot)}) \subset \mathbb{R}^{\mathbb{T}}$. Then since $\#\mathbb{T}^{\mathfrak{f}} = \mathbb{T} + 1$, the rank–nullity theorem yields $\hat{\mathbf{r}} = \mathbf{r}$. Finally, via linearity of the inner product, the Jacobi equations (29)–(31) hold for $u^{(\cdot,\cdot)}$ if, and only if, they hold for $\hat{\eta}^{(\cdot,\cdot)}$ and hence $\hat{u}^{(\cdot,\cdot)}$. 1200 \square

THE CASE WHERE Y HAS CARDINALITY 3, follows from the next two lemmas. 1205

Lemma 3.3. *If $Y = \{x, y, z\}$ and $\mathbf{r} = 2$, then 4-Jac and 4-stability hold on Y .*

Proof of lemma 3.3. Fix arbitrary $J \in \mathbb{J}$ such that \mathcal{R}_J is total. We first apply Zaslavski’s theorem to prove that the Y -generalization $\hat{\mathcal{R}}$ of lemma 3.2 (which is testworthy relative to J) satisfies $|\hat{\mathcal{G}}_{++}| = 6$. Via lemma 2.2, $|\hat{\mathcal{H}}_{++}| = 3$. And since every subarrangement of $\hat{\mathcal{H}}_{++}$ is central, for every $k = 0, 1, 2, 3$ there are $\binom{3}{k}$ ways to choose $|\mathcal{A}| = k$ hyperplanes from $\hat{\mathcal{H}}_{++}$. For $k < 3$, the rank of every 1210

subarrangement is k . For $k = 3$, the rank of the arrangement is \mathfrak{r} . So that, via lemma 3.2, since $\mathfrak{r} = \mathbf{r} = 2$,

$$|\mathcal{G}_{++}| = \binom{3}{3}(-1)^{3-\mathfrak{r}} + \binom{3}{2}(-1)^{2-2} + \binom{3}{1}(-1)^{1-1} + \binom{3}{0}(-1)^{0-0} = 6. \quad (15)$$

For both 4-Jac and 4-stability, we require that every member of \mathcal{G}_{++} is associated with a total ordering. This ensures that 3-diversity holds for $\hat{\mathcal{R}}$, so that, via lemma 2 of GS03, 3-Jac holds for $\hat{\mathcal{R}}$ and lemma 3.2 yields 3-Jac for \mathcal{R} . (When $|Y| = 3$, 3-Jac and 4-Jac coincide.) Also, for 4-stability, if $\hat{\mathcal{R}}$ is any novel, testable generalization that satisfies $\hat{\mathcal{R}}_{\mathfrak{f}} = \mathcal{R}_J^{-1}$, then $\hat{\mathcal{R}}$ is a diverse perturbation of $\hat{\mathcal{R}}$ that satisfies $\text{A0}^b\text{-A3}^b$. 1215

It therefore remains for us to show that every member of \mathcal{G}_{++} is associated with a total ordering of Y . Since \mathcal{H}_{++} is central, $\mathcal{G}_{++} \equiv \mathcal{G}$, we suppress reference to $++$. Let $\{\dot{G}_r : r = 1, \dots, 6\}$ be an enumeration of \mathcal{G} such that \dot{G}_r and $\dot{G}_{r+1} \pmod{6}$ are separated by a single hyperplane in \mathcal{H} . Moreover, let the first four members of this enumeration intersect $\mathbb{J} \times \{0\}$. Then, since $\mathcal{R} = \langle \mathcal{R}_I : I \in \mathbb{J} \rangle$ satisfies A0^b , 1220
 $\dot{G}_1, \dots, \dot{G}_4$ are associated with total orders. Take $L \in \dot{G}_5 \cap \mathbb{J}^{\mathfrak{f}}$ and consider the affine path $\lambda \mapsto \phi(\lambda) = (1 - \lambda)\dot{L} + \lambda\dot{J}$, where \dot{J} is defined in lemma 3.2. For some rational λ^* sufficiently close to, but greater than, 1, $L^* = \phi(\lambda^*)$ belongs to $\dot{G}_* \cap \mathbb{J}^{\mathfrak{f}}$ for some $\dot{G}_* \in \mathcal{G}$. Since $\phi(1) = \dot{J}$ belongs to the center of \mathcal{H} , L and L^* are separated by all three members of \mathcal{H} , so that $\hat{\mathcal{R}}_{L^*} = \hat{\mathcal{R}}_L^{-1}$. Thus, $\hat{\mathcal{R}}_L$ is transitive 1230
if, and only if, $\hat{\mathcal{R}}_{L^*}$ is. Since \dot{G}_5 is separated from \dot{G}_* by 3 hyperplanes, $\dot{G}_* = \dot{G}_2$. Thus $\hat{\mathcal{R}}_{L^*}$ is transitive. *Mutatis mutandis*, the same argument shows that \dot{G}_6 is also associated with a total ordering of Y . □

Lemma 3.4. *If $Y = \{x, y, z\}$ and $\mathbf{r} = 3$, then neither 4-stability nor 4-Jac hold.*

Proof of lemma 3.4. When $\mathbf{r} = 3$, $u^{(x,y)}, u^{(y,z)}$ and $u^{(x,z)}$ are linearly independent, 1235
so that 3-Jac fails to hold.

We now confirm that 3-stability also fails to hold. Via lemma 3.2, $\mathfrak{r} = \mathbf{r}$. The only term in eq. (15) that we adjust for the present lemma is $\mathbf{r} = 3$. Thus,

$$|\mathcal{G}| = (-1)^0 + 3(-1)^0 + 3(-1)^0 + (-1)^0 = 8. \quad \text{1240}$$

Then there are $3! = 6$ members of $\text{total}(\hat{\mathcal{R}})$, and the two additional regions of \mathcal{G} are associated with intransitive CAR rankings. It remains for us to show that every Y -generalization $\hat{\mathcal{R}}$ with $\#\text{total}(\hat{\mathcal{R}}) = 6$ is of this form. 1240




Figure 4: The intersection lattice of a central arrangement for $\#Y = 3$ and $\mathbf{r} = 3$.

In the Hasse diagram of fig. 4, an increase in level corresponds to a decrease in dimension: since $\hat{A}^{\{x,y,z\}}$ is nonempty, it is of dimension at least zero. Since \hat{J} belongs to the interior of $\mathbb{R}_{++}^{\mathbf{T}^f}$ and $\hat{A}^{\{x,y\}\{y,z\}}$ is at least one-dimensional, $\hat{A}^{\{x,y\}\{y,z\}}$ is one-dimensional. Since $\hat{A}^{\{x,y,z\}} \subset \hat{A}^{\{x,y\}\{y,z\}}$, the latter set is nonempty whenever $\hat{\mathcal{H}}_{++}$ is central. The same, of course, applies to other members at the same level. Conversely, if $\hat{A}^{\{x,y\}\{y,z\}}$ is empty, then so is $\hat{A}^{\{x,y,z\}}$. We now use this to show that there is a unique form of Y -generalization $\hat{\mathcal{R}}$ such that $\#\hat{\mathcal{G}}_{++} = 6$ and, moreover, that any such $\hat{\mathcal{R}}$ fails to satisfy $\mathbf{A0}^b$.

Recall from example 7 that $\mu(A^\emptyset) = 1$ and $\mu(A^{\{i,j\}}) = -\mu(A^\emptyset)$. Then, since $\hat{A}^{\{x,y\}\{y,z\}} \subset \hat{A}^{\{x,y\}}, \hat{A}^{\{y,z\}}$, we have $\mu(A^{\{x,y\}\{y,z\}}) = -\mu(A^\emptyset)(1 - 2) = 1$. Finally, by the same argument, since $\hat{A}^{\{x,y,z\}}$ belongs to every member of $\hat{\mathcal{L}}$, we have $\mu(A^{\{x,y,z\}}) = -\mu(A^\emptyset)(1 - 3 + 3) = -1$. Thus, when $\mathbf{r} = 3$, if $\#\hat{\mathcal{G}}_{++} = 6$, then, in addition to $\hat{A}_{++}^{\{x,y,z\}}$, at most one of the intersections $\hat{A}_{++}^{\{i,j\}\{k,l\}}$ is empty. Then, w.l.o.g., suppose $\hat{A}_{++}^{\{x,y\}\{y,z\}}$ is nonempty. We will show that $\mathbf{A0}^b$ fails to hold. Take $L \in \hat{A}_{++}^{\{x,y\}\{y,z\}}$. If $L \in \mathbb{J}^f$, then since $x \hat{\mathcal{I}}_L y \hat{\mathcal{I}}_L z$ and $\neg(x \hat{\mathcal{I}}_L z)$, the proof is complete. In any case, L belongs to one $\hat{G}_{++}^{(x,z)}$ or $\hat{G}_{++}^{(z,x)}$. W.l.o.g., suppose $L \in \hat{G}_{++}^{(x,z)}$. Then $\hat{H}^{\{x,y\}}$ and $\hat{H}^{\{y,z\}}$ split $\hat{G}_{++}^{(x,z)}$ into the four (nonempty) open regions formed by expanding

$$\hat{G}_{++}^{(x,z)} - \left(\hat{H}^{\{x,y\}} \cap \hat{H}^{\{y,z\}} \right) = \hat{G}_{++}^{(x,z)} \cap \left(\hat{G}_{++}^{(x,y)} \cup \hat{G}_{++}^{(y,x)} \right) \cap \left(\hat{G}_{++}^{(y,z)} \cup \hat{G}_{++}^{(z,y)} \right).$$

Take any K in the final member $\hat{G}_{++}^{(x,z)} \cap \hat{G}_{++}^{(y,x)} \cap \hat{G}_{++}^{(z,y)}$ of this expansion. Then, since $x \hat{\mathcal{P}}_K z \hat{\mathcal{P}}_K y \hat{\mathcal{P}}_K x$, we observe that $\hat{\mathcal{R}}$ fails to satisfy $\mathbf{A0}^b$. We conclude that only dogmatic perturbations of $\hat{\mathcal{R}}$ satisfy $\mathbf{A0}^b$. \square

THE CASE WHERE Y HAS CARDINALITY 4. Note that a failure of \mathcal{B} -Jac on $Z \subset Y$ such that $|Z| = 3$ implies a failure of \mathcal{A} -Jac on Y . And since the arguments for the case where $|Y| = 3$ account for the case where \mathcal{B} -Jac fails, we henceforth assume that \mathcal{B} -Jac holds on Y . That is, our conditionally \mathcal{B} -diverse representation $u^{(\cdot, \cdot)}$ will now satisfy equations (29)–(31) with $\hat{\beta} = \beta$ if, and only if, \mathcal{A} -Jac holds on Y . 1265

First some some useful results that exploit \mathcal{B} -Jac. 1270

Proposition 5. *If $Y = \{x, y, z, w\}$ and \mathcal{B} -Jac holds for $u^{(\cdot, \cdot)}$, then, for every $\hat{v}^{(\cdot, \cdot)} = u^{(\cdot, \cdot)} \times \hat{\eta}^{(\cdot, \cdot)}$ with rank $\hat{\mathbf{r}}$, that satisfies \mathcal{B} -Jac, $2 \leq \hat{\mathbf{r}} \leq 3$.*

Proof of proposition 5. Via proposition 4 and lemma 3.3, $\mathbf{r} \geq 2$. Indeed the span of $\{u^{(x, w)}, u^{(y, w)}\}$ is two. Let S denote the span of $\{u^{(x, w)}, u^{(y, w)}, u^{(z, w)}\}$. Since $u^{(y, w)} = -u^{(w, y)}$ and $u^{(\cdot, \cdot)}$ satisfies \mathcal{B} -Jac, equations (29)–(31) hold for $u^{(\cdot, \cdot)}$. (If \mathcal{A} -Jac fails to hold, then $\beta \neq \hat{\beta}$, but the equations still hold.) Thus $u^{(x, y)}$, $u^{(y, z)}$ and $u^{(x, z)}$ all belong to S and $\mathbf{r} \leq 3$. Now note that above argument does not depend on the cardinality of \mathbb{T} , thus take $\hat{\eta}^{(\cdot, \cdot)}$ to satisfy \mathcal{B} -Jac: indeed with the same parameters that feature in equations (29)–(31) for $u^{(\cdot, \cdot)}$. The preceding argument then generalizes *mutatis mutandis* to $\hat{v}^{(\cdot, \cdot)}$ and $2 \leq \hat{\mathbf{r}} \leq 3$. □ 1275

Proposition 6. *If $X = \{x, y, z, w\}$, then $4 \leq |\mathcal{H}| \leq 6$ and these bounds are tight. If, moreover, \mathcal{B} -Jac holds for $u^{(\cdot, \cdot)}$ and $|\mathcal{H}| < 6$, then $\mathbf{r} = 2$.*

Proof of proposition 6. The upper bound $|\mathcal{H}| \leq 6$ follows from the fact that there are $\binom{4}{2} = 6$ ways to choose distinct pairs of elements from the four-element set X . Via lemma 2.2 only the following equalities are feasible: $H^{\{x, y\}} = H^{\{z, w\}}$, $H^{\{x, z\}} = H^{\{y, w\}}$ and $H^{\{x, w\}} = H^{\{y, z\}}$. Via proposition 3, w.l.o.g., take $G^{(x, y, z, w)}$ and $G^{(w, z, y, x)}$ to belong to \mathcal{G} . We consider a convex path ϕ from (some member of) $G^{(x, y, z, w)}$ to $G^{(w, z, y, x)}$ such that $\#\{G \in \mathcal{G} : (\text{image } \phi) \cap G \neq \emptyset\}$ is maximal. Via $\mathbf{A0}^b$, one of $H^{\{x, y\}}$, $H^{\{y, z\}}$ and $H^{\{z, w\}}$ supports $G^{(x, y, z, w)}$. If it is $H^{\{y, z\}}$, then so does $H^{\{x, w\}}$ and, and since x and w lie at opposite ends of the ordering (x, y, z, w) , this would imply that $|\mathcal{H}| = 1$ in violation of lemma 2.2. Since we are seeking a greatest lower bound on $|\mathcal{H}|$, take $H_1 = H^{\{x, y\}} = H^{\{z, w\}}$ to be the first hyperplane to intersect image ϕ and the adjacent chamber is $G^{(y, x, w, z)}$. Via $\mathbf{A0}^b$, the second hyperplane to intersect image ϕ is $H_2 \stackrel{\text{def}}{=} H^{\{x, w\}}$. If $H^{\{x, w\}} = H^{\{y, z\}}$ then since y and z lie at opposite ends of (y, x, w, z) , this would once again lead to a violation 1280

of lemma 2.2. Then $G^{(y,w,x,z)}$ is next chamber to intersect image ϕ . Since we are seeking a greatest lower bound, via $A0^b$, take $H_3 \stackrel{\text{def}}{=} H^{\{y,w\}} = H^{\{x,z\}}$ to be the third hyperplane to intersect image ϕ . Then $G^{(w,y,z,x)}$ is the next chamber to intersect image ϕ . The final hyperplane to intersect image ϕ is $H_4 = H^{\{y,z\}}$ which brings us to the destination chamber $G^{(w,z,y,x)}$. We conclude that at most two pairs of the six hyperplanes coincide, so that $|\mathcal{H}| \geq 4$. 1300

We now prove that 3-Jac and $H^{\{x,y\}} = H^{\{z,w\}}$ together imply $\mathbf{r} = 2$. Consider equations (29)–(31) (so 3-Jac holds, but 4-Jac need not). Via (29), $S = \{u^{(x,w)}, u^{(w,y)}, u^{(x,y)}\}$ is 2-dimensional. Since $u^{(x,y)}$ and $u^{(z,w)}$ are collinear, $u^{(z,w)}$ belongs to S . Finally, equations (30) and (31) yield $u^{yz}, u^{(x,z)} \in S$. □ 1305

Lemma 3.5. *If $Y = \{x, y, z, w\}$, $\mathbf{r} = 3$ and 3-Jac holds on Y , then 4-stability and 4-Jac both hold on Y*

Proof of lemma 3.5. To see that 4-Jac holds, appeal to the proof of lemma 3 of GS03: if 3-Jac holds and 4-Jac does not, then $\{u^{(x,w)}, u^{(y,w)}, u^{(z,w)}\}$ is linearly dependent. This is in contradiction of $\mathbf{r} = 3$. 1310

We now verify that 4-stability also holds. We show that, for every $J \in \mathbb{J}$ such that \mathcal{R}_J is total, the Y -generalization $\hat{\mathcal{R}}$ of lemma 3.2 is maximal in the sense that $\text{total}(\hat{\mathcal{R}}) = |\hat{\mathcal{G}}_{++}| = 4!$. We then confirm that $\hat{\mathcal{R}}$ satisfies $A0^b$. Then, if $\tilde{\mathcal{R}}$ is any novel, testworthy generalization that satisfies $\tilde{\mathcal{R}}_{\mathfrak{f}} = \mathcal{R}_J^{-1}$, then $\tilde{\mathcal{R}}$ is a diverse perturbation of $\hat{\mathcal{R}}$ that satisfies $A0^b$ – $A3^b$. 1315

Since $\mathbf{r} = 3$, the contrapositive of proposition 6 implies that $|\mathcal{H}_{++}| = 6$. The proof that now follows is a simple generalization of lemma 3.3 to allow for the fact that $|Y| = 4$. Fix any $J \in \mathbb{J}$ such that \mathcal{R}_J is total and w.l.o.g. take $J > 0$. Via lemma 3.2, there exists a testworthy Y -generalization $\hat{\mathcal{R}}$ such that $\hat{\mathcal{R}}_{\mathfrak{f}} = \mathcal{R}_J^{-1}$ is total and $L = (1 - \lambda)J \times \lambda$ belongs to both $\mathbb{J}^{\mathfrak{f}}$ and the center of $\hat{\mathcal{H}}_{++}$ and $\hat{\nu}^{(\cdot, \cdot)}$ satisfies 4-Jac. Via remark 1, since $\hat{\mathcal{H}}$ is central, $\hat{\mathcal{G}}_{++}$ and $\hat{\mathcal{G}}$ are isomorphic, so we work with $\hat{\mathcal{G}}$. Since $\mathbf{r} = 3$, proposition 6 implies $|\mathcal{H}| = 6$, so that the same is true of $|\hat{\mathcal{H}}|$. The rank of subarrangements with cardinality 4 or more is \mathbf{r} . Let \mathfrak{r} denote the number of subarrangements \mathcal{A} that have cardinality 3 and rank 2. Each of the other $\binom{6}{3} - \mathfrak{r}$ subarrangements with cardinality 3 have rank \mathbf{r} . All 1320

other subarrangements have rank equal to their cardinality.

$$\begin{aligned}
 |\mathcal{G}| &= \binom{6}{6}(-1)^{6-\mathbf{r}} + \binom{6}{5}(-1)^{5-\mathbf{r}} + \binom{6}{4}(-1)^{4-\mathbf{r}} \\
 &\quad + \binom{6}{3}(-1)^{3-\mathbf{r}} - \mathfrak{r}(-1)^{3-\mathbf{r}} + \mathfrak{r}(-1)^{3-2} \\
 &\quad + \binom{6}{2}(-1)^{2-2} + \binom{6}{1}(-1)^{1-1} + \binom{6}{0}(-1)^{0-0}
 \end{aligned} \tag{16}$$

Via lemma 3.2, $\mathbf{r} = \mathbf{r}$, so that $\mathbf{r} = 3$. It remains for us to calculate the value of \mathfrak{r} . Each of the $\binom{4}{3} = 4$ subsets of Y that have cardinality 3 generates a subarrangement of cardinality $\binom{3}{2} = 3$. (For instance, $\mathcal{A}^{\{x,y,z\}} = \{\dot{H}^{\{x,y\}}, \dot{H}^{\{y,z\}}, \dot{H}^{\{x,z\}}\}$.)

1330

For such subarrangements, 4-Jac implies a rank of 2. Arguments from the final step in the proof of proposition 6 confirm that every other subarrangement with cardinality 3 has rank 3. Substituting into eq. (16), we obtain

$$|\mathcal{G}| = -1 + 6 - 15 + 20 - 4 - 4 + 15 + 6 + 1 = 24 = 4!.$$

The proof that every member of \mathcal{G} is associated with a transitive ordering is as follows. Via lemma 3.2, 4-Jac holds for $\hat{\mathcal{R}}$ if, and only if, it holds for \mathcal{R} . We have seen that 4-Jac holds. Then, via a straightforward application of the Jacobi identity and, for arbitrary $K \in \mathbb{J}^\dagger$, the definition of transitivity of $\hat{\mathcal{R}}_K$ yields $\mathbf{A}0^b$. Thus, every member of $\hat{\mathcal{G}}$ is total. \square

1335

IN THE REMAINING CASE, WHERE $Y = \{x, y, z, w\}$ AND $\mathbf{r} = 2$, the proof is complicated by the fact that maximally-diverse generalizations have centerless arrangements. We begin by choosing $\eta^{(\cdot, \cdot)}$ so as to construct $\acute{u}^{(\cdot, \cdot)} = u^{(\cdot, \cdot)} \times \eta^{(\cdot, \cdot)}$ with $\mathbf{r} = 3$.

1340

Since $\mathbf{r} = 2$, it follows that $u^{(x,z)}, u^{(y,z)}$ and $u^{(w,z)}$ form a linearly dependent set. Thus, for some $\pi, \rho \in \mathbb{R}$,

1345

$$\pi u^{(x,z)} + \rho u^{(y,z)} = u^{(w,z)}. \tag{17}$$

Fix arbitrary $J \in \mathbb{J}$ such that \mathcal{R}_J is total, and, as in lemma 3.2, w.l.o.g., we take $J \in G_{++}^{(x,y,z,w)}$. Then, for $\acute{\iota} = \frac{1}{2}$ and $\acute{J}^{\{x,y,z\}} = (1 - \acute{\iota})J \times \acute{\iota}$, let

$$\acute{\eta}^{(i,j)} = -\frac{1-\acute{\iota}}{\acute{\iota}} \langle u^{(i,j)}, J \rangle = -\langle u^{(i,j)}, J \rangle, \quad \text{for every } i, j \in \{x, y, z\}. \tag{18}$$

Note that $\dot{\eta}^{(y,z)} > 0$ is determined by eq. (18). Since $G_{++}^{(x,y,z,w)}$ is open and the inner product is continuous, there exists a compact neighbourhood $N_J \subset G_{++}^{(x,y,z,w)}$ of J such that, for every $L \in N_J$, $\langle u^{(y,z)}, L \rangle < -\langle u^{(w,z)}, L \rangle$ if, and only if, $\langle u^{(y,z)}, J \rangle < -\langle u^{(w,z)}, J \rangle$. For any such $L \neq J$, take $\dot{\lambda}$ and ϵ to be solutions to

$$\dot{\eta}^{(y,z)} = -\frac{1-\dot{\lambda}}{\dot{\lambda}} \langle u^{(y,z)}, L \rangle \quad \text{and} \quad \epsilon = \langle u^{(w,z)}, \frac{1-\dot{\lambda}}{\dot{\lambda}} L - J \rangle. \quad (19)$$

A rearrangement of the first equality in eq. (19) and, via eq. (18), a substitution for $\dot{\eta}^{(y,z)}$ yields $\dot{\lambda} = \frac{\langle u^{(y,z)}, L \rangle}{\langle u^{(y,z)}, J \rangle + \langle u^{(y,z)}, L \rangle}$ and $\frac{1-\dot{\lambda}}{\dot{\lambda}} = \frac{\langle u^{(y,z)}, J \rangle}{\langle u^{(y,z)}, L \rangle}$. Since $N_J \subset G^{(y,z)}$, $0 < \dot{\lambda} < 1$ and $\dot{L}^{\{y,z,w\}} \stackrel{\text{def}}{=} (1 - \dot{\lambda})L \times \dot{\lambda} \in \mathbb{R}_{++}^{\text{Tf}}$ for every such L . On N_J , the denominator of the map $L \mapsto \dot{L}^{\{y,z,w\}}$ is bounded away from zero and continuous (since it is the quotient of continuous functions). Furthermore, $\lim_{L \rightarrow J} \dot{L}^{\{y,z,w\}} = \dot{J}^{\{y,z,w\}}$. Similarly, on N_J , the fact that $\langle u^{(y,z)}, L \rangle > 0$ ensures that the map $L \mapsto \epsilon$ is well-defined, continuous and $\lim_{L \rightarrow J} \epsilon = 0$.²⁰

For every $i, j \in \{y, z, w\}$, let $\dot{\eta}^{(i,j)} = -\frac{1-\dot{\lambda}}{\dot{\lambda}} \langle u^{(i,j)}, L \rangle$. Now note that, via equations (17)–(19), we obtain

$$\pi \dot{\eta}^{(x,z)} + \rho \dot{\eta}^{(y,z)} = \dot{\eta}^{(w,z)} + \epsilon \neq \dot{\eta}^{(w,z)}, \quad (21)$$

so that, for every $\epsilon \neq 0$, $\{\dot{u}^{(i,j)} = u^{(i,j)} \times \dot{\eta}^{(i,j)} : (i, j) = (x, z), (y, z), (w, z)\}$ forms a linearly independent set. Let $\{\dot{u}^{(i,j)} : i, j \in \{x, y, z\} \cup \{y, z, w\}\}$ be defined similarly. To complete the definition of $\dot{u}^{(\cdot, \cdot)}$, we appeal to the fact that, via β -Jac, $u^{(\cdot, \cdot)}$ satisfies equations (29)–(31). In particular, from these equations, take parameters α , β , and γ and let $\dot{\eta}^{(x,w)}$ be the (unique) solution to the Jacobi identity

$$\alpha \dot{\eta}^{(x,w)} = \gamma \dot{\eta}^{(x,y)} + \beta \dot{\eta}^{(y,w)} = -\langle \gamma u^{(x,y)}, J \rangle - \frac{1-\dot{\lambda}}{\dot{\lambda}} \langle \beta u^{(y,w)}, L \rangle. \quad (22)$$

For the same set of parameter values, $\dot{u}^{(\cdot, \cdot)}$ also satisfies (29)–(31). That is, for $\{x, y, z\}$, via eq. (18) and (31), $\gamma \dot{\eta}^{(x,y)} + \tau \dot{\eta}^{(y,z)} = \phi \dot{\eta}^{(x,z)}$, so that $\dot{u}^{(\cdot, \cdot)}$ satisfies (31).

²⁰Substituting for $\frac{1-\dot{\lambda}}{\dot{\lambda}}$ in the equation for ϵ in (19) and rearranging yields a linear equation

$$k \langle u^{(y,z)}, L \rangle = \langle u^{(w,z)}, L \rangle \quad (20)$$

in L , where $k = \frac{\epsilon + \langle u^{(w,z)}, J \rangle}{\langle u^{(y,z)}, J \rangle}$. Note that, since $L \in G_{++}^{(z,w)} \cap G_{++}^{(y,z)}$, $\langle u^{(w,z)}, L \rangle < 0$ and $\langle u^{(y,z)}, L \rangle > 0$; moreover, the same is true of J . Thus, $k < 0$ or, equivalently, $\epsilon < -\langle u^{(w,z)}, J \rangle$. If $\langle u^{(y,z)}, L \rangle < -\langle u^{(w,z)}, L \rangle$, then, via eq. (20), $k < -1$. Moreover, our choice of N_J is such that $\frac{\langle u^{(w,z)}, J \rangle}{\langle u^{(y,z)}, J \rangle} < -1$. This ensures that, on N_J , ϵ lies in a neighbourhood of zero, as required.

For $\{y, z, w\}$, Via eq. (19) and (30), $\hat{\beta}\dot{\eta}^{(y,w)} + \sigma\dot{\eta}^{(w,z)} = \tau\dot{\eta}^{(y,z)}$, so that $\dot{u}^{(\cdot,\cdot)}$ satisfies (30). For $\{x, y, w\}$, via eq. (22), $\dot{u}^{(\cdot,\cdot)}$ satisfies (29). We now demonstrate that for the final triple $\{x, z, w\}$, the Jacobi identity holds if $\hat{\beta} = \beta$, and $\{\dot{u}^{(x,w)}, \dot{u}^{(w,z)}, \dot{u}^{(x,z)}\}$ has rank 3 otherwise. 1375

First extract the parameters from equations (29)–(31) to obtain the matrix form

$$\begin{array}{ccccc} (x, w) & (y, w) & (x, y) & (w, z) & (y, z) & (x, z) \\ \left[\begin{array}{ccc|ccc} \alpha & -\beta & -\gamma & 0 & 0 & 0 \\ 0 & \hat{\beta} & 0 & \sigma & -\tau & 0 \\ 0 & 0 & \gamma & 0 & \tau & -\phi \end{array} \right] \end{array} \begin{array}{l} (29) \\ (30) \\ (31) \end{array} \quad (23)$$

Since the triple $\{\dot{u}^{(i,z)} : i = x, y, w\}$ provides a basis for $\text{span}(\dot{u}^{(\cdot,\cdot)})$, we will write all vectors in terms of this basis. To this end, we derive the reduced row echelon form of eq. (23). In particular, letting r_i denote the rows of the matrix, we perform the operation $r_1 \mapsto r_1 + \frac{\beta}{\hat{\beta}}r_2 + r_3$ to obtain 1380

$$\begin{array}{ccccc} (x, w) & (y, w) & (x, y) & (w, z) & (y, z) & (x, z) \\ \left[\begin{array}{ccc|ccc} \alpha & 0 & 0 & \frac{\beta}{\hat{\beta}}\sigma & (1 - \frac{\beta}{\hat{\beta}})\tau & -\phi \\ 0 & \hat{\beta} & 0 & \sigma & -\tau & 0 \\ 0 & 0 & \gamma & 0 & \tau & -\phi \end{array} \right] \end{array} \begin{array}{l} (29) \\ (30) \\ (31) \end{array} \quad (24)$$

In eq. (24), the fact that $\hat{\beta}$ (instead of β) that appears as a pivot in column 2 indicates that, in this derivation, we are, w.l.o.g., choosing $\dot{v}^{(y,w)} = \hat{\beta}\dot{u}^{(y,w)}$. *Mutatis mutandis*, the conclusions we will draw are the same if instead we chose to define $\dot{v}^{(y,w)}$ using β . The other (relevant) rows of $\dot{v}^{(\cdot,\cdot)} : Y^2 \times \mathbb{T}^f \rightarrow \mathbb{R}$ are $\dot{v}^{(x,w)} = \alpha\dot{u}^{(x,w)}$, $\dot{v}^{(x,y)} = \gamma\dot{u}^{(x,y)}$, $\dot{v}^{(z,w)} = \sigma\dot{u}^{(z,w)}$, $\dot{v}^{(y,z)} = \tau\dot{u}^{(y,z)}$ and $\dot{v}^{(x,z)} = \phi\dot{u}^{(x,z)}$. The matrix of the equation that now follows, is invertible if, and only if, $(1 - \frac{\beta}{\hat{\beta}}) \neq 0$. 1385 1390

$$\begin{bmatrix} \dot{v}^{(x,w)} \\ \dot{v}^{(x,z)} \\ \dot{v}^{(z,w)} \end{bmatrix} = \begin{bmatrix} -\frac{\beta}{\hat{\beta}} & -(1 - \frac{\beta}{\hat{\beta}}) & 1 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix} \begin{bmatrix} \dot{v}^{(w,z)} \\ \dot{v}^{(y,z)} \\ \dot{v}^{(x,z)} \end{bmatrix} \quad (25)$$

Thus, unless $\hat{\beta} = \beta$, we conclude that $\{\dot{v}^{(x,w)}, \dot{v}^{(x,z)}, \dot{v}^{(z,w)}\}$ has the same rank as $\{\dot{v}^{(w,z)}, \dot{v}^{(y,z)}, \dot{v}^{(x,z)}\}$ which, by construction, has rank 3 for every choice of $\epsilon \neq 0$.




Figure 5: The intersection semilattices $\hat{\mathcal{L}}$ and $\hat{\mathcal{L}}_{++} = \hat{\mathcal{L}} - \{A^Y, \hat{A}^{\{x,z\}\{y,w\}}\}$ when $\#\mathbb{T} = 2$, $\hat{\beta} = \beta$ and ϵ is sufficiently small but distinct from zero.

Since $\hat{\beta} = \beta$ if, and only if, 4-Jac holds for \mathcal{R} , we conclude that 4-Jac holds for \mathcal{R} if, and only if, it holds for $\hat{\mathcal{R}}$. It remains for us to show that, for ϵ sufficiently small $\hat{\mathcal{G}}$ is maximal. For if $\hat{\mathcal{G}}$ is maximal, then via lemma 3.3 and lemma 3.4, A_0^b holds if, and only if, $\text{rank}\{\hat{v}^{(x,w)}, \hat{v}^{(x,z)}, \hat{v}^{(z,w)}\} = 2$. 1395

By Zaslavski's theorem, it suffices to show that every member of the intersection lattice $\hat{\mathcal{L}}$, other than the center $\hat{A}^{\{x,y,z,w\}}$, has nonempty intersection with $\mathbb{R}_{++}^{\mathbb{T}^f}$. We now make explicit the dependence of the generalization $\hat{\mathcal{R}}$ on our choice of $L \in N_J$, though we do so indirectly via ϵ . Let $d^\epsilon = \max\{d(\hat{J}, \hat{A}) : \hat{A} \in \hat{\mathcal{L}}^\epsilon\}$, where $d(\hat{J}, \hat{A})$ is the minimum (Euclidean) distance between \hat{J} and the (closed) linear subspace \hat{A} of $\mathbb{R}^{\mathbb{T}^f}$. Note that for $\epsilon = 0$, we obtain a central arrangement of the form of lemma 3.2 with $d^\epsilon = 0$. Moreover, since the Euclidean metric is continuous in its arguments, and, for every ϵ , $\hat{\mathcal{L}}^\epsilon$ is finite, the map $\epsilon \mapsto d^\epsilon$ is continuous and $\lim_{\epsilon \rightarrow 0} d^\epsilon = 0$. Thus, for sufficiently small $\epsilon \neq 0$, every $\hat{A} \in \hat{\mathcal{L}}^\epsilon$ intersects $\mathbb{R}_{++}^{\mathbb{T}^f}$. 1400
1405

Remark. We note that the above arguments apply without modification to the case where $\#\mathcal{H} = 4, 5$. Consider, for example, the Hasse diagram of fig. 5. That case arises when $u^{(x,z)}$ and $u^{(y,w)}$ are collinear, so that $\hat{A}^{\{x,z\}\{y,w\}}$ is a hyperplane of dimension $\#\mathbb{T} - 1$. Assuming the same construction, with $\epsilon \neq 0$, so that $\hat{u}^{(\cdot, \cdot)}$ has rank 3 and, via proposition 6, $\hat{u}^{(x,z)}$ and $\hat{u}^{(y,w)}$ are linearly independent. Thus $\hat{A}^{\{x,z\}\{y,w\}}$ is of dimension $\#\mathbb{T}^f - 2 = \#\mathbb{T} - 1$. Thus, $\hat{A}^{\{x,z\}\{y,w\}} = A^{\{x,z\}\{y,w\}} \times \{0\}$ which belongs to the boundary of $\mathbb{R}_{++}^{\mathbb{T}^f}$. At $\epsilon = 0$, $\hat{A}^{\{x,z\}\{y,w\}}$ increases by one 1410

dimension and the upper two levels of the Hasse diagram collapse to equal $A^{\{x,y,z,w\}}$.

ONLINE APPENDIX D. PROOFS

1415

Proof of Observation 1. Fix $Y \subseteq X$ nonempty and $\hat{\mathcal{R}}$ regular. W.l.o.g., take $C \in \mathbb{D}^f - \mathbb{D}$, so that C contains at least one copy of \mathfrak{f} . For any $c \in C \cap [\mathfrak{f}]$, the fact that $\hat{\mathcal{R}}$ is regular implies that $c \sim^{\hat{\mathcal{R}}} c_1$ for some $c_1 \in \mathbb{C}$. The richness assumption ensures that we may choose c_1 from the complement of C . Then, since neither c nor c_1 belong to $C_1 \stackrel{\text{def}}{=} C - \{c\}$, $c \sim^{\hat{\mathcal{R}}} c_1$ implies $\hat{\mathcal{R}}_C = \hat{\mathcal{R}}_{C_1 \cup \{c_1\}}$. If c is the unique member of $C \cap [\mathfrak{f}]$, then the proof is complete. Otherwise, using the fact that C is finite, we may proceed by induction until we obtain a set C_n such that $C_n \cap [\mathfrak{f}]$ is empty and $D \stackrel{\text{def}}{=} C_n \cup \{c_1, \dots, c_n\}$ belongs to \mathbb{D} . Part 1 of definition 1 then implies $\hat{\mathcal{R}}_D = \leq_D \cap Y^2$, so that, since \mathcal{R} is improper, $\hat{\mathcal{R}}_D = \mathcal{R}_D$. Finally, since $C \sim^{\hat{\mathcal{R}}} D$, $\hat{\mathcal{R}}_C = \hat{\mathcal{R}}_D$, as required. \square

1420

1425

Proof of Corollary 1. This follows from theorem 3, lemma 2.4 and the fact that, via lemma 3.1, 4-stability implies A0–A3 when the number of case types is finite. \square

Proof of Lemma 1.1. We show that there exists a canonical embedding (a structure preserving injection) of $\text{nov}(Y, \leq_{\mathbb{J}})$ into $\text{nov}(Y, \leq_{\mathbb{D}})$. The fact that this map is also surjective follows from the fact that $\text{nov}(Y, \leq_{\mathbb{D}})$ can be embedded in $\text{nov}(Y, \leq_{\mathbb{J}})$ in precisely the same way. The proof that the two sets of regular generalizations are isomorphic follows via a similar argument plus the observation that every Y -generalization is either regular or novel.

1430

Take $\mathcal{R} \in \text{nov}(Y, \leq_{\mathbb{J}})$ and define $\hat{\mathcal{R}} = \langle \hat{\mathcal{R}}_C : C \in \mathbb{D}^f \rangle$ via the property: for each $C \in \mathbb{D}^f$, $\hat{\mathcal{R}}_C \stackrel{\text{def}}{=} \mathcal{R}_J$ if, and only if, $L_C = L_J$, where, as before, $t \mapsto L_C(t)$ counts the number of cases of type t in C and $L_J = \kappa_J J \in \mathbb{L}^f$ for some minimal $\kappa_J \in \mathbb{Z}_+$. Now, for any $\mathcal{R}' \neq \mathcal{R}$ in $\text{nov}(Y, \leq_{\mathbb{J}})$, there exists $J \in \mathbb{J}^f$ such that $\mathcal{R}'_J \neq \mathcal{R}_J$. If we define $\hat{\mathcal{R}}'$ analogously, so that it is equivalent to \mathcal{R}' , then $\hat{\mathcal{R}}' \neq \hat{\mathcal{R}}$. As a consequence, the canonical mapping $\mathcal{R} \mapsto \hat{\mathcal{R}}$ is injective. If we can show that $\hat{\mathcal{R}}$ does in fact belong to $\text{nov}(Y, \leq_{\mathbb{D}})$, then we have constructed the required embedding. The fact that $\hat{\mathcal{R}}$ satisfies 2 and 1 of definition 1 follows immediately from definition 2. The proof that item 3 of definition 1 holds is as follows. Take any $c, c' \in \mathbb{C}^f$ and $D \in \mathbb{D}^f$ such that $c \sim^{\star} c'$ and $c, c' \notin D$. First, observe that $D \cup \{c\} \sim^{\star} D \cup \{c'\}$, and moreover, for some $t \in \mathbb{T}^f$ we have $c, c' \in t$. Then, for every $t \in \mathbb{T}^f$, $|D \cup \{c\}| = |D \cup \{c'\}| = L$ for

1435

1440

1445

some $L \in \mathbb{L}^f \cap \mathbb{J}^f$. Thus $\hat{\mathcal{R}}_{D \cup \{c\}} = \hat{\mathcal{R}}_{D \cup \{c'\}}$, as required for $\hat{\mathcal{R}}$ to be a generalization of $\leq_{\mathbb{D}}$. Finally, via definition 2, the definition of a novel generalization ensures that the induced equivalence relation $\sim^{\mathcal{R}}$ on \mathbb{C}^f satisfies $c \not\sim^{\mathcal{R}} f$ for every $c \in \mathbb{C}$. Since $\sim^{\hat{\mathcal{R}}}$ inherits this property, $\hat{\mathcal{R}}$ is novel. \square

Proof of Lemma 1.3. Fix $\mathcal{R}_{\mathfrak{J}} \equiv \hat{\mathcal{R}}_{\mathfrak{D}}$ and assume that $\hat{\mathcal{R}}_{\mathfrak{D}}$ satisfies A2. We show that $\mathcal{R}_{\mathfrak{J}}$ satisfies A2^b. Fix $x, y \in Y$ and $J \in \mathfrak{J}$ such that $x \mathcal{R}_J y$ and $x \mathcal{R}_{J'} y$. Fix $\lambda, \mu \in \mathbb{Q}_{++}$ and let κ be the smallest positive integer such that both $L \stackrel{\text{def}}{=} \kappa \lambda J$ and $L' \stackrel{\text{def}}{=} \kappa \mu J'$ belong to \mathfrak{L} . Then, by lemma 1.2, we have both $x \mathcal{R}_L y$ and $x \mathcal{R}_{L'} y$. Moreover, for D, D' such that $L_D = L$ and $L_{D'} = L'$, we have $x \hat{\mathcal{R}}_D y$ and $x \hat{\mathcal{R}}_{D'} y$ and, by A2, $x \hat{\mathcal{R}}_{D \cup D'} y$. Finally, since $L_D + L_{D'} = \kappa(\lambda J + \mu J')$, one further application of lemma 1.2 yields $x \mathcal{P}_{\lambda J + \mu J'} y$, as required for A2^b. 1450

The proof that “A2 implies A2^b” is *mutatis mutandis* a special case of the above argument and omitted. We now assume $\hat{\mathcal{R}}_{\mathfrak{D}}$ satisfies A2 and A3 and prove that $\mathcal{R}_{\mathfrak{J}}$ satisfies A3^b. Fix $x, y \in X$ such that $x \mathcal{P}_J y$ for some $J \in \mathbb{J}$ and take any $J' \in \mathfrak{J}$. Then, by the construction of $\mathcal{R}_{\mathfrak{J}}$, there exists $L, L' \in \mathbb{L}$ such that $jJ = L$ and $j'J' = L'$ for some $j, j' \in \mathbb{Z}_{++}$. By lemma 1.2, $\mathcal{R}_L = \mathcal{R}_J$ and $\mathcal{R}_{L'} = \mathcal{R}_{J'}$. Moreover, by construction, for some D and D' such that $L_D = L$ and $L_{D'} = L'$, $\hat{\mathcal{R}}_D = \mathcal{R}_J$ and $\hat{\mathcal{R}}_{D'} = \mathcal{R}_{J'}$. We therefore conclude that $x \hat{\mathcal{P}}_D y$, so that A3 implies the existence of $\kappa \in \mathbb{Z}_{++}$ and $\{D_l : D_l \sim^{\hat{\mathcal{R}}} D\}_1^{\kappa}$ such that $x \hat{\mathcal{P}}_{D_1 \cup \dots \cup D_{\kappa} \cup D'} y$. Then, by the construction of $\mathcal{R}_{\mathfrak{J}}$, $x \mathcal{P}_{\kappa L_D + L_{D'}} y$. Let $\nu \stackrel{\text{def}}{=} \frac{1}{\kappa j + j'}$ and take $\lambda = \nu j'$, so that $0 < \lambda < 1$ and $1 - \lambda = \nu \kappa j$. In fact, since $\lambda \in \mathbb{Q}$, we have 1460

$$K \stackrel{\text{def}}{=} (1 - \lambda)J + \lambda J' \in \mathfrak{J}.$$

Simplifying, we obtain $K = \nu(\kappa L + L')$. Since $\nu \in \mathbb{Q}_{++}$ and $\kappa L + L' \in \mathfrak{J}$, lemma 1.2 implies $\mathcal{R}_K = \mathcal{R}_{\kappa L + L'}$. This allows us to conclude that $x \mathcal{P}_K y$. Finally, take any $\mu \in \mathbb{Q} \cap (0, \lambda)$. From basic properties of the real numbers, there exists $\xi < 1$ such that $\mu = \xi \lambda$ and, moreover, ξ is rational. Next, note that the definition of K implies $\xi(K - J) = \xi \lambda(J' - J)$. Adding J to each side of the latter and applying the definition of μ yields 1470

$$(1 - \xi)J + \xi K = (1 - \mu)J + \mu J'.$$

Then, since $x \mathcal{P}_J y$ and $x \mathcal{P}_K y$, A2^b implies $x \mathcal{P}_{(1-\mu)J + \mu J'} y$, as required for A3^b. 1475

Conversely, we now assume that \mathcal{R}_J satisfies $A2^b$ and $A3^b$ and prove that $A3$ holds. Take $D, D' \in \mathbb{D}$ such that $x \hat{\mathcal{P}}_D y$ and any other $D' \in \mathbb{D}$. Let $L = L_D$ and $L' = L_{D'}$. Then, by construction, $x \mathcal{P}_L y$ and, by $A3^b$, there exists $\lambda \in \mathbb{Q} \cap (0, 1)$ such that $x \mathcal{P}_{(1-\mu)L+\mu L'} y$. Then, since μ is rational, $\mu = j/k$ for some $j, k \in \mathbb{Z}_{++}$. Let $q := (1-\mu)/\mu = (k-j)/j$ and let $\kappa = jq$, so that $\kappa = k-j$. The fact that $0 < \mu < 1$ ensures that $\kappa \in \mathbb{Z}_{++}$. To complete the proof, we show that $x \mathcal{P}_{\kappa L+L'} y$, for then the existence of D_1, \dots, D_κ such that $x \hat{\mathcal{P}}_{D_1 \cup \dots \cup D_\kappa \cup D'} y$ immediately follows. Together $x \mathcal{P}_{(1-\mu)L+\mu L'} y$ and lemma 1.2 imply $x \mathcal{P}_{qL+L'} y$. Similarly, together $x \mathcal{P}_L y$ and lemma 1.2 imply $x \mathcal{P}_{(j-1)qL} y$. Then, since $(j-1)qL + (qL + L') = jqL + L'$ and $\kappa = jq$, an application of $A2^b$ yields $x \mathcal{P}_{\kappa L+L'} y$, as required. \square

Proof of Proposition 2. Let $\mathcal{Y} = \{Y_\alpha \subseteq Y : \#Y_\alpha = 2\}$ be the collection of distinct (unordered) pairs in Y . For every $Y_\alpha \in \mathcal{Y}$, $A0$ holds simply because Y_α is of cardinality two; moreover, on Y_α , 4-diversity is equivalent to 2-diversity. This allows us to apply theorem 2 of GS03.²¹ (Part (ii) of the present lemma follows from claim 5 of the proof of lemma 1 of GS03.) Thus, on Y_α , $A1^b$ – $A3^b$ and 2-diversity hold if, and only if, there exists of a matrix $v_\alpha^{(\cdot, \cdot)} : Y_\alpha^2 \times \mathfrak{T} \rightarrow \mathbb{R}$ with rows satisfying condition (i) of the present lemma. In particular, GS03 yields a matrix representation $v_\alpha : Y_\alpha \times \mathfrak{T} \rightarrow \mathbb{R}$. For $x, y \in Y_\alpha$, take $v_\alpha^{(x, y)} = -v_\alpha(x, \cdot) + v_\alpha(y, \cdot)$. For condition (ii), note that via theorem 2 of GS03, the matrix v_α is 2-diversified if, and only if, 4-diversity holds for $\acute{\mathcal{R}}$ on Y_α . Thus, for every $\lambda \in \mathbb{R}$, $v_\alpha(x) \not\leq \lambda v_\alpha(y)$ if, and only if, 2-diversity. Take $\lambda = 1$ and observe that this implies $v_\alpha^{(x, y)}(s) < 0 < v_\alpha^{(x, y)}(t)$ for some $s, t \in \mathfrak{T}$.

To extend this to arbitrary $Y \subseteq X$, note that, for some A' , $\bigcup \{\mathcal{Y}_\alpha : \alpha \in A'\} = Y$. For an arbitrary function f , let $\text{gr } f$ denote the graph of that function. To obtain the desired matrix $\acute{v}^{(\cdot, \cdot)} : Y^2 \times \mathbb{T} \rightarrow \mathbb{R}$, take

$$\text{gr } \acute{v}^{(\cdot, \cdot)} = \bigcup \{\text{gr } v_\alpha^{(\cdot, \cdot)} : \alpha \in A'\}. \quad (26)$$

\square

²¹Gilboa and Schmeidler explicitly prove their theorem 1 holds for the case of arbitrary X and \mathbb{T} . But although their theorem 2 is stated and proved in the first step (the case where $X, \mathbb{T} < \infty$) of the proof of their theorem 1, steps 2 and 3 of that proof apply equally to their theorem 2.

Proof of Lemma 2.1. In the remainder of the proof of the present lemma, we suppress reference to the acute accent. It remains for us to prove the characterisation of novel generalizations. Let \mathcal{R} be a novel Y -generalization with matrix representation $v^{(\cdot, \cdot)}$ satisfying parts (i)–(iv) of the lemma. Fix arbitrary $t \neq \mathfrak{f}$. Then definition 2 implies the existence of $J \in \mathbb{J}$ and $L = J \times 0 \in \mathbb{J}^\mathfrak{f}$ such that $\mathcal{R}_{L+\delta_t} \neq \mathcal{R}_{L+\delta_\mathfrak{f}}$. W.l.o.g., consider the case where, for some $x, y \in Y$, it holds that both $y \mathcal{R}_{L+\delta_t} x$ and $x \mathcal{P}_{L+\delta_\mathfrak{f}} y$. Equivalently,

$$\langle v^{(x,y)}, L + \delta_t \rangle \leq 0 < \langle v^{(x,y)}, L + \delta_\mathfrak{f} \rangle$$

which, via linearity of $\langle v^{(x,y)}, \cdot \rangle$, we may rearrange to obtain

$$v^{(x,y)}(t) \leq -\langle v^{(x,y)}, L \rangle < v^{(x,y)}(\mathfrak{f}). \quad (27)$$

Thus, $v^{(x,y)}(t) \neq v^{(x,y)}(\mathfrak{f})$, as required for the lemma.

For the converse argument, fix arbitrary $t \in \mathbb{T}$. Then $\acute{v}^{(\cdot, \cdot)}(t) \neq \acute{v}^{(\cdot, \cdot)}(\mathfrak{f})$ implies the existence of distinct $x, y \in Y$ such that $\acute{v}^{(x,y)}(t) \neq \acute{v}^{(x,y)}(\mathfrak{f})$. We show that there exists $J \in \mathbb{J}$ and $L = J \times 0 \in \mathbb{J}^\mathfrak{f}$ satisfying eq. (27). For then, by retracing (in reverse order) the arguments that lead to eq. (27), we arrive at the conclusion that $\mathcal{R}_{L+\delta_t} \neq \mathcal{R}_{L+\delta_\mathfrak{f}}$, as required for \mathcal{R} to be novel.

Take $\mu = v^{(x,y)}(t)$ and $\xi = v^{(x,y)}(\mathfrak{f})$ and consider the case where $\mu < \xi < 0$. Since $\mu < 0$, 2-diversity implies that there exists $s \in \mathbb{T}$ such that $v^{(x,y)}(s)$ is positive. Then, for some $\lambda \in \mathbb{Q}_{++}$, $-\lambda v^{(x,y)}(s) \in (\mu, \xi)$. Let $L = \lambda \delta_s^\mathfrak{f}$ and observe that

$$\mu < -\langle v^{(x,y)}, L \rangle < \xi,$$

as required. *Mutatis mutandis*, the case where both μ and ξ are positive is the same. If $\mu \leq 0 \leq \xi$, then take $L = 0$, so that $\mu \neq \xi$ yields $\mathcal{R}_{L+\delta_t^\mathfrak{f}} \neq \mathcal{R}_{L+\delta_\mathfrak{f}^\mathfrak{f}}$. \square

Proof of Proposition 3. Fix $\#Y = 3$ or 4 , via lemma 2.1, let $v^{(\cdot, \cdot)}$ denote the 2-diverse matrix representation of the improper Y -generalization \mathcal{R} . Let \mathcal{H}_{++} denote the associated arrangement of hyperplanes. For every distinct $x, y \in X$, lemma 2.1 implies that $H_{++}^{\{x,y\}}$ intersects $\mathbb{R}_{++}^\mathbb{T}$. Then, similar to example 7, the $1 \leq n \leq \binom{\#Y}{2}$ distinct hyperplanes of \mathcal{H}_{++} cut $\mathbb{R}_{++}^\mathbb{T}$ into at least $n + 1$ regions. At least one pair G and G^* in \mathcal{G}_{++} are therefore separated by all n distinct members of \mathcal{H}_{++} . Take $J \in G$, so that, for every distinct $x, y \in Y$, $\langle u^{(x,y)}, J \rangle \neq 0$. Thus \mathcal{R}_J is

antisymmetric, complete and, via $A0^b$, total. Next, take $L \in G^*$, so that since J and L are separated by every hyperplane in \mathcal{H}_{++} , $\mathcal{R}_J = \mathcal{R}_L^{-1}$. \square

Proof of Lemma 2.2. Let $A4^b$ hold on Y . Since $v^{(\cdot, \cdot)}$ is a 2-diverse pairwise representation, $v^{(x,z)}, v^{(y,z)} \not\leq 0$. By $A4^b$, one of $G_+^{(x,z)}$ and $G_+^{(z,x)}$ contains both J, L such that 1535

$$\langle v^{(y,z)}, L \rangle < 0 < \langle v^{(y,z)}, J \rangle.$$

W.l.o.g., suppose J, L belongs to $G_+^{(x,z)}$. Then $\langle v^{(x,z)}, \cdot \rangle$ is positive on $\{L, J\}$, so that, for every $\lambda \in \mathbb{R}$ $v^{(x,z)} \neq \lambda v^{(y,z)}$, as required.

Conversely, let $x, y, z \in Y$ be such that $v^{(x,z)}$ and $v^{(y,z)}$ are noncollinear. Then 1540
 $H_{++}^{\{x,z\}} \neq H_{++}^{\{y,z\}}$, and there exists $L \in H_{++}^{\{x,z\}} - H_{++}^{\{y,z\}}$. W.l.o.g., therefore, suppose $L \in H_{++}^{\{x,z\}} \cap G_{++}^{(y,z)}$. Since $v^{(\cdot, \cdot)}$ is 2-diverse, there exists $s, t \in \mathbb{T}$ such that $v^{(x,z)}(s) < 0 < v^{(x,z)}(t)$. Noting that $L \in \mathbb{R}_{++}^{\mathfrak{T}}$, so that $v^{(x,z)} \neq \delta_s, \delta_t$, let ψ_s and ψ_t be the convex paths from L to δ_s and δ_t respectively. For sufficiently small $\lambda > 0$, $\langle v^{(y,z)}, \psi_{s'}(\lambda) \rangle$ remains positive for $s' = s, t$ and, moreover, since $L \in H_{++}^{\{x,z\}}$, 1545

$$\langle v^{(x,z)}, \psi_s(\lambda) \rangle < 0 < \langle v^{(x,z)}, \psi_t(\lambda) \rangle.$$

Finally, since L has finite support, a finite sequence of perturbations of the elements of $\psi_s(\lambda)$ and $\psi_t(\lambda)$ yields (rational-valued) members of \mathfrak{J} with the same properties, as required for $A4^b$. \square

Proof of Proposition 4. When $X = 2$, $A4^b$ and $A4'^b$ are identical to 2-diversity. 1550
 Let $Y = \{x, y, z\} \subseteq X$ and let \mathcal{R} denote the improper Y -generalization of $\leq_{\mathbb{J}}$. We begin by assuming $A4^b$ and showing that $\#Y + 1 = 4 \leq \#\text{total}(\mathcal{R})$. Via lemma 2.2, there are three distinct hyperplanes $H_{++}^{\{x,y\}}, H_{++}^{\{y,z\}}$ and $H_{++}^{\{x,z\}}$ in the associated arrangement \mathcal{H}_{++} . Then, as in example 7, $\emptyset = \mathbb{R}_{++}^{\mathbb{T}}$ is the unique element of \mathcal{L}_{++} that lies below each member of \mathcal{H}_{++} . Thus, via eq. (14), $\mu(A^\emptyset) = 1$ 1555
 and $\mu(A) = -\mu(A^\emptyset)$ for all three hyperplanes $A \in \mathcal{H}_{++}$. Thus, Zaslavski's theorem implies that $\#\mathcal{G}_{++}$ is bounded below by 4. Thus $\text{total}(\mathcal{R}) \geq 4$, and since, for every Y -generalization $\hat{\mathcal{R}}$, $\#\text{total}(\hat{\mathcal{R}}) \geq \#\text{total}(\mathcal{R})$, $A4'^b$ holds.

Conversely, suppose $A4'^b$ holds and, once again let \mathcal{R} denote the improper Y -generalization of $\leq_{\mathbb{J}}$, so that $\text{total}(\mathcal{R}) \geq \#Y = 3$. Now $A4'^b$ implies 2-diversity, so 1560
 that, via lemma 2.1, there exists a 2-diverse matrix representation with associated

arrangement \mathcal{H}_{++} . It is not the case that $|\mathcal{H}_{++}| = 1$, for this would imply that $\#\text{total}(\mathcal{R}) = 2$. W.l.o.g., suppose $H_{++}^{\{x,y\}} \neq H_{++}^{\{y,z\}}$. Observe that $A0^b$ then implies $H_{++}^{\{x,z\}} \neq H_{++}^{\{x,y\}}$ and $H_{++}^{\{x,z\}} \neq H_{++}^{\{y,z\}}$. This implies that $v^{(x,y)}, v^{(y,z)}$ and $v^{(x,z)}$ are pairwise noncollinear. Finally, an application of lemma 2.2 then yields $A4^b$. \square 1565

Proof of Lemma 2.4. Note that, when $1 \leq |X| \leq 2$, 4-Jac holds vacuously and $\leq_{\mathbb{J}}$ has a Jacobi representation via lemma 2.1. For general X , the fact that 4-Jac is necessary for $\leq_{\mathbb{J}}$ to have a Jacobi representation follows simply because if the Jacobi identity holds on X , then it holds on every $Y \subseteq X$. For the sufficiency of 4-Jac, we proceed by induction. As in lemma 3 and claim 9 of GS03, we assume 1570 that X is well-ordered.

In the case that $|X| \leq 4$, we only need to show that $v^{(\cdot, \cdot)}$ is unique. W.l.o.g., we take the initial step in our induction argument to satisfy $|X| = 4$. Let $\mathbf{v}^{(\cdot, \cdot)}$ denote any other Jacobi representation of $\leq_{\mathbb{J}}$. By lemma 2.1, for every distinct $x, y \in Y^2$, there exists $\lambda^{\{x,y\}} > 0$ such that $\mathbf{v}^{(x,y)} = \lambda^{\{x,y\}} v^{(x,y)}$. We need to show 1575 that $\lambda^{\{x,y\}} = \lambda$ for every distinct $x, y \in Y$. Let $Y = \{x, y, z, w\}$. By lemma 2.2, the set $\{v^{(x,y)}, v^{(x,z)}, v^{(x,w)}\}$ is pairwise noncollinear. Then, since the Jacobi identity holds for both $v^{(\cdot, \cdot)}$ and $\mathbf{v}^{(\cdot, \cdot)}$, we derive the equation

$$(1 - \lambda^{\{x,y\}})v^{(x,y)} + (1 - \lambda^{\{y,z\}})v^{(y,z)} = (1 - \lambda^{\{x,z\}})v^{(x,z)} \quad (28)$$

Suppose that $1 - \lambda^{\{y,z\}} = 0$. Then, either the other coefficients in eq. (28) are 1580 both equal to zero (and our proof is complete), or we obtain a contradiction of lemma 2.2. Thus, $1 - \lambda^{\{y,z\}}$ is nonzero and we may divide through by this term and solve for $v^{(y,z)}$. First note that, since $v^{(\cdot, \cdot)}$ is a Jacobi representation, $v^{(y,x)} + v^{(x,y)} = v^{(y,y)} = 0$. Then, since $v^{(y,x)} = -v^{(x,y)}$,

$$v^{(y,z)} = \frac{1 - \lambda^{xy}}{1 - \lambda^{yz}} v^{(y,x)} + \frac{1 - \lambda^{xy}}{1 - \lambda^{yz}} v^{(x,z)}. \quad 1585$$

We then conclude that both of the coefficients in the latter equation are equal to one. (This follows from linear independence of $v^{(y,x)}$ and $v^{(x,z)}$ together with the Jacobi identity $v^{(y,z)} = v^{(y,x)} + v^{(x,z)}$.) Thus, $\lambda^{\{x,y\}} = \lambda^{\{y,z\}} = \lambda^{\{x,z\}}$, as required. Repeated application of the same argument to the remaining Jacobi identities yields the desired conclusion, $\mathbf{v}^{(\cdot, \cdot)} = \lambda v^{(\cdot, \cdot)}$. 1590

For the inductive step, take Y to be an initial segment of X . By the induction hypothesis, there exists a Jacobi representation $\mathbf{v}^{(\cdot, \cdot)} : Y^2 \times \mathbb{T} \rightarrow \mathbb{R}$ of the improper Y -generalization $\mathcal{R} = \leq_{\mathbb{J}} \cap Y^2$ that is suitably unique.

Claim 3.5.1. *For every $w \in X - Y$ and $W \stackrel{\text{def}}{=} Y \cup \{w\}$, there exists a Jacobi representation $\hat{\mathbf{v}}^{(\cdot, \cdot)} : W^2 \times \mathbb{T} \rightarrow \mathbb{R}$ of the improper W -generalization $\hat{\mathcal{R}}$.*

1595

Proof of claim 3.5.1. Via lemma 2.2, there exists a conditionally 2-diverse pairwise representation $u^{(\cdot, \cdot)}$ of $\leq_{\mathbb{J}}$. Fix any four distinct elements x, x', y, z in Y . Lemma 2.1 implies the existence of $\phi, \phi' \in \mathbb{R}_{++}$ such that $\phi u^{(x, z)} = \mathbf{v}^{(x, z)}$ and $\phi' u^{(x', z)} = \mathbf{v}^{(x', z)}$. Let $Z = \{x, y, z, w\}$ and $Z' = \{x', y, z, w\}$. Since β -Jac holds, there exist positive scalars $\alpha, \beta, \hat{\beta}, \gamma, \sigma$ and τ such that

1600

$$\alpha u^{(x, w)} + \beta u^{(w, y)} = \gamma u^{(x, y)}, \quad (29)$$

$$\hat{\beta} u^{(y, w)} + \sigma u^{(w, z)} = \tau u^{(y, z)}, \text{ and} \quad (30)$$

$$\gamma u^{(x, y)} + \tau u^{(y, z)} = \mathbf{v}^{(x, z)}. \quad (31)$$

Moreover, β -Jac ensures that we may take $\beta = \hat{\beta}$. Since $u^{(\cdot, \cdot)}$ is conditionally 2-diverse, $\{u^{(x, y)}, u^{(y, z)}\}$ is linearly independent, and the linear system eq. (31) in the unknowns γ and τ has a unique solution. This, together with the induction hypothesis (which yields $\mathbf{v}^{(x, y)} + \mathbf{v}^{(y, z)} = \mathbf{v}^{(x, z)}$) implies that $\gamma u^{(x, y)} = \mathbf{v}^{(x, y)}$ and $\tau u^{(y, z)} = \mathbf{v}^{(y, z)}$. Similarly, for Z' , β -Jac yields $\alpha', \beta', \sigma', \gamma', \tau' > 0$ such that

1605

$$\alpha' u^{(x', w)} + \beta' u^{(w, y)} = \gamma' u^{(x', y)}, \quad (32)$$

$$\beta' u^{(y, w)} + \sigma' u^{(w, z)} = \tau' u^{(y, z)}, \text{ and} \quad (33)$$

1610

$$\gamma' u^{(x', y)} + \tau' u^{(y, z)} = \mathbf{v}^{(x', z)}. \quad (34)$$

As in the arguments involving γ and τ , the induction hypothesis yields $\gamma' u^{(x', y)} = \mathbf{v}^{(x', y)}$ and $\tau' u^{(y, z)} = \mathbf{v}^{(y, z)}$. We conclude that $\tau = \tau'$. Substituting for τ' in eq. (33) and appealing to linear independence of $\{u^{(y, w)}, u^{(w, z)}\}$ then yields the desired equalities $\beta = \beta'$ and $\sigma = \sigma'$.

1615

As a consequence of the above argument, for every $y, z \in Y$, take $\hat{\mathbf{v}}^{(y, w)}$ and $\hat{\mathbf{v}}^{(w, z)}$ to be the unique vectors in $\mathbb{R}^{\mathbb{T}}$ that solve the equation $\hat{\mathbf{v}}^{(y, w)} + \hat{\mathbf{v}}^{(w, z)} = \mathbf{v}^{(y, z)}$. For every $y, z \in Y$, let $\hat{\mathbf{v}}^{(y, z)} = \mathbf{v}^{(y, z)}$ and $\hat{\mathbf{v}}^{(w, w)} = 0$. Then the matrix $\hat{\mathbf{v}}^{(\cdot, \cdot)}$ with row vectors $\{\hat{\mathbf{v}}^{(x, y)} : x, y \in W\}$ is a Jacobi representation of $\hat{\mathcal{R}}$. \square

Our proof of claim 3.5.1 shows that the generalization to W holds for any 1620 initial subsegment of Y consisting of four elements. Our proof thereby accounts for the case where X is infinite and w is a limit ordinal. \square