

A
Difference in subjective SES coefficient
(weighted minus unweighted, same sample)
-0.01
0.00
0.01
0.02
Outcome
Meaningfulness of life
Evaluative wellbeing
Hedonic wellbeing
Loneliness (rev.)
Depression (rev.)
Anxiety (rev.)
Frailty (rev.)
Self-rated health
Mobility limitations (rev.)
IADL difficulties (rev.)
Fibrinogen (rev.)
ADL difficulties (rev.)
HDL/total cholesterol ratio
Physical activity
C-reactive protein (rev.)
Executive function
Memory
Gait speed
Triglycerides (rev.)
Allostatic load (rev.)
Processing speed
Glycated hemoglobin (rev.)
Grip strength
Cross-sectional weights
Longitudinal weights
B
Difference in subjective SES coefficient
(weighted minus unweighted, same sample)
-0.010
0.000
0.010
0.020
0.030
Outcome
Low peak expiratory flow (rev.)
Limiting longstanding illness (rev.)
Good hearing
Low forced vital capacity (rev.)
Low forced expiratory volume (rev.)
Pain (rev.)
Good eyesight
High blood pressure: diastolic (rev.)
Smoking (rev.)
High fasting blood glucose (rev.)
Central obesity (rev.)
High blood pressure: systolic (rev.)
Drinking alcohol (rev.)
BMI obesity (rev.)