

Notes for Math 8302-01
Fall 2014
Introduction to Geometric Measure
Theory and Quantitative Rectifiability

Patrick Spencer
Last updated August 28, 2014

Contents

Chapter 1. Intro and Syllabus	5
Chapter 2. Review of measure theory	7
Bibliography	11

CHAPTER 1

Intro and Syllabus

The following is basic information about the class from the syllabus.

Institution: University of Missouri at Columbia

Semester: Fall 2014

Instructor: Steve Hofmann

Course title: Math 8302 - Introduction to Geometric Measure Theory and Quantitative Rectifiability.

Description: Geometric Measure Theory (GMT) is largely concerned with the notion of “rectifiability”, which describes a sense in which a general set of points (in \mathbb{R}^n , say) is approximated by “nice” surfaces. In the past couple of decades, this concept has been sharpened: “uniform (i.e., quantitative) rectifiability” entails approximation of a set by nice surfaces, but in a quantitatively precise way, which turns out to have deep connections with the behavior, on the set, of singular integral operators, square functions, and harmonic measure. We plan to cover the following topics, to the extent that time permits

- (1) Review of Hausdorff measure
- (2) Bounded variation, sets of finite perimeter, isoperimetric inequality, reduced boundary, measure theoretic boundary, Hausdorff-Green theorem.
- (3) Ahlfors-David Regularity, “Spaces of homogeneous type” and M. Christ’s dyadic cube reconstruction.
- (4) NTA domains and “big pieces of Lipschitz graphs”.
- (5) Uniform rectifiability, Corona decomposition, “Geometric Lemma” and the “Bilateral Weak Geometric Lemma”, singular integrals and square functions.

Prerequisites Familiarity with the elementary theory of measure and integration and with the basic subject matter of harmonic analysis: Fourier transform, Hardy-Littlewood maximal function, approximate identities, Whitney decomposition, Littlewood-Paley theory, classical Calderon-Zygmund theory, BMO, Carleson measures, theory of singular integrals on Lipschitz graphs.

Text: There is no official textbook for the course, however, much of the material presented in the course will follow [1],[2], [3], and [4]. Other material may be taken directly from the literature.

CHAPTER 2

Review of measure theory

Most of the material in this chapter is from the book “Measure Theory and Fine Properties of Functions” by Evans and Gariepy [3]

DEFINITION 0.1. A *set function* f is a mapping $f : P(X) \rightarrow Y$ where Y is a Banach space. Y is usually \mathbb{R} or \mathbb{C} .

DEFINITION 0.2. Let X be a set. We let $P(X)$ denote the power set of X i.e. the set of all subsets of X .

DEFINITION 0.3. Let X be set. A non-empty subset $\mathfrak{M} \subseteq P(X)$ is called an *algebra* on X if \mathfrak{M} has the following properties:

- (1) $A \in \mathfrak{M}$ implies $A^c \in \mathfrak{M}$.
- (2) If $n \in \mathbb{N}$ and $A_1, \dots, A_n \in \mathfrak{M}$ then $\bigcup_{i=1}^n A_i \in \mathfrak{M}$

A σ -*algebra* is an algebra which is closed under countable unions instead of finite unions.

REMARK 0.4. Notice that if \mathfrak{M} is an algebra or a σ -algebra on X then $\emptyset \in \mathfrak{M}$ and $X \in \mathfrak{M}$ because if $E \in \mathfrak{M}$ then $\emptyset = E \cap E^c$ and $X = E \cup E^c$.

DEFINITION 0.5. Suppose X is a set and \mathfrak{M} is a σ -algebra on X . A set function $\mu : \mathfrak{M} \rightarrow [0, \infty]$ is called a *measure* on (X, \mathfrak{M}) if μ has the following properties:

- (1) $\mu(\emptyset) = 0$
- (2) $\mu(\sum_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \mu(A_i)$ whenever $\{A_i\}_{i=1}^{\infty}$ is a countable collection of disjoint sets in \mathfrak{M} .

A slightly more relaxed notion of a measure is what is called an outer measure.

DEFINITION 0.6. A set function $\tilde{\mu} : P(X) \rightarrow [0, \infty]$ is called an *outer measure* if

- (1) $\tilde{\mu}(\emptyset) = 0$
- (2) $\tilde{\mu}(A) \leq \tilde{\mu}(B)$ if $A \subseteq B \subseteq X$,
- (3) $\tilde{\mu}(\bigcup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \tilde{\mu}(A_i)$ whenever $\{A_i\}_{i=1}^{\infty} \subseteq P(X)$.

DEFINITION 0.7. Given an outer measure $\tilde{\mu}$ a subset $E \subseteq X$ is $\tilde{\mu}$ -measurable if it satisfies the following *Caratheodory Criterion*:

$$(0.1) \quad \tilde{\mu}(A) = \tilde{\mu}(A \cap E) + \tilde{\mu}(A \setminus E), \quad \text{for all } A \subseteq X.$$

Equivalently we have that E is $\tilde{\mu}$ -measurable iff $\tilde{\mu}(A_1 \cap A_2) = \tilde{\mu}(A_1) + \tilde{\mu}(A_2)$ whenever $\tilde{\mu}(A_2 \subseteq E)$ and $A_2 \cap E = \emptyset$.

PROPOSITION 0.8. (exercise) Let $\tilde{\mu}$ be an outer measure on X . If $\tilde{\mu}(Z) = 0$ then Z is $\tilde{\mu}$ -measurable.

NOTE 0.9. This says \emptyset is measurable.

PROPOSITION 0.10. (exercise) Given an outer measure $\tilde{\mu}$, the collection of all $\tilde{\mu}$ -measurable sets forms a σ -algebra.

DEFINITION 0.11. Denote the σ -algebra of $\tilde{\mu}$ -measurable sets as $\mathfrak{M}_{\tilde{\mu}}$.

PROPOSITION 0.12. (exercise) Any outer measure $\tilde{\mu}$ restricted to its measurable sets is a measure.

THEOREM 0.13. (exercise) Suppose $\tilde{\mu}$ is an outer measure and that $\{A_k\}_{k=1}^{\infty}$ is a collection of $\tilde{\mu}$ -measurable sets.

- (1) If $\{A_k\}_{k=1}^{\infty}$ are pairwise disjoint then $\tilde{\mu}(\bigcup_{k=1}^{\infty} A_k) = \sum_{k=1}^{\infty} \tilde{\mu}(A_k)$.
- (2) If $A_1 \subseteq A_2 \subseteq \dots$ then $\lim_{k \rightarrow \infty} \tilde{\mu}(A_k) = \tilde{\mu}(\bigcup_{k=1}^{\infty} A_k)$.
- (3) If $A_1 \supseteq A_2 \supseteq \dots$ and $\tilde{\mu}(A_1) < \infty$ then $\lim_{k \rightarrow \infty} \tilde{\mu}(A_k) = \tilde{\mu}(\bigcap_{k=1}^{\infty} A_k)$.

DEFINITION 0.14. Suppose X is a topological space. The *Borel σ -algebra*, denoted \mathfrak{B}_X , is the smallest σ -algebra that contains all the open set. This can be constructed by taking the intersection of all σ -algebras on X which contain the all open sets of X .

DEFINITION 0.15. A subset $A \subseteq X$ is called σ -finite w.r.t. $\tilde{\mu}$ if $A = \bigcup_{k=1}^{\infty} B_k$ where $\{B_i\}_{i=1}^{\infty} \subseteq \mathfrak{M}_{\tilde{\mu}}$ and $\tilde{\mu}(B_i) < \infty, \forall i$.

DEFINITION 0.16. Let X be an arbitrary space and $\tilde{\mu}$ an outer measure on X .

- (1) $\tilde{\mu}$ is *regular* if $\forall A \subseteq X$ there exists a $B \in \mathfrak{M}_{\tilde{\mu}}$ so that $A \subseteq B$ and $\tilde{\mu}(A) = \tilde{\mu}(B)$.

- (2) $\tilde{\mu}$ is a *Borel* measure if every Borel set is $\tilde{\mu}$ -measurable.
- (3) $\tilde{\mu}$ is *Borel regular* if $\tilde{\mu}$ is Borel and $\forall A \subseteq \mathbb{R}$ (or X) there exists a Borel set B with $A \subseteq B$ and $\tilde{\mu}(A) = \tilde{\mu}(B)$.
- (4) $\tilde{\mu}$ is a *Radon* measure if $\tilde{\mu}$ is Borel regular and if $\tilde{\mu}(K) < \infty$ for every compact set K .

THEOREM 0.17. [3, thm 2] Let $\tilde{\mu}$ be a regular outer measure on X . If $A_1 \subseteq A_2 \subseteq \dots$ then $\lim_{k \rightarrow \infty} \tilde{\mu}(A_k) = \tilde{\mu}(\cup_{k=1}^{\infty} A_k)$.

REMARK 0.18. The significance of this theorem is that we are not requiring the A_k to be $\tilde{\mu}$ -measurable.

DEFINITION 0.19. Let $\tilde{\mu}|_A$ be defined as $\tilde{\mu}|_A(B) = \tilde{\mu}(A \cap B)$ for ever $B \subseteq X$.

THEOREM 0.20. [3, thm 3] Suppose $\tilde{\mu}$ is a Borel regular outer measure on \mathbb{R}^n and $A \subseteq \mathbb{R}^n$ is $\tilde{\mu}$ -measurable with $\tilde{\mu}(A) < \infty$. Then $\tilde{\mu}|_A$ is a Radon measure.

LEMMA 0.21. [3, lemma 1] Let $\tilde{\mu}$ be a Borel regular outer measure on \mathbb{R}^n . and let $B \in \mathfrak{B}_X$.

- (1) If $\tilde{\mu} < \infty$ then for every $\varepsilon > 0$ there exists a closed set C_ε so that $C_\varepsilon \subseteq B$ and $\tilde{\mu}(B \setminus C_\varepsilon) < \varepsilon$.
- (2) If $\tilde{\mu}$ is a Radon outer measure, then for ever $\varepsilon > 0$ there exists an open set U_ε so that $B \subseteq U_\varepsilon$ and $\tilde{\mu}(B \setminus U_\varepsilon) < \varepsilon$.

THEOREM 0.22. [3, thm 4](Approximation by open and compact sets)
Let $\tilde{\mu}$ be a Radon outer measure on \mathbb{R}^n . Then

- (1) for each set $A \subseteq \mathbb{R}^n$

$$\tilde{\mu}(A) = \inf \{ \tilde{\mu}(U) : A \subseteq U, U \text{ is open} \},$$

and

- (2) for each $\tilde{\mu}$ -measurable set $A \subseteq \mathbb{R}^n$

$$\tilde{\mu}(A) = \sup \{ \tilde{\mu}(K) : K \subseteq A, K \text{ is compact} \}.$$

THEOREM 0.23. [3, thm 5](Caratheodory's Criterion) Let $\tilde{\mu}$ be an outer measure on \mathbb{R}^n . If $\tilde{\mu}(A \cup B) = \tilde{\mu}(A) + \tilde{\mu}(B)$ for all sets $A, B \subseteq \mathbb{R}^n$ with $\text{dist}(A, B) > 0$, then $\tilde{\mu}$ is a Borel outer measure.

Bibliography

1. Guy David and Stephen Semmes, *Singular integrals and rectifiable sets in \mathbb{R}^n : Beyond lipschitz graphs*, vol. 193, Société mathématique de France, 1991.
2. ———, *Analysis of and on uniformly rectifiable sets*, vol. 38, American Mathematical Soc., 1993.
3. Lawrence Craig Evans and Ronald F Gariepy, *Measure theory and fine properties of functions*, vol. 5, CRC press, 1991.
4. Richard Wheeden, Richard L Wheeden, and Antoni Zygmund, *Measure and integral: an introduction to real analysis*, CRC Press, 1977.