

# Adverse Health Effects of Frailty in CKD Patients

Cause .....	4
Lifestyle .....	4
Ethnicity .....	4
Biological .....	4
Cardiovascular .....	4
Vascular access.....	4
Cerebrovascular .....	4
Pulmonary .....	4
Inflammatory .....	4
Endocrinologic/ Metabolic.....	5
Body Composition .....	5
Cancer .....	5
Arthritis .....	5
Laboratory Data .....	5
Non-adjusted differences in comorbidity and system-based influences.....	8
Biological .....	8
Cardiovascular .....	8
Cerebrovascular .....	8
Vascular Access .....	8
Endocrinologic/ Metabolic.....	8
Cognitive .....	9

Immunological .....	9
Body Composition .....	9
Laboratory Data .....	9
Miscellaneous .....	10
Composite .....	11
Functional Status.....	11
Microbiota .....	12
Gut microbiota .....	12
Neurological .....	12
EEG .....	13
Psychological .....	13
Mood.....	13
Mental Health .....	14
Mental Function.....	14
Sociological.....	14
Interaction.....	14
Physical activity .....	14
Nutritional Status.....	15
Quality of Life .....	15
Independence .....	16
Health-care utilization.....	16
Complications.....	18
Biological .....	18
Cardiovascular.....	18
Renal Function Decline .....	18

Immunological .....	18
Cognitive .....	18
Functional status .....	19
Diabetes .....	19
Body composition .....	19
Laboratory data .....	23
Psychological .....	24
Delirium .....	24
Distress .....	24
Quality of Life .....	24
Graft Loss .....	28
Health-care utilization .....	28
Mortality .....	30
Composite .....	33
Bibliography .....	35

## Cause

	Effect (descriptions)	Risk Difference	CKD severity	Frailty Assessment	Sample Size	Reference
<b>Lifestyle</b>	Smoking	RR 1.18 (1.04-1.34)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	<sup>1</sup>
<b>Ethnicity</b>	Hispanic	Frailty scores +0.6 points per year	CKD stage 5D (hemodialysis)	Fried Phenotypes	762	<sup>2</sup>
<b>Biological</b>						
Cardiovascular	Hypertension	RR 1.6 (1.26-2.04)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	<a href="#"><sup>1</sup></a>
	Peripheral vascular disease	RR 1.58 (1.34-1.8)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	<a href="#"><sup>1</sup></a>
	Left ventricular dysfunction	RR 1.18 (1.03-1.36)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	<a href="#"><sup>1</sup></a>
	Endothelial dysfunction	$r = -0.367$ (p= 0.004) OR 3.86 (1.00-14.88)	CKD stages 3-5	Fried Phenotypes	61	<sup>3</sup>
Vascular access	Use of a permanent vascular access (fistula or graft)	HR 0.71 (0.51-0.98)	CKD stage 5D (hemodialysis)	Fried Phenotypes	2275	<sup>4</sup>
Cerebrovascular	Cerebrovascular Accident	RR 1.34 (1.19-1.5)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	<a href="#"><sup>1</sup></a>
Pulmonary	COPD	OR 1.68 (1.16-2.45)	CKD stages 1-5	Fried Phenotypes	10256	<sup>5</sup>
Inflammatory	IL-6	Worse frailty	CKD stage 5D	Fried	762	<sup>2</sup>

			(hemodialysis)	Phenotypes		
	CRP	After adjustment, OR	Chronic kidney	Fried	5888	<sup>6</sup>
	Fibrinogen	1.76 (1.28-2.41) to 1.50 (1.07-2.09)	insufficiency, serum creatinine ≥ 1.3 mg/dL	Phenotypes		
Endocrinologic/ Metabolic	Diabetes	Frailty scores +0.7 points per year	CKD stage 5D (hemodialysis)	Fried	762	<sup>2</sup>
		OR 1.68 (1.16-2.45)	CKD stages 1-5	Fried	10256	<sup>5</sup>
				Phenotypes		
	Obesity (IMC ≥ 30 kg/m <sup>2</sup> )	OR 6.63 (1.16-36.77)	CKD stages 3-5	Fried	61	<sup>3</sup>
				Phenotypes		
	Higher parathyroid hormones (PTH)	r= 0.30 (p= 0.01)	CKD stages 3-5	Fried	61	<sup>3</sup>
				Phenotypes		
Body Composition	Higher fat mass	r= 0.25 (p= 0.04)	CKD stages 3-5	Fried	61	<sup>3</sup>
				Phenotypes		
Cancer	Cancer	OR 1.89 (1.19-2.99)	CKD stages 1-5	Fried	10256	<sup>5</sup>
				Phenotypes		
Arthritis	Arthritis	OR 3.34 (2.08-5.38)	CKD stages 1-5	Fried	10256	<sup>5</sup>
				Phenotypes		
Laboratory Data	eGFR (mL/min/1.72m <sup>2</sup> ) reference: > 60					
	eGFRcys < 30	Frailty prevalence 2.8	CKD stages 1-4	Fried	336	<sup>7</sup>
				Phenotypes		
	eGFRcys 30-44	Frailty prevalence 2.1				

Serum Albumin Concentrations (g/dL)	Frailty scores -1.1 points per g/dL	CKD stage 5D (hemodialysis)	Fried Phenotypes	762	<sup>2</sup>
Serum Creatinine <4 mg/dL*	RR 1.46 (1.22-1.71)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	<sup>1</sup>
Testosterone, every 50% lower free form		CKD stage 5D (hemodialysis), men	Fried Phenotypes	440	<sup>8</sup>
being frail	OR 1.40 (1.05-1.53)				
becoming frail over 12 months	OR 1.40 (1.07-1.73)				
25-hydroxy vitamin D (ng/mL)	Pearson correlation coefficient r = -0.363 (p = 0.002) 11.58 vs. 17.09 (p = 0.005)	CKD stage 5D (hemodialysis)	Fried Phenotypes	74	<sup>9</sup>
Hemoglobin	Pearson correlation coefficient r = -0.336 (p = 0.004) 6.49 g/dL vs. 8.07 g/dL (p = 0.037)	CKD stage 5D (hemodialysis)	Fried Phenotypes	74	<sup>9</sup>
	Adjusted, OR 1.76 (1.28-2.41) to 1.50 (1.07-2.09)	Chronic kidney insufficiency, serum creatinine ≥1.3mg/dL	Fried Phenotypes	5888	<sup>6</sup>
LDL, HDL	Adjusted, OR 1.76 (1.28-2.41) to 1.50	Chronic kidney insufficiency, serum	Fried Phenotypes	5888	<sup>6</sup>

(1.07-2.09)	creatinine $\geq 1.3$ mg/dL
-------------	-----------------------------

## Non-adjusted differences in comorbidity and system-based influences

		Effect (descriptions)	Prevalence	CKD Severity	Frailty Assessment	Sample Size	Reference
Biological							
	Cardiovascular	Heart Failure (%)	30% vs 12%	CKD stages 1-4	Fried Phenotypes	336	<sup>7</sup>
		Angina (%)	34% vs. 22%	CKD stages 1-4	Fried Phenotypes	336	<sup>7</sup>
	Cerebrovascular	Cerebrovascular Disease (%)	26.4 vs. 12.0	CKD stage 5D (hemodialysis)	Fried Phenotypes	324	<sup>10</sup>
	Vascular Access	Catheter vs. Native/Prosthetic FAVI (%)	61.6% vs. 27.78/11.11%	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	<sup>11</sup>
	Endocrinologic/ Metabolic	Diabetes (%)	F vs. NF				
			64% vs. 49%	CKD stages 1-4	Fried Phenotypes	336	<sup>7</sup>
		Obesity (%)	F vs. NF				
			64% vs. 50%	CKD stages 1-4	Fried Phenotypes	336	<sup>7</sup>
			51.8% vs. 23.9%	CKD stage 5D (hemodialysis)	Fried Phenotypes	324	<sup>10</sup>
		BMI based on dry weight	31.5 vs. 27.6	CKD stage 5D (hemodialysis)	Fried Phenotypes	324	<sup>10</sup>
		BMI	22.53 vs. 26.16	CKD stage 5D	Fried	320	<sup>11</sup>



				(hemodialysis)	Phenotypes		
Cognitive	Executive Function		F vs. NF				
		Trail Making Tests A (TMTA) scores	+12.08	CKD stage 5D (hemodialysis)	Fried Phenotypes	324	<sup>10</sup>
		Trail Making Tests B (TMTB) scores	+33.15	CKD stage 5D (hemodialysis)	Fried Phenotypes	324	<sup>10</sup>
Immunological	Inflammatory						
		CRP (ln CRP) (mg/dL)	1.12 vs 0.28	CKD stage 5D (peritoneal dialysis)	Clinical Frailty Scale (CFS)	119	<sup>12</sup>
		IL6 (ln IL6) (mg/dL)	2.45 vs. 1.58				
	Mycophenolate mofetil (MMF) dose reduction (MDR)		F vs. NF	CKD stage 5T	Fried Phenotypes	525	<sup>13</sup>
		1 year since KT (%)	44 vs 40				
		2 years since KT (%)	54 vs. 45				
		3 years since KT (%)	67 vs. 51				
	Viral infection		F vs. NF				
		HCV (n=37) (capitals)	36 vs. 1	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	<sup>1</sup>
Body Composition	Lower lean mass						
		Lean body mass (in frail vs. nonfrail) (%)	57.1% vs .14.7%	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup>
Laboratory Data	Prealbumin (PRAB) (mg/dL)		28.9 vs. 38.3	CKD stage 1-5	Edmonton Frail Scale	41	<sup>12</sup>

					(EFS)		
		Serum albumin (mg/L)	Pearson correlation coefficient $r = -0.263$ ( $p = 0.025$ )	CKD stage 5D (hemodialysis)	Fried Phenotypes	74	<sup>9</sup>
			38 vs. 41	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup>
			29.2 vs. 34.8	CKD stage 5D (peritoneal dialysis)	Clinical Frailty Scale (CFS)	119	<sup>12</sup>
			36.1 vs. 38.5	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	<sup>11</sup>
		Frail with depression vs. Frail without depression vs. Nonfrail	32.9 vs. 34.9 vs. 35.8 ( $p=0.025$ )	CKD stage 5D (peritoneal dialysis)	In-house Chinese questionnaire	178	<sup>15</sup>
		Calcium (mmol/L)	2.24 vs. 2.36	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup>
		Hemoglobin (g/dL)	10.35 vs. 10.97	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	<sup>11</sup>
	Miscellaneous	Dialysis clearance rate (Kt/V and urea	↑	CKD stage 5D	FRAIL scale	46	<sup>16</sup>

		reduction ratio)		(chronic dialysis)			
Composite	Number of complications (complications identified at data collection: high pressure cramping, anemia, weight loss pain, weakness, weight gain constipation, heart arrhythmia, headache, itch, recurrent infections, arterial hypertension)		Spearman's correlation 0.666 (p = 0.000 in table) (p < 0.05 in text)	Elderly ( $\geq 60$ yo), with diagnosis of CKD	Edmonton Frail Scale (EFS)	35	<sup>17</sup>
	Higher number of comorbid conditions		6 vs. 4 (p = 0.03)	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup>
	Charlson's comorbidity score		Spearman's rank correlation coefficient r = 0.40 (p < 0.0001)	CKD stage 5D (peritoneal dialysis)	Chinese questionnaire	193	<sup>18</sup>
Functional Status	Disability		F vs. NF				
		$\geq 1$ disability in activities of daily Living (ADLs)	15% vs. 5%	CKD stages 1-4	Fried Phenotypes	336	<sup>7</sup>
		$\geq 1$ disability in instrumental activities of daily living (IADLs)	60% vs. 28%				

			≥1 disability in mobility tasks	40% vs. 18%				
			Less ADL-independent	55% vs. 91%	≥ 65 yo, predialysis, eGFR < 20 mL/min	Groningen frailty indicator (GFI)	65	19
		Ability to perform basic activities of daily living		33.33% vs 76.4%	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	11
		Ability to perform transfers		38.8% vs. 84.7%				
Microbiota								
	Gut microbiota	Malnutrition-Inflammation-Score (MIS)		7.6 vs. 3.9	Stage 3b-4, eGFR 15-45ml/min	Fried Phenotype score	64 (and 15 control subjects)	20
			Abundance of unclassified Mogibacteriaceae and Oscillospira	Directly proportional to MIS				
			Abundance of Akkermansia, Ruminococcus, and Eubacterium	Inversely proportional to MIS				
		Bacterial Abundance of some genera (Mogibacteriacee, Coriobacteriacee, Eggerthella, Erwinia, Coprobacillus, Anaerotruncus, etc)		↑				
Neurological								

EEG	Brain Wave		F vs. NF	ESRD, under chronic dialysis	FRAIL scale	46	16
		Global DAR	283 ± 679 vs. 2971 ± 4859				
		DARs (left frontal)	135 ± 250 vs. 3073 ± 4702				
		DAR (left TO)	197 ± 318 vs. 3708 ± 6398				
		DAR (central)	55 ± 96 vs. 1773 ± 3262				
		DAR (right TO)	187 ± 261 vs. 4400 ± 7763				
		Global DTABR	191 ± 469 vs. 1781 ± 2793				
		DTABR (left frontal)	86 ± 158 vs. 1680 ± 2388				
		DTABR (left TO)	130 ± 210 vs. 1884 ± 2828				
		DTABR (central)	39 ± 65 vs. 1132 ± 1957				
		DTABR (right TO)	126 ± 178 vs. 2960 ± 5271				
Psychological							
	Mood	Mood Change	Negative change	CKD stage 5D (hemodialysis)	Edmonton Frail Scale	N/A	21

					(EFS)		
	<b>Mental Health</b>						
	Anxiety	Hospital Anxiety and Depression Scale (HADS)	<u>Women</u> : ↑ in global, psychological, social frailty <u>Men</u> : ↑ in Physical frailty	ESRD, under online-haemodiafiltration (OL-HDF)	N/A	97	<sup>22</sup>
	Depression	Hospital Anxiety and Depression Scale (HADS)	<u>Men</u> ↑ in global, psychological, physical frailty	ESRD, under online-haemodiafiltration (OL-HDF)	N/A	97	<sup>22</sup>
		Incidence (%) (Self-reported Major Depression Inventory)	83 vs. 6	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup>
	<b>Mental Function</b>	Post-KT delirium	9.0% vs. 3.9%	CKD stage 5T	Fried Phenotypes	893	<sup>23</sup>
	<b>Sociological</b>						
	Interaction	Interaction with family	Good				<sup>24</sup>
	<b>Physical activity</b>	Minnesota Leisure Time Activity (LTA)	95 vs. 735 (p<0.001)	CKD stage 5D (hemodialysis)	Fried Phenotypes	68	<sup>25</sup>
		Low Physical Activity Questionnaire (LoPAQ)	280 vs. 798 (p=0.003)				
		Sitting (hours/day)	6.5 vs. 5				

			(p=0.04)				
<b>Nutritional Status</b>	Overall subjective global assessment (SGA) (weight loss, anorexia, subcutaneous fat, muscle mass) (Frail with depression vs. Frail without depression vs. Nonfrail)		5.04 vs. 5.41 vs. 5.75 (p < 0.0001)	CKD stage 5D (peritoneal dialysis)	In-house Chinese questionnaire	178	<sup>15</sup>
			Spearman's rank correlation coefficient r = - 0.44, p < 0.0001	CKD stage 5D (peritoneal dialysis)	Chinese questionnaire	193	<sup>18</sup>
	Malnutrition inflammation score (MIS) (frail with depression vs. frail without depression vs. nonfrail)		9.48 vs. 7.13 vs. 5.12 (p < 0.0001)	CKD stage 5D (peritoneal dialysis)	In-house Chinese questionnaire	178	<sup>15</sup>
			Spearman's rank correlation coefficient r = 0.40, p < 0.0001	CKD stage 5D (peritoneal dialysis)	Chinese questionnaire	193	<sup>18</sup>
<b>Quality of Life</b>	HRQoL						
		SF-36					
		Scores in physical functioning, blood pressure, role physical,	↓	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup>

		and physical component summary domains					
	Kidney Disease Quality of Life (KDQoL)		F vs. NF				
		Physical health	33.7 vs. 40.7	ESRD CKD stage 5D (conventional hemodialysis)	Fried Phenotypes	151	26
		Kidney disease effects	51.6 vs. 66.8				
<b>Independence</b>	Functional Independence Measure (FIM)			Elderly, ≥65y/o	Edmonton Frail Scale (EFS)	137	27
	Spearman's correlation coefficient	Frailty diagnosis with global FIM	-0.703 (p<0.001)				
		Frailty diagnosis with motor FIM	-0.714 (p<0.001)				
		Frailty diagnosis with cognitive FIM	-0.575 (p<0.001)				
		EFS scores with gross FIM	-0.53 (p<0.01)				
<b>Health-care utilization</b>	Hospitalization		90% vs. 53% (p = 0.04)	≥ 65 yo, predialysis, eGFR < 20 mL/min	Groningen frailty indicator (GFI)	65	19
		Cumulative number of inpatient health-care visits	↑	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	14
		Cumulative number of emergency health-care visits					



		Cumulative number of total health-care visits					
		Admissions/year	0.77727 vs. 0.2838	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	<sup>11</sup>

## Complications

	Effect (descriptions)	Risk Difference	CKD Severity	Frailty Assessment	Sample Size	Reference
Biological						
Cardiovascular	Cerebrovascular Accident	OR 1.55 (1.05-2.99)	CKD stage 5D (hemodialysis)	Fried Phenotypes	2275	<sup>4</sup>
	QRS duration	$\beta$ coefficient = -0.29, t = -2.03 (p = 0.048)	CKD stage 5D (chronic hemodialysis)	Edmonton frailty scale (EFS)	41	<sup>28</sup>
		$\beta$ coefficient = -0.27, t = -1.84 (p = 0.05)		FRAIL scale		
	Vascular Access failure	HR 2.63 (1.03-6.71)	ESRD (CKD stage 5D)	Self-reported FRAIL scale	51	<sup>29</sup>
Renal Function Decline	Risk for death or dialysis therapy	2.5 (1.4-4.4)-fold greater	CKD stages 1-4	Fried Phenotypes	336	<sup>7</sup>
Immunological	Mycophenolate mofetil (MMF) dose reduction (MDR)	HR 1.29 (1.01-1.66)	CKD stage 5T	Fried Phenotypes	525	<sup>13</sup>
Cognitive	Modified Mini-Mental State (3MS)	-2.37 to -2.80 (1 year) (p=0.03)	ESRD	Fried Phenotypes	324	<sup>10</sup>
	Declined, 1-4 years post-KT (points/week)	Slope = -0.04 vs. 0.005	CKD stage 5T	Fried physical frailty phenotypes (PFP)	665	<sup>30</sup>
	At 4 year post-KT (points)	-5.5 (87.4 vs.				

		92.9)				
Functional status	Disability	F vs. Prefrail vs. NF				
	Need for activities of daily livings (ADL) assistance	OR 11.32 (5.49-23.32) vs. 1.93 (1.01-3.68) vs. 1.00	CKD stage 5D (hemodialysis)	Fried Phenotypes	742	<sup>31</sup>
Diabetes	Diabetes	OR 1.35 (1.10-1.65)	CKD stage 5D	Fried Phenotypes	2275	<sup>4</sup>
Body composition	Lower lean mass	F/PF vs. NF				
	Whole body (kg)	34.7 vs. 43.1	CKD stage 5D (hemodialysis)	Self- report instrument evaluating five dimensions of frailty (fatigue, resistance, ambulation, illnesses, and weight loss)	44	<sup>32</sup>
	Cephalic/trunk (g)	3059 vs. 3288 (cephalic); 17.4 vs. 22.1 (trunk)				
	Right/Left upper limb (g)	1831 vs. 2493 (right); 1869 vs. 2515 (left)				
	Right/Left lower limb (g)	4920 vs. 6114 (right); 4650 vs. 6349 (left)				
	Bones					

Bone Mineral Density (BMD)	One year follow-up, with frailty	ESRD	FRAIL Scale	43	33
L1	$\beta = -0.4$ , t = -2.18, p=0.04	CKD stage 5D (chronic hemodialysis)			
L4	$\beta = -0.39$ , t = -2.1, p=0.046				
Femur Neck (FN)	$\beta = -0.5$ , t = -2.96, p<0.01				
Total	$\beta = -4$ , t = -3.17, p = 0.004				
	$\beta = -0.53$ , t = -3.27, p < 0.01	ESRD	FRAIL Scale	43	33
		CKD stage 5D (chronic hemodialysis)			
Areas	One year follow-up, with frailty	ESRD	FRAIL Scale	43	33
Average L-spine areas	$\beta = -0.48$ , t = -2.84, p < 0.01	CKD stage 5D (chronic hemodialysis)			
Changes of average L-spine areas	$\beta = -0.5$ , t = -3.02, p<0.01				
Z-score	One year follow-up, with frailty				

Percentage change of L1 Z-score	$\beta = -0.45$ , t = -2.11, p=0.049				
Vertebral Compression Fracture (VCF)	OR 1.8 (p = 0.01)	ESRD CKD stage 5D (chronic hemodialysis)	FRAIL Scale	43	34
Quantitative ultrasound (QUS) parameters of calcaneus					
Speed of sound (SOS)	Standardized $\beta$ (range, p value)	CKD stage 5D (maintenance hemodialysis)	Fried Phenotypes	214	35
Female (Five frailty criteria)	Negative (-0.253 to -0.439, p $\leq$ 0.034)				
Male (All criteria significant except weight loss)	Negative (-0.277 to -0.402, p $\leq$ 0.003)				
Broadband ultrasound attenuation (BUA)	Standardized $\beta$ (range, p value)				
Female (All criteria significant except weakness and weight loss)	Negative (-0.209 to -0.354, p $\leq$ 0.045)				

Male (All criteria significant except weight loss)	Negative (-0.171 to -0.371, $p \leq$ 0.045)				
Stiffness index	Standardized $\beta$ (range, $p$ value)				
Female (All criteria significant except weight loss)	Negative (-0.271 to -0.461, $p \leq$ 0.018)				
Male (Five frailty criteria)	Negative (-0.183 to -0.461, $p \leq$ 0.048)				
Muscles					
Quadriceps muscle area (magnitude of association with PbF vs. 10 years of age)	Multivariable coefficient -30.3 $\text{cm}^2$ ( $p = 0.02$ ) vs. -6.6 $\text{cm}^2$ ( $p =$ 0.0001)	CKD stage 5D (hemodialysis)	Performance-based frailty (PbF)	80	36
Appendicular skeletal muscle mass index (ASMI)	Unadjusted 6.8 vs. 7.7; adjusted $p < 0.05$	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	14
Fat					

	Appendicular fat percentage	FRAIL scale				
	Left/Right lower extremity	t = 2.32; p = 0.03 (left); t = 2.05; p = 0.048 (right)	CKD stage 5D (hemodialysis)	FRAIL scale	44	32
	Left/Right upper extremity	t = 2.66; p = 0.01 (left); t = 3.09; p = <0.01 (right)				
	Appendicular fat percentage	Frail/Prefrail vs. Nonfrail				
	Left/Right lower extremity	t = 2.31; p = 0.03 (left); t = 2.28; p = 0.03 (right)	CKD stage 5D (hemodialysis)	Self- report instrument evaluating five dimensions of frailty (fatigue, resistance, ambulation, illnesses, and weight loss)	44	32
	Right upper extremity	t = 2.35; p = 0.03				
	Sarcopenia	aOR 12.2 (2.27-65.5)	CKD stage 5D (peritoneal dialysis)	Clinical Frailty Scale (CFS)	11912	12
Laboratory data	Serum Albumin Concentrations (g/dL)					
	<3.2 vs. ≥3.9	OR 1.89 (1.30-2.59)	CKD stage 5D	Fried Phenotypes	2275	4
	Hypoalbuminemia	Negative association (p =	CKD stage 5D (maintenance	Frail Scale	46	37

		0.01)	hemodialysis) (ESRD)			
Psychological						
Delirium	Post-KT delirium	OR 2.05 (1.02-4.13)	CKD stage 5T	Fried Phenotypes	893	23
Distress	Distress Thermometer	$\beta = 0.35$ , $t = 3.0$ (95% CL = 0.12-0.58) ( $p = 0.003$ )	CKD stage 5D (hemodialysis)	Canadian frailty score	382	38
Quality of Life						
	HRQoL					
	Fair/Poor HRQOL at follow-up (median 9.4 mo)	aOR 2.79 (1.32-5.90)	ESRD CKD stage 5T	Fried Phenotypes	233	39
	Worsening HRQOL at follow-up (median 9.4 mo)	aRR 2.91 (1.08-7.80)				
	SF-36					
	Hierarchical regression R <sup>2</sup> change (effects of frailty on HRQoL) in Physical Component Summary (PCS)	29% ( $p < 0.001$ )	CKD stage 2-4	Fried Phenotypes	168	40
	Hierarchical regression R <sup>2</sup> change (effects of frailty on HRQoL) in Mental Component Summary (MCS)	21.3% ( $p < 0.001$ )				



Physical components	Simple linear regression coefficient = -1.12 (-1.47 to -0.76) (p < 0.001)	CKD stages 3-5 (predialysis treatment)	Fried Phenotypes	61	41
Mental components	Simple linear regression coefficient = -0.75 (-1.40 to -.016)				
SF-12					
MCS	Effect estimate 0.94 (0.91-0.97) (p<0.01)	CKD stage 5D (peritoneal dialysis, n=129;	The Canadian Study of Health and Aging Clinical Frailty Scale (CFS)	251	42
PCS	Effect estimate 0.88 (0.84-0.91) (p<0.01)	hemodialysis, n=122)			
KDQOL-SF scores within 3 months post-KT	F vs. NF				
At KT	↓	CKD stage 5T	Fried Phenotypes	443	43
Physical HRQoL	-6.31 points (95% CI -8.16 to -4.46)				
Kidney disease-	-6.53 points				

specific HRQoL	(95% CI -9.17 to -3.89)
Post-KT	Greater improvement
Physical HRQoL	1.35 points/month (0.65 to 2.05) vs. 0.34 points/month (-0.17 to 0.85)
Kidney disease-specific HRQoL	3.75 points/month (2.89 to 4.60) vs. 2.41 points/month (1.78 to 3.04)
Constituent domains	Greater improvement
General health	4.93 points/month (3.51 to 6.35) vs. 2.87 points/month (1.82 to 3.92)

Effects of ESRD on daily living	7.10 points/month (5.68 to 8.51) vs. 4.01 points/month (2.99 to 5.03)				
Cognitive function	2.88 points/month (1.80 to 3.96) vs. 1.28 points/month (0.50 to 2.07)				
Social interaction	1.18 points/month (-0.06 to 2.43) vs. -0.57 points/month (-1.47 to 0.33)				
Illness Intrusiveness Rating Scale	Effect estimate 1.14 (1.09-1.20)	CKD stage 5D (peritoneal dialysis, n=129;	The Canadian Study of Health and Aging Clinical Frailty Scale (CFS)	251	42
Barthel Index	Effect estimate 0.89 (0.86-.093)	hemodialysis, n=122)			
Symptom score	Effect estimate 1.23 (1.13-1.34)				

	Hospital Anxiety and Depression Scale	Effect estimate 1.21 (1.11-1.31)				
	Falls	HR 2.1 (1.21-3.92)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	<sup>1</sup>
		OR 2.39 (1.22-4.71)	CKD stage 5D (maintenance hemodialysis)	Fried frailty index	762	<sup>44</sup>
	Higher numbers of falls	HR 3.09 (1.38-6.90)	CKD stage 5D (hemodialysis)		95	<sup>45</sup>
	Time to first fall or fracture requiring medical attention	HR 1.60 (1.16-2.20)	CKD stage 5D (maintenance hemodialysis)	Modified Fried Phenotypes by Bao Y <sup>46</sup> .	1646	<sup>47</sup>
Graft Loss	Death-censored graft loss					
	F vs. NF (in patients with depressive symptoms)	aHR 6.20 (1.67, 22.95) vs. 3.16 (0.90, 11.04)	CKD stage 5T	Fried Phenotypes	773	<sup>48</sup>
Health-care utilization	Hospitalization/Death	HR 1.56 (1.36-1.79)	CKD stage 5D	Fried Phenotypes	2275	<sup>4</sup>
	Hospitalization	HR 2.06 (1.18-3.58)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	<sup>1</sup>
		aHR 1.80 (1.4-2.3)	CKD stage 5D (maintenance hemodialysis &	Adopted	1658	<sup>49</sup>

		peritoneal dialysis)			
	Relative risk = 1.43 (1.00-2.03)	CKD stage 5D (hemodialysis)	Fried Phenotypes	146	50
Number of hospitalizations for all causes	beta = 0.29 (p < 0.0001)	CKD stage 5D (peritoneal dialysis)	Chinese questionnaire	193	18
Number of hospitalizations related to cardiovascular events	beta = 0.37 (p < 0.0001)				
Non-vascular access- related hospitalizations	aHR 1.98 (1.41- 1.87) (內文應該 寫錯，無勘誤)	CKD stage 5D	Fried Phenotypes	2275	4
Time to first hospitalization	HR 1.26 (1.09- 1.45)	CKD stage 5D (maintenance dialysis)	Earlier modification of Fried Phenotypes by Johansen et al <sup>4</sup> , but without weight loss.	1576	46
Early Hospital Readmission (EHR)	aRR 1.61 (1.81- 2.19) (p=0.002)	CKD stage 5T	Fried Phenotypes	383	51
Duration of hospitalization					
Hospital stay (days per year of follow up) (frail with depression vs. frail without depression vs. nonfrail)	26.62 (IQR 10.65-61.18) vs. 14.05 (IQR 3.57- 37.27) vs. 8.04 (IQR 0.91-19.42) (p<0.0001)	CKD stage 5D (peritoneal dialysis)	In-house Chinese questionnaire	178	15

	Total length of hospital stay	beta = 0.34 (p < 0.0001)	CKD stage 5D (peritoneal dialysis)	Chinese questionnaire	193	18
	Longer Length of Stay (LOS)					
	with delayed graft function (DGF), LOS	Relative Risk 1.15 (1.03-1.29)	CKD stage 5T	Fried Phenotypes	589	52
	With DGF, LOS ≥2 weeks	OR 1.57 (1.06-2.33)				
	≥2 weeks		CKD stage 5 to 5T	Fried Phenotypes	569	53
	Change in 3 categories (more frail)	OR 2.02 (1.20-3.40)				
	Change in frailty scores (more frail)	OR 1.92 (1.13-3.25)				
	With depressive symptoms (aRR difference between F and NF)	aRR 1.88 (1.70-2.08) vs. 1.38 (1.27-1.52)	CKD stage 5T	Fried Phenotypes	773	48
	CES-D score (10-point increase) (aRR increase between F and NF)	aRR 1.23 (1.16-1.31) vs. 1.17 (1.08-1.27)				
Mortality	Mortality	2.17 fold	CKD stage 5T	Fried Phenotypes	537	54
		HR 1.57 (1.25-1.97)	CKD stage 5D (maintenance dialysis)	Earlier modification of Fried Phenotypes by Johansen et al <sup>4</sup> , but without weight loss.	1576	46
		HR 2.24 (1.60-	CKD stage 5D	Fried Phenotypes	2275	4

	3.15)				
	HR 1.22 (1.04-1.43)	CKD stage 5D (incident chronic dialysis)	CFS	390	55
	HR 4.28 (1.22-14.98)	Predialysis (eGFR ≤ 25 mL)	PRISMA questionnaire & Timed up and Go test	104	56
	aHR 9.83 (1.80-53.7)	CKD stage 5D (peritoneal dialysis)	CFS	119	12
	HR 2.60 (1.04-6.49)	CKD stage 5D (hemodialysis)	Fried Phenotypes	146	50
	HR 2.37 (1.11-5.02)	CKD stage 5D (maintenance hemodialysis & peritoneal dialysis)	Adopted	1658	49
	20.45% vs. 12.36% (p<0.005)	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	11
Performance-based frailty	HR 2.16 (1.41-3.29)	CKD stage 5D (hemodialysis)	Fried Phenotypes & a definition that substitutes self-reported measures available on the Medical Outcomes Study 36-Item Short Form (SF-36) for the physical performance and	771	57
Self-reported function-based frailty	HR 1.93 (1.24-3.00)				
Patients who met both performance- and self-reported function-based	HR 2.46 (1.51-4.01)				

frailty			exhaustion criteria.		
F vs. NF (in patients with depressive symptoms)	aHR 2.62 (1.03, 6.70) vs. 1.92 (0.68, 5.38)	CKD stage 5T	Fried Phenotypes	773	48
At 24-month follow up, frail with depression vs. frail without depression vs. nonfrail	62.5% vs. 71.4% vs 86.6% (p=0.001)	CKD stage 5D (peritoneal dialysis)	In-house Chinese questionnaire	178	15
Prediction ability of comorbidities in F vs. NF	HR 0.75 (0.44-1.29) vs. 1.66 (1.17-2.35)	CKD stage 5T (KT candidates, on waitlist)	Fried Phenotypes	2086	58
Out of 10 deceased within 1 year of initiation (percentage of F vs. NF)	30% vs. 9%	≥ 65 yo, predialysis, eGFR < 20 mL/min	Groningen frailty indicator (GFI)	65	19
Risk for death or dialysis therapy	2.5 (1.4-4.4)-fold greater	CKD stages 1-4	Fried Phenotypes	336	7
All-cause mortality					
Adjusted	HR 1.66 (1.03-2.67)	CKD stage 5D (incident chronic dialysis)	Fried Phenotypes	370	59
Among BMI ≥30 kg/m <sup>2</sup>	HR 3.77 (1.10-12.92)				
Above median Waist-Hip Ratio (WHR)	HR 2.38 (1.17-4.82)				
Anti-neutrophil cytoplasmic	HR 2.43 (1.48-	CKD stage 5D to 5T	Inability to walk without	425	60



	antibody (ANCA)-associated vasculitides (AAV) patients' mortality	3.99)	(RRT [hemodialysis, peritoneal dialysis, transplantation])	help		
	Mortality of patients with monoclonal gammopathy and ESRD caused by myeloma cast nephropathy (MCN), immunoglobulin light chain amyloidosis (ALA), or light-chain deposition disease (LCDD)	HR, 1.93 (1.58-2.36)	CKD stage 5D to 5T (RRT [hemodialysis, peritoneal dialysis, transplantation])	Inability to walk without help	1462	<sup>61</sup>
	Post-KT mortality					
	Change in 3 categories (more frail)	HR 2.27 (1.11-4.65)	CKD stage 5 to 5T	Fried Phenotypes	569	<sup>53</sup>
	Change in frailty scores (more frail)	HR 2.36 (1.12-4.99)				
Composite	Composite outcomes of all-cause death or cardiovascular hospitalization	HR 23.58 (1.61-346.03)	CKD stage 5D (hemodialysis)	Multidimensional frailty score based on comprehensive geriatric assessment (CGA) protocol	46	<sup>62</sup>
	30-day postoperative (KT) complications according to Comprehensive Complication	$\beta=13.31$ , 95% CI 5.72-20.89 (p = 0.0007)	CKD stage 5T	Groningen Frailty Indicator	150	<sup>63</sup>

---

Index (CCI)

---

## Bibliography

1. YadlaM, JohnJ, MummadiM. A study of clinical assessment of frailty in patients on maintenance hemodialysis supported by cashless government scheme. *Saudi J Kidney Dis Transplant*. 2017. doi:10.4103/1319-2442.198102
2. JohansenKL, DalrympleLS, DelgadoC, et al. Factors Associated with Frailty and Its Trajectory among Patients on Hemodialysis. *Clin J Am Soc Nephrol*. 2017;12(7):1100-1108. doi:10.2215/CJN.12131116
3. MansurHN. Fragilidade na doença renal crônica: prevalência e fatores associados. 2012.
4. JohansenKL, ChertowGM, JinC, KutnerNG. Significance of frailty among dialysis patients. *J Am Soc Nephrol*. 2007;18(11):2960-2967.
5. Wilhelm-LeenER, HallYN, MKT, ChertowGM. Frailty and chronic kidney disease: the Third National Health and Nutrition Evaluation Survey. *Am J Med*. 2009;122(7):664-71 e2. doi:10.1016/j.amjmed.2009.01.026
6. ShlipakMG, Stehman-BreenC, FriedLF, et al. The Presence of Frailty in Elderly Persons with Chronic Renal Insufficiency. *Am J Kidney Dis*. 2004;43(5):861-867. doi:10.1053/j.ajkd.2003.12.049
7. RoshanravanB, KhatriM, Robinson-CohenC, et al. A prospective study of frailty in nephrology-referred patients with CKD. *Am J Kidney Dis*. 2012;60(6):912-921. doi:10.1053/j.ajkd.2012.05.017
8. ChiangJM, KaysenGA, SegalM, ChertowGM, DelgadoC, JohansenKL. Low testosterone is associated with frailty, muscle wasting and physical dysfunction among men receiving hemodialysis: a longitudinal analysis. *Nephrol Dial Transplant*. 2018. doi:10.1093/ndt/gfy252
9. DemirciogluD. The Association of Vitamin D Levels and the Frailty Phenotype Among Non-geriatric Dialysis Patients: A Cross-sectional Study. *Clinics*. 2018. doi:10.6061/clinics/2018/e116
10. McAdams-DemarcoMA, TanJ, SalterML, et al. Frailty and cognitive function in incident hemodialysis patients. *Clin J Am Soc Nephrol*. 2015;10(12):2181-2189. doi:10.2215/CJN.01960215
11. Bancul, GrateroIF, BonalJ, et al. Frail Patient in Hemodialysis: A New Challenge in Nephrology-Incidence in Our Area, Barcelones Nord and Maresme. *J Aging Res*. 2017;2017:7624139. doi:10.1155/2017/7624139
12. KamijoY, KandaE, IshibashiY, YoshidaM. Sarcopenia and Frailty in PD: Impact on Mortality, Malnutrition, and Inflammation. *Perit*

- Dial Int.* 2018;38(6):447-454. doi:10.3747/pdi.2017.00271
13. McAdams-DemarcoMA, LawA, TanJ, et al. Frailty, mycophenolate reduction, and graft loss in kidney transplant recipients. *Transplantation.* 2015;99(4):805-810. doi:10.1097/TP.0000000000000444
  14. Adame PerezSI, SeniorPA, FieldCJ, JindalK, MagerDR. Frailty, Health-Related Quality of Life, Cognition, Depression, Vitamin D and Health-Care Utilization in an Ambulatory Adult Population with Type 1 or Type 2 Diabetes Mellitus and Chronic Kidney Disease: A Cross-Sectional Analysis. *Can J Diabetes.* June2018. doi:10.1016/j.jcjd.2018.06.001
  15. SzetoC-C, ChanGC-K, NgJK-C, et al. Depression and Physical Frailty Have Additive Effect on the Nutritional Status and Clinical Outcome of Chinese Peritoneal Dialysis. *Kidney Blood Press Res.* 2018;43(3):914-923. doi:10.1159/000490470
  16. ChaoC-T, LaiH-J, TsaiH-B, YangS-Y, HuangJ-W. Frail phenotype is associated with distinct quantitative electroencephalographic findings among end-stage renal disease patients: an observational study. *BMC Geriatr.* 2017;17(1):277. doi:10.1186/s12877-017-0673-3
  17. deSousa MeiraA, Aparecido BatistaM, dePina PereiraRM, Partezani RodriguesRA, Silva FhonJR, KusumotaL. Frailty in elderly patients with chronic kidney disease under conservative treatment. *Rev da Rede Enferm do Nord.* 2016;17(3):386-392. doi:10.15253/2175-6783.2016000300012
  18. NgJK-C, KwanBC-H, ChowK-M, et al. Frailty in Chinese Peritoneal Dialysis Patients: Prevalence and Prognostic Significance. *Kidney Blood Press Res.* 2016;41(6):736-745. doi:10.1159/000450563
  19. MeulendijksFG, HamakerME, BoereboomFTJ, KalfA, VögtlanderNPJ, vanMunsterBC. Groningen frailty indicator in older patients with end-stage renal disease. *Ren Fail.* 2015;37(9):1419-1424.
  20. MargiottaE, Caldirolil, VettorettiS, et al. SuO004GUT MICROBIOTA COMPOSITION AND FRAILITY IN ELDERLY PATIENTS WITH CHRONIC KIDNEY DISEASE. *Nephrol Dial Transplant.* 2018;33(suppl\_1):i618-i618. doi:10.1093/ndt/gfy104.SuO004
  21. DeSouza OrlandiF, GesualdoGD. Assessment of the frailty level of elderly people with chronic kidney disease undergoing hemodialysis. *ACTA Paul Enferm.* 2014;27(1):29-34. doi:10.1590/1982-01942014000007
  22. SalesC, TavaresR, AmadoL, et al. SP651ANXIETY AND DEPRESSION IN END STAGE RENAL DISEASE PATIENTS AND ITS ASSOCIATION WITH CLINICAL AND LABORATORIAL DATA. *Nephrol Dial Transplant.* 2017;32(suppl\_3):iii355-iii355. doi:10.1093/ndt/gfx154.SP651

23. HaugenCE, MountfordA, WarsameF, et al. Incidence, Risk Factors, and Sequelae of Post-kidney Transplant Delirium. *J Am Soc Nephrol*. 2018;29(6):1752-1759. doi:10.1681/ASN.2018010064
24. MoffattH, MoorhouseP, MalleryL, LandryD, TennankoreK. Using the Frailty Assessment for Care Planning Tool (FACT) to screen elderly chronic kidney disease patients for frailty: the nurse experience. *Clin Interv Aging*. 2018;13:843.
25. JohansenKL, PainterP, DelgadoC, DoyleJ. Characterization of physical activity and sitting time among patients on hemodialysis using a new physical activity instrument. *J Ren Nutr*. 2015;25(1):25-30. doi:10.1053/j.jrn.2014.06.012
26. NooriN, Sharma ParpiaA, LakhaniR, JanesS, GoldsteinMB. Frailty and the Quality of Life in Hemodialysis Patients: The Importance of Waist Circumference. *J Ren Nutr*. 2018;28(2):101-109. doi:10.1053/j.jrn.2017.07.007
27. Fabrício-WehbeSCC, SchiavetoFV, VendrusculoTRP, HaasVJ, DantasRAS, RodriguesRAP. Cross-cultural adaptation and validity of the “Edmonton Frail Scale - EFS” in a Brazilian elderly sample. *Rev Lat Am Enfermagem*. 2009;17(6):1043-1049. doi:10.1590/S0104-11692009000600018
28. ChaoC-T, HuangJ-W. Frailty severity is significantly associated with electrocardiographic QRS duration in chronic dialysis patients. *PeerJ*. 2015;3:e1354. doi:10.7717/peerj.1354
29. ChaoC-T, ChiangC-K, HuangJ-W, HungK-Y. Self-reported frailty among end-stage renal disease patients: A potential predictor of dialysis access outcomes. *Nephrology (Carlton)*. 2017;22(4):333-334. doi:10.1111/nep.12961
30. ChuNM, GrossAL, ShafferAA, et al. Frailty and Changes in Cognitive Function after Kidney Transplantation. *J Am Soc Nephrol*. 2019;30(2):336-345. doi:10.1681/ASN.2018070726
31. KutnerNG, ZhangR, AllmanRM, BowlingCB. Correlates of ADL difficulty in a large hemodialysis cohort. *Hemodial Int*. 2014;18(1):70-77. doi:10.1111/hdi.12098
32. ChaoC-T, ChanD-C, HuangJ-W. Frail Phenotype Might Be Associated With Higher Appendicular but Not Truncal Fat Among End-Stage Renal Disease Patients. *J Pain Symptom Manage*. 2017;53(4):e1-e4. doi:10.1016/j.jpainsymman.2017.01.004
33. ChaoC-T, HuangJ-W, ChanD-C. Frail phenotype might herald bone health worsening among end-stage renal disease patients. *PeerJ*. 2017;5:e3542. doi:10.7717/peerj.3542
34. ChaoC-T, ChiangC-K, HuangJ-W, ChanD-C. Effect of Frail Phenotype on Bone Mass and Vertebral Compression Fracture in Individuals Undergoing Dialysis. *J Am Geriatr Soc*. 2016;64(9):e19-21. doi:10.1111/jgs.14296

35. YonekiK, KitagawaJ, HoshiK, et al. Association between frailty and bone loss in patients undergoing maintenance hemodialysis. *J Bone Miner Metab*. January 2018. doi:10.1007/s00774-017-0898-4
36. DelgadoC, DoyleJW, JohansenKL. Association of frailty with body composition among patients on hemodialysis. *J Ren Nutr*. 2013;23(5):356-362. doi:10.1053/j.jrn.2013.02.010
37. ChaoC-T, HsuY-H, ChangP-Y, et al. Simple self-report FRAIL scale might be more closely associated with dialysis complications than other frailty screening instruments in rural chronic dialysis patients. *Nephrology (Carlton)*. 2015;20(5):321-328. doi:10.1111/nep.12401
38. CamilleriS, ChongS, TangvoraphonkchaiK, YoowannakulS, DavenportA. Effect of Self-Reported Distress Thermometer Score on the Maximal Handgrip and Pinch Strength Measurements in Hemodialysis Patients. *Nutr Clin Pract*. 2017;32(5):682-686. doi:10.1177/0884533617697936
39. McAdams-DeMarcoMA, YingH, OlorundareI, et al. Frailty and Health-Related Quality of Life in End Stage Renal Disease Patients of All Ages. *J frailty aging*. 2016;5(3):174-179.
40. LeeSJ, SonH, ShinSK. Influence of frailty on health-related quality of life in pre-dialysis patients with chronic kidney disease in Korea: a cross-sectional study. *Health Qual Life Outcomes*. 2015;13(1):70.
41. MansurHN, ColugnatiFA, GrincenkovFR, BastosMG. Frailty and quality of life: a cross-sectional study of Brazilian patients with pre-dialysis chronic kidney disease. *Heal Qual Life Outcomes*. 2014;12:27. doi:10.1186/1477-7525-12-27
42. IyasereOU, BrownEA, JohanssonL, et al. Quality of Life and Physical Function in Older Patients on Dialysis: A Comparison of Assisted Peritoneal Dialysis with Hemodialysis. *Clin J Am Soc Nephrol*. 2016;11(3):423-430. doi:10.2215/CJN.01050115
43. McAdams-DeMarcoMA, OlorundareIO, YingH, et al. Frailty and Postkidney Transplant Health-Related Quality of Life. *Transplantation*. 2018;102(2):291-299. doi:10.1097/TP.0000000000001943
44. KutnerNG, ZhangR, HuangY, WasseH. Falls among hemodialysis patients: potential opportunities for prevention? 2014. doi:10.1093/ckj/sfu034
45. McAdams-DeMarcoMA, SureshS, LawA, et al. Frailty and falls among adult patients undergoing chronic hemodialysis: a prospective cohort study. *BMC Nephrol*. 2013;14:224. doi:10.1186/1471-2369-14-224
46. BaoY, DalrympleL, ChertowGM, KaysenGA, JohansenKL. Frailty, dialysis initiation, and mortality in end-stage renal disease. *Arch*

- Intern Med.* 2012;172(14):1071-1077. doi:10.1001/archinternmed.2012.3020
47. DelgadoC, ShiehS, GrimesB, et al. Association of Self-Reported Frailty with Falls and Fractures among Patients New to Dialysis. *Am J Nephrol.* 2015;42(2):134-140. doi:10.1159/000439000
  48. KonelJM, WarsameF, YingH, et al. Depressive symptoms, frailty, and adverse outcomes among kidney transplant recipients. *Clin Transplant.* 2018;32(10):e13391. doi:10.1111/ctr.13391
  49. LeeS-Y, YangDH, HwangE, et al. The prevalence, association, and clinical outcomes of frailty in maintenance dialysis patients. *J Ren Nutr.* 2017;27(2):106-112.
  50. McAdams-DeMarcoMA, LawA, SalterML, et al. Frailty as a novel predictor of mortality and hospitalization in individuals of all ages undergoing hemodialysis. *J Am Geriatr Soc.* 2013;61(6):896-901. doi:10.1111/jgs.12266
  51. McAdams-DeMarcoMA, LawA, SalterML, et al. Frailty and Early Hospital Readmission After Kidney Transplantation. *Am J Transplant.* 2013;13(8):2091-2095. doi:10.1111/ajt.12300
  52. McAdams-DeMarcoMA, KingEA, LuoX, et al. Frailty, Length of Stay, and Mortality in Kidney Transplant Recipients: A National Registry and Prospective Cohort Study. *Ann Surg.* 2017;266(6):1084-1090. doi:10.1097/SLA.0000000000002025
  53. ChuNM, DengA, YingH, et al. Dynamic Frailty Before Kidney Transplantation-Time of Measurement Matters. *Transplantation.* February2019. doi:10.1097/TP.0000000000002563
  54. McAdams-DeMarcoMA, LawA, KingE, et al. Frailty and mortality in kidney transplant recipients. *Am J Transplant.* 2015;15(1):149-154. doi:10.1111/ajt.12992
  55. AlfaadhelTA, SorokaSD, KiberdBA, LandryD, MoorhouseP, TennankoreKK. Frailty and mortality in dialysis: evaluation of a clinical frailty scale. *Clin J Am Soc Nephrol.* 2015;10(5):832-840. doi:10.2215/CJN.07760814
  56. AliH, AbdelazizT, AbdelaalF, BaharaniJ. Assessment of prevalence and clinical outcome of frailty in an elderly predialysis cohort using simple tools. *Saudi J Kidney Dis Transpl.* 2018;29(1):63-70. doi:10.4103/1319-2442.225175
  57. JohansenKL, DalrympleLS, GliddenD, et al. Association of Performance-Based and Self-Reported Function-Based Definitions of Frailty with Mortality among Patients Receiving Hemodialysis. *Clin J Am Soc Nephrol.* 2016;11(4):626-632. doi:10.2215/CJN.03710415
  58. Pérez FernándezM, Martínez MiguelP, YingH, et al. Comorbidity, Frailty, and Waitlist Mortality among Kidney Transplant

- Candidates of All Ages. *Am J Nephrol*. 2019;49(2):103-110. doi:10.1159/000496061
59. FitzpatrickJ, SozioSM, JaarBG, et al. Frailty, body composition and the risk of mortality in incident hemodialysis patients: the Predictors of Arrhythmic and Cardiovascular Risk in End Stage Renal Disease study. *Nephrol Dial Transplant*. 2019;34(2):346-354. doi:10.1093/ndt/gfy124
60. RomeuM, CouchoudC, DelarozierJC, et al. Survival of patients with ANCA-associated vasculitis on chronic dialysis: data from the French REIN registry from 2002 to 2011. *QJM*. 2014;107(7):545-555. doi:10.1093/qjmed/hcu043
61. DecourtA, GondouinB, DelarozierJC, et al. Trends in Survival and Renal Recovery in Patients with Multiple Myeloma or Light-Chain Amyloidosis on Chronic Dialysis. *Clin J Am Soc Nephrol*. 2016;11(3):431-441. doi:10.2215/CJN.06290615
62. LeeSW, LeeA, YuMY, et al. Is Frailty a Modifiable Risk Factor of Future Adverse Outcomes in Elderly Patients with Incident End-Stage Renal Disease? *J Korean Med Sci*. 2017;32(11):1800-1806. doi:10.3346/jkms.2017.32.11.1800
63. SchopmeyerL, ElMoumniM, Nieuwenhuijs-MoekeGJ, BergerSP, BakkerSJM, PolRA. Frailty has a significant influence on postoperative complications after kidney transplantation-a prospective study on short-term outcomes. *Transpl Int*. 2019;32(1):66-74. doi:10.1111/tri.13330