

# Aortic pulse wave velocity in haemodialysis patients is associated with the prescription of active vitamin D analogues

Evangelina Charitaki · Andrew Davenport

Received: 18 September 2013 / Accepted: 3 November 2013  
© Italian Society of Nephrology 2014

## Abstract

**Background** Cardiovascular disease remains the most common cause of death for haemodialysis patients. In addition to traditional cardiovascular risk factors, haemodialysis patients have additional risk factors, including vascular calcification. Pulse wave velocity (PWV) is a measurement of arterial stiffness, and we wished to determine whether PWV is affected by different factors in haemodialysis patients compared to the general population. **Methods** Aortic PWV was measured in 303 adult patients attending for routine outpatient dialysis.

**Results** 303 patients, 63.4 % male, mean age  $68.5 \pm 15.8$  years, 47.5 % diabetic with a body mass index of  $25.8 \pm 5.3 \text{ kg/m}^2$ , were studied. Systolic blood pressure (SBP) was  $148.7 \pm 28.6 \text{ mmHg}$  and diastolic  $80.4 \pm 15.3 \text{ mmHg}$ . Aortic PWV was  $9.73 \pm 2.08 \text{ m/s}$ , and was correlated with SBP ( $\beta$  0.015,  $F$  5.29,  $p = 0.023$ ), log serum parathyroid hormone (PTH) ( $\beta$  1.58,  $F$  13.85,  $p < 0.001$ ) and prescription of alfacalcidol ( $\beta$  -1.11,  $F$  6.81,  $p = 0.010$ ). 197 patients had corresponding ECHO cardiograms, and in this cohort PWV was associated with SBP ( $\beta$  0.017,  $F$  7.49,  $p = 0.006$ ), log serum parathyroid hormone ( $\beta$  0.85,  $F$  5.99,  $p < 0.015$ ) and prescription of alfacalcidol ( $\beta$  -0.8,  $F$  4.18,  $p = 0.042$ ), left ventricular mass index (LVMI) ( $\beta$  0.01,  $F$  11.4,  $p = 0.001$ ), and log serum triglycerides ( $\beta$  1.43,  $F$  4.79,  $p = 0.03$ ).

**Conclusions** We found that PWV, a measurement of arterial stiffness, was associated with both traditional cardiovascular risk factors, including SBP and LVMI, but also non-traditional risk factors such as prescription of active vitamin D analogues, suggesting a potential link between vascular calcification and arterial stiffness in haemodialysis patients.

**Keywords** Hypertension · Haemodialysis · Parathyroid hormone · Vitamin D · Pulse wave velocity

## Background

The mortality of haemodialysis patients remains high, with survival rates similar to those of some solid organ malignancies. Cardiovascular disease continues to be the major cause of this excess mortality. Whereas atheromatous coronary artery disease with cholesterol rich plaque formation is the predominant cardiovascular risk factor for the general population, haemodialysis patients typically develop arteriosclerosis with medial vascular calcification and increased risk of death due to heart failure, stroke and cardiac arrhythmias [1, 2].

Arterial stiffness is one of the first detectable manifestations of arteriosclerosis. Pulse wave velocity (PWV) is the most validated non-invasive method of assessing arterial stiffness, and as such is considered the gold standard measurement [3, 4]. Aortic stiffness is recognised as an important risk factor for cardiovascular mortality in both the general population and also patients with chronic kidney disease [5]. In order to determine if there were different factors associated with aortic stiffness in patients on haemodialysis compared to the general population, we

E. Charitaki · A. Davenport (✉)  
UCL Centre for Nephrology, Royal Free Hospital, University  
College London Medical School, Rowland Hill Street,  
London NW3 2PF, UK  
e-mail: andrewdavenport@nhs.net

E. Charitaki  
e-mail: evacharitaki@hotmail.com

reviewed aortic PWV measurements in a cohort of established haemodialysis outpatients.

## Patients and methods

The records of 303 adult patients with chronic kidney disease stage 5 who were undergoing outpatient haemodialysis treatment thrice weekly and had reliable pulse wave velocity (PWV) measurements taken before the dialysis session were reviewed. PWV measurement was attempted in all outpatient haemodialysis patients under the care of the Royal Free Hospital, but it was not possible to have reliable PVW measurements in patients with atrial fibrillation, other cardiac arrhythmias, those patients with fistulae in both arms, and those in whom no recordable upper arm blood pressure recording was possible. Aortic-brachial pulse wave velocity was measured using the Tensio Clinic Ateriograph which has been validated against direct invasive measurements [6], and also been shown to be the most reproducible of the currently available devices for measuring PWV [7]. The distance between jugular notch and symphysis pubis was measured with a specially designed measuring device, and after patients had rested pulse wave velocity measurements were made in the recumbent position in the non-fistula arm. The upper arm circumference was measured and the appropriately sized blood pressure arm cuff chosen. Aortic augmentation index (AoAXi), brachial artery augmentation index (BrAXi), and diastolic relaxation index (DRA) were corrected for heart rate.

Patients were dialysed thrice weekly, targeted to achieve an on-line clearance of  $\geq 1.4$  (Fresenius 4008/5008, Fresenius, Bad Homburg, Germany; Dialog+, B Braun Medical Inc, Melsungen, Germany), mean session time  $3.96 \pm 0.38$  h, using polysulfone dialyzers (Nipro, Osaka, Japan) [8] with ultrapure quality dialysate, and anticoagulated with tinzaparin (Leo Laboratories, Market Harborough, UK), median dose 2,500 IU [9]. Regarding the vascular access, 86 (28.4 %) patients dialysed using central venous catheters [10], 205 (67.7 %) by arteriovenous fistulae and 12 (3.9 %) by arteriovenous grafts. Dialysate temperature was set at 35–35.5 °C, and dialysate sodium regularly checked by flame photometry (Flame photometer 943, Instrumentation Laboratory, Warrington, UK) with appropriate aqueous standards, to ensure quality control [11]. Mean dialysis session time was  $3.96 \pm 0.38$  h, using a mean dialysate sodium of  $137.3 \pm 1.2$  mmol/l, potassium  $1.9 \pm 0.5$  mmol/l, and calcium  $1.3 \pm 0.2$  mmol/l, with a set dialysate bicarbonate of 32 mmol/l, 3 mmol/l of acetate and 0.5 mmol/l of magnesium.

Standard biochemical investigations were measured using an automated multichannel analyser, with albumin

measured by the bromocresol green method (Roche Integra, Roche diagnostics, Lewes, UK), and C reactive protein used the same assay as the UK National Amyloid Centre [12]. Standard 2 dimensional echocardiograms were performed on a non-dialysis day (Philips IE33, Philips Medical Systems, Eindhoven, The Netherlands) [13] in a subset of 197 patients and 137 (45.2 %) patients had attended for cardiac stress testing using adenosine triphosphate myocardial perfusion scintigraphy on a non-dialysis day [14]. Echocardiograms and isotope heart stress tests were performed as part of patient assessment for transplantation and introduction of the Kidney Disease: Improving Global Outcomes (KDIGO) clinical practice guidelines [15].

This retrospective audit complied with the UK National Health Service (NHS) guidelines for audit and clinical service development. Measurement of pulse wave velocity was introduced into clinical practice in 2011 and measured as part of an audit of blood pressure control in haemodialysis patients.

## Statistical analysis

Results are expressed as mean  $\pm$  standard deviation, median and interquartile range, or percentage. Simple correlation analysis was used to determine associates of PVW, with non-parametric variables log transformed when entered into a step back multiple regression model if  $p < 0.1$ , and retained if significant and confidence limits did not cross the line of unity, unless variables improved the model fit. Statistical analysis was performed using Graph Pad Prism (version 6.0, Graph Pad, San Diego, CA, USA) and SPSS (version 17, University Chicago, USA). Results are expressed as mean and standard deviation or, median and interquartile range, unless otherwise stated. Statistical significance was taken at or below the 5 % level.

## Results

Data was available on 303 patients (male 63.4 %, mean age  $68.5 \pm 15.8$  years, weight  $71.4 \pm 16.7$  kg, body mass index  $25.8 \pm 5.3$  kg/m<sup>2</sup>, with a median dialysis vintage of 32.5 (14–60) months). 47.5 % of the patients were coded as being diabetic, with either a fasting plasma glucose  $\geq 7.0$  mmol/l (126 mg/dl) or 2-h plasma glucose  $\geq 11.1$  mmol/l (200 mg/dl) being recorded in their medical records. 79.5 % had a past medical history of hypertension, 29.0 % ischaemic heart disease, 19.8 % peripheral vascular disease, 14.9 % cerebrovascular disease and 5.3 % had undergone previous parathyroid surgery. The mean systolic blood pressure (SBP) was  $148.7 \pm 28.6$  mmHg, diastolic blood pressure (DBP)  $80.4 \pm 15.3$  mmHg, and pulse pressure (PP)  $68.2 \pm 20.3$  mmHg. As for antihypertensive

medications, 27.4 % were prescribed angiotensin converting enzyme inhibitors (ACEIs) or angiotensin receptor blockers (ARBs), 26.4 % beta-blockers and 21.5 % calcium channel blockers. In addition, 60.1 % were prescribed aspirin, 13.2 % clopidogrel and 2.3 % were anticoagulated with coumarins.

Mean haemoglobin was  $111.7 \pm 14.5$  g/l, with a serum albumin of  $39.6 \pm 4.1$  g/l, median C reactive protein 4 (2–13) mg/l, calcium  $2.27 \pm 0.17$  mmol/l, corrected calcium  $2.44 \pm 0.15$  mmol/l, phosphate  $1.48 \pm 0.47$  mmol/l, alkaline phosphatase 97 (72–138) IU/l, CRP mg/l and parathyroid hormone (PTH) 27.0 (13.9–53.4) pmol/l. The median dose of alfacalcidol prescribed was 2 (0.75–4.5) µg/week. 53.1 % of patients were prescribed calcium containing phosphate binders [median dose 1,000 (0–1,500) g/day], 16.2 % sevelamer, 13.5 % lanthanum carbonate and 8.5 % cinacalcet [all median dose 0 (0–0)]. Vitamin D (25 hydroxy vitamin D3) levels were measured and 181 (59.7 %) patients with reduced concentrations were prescribed 20,000 IU/week of cholecalciferol.

Serum total cholesterol (TChol) was  $3.98 \pm 1.1$  mmol/l, high density lipoprotein (HDL)  $1.27 \pm 0.48$  mmol/l, low density lipoprotein (LDL)  $2.83 \pm 0.9$  mmol/l and triglycerides  $1.3$  (0.9–1.9) mmol/l, with 58.4 % prescribed HMG CoA 3 reductase inhibitors (statins).

Aortic PWV was  $9.73 \pm 2.08$  m/s, with a heart rate of  $73.3 \pm 13.6$  beats/min, giving a corrected aortic PWV  $9.67 \pm 2.99$  (for a heart rate of 70). Aortic systolic blood pressure was  $149.2 \pm 46.8$  mmHg, and augmentation indices corrected for heart rate were  $3.77 \pm 31.9$  and  $39.1 \pm 20.7$  % for the brachial and aortic arteries respectively, with a diastolic relaxation area of  $40.2 \pm 24.6$  %.

Left ventricular (LV) mass was calculated in 197 patients and corrected for body surface area, using standard equations [15, 16], with a mean LV mass of  $265.7 \pm 93.6$  g, and LVmass index of  $147.5 \pm 48.6$  g/cm<sup>2</sup>, with a relative wall thickness of  $0.52 \pm 0.15$ . No patient was reported to have severe valvular calcification, but minor degrees of valvular calcification were not reported. 137 (45.2 %) patients had attended for cardiac stress testing using adenosine triphosphate myocardial perfusion scintigraphy, and 39.4 % of these tests were reported to show reversible cardiac ischaemia. Dialysis adequacy was targeted to achieve a minimum on-line Kt/V of >1.4 [17].

Simple correlation analysis showed that there was no correlation between pulse wave velocity and haemoglobin, serum albumin or C reactive protein, although a number of variables were statistically associated with PWV (Table 1). Nonparametric variables were log transformed as appropriate and a linear regression model was created for all patients, using all variables with a simple correlation of  $p < 0.01$ . A stepwise regression analysis was then performed with elimination of variables that were not

**Table 1** Simple Pearson correlation analysis for pulse wave velocity

Variable	Pearson r	p
Left ventricular mass	0.236	0.001
Systolic blood pressure	0.229	0.000
Left ventricular mass index	0.206	0.004
Pulse pressure	0.205	0.000
Body mass index	0.172	0.003
Prescription of HMG-CoA 3 reductase inhibitors (statins)	0.164	0.004
History of peripheral vascular disease	0.160	0.005
History of diabetes mellitus	0.158	0.006
Diastolic blood pressure	0.157	0.006
Log serum triglycerides	0.155	0.007
Prescription of alfacalcidol	−0.150	0.009
Age	0.149	0.010
Weight	0.147	0.010
Prescription of cholecalciferol	−0.131	0.023
History of hypertension	0.128	0.026
Sex (female vs. male)	−0.124	0.031

**Table 2** Linear regression models for pulse wave velocity (PWV) measured in 303 patients

	β	SEM	F	95 % CL	p
log PTH	1.58	0.42	13.9	0.74 to 2.42	<0.001
SBP	0.02	0.01	5.3	0.03 to 0.023	0.023
1α	−1.11	0.04	6.81	−1.94 to −0.27	0.01
log TG	1.37	0.69	4.0	1.99 to 01	0.048

Model fit  $r^2 = 0.47$ , adjusted  $r^2 = 0.18$

statistically significant, unless they improved the fit of the model. PWV was associated with SBP, serum PTH and prescription of alfacalcidol (Tables 2, 3). Another model was created by restricting the analysis to only those patients with echocardiography-derived LVmass index (Tables 2, 3). Co-linearity was checked by simple plotting of variables and checking correlations; the correlation between serum PTH and alfacalcidol prescription was somewhat weak with an  $r^2$  value of 0.023. As both alfacalcidol and PTH were present in both models, patients were re-analysed according to alfacalcidol prescription and quartiles of serum PTH concentrations (Tables 4, 5). PWV was greater in those patients not prescribed alfacalcidol (Fig. 1), despite similar patient demographics, body size, blood pressure and left ventricular mass (Table 4), prescription of antihypertensive medications (ACEI/ARBs 30 vs. 26.6 %, β blockers 33.3 vs. 24.6 %), and prescription of calcium based binders (41.7 vs. 51.5 %), all  $p > 0.05$ . Although PTH was lower in the group not prescribed alfacalcidol, there was no difference in PVW, blood

**Table 3** Linear regression models for pulse wave velocity (PWV) restricted to 197 patients with echocardiography-derived left ventricular mass index (LVMI)

	$\beta$	SEM	F	95 % CL	p value
log PTH	0.85	0.35	6.0	0.16 to 1.53	0.005
SBP	0.02	0.01	7.5	0.01 to 0.03	0.006
LVMI	0.01	0.00	11.4	0.01 to 0.07	0.001
1 $\alpha$	−0.8	0.39	4.2	−1.56 to −0.27	0.04
logTG	1.43	0.65	4.79	0.14 to 2.72	0.03

Model fit  $r^2 = 0.37$ , adjusted  $r^2 = 0.19$

CL confidence limits, SBP systolic blood pressure, PTH serum parathyroid hormone, TG triglycerides, 1 $\alpha$  prescription of calcitriol

**Table 4** Demographics of 60 patients not prescribed alfacalcidol vs. 243 prescribed alfacalcidol

	No alfacalcidol	Alfacalcidol	p
Age, years	69.6 $\pm$ 1.9	65.8 $\pm$ 1.1	0.09
Male sex (%)	63.3	63.1	1.00
Diabetes (%)	51.7	46.3	0.56
Dialysis vintage months	1.33 $\pm$ 0.07	1.46 $\pm$ 0.04	0.10
Weight, kg	69.5 $\pm$ 1.8	71.9 $\pm$ 1.1	0.33
Body mass index, kg/m <sup>2</sup>	25.2 $\pm$ 0.6	25.9 $\pm$ 0.4	0.33
Ejection fraction (%)	53.3 $\pm$ 1.7	54.4 $\pm$ 0.9	0.16
Left ventricular mass, g	262.5 $\pm$ 15.5	266.5 $\pm$ 7.5	0.82
Left ventricular mass index, g/cm <sup>2</sup>	147.5 $\pm$ 8.1	147.5 $\pm$ 3.9	0.99
Heart rate, beats/min	71.4 $\pm$ 1.5	73.7 $\pm$ 0.9	0.24
Systolic blood pressure, mmHg	147.2 $\pm$ 3.9	149.0 $\pm$ 1.8	0.66
Diastolic blood pressure, mmHg	79.2 $\pm$ 2.1	80.7 $\pm$ 1.0	0.52
Albumin, g/l	39.1 $\pm$ 0.6	39.8 $\pm$ 0.3	0.29
Cholesterol, mmol/l	4.0 $\pm$ 0.1	4.0 $\pm$ 0.1	0.77
Parathyroid hormone, pmol/l	16.9 (4–30.5)	30.9 (17.2–56.2)	0.005*
Calcium, mmol/l	2.43 $\pm$ 0.02	2.44 $\pm$ 0.01	0.21
Phosphate, mmol/l	1.44 $\pm$ 0.06	1.49 $\pm$ 0.03	0.45

Dialysis vintage was log transformed

Results expressed as percentage, mean  $\pm$  standard error of the mean (SEM), median and interquartile range

Uncorrected p values displayed and \* p < 0.05 after correction for multiple testing

pressure or left ventricular mass between patient quartiles according to serum PTH (Table 5). There was a trend for the highest PTH quartile to be younger, but heavier and have lower serum phosphate. However after correcting for multiple statistical analyses, these differences were no longer significant.

## Discussion

Patients with chronic kidney disease develop arterial stiffening well before progression to CKD stage 5 [5]. Pulse wave velocity is currently considered the “gold” standard non-invasive measurement of arterial stiffness, due to the reliability of the measurements [3, 4]. We measured aortic PWV, as this is more reproducible compared to carotid-femoral measurements [7, 18].

In the general population, PWV has been shown to increase with age, while there is no significant difference of PWV between men and women [19]. In our cohort, we only found a simple univariate association between age and increasing PWV. Although several smaller studies in dialysis patients have observed increasing PWV with age [1, 20, 21], this finding is not universal, particularly with larger cross sectional studies [22]. As expected, PWV was associated with systolic and diastolic blood pressures and pulse pressure, and left ventricular mass. In addition PWV was also, on univariate analysis, associated with past medical history of hypertension and peripheral vascular disease. While some studies in dialysis patients have reported increased PWV in women [23], our measurements initially suggested a higher PWV in women but after adjustment for heart rate and body size there were no differences.

Some reports have noted increased PWV in diabetic dialysis patients [20, 22], but although we found diabetes to be associated with PWV on univariate analysis we could not confirm this association on multivariate analysis. This may reflect the greater prevalence of impaired glucose metabolism in our inner city catchment population, and the greater prevalence of type 2 diabetes in younger dialysis patients from ethnic minority groups. Our cohort did not show an effect of the type of vascular access on PWV, although there are studies suggesting that PWV is reduced by the presence of an arteriovenous fistula [24]. Other studies have also reported reduced PWV with use of angiotensin converting enzyme inhibitors, angiotensin receptor blockers and calcium channel blockers compared to beta blockers [3, 25]. In our cross sectional analysis, there was no observed effect of any antihypertensive agent, or of the number of antihypertensive medications on PWV. The association of PWV with dyslipidaemia [3, 26] was also not confirmed by our results, since we did not find any association with total cholesterol, high density or low density lipoprotein. This may have been due to the relatively large number of our patients prescribed statins, and on simple correlation there was an affect with statin prescription. However on further analysis there was an association with triglycerides, which supports the observations of an earlier study [27].

**Table 5** Demographics of patients characterised by serum parathyroid hormone (PTH) concentration quartiles


PTH, pmol/l	<13.9	13.9–27.0	27.1–53.4	>53.4	p value
Age, years	68.8 ± 1.7	69.3 ± 1.9	70.3 ± 1.8	58.0 ± 1.7	<0.001*
Male sex (%)	62.7	65.4	62.3	63.5	0.98
Diabetes (%)	41.3	53.8	53.6	41.9	0.22
Dialysis vintage, months	1.39 ± 0.07	1.37 ± 0.02	1.5 ± 0.06	1.51 ± 0.06	0.30
Weight, kg	67.2 ± 1.8	72.3 ± 1.9	70.2 ± 1.8	76.3 ± 2.1	0.015
BMI, kg/m <sup>2</sup>	24.9 ± 0.6	25.1 ± 0.6	25.2 ± 0.6	27.2 ± 0.6	0.068
ej (%)	55.1 ± 1.3	54.8 ± 1.6	54.6 ± 1.9	56.2 ± 1.6	0.94
LVmass, g	256.4 ± 13.5	282.5 ± 14	257.8 ± 14.3	261.7 ± 12.4	0.44
LVMI, g/cm <sup>2</sup>	148.2 ± 7.0	154.3 ± 6.8	145.5 ± 7.6	140.0 ± 6.7	0.52
HR, beats/min	73.3 ± 1.4	70.7 ± 1.5	74.5 ± 1.7	74.5 ± 1.7	0.26
SBP, mmHg	147.1 ± 3.6	150.7 ± 2.9	148.1 ± 3.1	148.4 ± 3.7	0.82
DBP, mmHg	79.0 ± 1.9	80.4 ± 1.6	77.5 ± 1.8	84.5 ± 1.9	0.039
Albumin, g/l	39.5 ± 0.5	38.7 ± 0.4	39.9 ± 0.2	40.6 ± 0.4	0.042
Cholesterol, mmol/l	3.87 ± 0.11	3.91 ± 0.13	4.0 ± 0.13	3.3 ± 0.14	0.38
PWV, m/s	9.62 ± 0.25	9.58 ± 0.23	9.97 ± 0.24	9.79 ± 0.24	0.66
Calcium, mmol/l	2.26 ± 0.02	2.26 ± 0.02	2.28 ± 0.02	2.16 ± 0.02	0.72
Phosphate, mmol/l	1.38 ± 0.5	1.47 ± 0.05	1.45 ± 0.06	1.28 ± 0.05	0.029

Dialysis vintage in months log transformed

BMI body mass index, ej cardiac echo derived ejection fraction, LVMI left ventricular mass index, HR heart rate, SBP systolic blood pressure, DBP diastolic blood pressure, PWV pulse wave velocity

Results expressed as percentage, mean ± SEM, median and interquartile range

p values for one way anova with Holm-Sidak correction and then \* p < 0.05 after correction for multiple testing



**Fig. 1** Measured pulse wave velocity, and pulse wave velocity corrected for heart rate in patients prescribed and not prescribed alfalcidol. Data expressed as mean ± SEM. \*p < 0.05 after correction for multiple analyses

Chronic dialysis patients are at risk of medial vascular calcification, and arterial stiffness [20, 28]. Although the biology of vascular calcification remains complex [29], the risks appear to be increased both when bone turnover is suppressed and also at high turnover states [30]. We did not systematically perform X-rays or computerized tomography (CT) scans on our patients and as such cannot comment on radiological calcification scores and PWV. Due to the limited number of patients with valvular calcification

on echocardiography we did not observe any association between PWV and valvular calcification. Previous reports have suggested an association between PWV and PTH [31, 32]. In our cohort there was no relationship with PTH on simple univariate analysis, and no difference between PTH quartiles. There was a trend for the highest PTH quartile group to be heavier, on one hand, which is reported to be associated with increased PVW, but on the other hand they were younger, which is associated with lower PVW. As such there was a trend for PVW to be higher for those patients in the higher PTH quartiles compared to those in the lower quartiles, and PTH did achieve significance in our multiple linear regression models. However in every day clinical practice alfalcidol dosing is based on review of the trend in serum PTH, calcium and phosphate. Hence alfalcidol therapy may be lowered or withdrawn in patients with low PTH values in an attempt to prevent further PTH suppression and low bone turnover, and this may explain the differences in serum PTH levels in those patients who were or were not prescribed calcitriol. Previous studies suggesting a relationship with PTH and PVW did not analyse the effect of medications. Patients not prescribed active vitamin D analogues had increased pulse wave velocity, and several large cohort studies have observed increased mortality in those patients not prescribed active vitamin D analogues [33, 34]. Gla-matrix



protein which inhibits vascular calcification is regulated by vitamin K, and possibly vitamin D3. Many dialysis patients in the UK are deficient in 25 hydroxy-vitamin D3, as sunlight levels are generally inadequate for most of the year, particularly for ethnic minority groups. As such our policy is to provide replacement cholecalciferol, and we could therefore not demonstrate any association between 25 hydroxy vitamin D3 serum concentrations and PWV. Increased arterial vessel stiffness due to vascular calcification could therefore link our results with previous reports of increased mortality in those patients not prescribed active vitamin D analogues. As such pulse wave velocity in haemodialysis patients is associated with non-traditional risk factors associated with bone-mineral disorders of dialysis patients. Further studies are required to determine whether newer markers of vascular calcification are associated with the increased PWV observed in dialysis patients.

**Acknowledgments** Funding—Royal Free hospital.

**Conflict of interest** The authors have no conflict of interest.

## References

- Green D, Roberts PR, New DI, Kalra PA (2011) Sudden cardiac death in haemodialysis patients: an in-depth review. *Am J Kidney Dis* 57(6):921–929
- Seliger SL, Gillen DL, Tirschwell D, Wasse H, Kestenbaum BR, Stehman-Breen CO (2003) Risk factors for incident stroke among patients with end-stage renal disease. *J Am Soc Nephrol* 14(10):2623–2631
- Cavalcante JL, Lima JA, Redheuil A, Al-Mallah MH (2011) Aortic stiffness: current understanding and future directions. *J Am Coll Cardiol* 57(14):1511–1522
- Sigrist MK, Chiarelli G, Levin A, Romann A, Weber C (2010) Pulse wave velocity measurements are reproducible in multiple trained observers: a short report. *Nephron Clin Pract* 116(1):c60–c64
- Covic A, Gusbeth-Tatomir P, Goldsmith DJ (2005) Arterial stiffness in renal patients: an update. *Am J Kidney Dis* 45(6):965–977
- Horváth IG, Németh A, Lenkey Z, Alessandri N, Tufano F, Kis P, Gaszner B, Cziráki A (2010) Invasive validation of a new oscillometric device (Arteriograph) for measuring augmentation index, central blood pressure and aortic pulse wave velocity. *J Hypertens* 28(10):2068–2075
- Davies JM, Bailey MA, Griffin KJ, Scott DJ (2012) Pulse wave velocity and the non-invasive methods used to assess it: Complior, SphygmoCor, Arteriograph and Vicorder. *Vascular* 20(6):342–349
- Vernon K, Peasegood J, Riddell A, Davenport A (2010) Dialyzers designed to increase internal filtration do not result in significantly increased platelet activation and thrombin generation. *Nephron Clin Pract* 117(4):c403–c408
- Davenport A (2009) Low-molecular-weight heparin as an alternative anticoagulant to unfractionated heparin for routine outpatient haemodialysis treatments. *Nephrology (Carlton)* 14: 455–461
- Willicombe MK, Vernon K, Davenport A (2010) Embolic complications from central venous haemodialysis catheters used with hypertonic citrate locking solution. *Am J Kidney Dis* 55(2):348–351
- Sandhu E, Crawford C, Davenport A (2012) Weight gains and increased blood pressure in outpatient hemodialysis patients due to change in acid dialysate concentrate supplier. *Int J Artif Organs* 35(9):642–647
- Booth J, Pinney J, Davenport A (2010) N-terminal proBNP—marker of cardiac dysfunction, fluid overload, or malnutrition in hemodialysis patients? *Clin J Am Soc Nephrol* 5(6):1036–1040
- Davenport A (2012) Changes in N-terminal pro-brain natriuretic peptide correlate with fluid volume changes assessed by bioimpedance in peritoneal dialysis patients. *Am J Nephrol* 36(4):371–376
- Moran N, Buscombe J, Davenport A (2008) Predicting cardiac events using adenosine triphosphate myocardial perfusion scintigraphy in renal transplant candidates. *J Nephrol Renal Transplant* 1:41–52
- Levin NW, Kotanko P, Eckardt KU, Kasiske BL, Chazot C, Cheung AK, Redon J, Wheeler DC, Zoccali C, London GM (2010) Blood pressure in chronic kidney disease stage 5D-report from a kidney disease: improving global outcomes controversies conference. *Kidney Int* 77(4):273–284
- Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise J, Solomon S, Spencer KT, St John Sutton M, Stewart W (2006) American Society of Echocardiography's Nomenclature and Standards Committee; Task Force on Chamber Quantification; American College of Cardiology Echocardiography Committee; American Heart Association; European Association of Echocardiography, European Society of Cardiology. Recommendations for chamber quantification. *Eur J Echocardiogr* 7(2):79–108
- Davenport A. Differences in prescribed Kt/V and delivered haemodialysis dose—why obesity makes a difference to survival for haemodialysis patients when using a 'one size fits all' Kt/V target. *Nephrol Dial Transplant*. 2013. PMID: 23787543
- Nemcsik J, Egresits J, El Hadj Othmane T, Fekete BC, Fodor E, Szabó T, Járó Z, Jekkel C, Kiss I, Tislér A (2009) Validation of arteriograph—a new oscillometric device to measure arterial stiffness in patients on maintenance haemodialysis. *Kidney Blood Press Res* 32(3):223–229
- The Reference Values for Arterial Stiffness' Collaboration (2010) Determinants of pulse wave velocity in healthy people and in the presence of cardiovascular risk factors: 'establishing normal and reference values'. *Eur Heart J* 31(19):2338–2350
- Toussaint ND, Lau KK, Strauss BJ, Polkinghorne KR, Kerr PG (2009) Relationship between vascular calcification, arterial stiffness and bone mineral density in a cross-sectional study of prevalent Australian haemodialysis patients. *Nephrology (Carlton)* 14(1):105–112
- Utescu MS, Couture V, Mac-Way F, De Serres SA, Marquis K, Larivière R, Desmeules S, Lebel M, Boutouyrie P, Agharazii M (2013) Determinants of progression of aortic stiffness in haemodialysis patients: a prospective longitudinal study. *Hypertension* 62(1):154–160
- Celik G, Demirci MS, Tumuklu M, Asci G, Sipahi S, Toz H, Basci A, Ok E (2011) Factors related to pulse wave velocity and augmentation index in chronic haemodialysis patients. *Ren Fail* 33(10):957–963
- Kuang DW, Li CL, Kuok UI, Cheung K, Lio WI, Xin J (2012) Risk factors associated with brachial-ankle pulse wave velocity among peritoneal dialysis patients in Macao. *BMC Nephrol* 13:143. doi:10.1186/1471-2369-13-143
- Korsheed S, Eldehni MT, John SG, Fluck RJ, McIntyre CW (2011) Effects of arteriovenous fistula formation on arterial

- stiffness and cardiovascular performance and function. *Nephrol Dial Transplant* 26(10):3296–3302
25. Shigenaga A, Tamura K, Dejima T, Ozawa M, Wakui H, Masuda S, Azuma K, Tsurumi-Ikeya Y, Mitsuhashi H, Okano Y, Kokuho T, Sugano T, Ishigami T, Toya Y, Uchino K, Tokita Y, Umemura S (2009) Effects of angiotensin II type 1 receptor blocker on blood pressure variability and cardiovascular remodelling in hypertensive patients on chronic peritoneal dialysis. *Nephron Clin Pract* 112(1):c31–c40
  26. Jung JY, Hwang YH, Lee SW, Lee H, Kim DK, Kim S, Oh YG, Yang J, Joo KW, Ahn C, Oh KH (2010) Factors associated with aortic stiffness and its change over time in peritoneal dialysis patients. *Nephrol Dial Transplant* 25(12):4041–4048
  27. Noordzij M, Cranenburg EM, Engelsman LF, Hermans MM, Boeschoten EW, Brandenburg VM, Bos WJ, Kooman JP, Dekker FW, Ketteler M, Schurgers LJ, Krediet RT, Korevaar JC, NECOSAD Study Group (2011) Progression of aortic calcification is associated with disorders of mineral metabolism and mortality in chronic dialysis patients. *Nephrol Dial Transplant* 26(5):1662–1669
  28. Shroff RC, Shanahan CM (2007) The vascular biology of calcification. *Semin Dial* 20(2):103–109
  29. London GM, Marchais SJ, Guérin AP, Boutouyrie P, Métivier F, de Vernejoul MC (2008) Association of bone activity, calcium load, aortic stiffness, and calcifications in ESRD. *J Am Soc Nephrol* 19(9):1827–1835
  30. Townsend RR, Chirinos JA, Parsa A, Weir MA, Sozio SM, Lash JP, Chen J, Steigerwalt SP, Go AS, Hsu CY, Rafey M, Wright JT Jr, Duckworth MJ, Gadegbeku CA, Joffe MP, Chronic Renal Insufficiency Cohort Investigators (2010) Central pulse pressure in chronic kidney disease: a chronic renal insufficiency cohort ancillary study. *Hypertension* 56(3):518–524
  31. Jung JY, Hwang YH, Lee SW, Lee H, Kim DK, Kim S, Oh YG, Yang J, Joo KW, Ahn C, Oh KH. Factors associated with aortic stiffness and its change over time in peritoneal dialysis patients
  32. Shoji T, Shinohara K, Kimoto E, Emoto M, Tahara H, Koyama H, Inaba M, Fukumoto S, Ishimura E, Miki T, Tabata T, Nishizawa Y (2004) Lower risk for cardiovascular mortality in oral 1 $\alpha$ -hydroxy vitamin D3 users in a haemodialysis population. *Nephrol Dial Transplant* 19:179–184
  33. Teng M, Wolf M, Lowrie E, Ofsthun N, Lazarus JM, Thadhani R (2003) Survival of patients undergoing hemodialysis with paricalcitol or calcitriol therapy. *N Engl J Med* 349(5):446–456
  34. Cozzolino M, Mehmeti F, Ciceri P, Volpi E, Stucchi A, Brenna I, Cusi D (2011) The effect of paricalcitol on vascular calcification and cardiovascular disease in uremia: beyond PTH control. *Int J Nephrol* 2011:269060. doi:[10.4061/2011/269060](https://doi.org/10.4061/2011/269060)