

Adverse Health Effects of Frailty in CKD Patients

Cause	2
Non-adjusted differences in comorbidity and system-based influences	5
Complications	17
Bibliography	32

Cause

	Effect (descriptions)	Risk Difference	CKD severity	Frailty Assessment	Sample Size	Reference
Lifestyle	Smoking*	RR 1.18 (1.04-1.34)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	¹
Ethnicity	Hispanic*	Frailty scores +0.6 points per year	CKD stage 5D (hemodialysis)	Fried Phenotypes	762	²
Biological						
Cardiovascular	Hypertension*	RR 1.6 (1.26-2.04)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	¹
	Peripheral vascular disease*	RR 1.58 (1.34-1.8)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	¹
	Left ventricular dysfunction*	RR 1.18 (1.03-1.36)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	¹
	Endothelial dysfunction	r = -0.367 (p= 0.004)	CKD stages 3-5	Fried Phenotypes	61	³
		OR 3.86 (1.00-14.88)				
Vascular access	Use of a permanent vascular access (fistula or graft)	HR 0.71 (0.51-0.98)	CKD stage 5D (hemodialysis)	Fried Phenotypes	2275	⁴
Cerebrovascular	Cerebrovascular Accident	RR 1.34 (1.19-1.5)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	¹
Pulmonary	COPD	OR 1.68 (1.16-2.45)	CKD stages 1-5	Fried Phenotypes	10256	⁵

	Inflammatory	IL-6*	Worse frailty	CKD stage 5D (hemodialysis)	Fried Phenotypes	762	²
		CRP	After adjustment, OR 1.76 (1.28-2.41) to 1.50 (1.07-2.09)	Chronic kidney insufficiency, serum creatinine \geq 1.3 mg/dL	Fried Phenotypes	5888	⁶
		Fibrinogen					
	Endocrinologic/ Metabolic	Diabetes	Frailty scores +0.7 points per year	CKD stage 5D (hemodialysis)	Fried Phenotypes	762	²
			OR 1.68 (1.16-2.45)	CKD stages 1-5	Fried Phenotypes	10256	⁵
		Obesity (IMC \geq 30 kg/m ²)	OR 6.63 (1.16-36.77)	CKD stages 3-5	Fried Phenotypes	61	³
		Higher parathyroid hormones (PTH)	r= 0.30 (p= 0.01)	CKD stages 3-5	Fried Phenotypes	61	³
	Body Composition	Higher fat mass	r= 0.25 (p= 0.04)	CKD stages 3-5	Fried Phenotypes	61	³
	Cancer	Cancer	OR 1.89 (1.19-2.99)	CKD stages 1-5	Fried Phenotypes	10256	⁵
	Arthritis	Arthritis	OR 3.34 (2.08-5.38)	CKD stages 1-5	Fried Phenotypes	10256	⁵
	Laboratory Data	eGFR (mL/min/1.72m ²) reference: > 60					
		eGFRcys < 30	Frailty prevalence 2.8	CKD stages 1-4	Fried	336	⁷

		eGFRcys 30-44	Frailty prevalence 2.1		Phenotypes		
		Serum Albumin Concentrations (g/dL)	Frailty scores -1.1 points per g/dL	CKD stage 5D (hemodialysis)	Fried Phenotypes	762	²
		Serum Creatinine <4 mg/dL*	RR 1.46 (1.22-1.71)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	¹
		Testosterone, every 50% lower free form		CKD stage 5D (hemodialysis), men	Fried Phenotypes	440	⁸
		being frail	OR 1.40 (1.05-1.53)				
		becoming frail over 12 months	OR 1.40 (1.07-1.73)				
		Hemoglobin	Adjusted, OR 1.76 (1.28-2.41) to 1.50 (1.07-2.09)	Chronic kidney insufficiency, serum creatinine ≥1.3mg/dL	Fried Phenotypes	5888	⁶
		LDL, HDL					

Non-adjusted differences in comorbidity and system-based influences

	Effect (descriptions)	Prevalence	CKD Severity	Frailty Assessment	Sample Size	Reference
Biological						
Cardiovascular	Heart Failure (%)	30% vs 12%	CKD stages 1-4	Fried Phenotypes	336	⁷
	Angina (%)	34% vs. 22%	CKD stages 1-4	Fried Phenotypes	336	⁷
Cerebrovascular	Cerebrovascular Disease (%)	26.4 vs. 12.0	CKD stage 5D (hemodialysis)	Fried Phenotypes	324	⁹
Endocrinologic/ Metabolic	Diabetes (%)					
		64% vs. 49%	CKD stages 1-4	Fried Phenotypes	336	⁷
	Obesity (%)	F vs. NF				
		64% vs. 50%	CKD stages 1-4	Fried Phenotypes	336	⁷
		51.8% vs. 23.9%	CKD stage 5D (hemodialysis)	Fried Phenotypes	324	⁹
	BMI based on dry weight	31.5 vs. 27.6	CKD stage 5D (hemodialysis)	Fried Phenotypes	324	⁹
	BMI	22.53 vs. 26.16	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	¹⁰
Cognitive	Mini-Mental State Examination		Elderly, ≥65y/o	Edmonton Frail	137	¹¹

		(MMSE)			Scale (EFS)		
			Spearman's correlation coefficient of EFS scores with gross MMSE scores	-0.607 (p<0.01)			
		Executive Function		F vs. NF			
			Trail Making Tests A (TMTA) scores	+12.08	CKD stage 5D (hemodialysis)	Fried Phenotypes	324 ⁹
			Trail Making Tests B (TMTB) scores	+33.15	CKD stage 5D (hemodialysis)	Fried Phenotypes	324 ⁹
Immunological		Inflammatory					
			CRP (ln CRP) (mg/dL)	1.12 vs 0.28	CKD stage 5D (peritoneal dialysis)	Clinical Frailty Scale (CFS)	119 ¹²
			IL6 (ln IL6) (mg/dL)	2.45 vs. 1.58			
		Mycophenolate mofetil (MMF) dose reduction (MDR)		F vs. NF	CKD stage 5T	Fried Phenotypes	525 ¹³
			1 year since KT (%)	44 vs 40			
			2 years since KT (%)	54 vs. 45			
			3 years since KT (%)	67 vs. 51			
		Viral infection		F vs. NF			
			HCV (n=37) (capitals)	36 vs. 1	CKD stage 5D (hemodialysis)	Fried Phenotypes	205 ¹
Body Composition		Appendicular					
			Appendicular skeletal muscle mass index	6.8 vs. 7.7	CKD stage 1-5	Edmonton Frail Scale (EFS)	41 ¹⁴

		(ASMI)					
		Appendicular fat percentage	FRAIL scale				
		Left/Right lower extremity	t = 2.32; p = 0.03 (left); t = 2.05; p = 0.048 (right)	CKD stage 5D (hemodialysis)	FRAIL scale	44	15
		Left/Right upper extremity	t = 2.66; p = 0.01 (left); t = 3.09; p = <0.01 (right)				
		Appendicular fat percentage	Frail/Prefrail vs. Nonfrail				
		Left/Right lower extremity	t = 2.31; p = 0.03 (left); t = 2.28; p = 0.03 (right)	CKD stage 5D (hemodialysis)	Self- report instrument evaluating five dimensions of frailty (fatigue, resistance, ambulation, illnesses, and weight loss)	44	15
		Right upper extremity	t = 2.35; p = 0.03				
	Lower lean mass						
			F/PF vs. NF				

			Whole body (kg)	34.7 vs. 43.1	CKD stage 5D (hemodialysis)	Self- report instrument evaluating five dimensions of frailty (fatigue, resistance, ambulation, illnesses, and weight loss)	44	15
			Cephalic/trunk (g)	3059 vs. 3288 (cephalic); 17.4 vs. 22.1 (trunk)				
			Right/Left upper limb (g)	1831 vs. 2493 (right); 1869 vs. 2515 (left)				
			Right/Left lower limb (g)	4920 vs. 6114 (right); 4650 vs. 6349 (left)				
			Lean body mass (in frail vs. nonfrail) (%)	57.1% vs .14.7%	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	14
	Laboratory Data	Prealbumin (PRAB) (mg/dL)		28.9 vs. 38.3	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	14
		Serum albumin (g/L)		Pearson correlation coefficient r = - 0.263 (p = 0.025)	CKD stage 5D (hemodialysis)	Fried Phenotypes	74	16
				38 vs. 41	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	14
				29.2 vs. 34.8	CKD stage 5D (peritoneal	Clinical Frailty Scale (CFS)	119	12

					dialysis)			
				36.1 vs. 38.5	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	10
			Frail with depression vs. Frail without depression vs. Nonfrail	32.9 vs. 34.9 vs. 35.8 (p=0.025)	CKD stage 5D (peritoneal dialysis)	In-house Chinese questionnaire	178	17
		Calcium (mmol/L)		2.24 vs. 2.36	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	14
		Hemoglobin (g/dL)		Pearson correlation coefficient r = - 0.336 (p = 0.004)	CKD stage 5D (hemodialysis)	Fried Phenotypes	74	16
				6.49 vs. 8.07 (p = 0.037)				
				10.35 vs. 10.97	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	10
		25-hydroxy vitamin D (ng/mL)		Pearson correlation coefficient r = - 0.363 (p = 0.002)	CKD stage 5D (hemodialysis)	Fried Phenotypes	74	16
				11.58 vs. 17.09				

			(p = 0.005)				
	Miscellaneous	Dialysis clearance rate (Kt/V and urea reduction ratio)	↑	CKD stage 5D (chronic dialysis)	FRAIL scale	46	18
	Composite	Number of complications (complications identified at data collection: High Pressure Cramping, Anemia, Weight loss Pain, Weakness, Weight gain Constipation, Heart Arrhythmia, Headache, Itch, Recurrent infections, Arterial hypertension)	Spearman's correlation 0.666 (p = 0.000 in table) (p < 0.05 in text)	Elderly (≥ 60 yo), with diagnosis of CKD	Edmonton Frail Scale (EFS)	35	28
		Higher number of comorbid conditions	6 vs. 4 (p = 0.03)	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	14
		Charlson's comorbidity score	Spearman's rank correlation coefficient r = 0.40 (p < 0.0001)	CKD stage 5D (peritoneal dialysis)	Chinese questionnaire	193	26
	Functional Status	Disability	F vs. NF				
		≥1 disability in activities of daily Living (ADLs)	15% vs. 5%	CKD stages 1-4	Fried Phenotypes	336	7
		≥1 disability in instrumental	60% vs. 28%				

		activities of daily living (IADLs)					
		≥1 disability in mobility tasks	40% vs. 18%				
		Less ADL-independent	55% vs. 91%	≥ 65 yo, predialysis, eGFR < 20 mL/min	Groningen frailty indicator (GFI)	65	¹⁹
	Ability to perform basic activities of daily living		33.33% vs 76.4%	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	¹⁰
	Ability to perform transfers		38.8% vs. 84.7%				
Microbiota							
	Gut microbiota	Malnutrition-Inflammation-Score (MIS)	7.6 vs. 3.9	Stage 3b-4, eGFR 15-45ml/min	Fried Phenotype score	64 (and 15 control subjects)	²⁰
		Abundance of unclassified Mogibacteriaceae and Oscillospira	Directly proportional to MIS				
		Abundance of Akkermansia, Ruminococcus, and Eubacterium	Inversely proportional to MIS				
		Bacterial Abundance of some genera (Mogibacteriaceae, Coriobacteriaceae, Eggerthella, Erwinia, Coprobacillus,	↑				

		Anaerotruncus, etc)					
Neurological							
EEG	Brain Wave		F vs. NF	ESRD, under chronic dialysis	FRAIL scale	46	18
		Global DAR	283 ± 679 vs. 2971 ± 4859				
		DARs (left frontal)	135 ± 250 vs. 3073 ± 4702				
		DAR (left TO)	197 ± 318 vs. 3708 ± 6398				
		DAR (central)	55 ± 96 vs. 1773 ± 3262				
		DAR (right TO)	187 ± 261 vs. 4400 ± 7763				
		Global DTABR	191 ± 469 vs. 1781 ± 2793				
		DTABR (left frontal)	86 ± 158 vs. 1680 ± 2388				
		DTABR (left TO)	130 ± 210 vs. 1884 ± 2828				
		DTABR (central)	39 ± 65 vs. 1132 ± 1957				
		DTABR (right TO)	126 ± 178 vs. 2960 ± 5271				
Psychological							

	Mood	Mood Change	Negative change	CKD stage 5D (hemodialysis)	Edmonton Frail Scale (EFS)	N/A	21
	Mental Health						
	Anxiety	Hospital Anxiety and Depression Scale (HADS)	<u>Women</u> : ↑ in global, psychological, social frailty <u>Men</u> : ↑ in Physical frailty	ESRD, under online-haemodiafiltration (OL-HDF)	N/A	97	22
	Depression	Hospital Anxiety and Depression Scale (HADS)	<u>Men</u> ↑ in global, psychological, physical frailty	ESRD, under online-haemodiafiltration (OL-HDF)	N/A	97	22
		Incidence (%) (Self-reported Major Depression Inventory)	83 vs. 6	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	14
	Mental Function	Post-KT delirium	9.0% vs. 3.9%	CKD stage 5T	Fried Phenotypes	893	23
	Sociological						
	Interaction	Interaction with family	Good				24
	Physical activity	Minnesota Leisure Time Activity (LTA)	95 vs. 735 (p<0.001)	CKD stage 5D (hemodialysis)	Fried Phenotypes	68	25
		Low Physical Activity Questionnaire (LoPAQ)	280 vs. 798 (p=0.003)				
		Sitting (hours/day)	6.5 vs. 5				

			(p=0.04)				
Nutritional Status	Overall subjective global assessment (SGA) (weight loss, anorexia, subcutaneous fat, muscle mass) (Frail with depression vs. Frail without depression vs. Nonfrail)		5.04 vs. 5.41 vs. 5.75 (p < 0.0001)	CKD stage 5D (peritoneal dialysis)	In-house Chinese questionnaire	178	17
			Spearman's rank correlation coefficient r = - 0.44, p < 0.0001	CKD stage 5D (peritoneal dialysis)	Chinese questionnaire	193	26
	Malnutrition inflammation score (MIS) (frail with depression vs. frail without depression vs. nonfrail)		9.48 vs. 7.13 vs. 5.12 (p < 0.0001)	CKD stage 5D (peritoneal dialysis)	In-house Chinese questionnaire	178	17
			Spearman's rank correlation coefficient r = 0.40, p < 0.0001	CKD stage 5D (peritoneal dialysis)	Chinese questionnaire	193	26
Quality of Life	HRQoL						
		SF-36					
		Scores in physical functioning, blood pressure, role	↓	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	14

		physical, and physical component summary domains					
	Kidney Disease Quality of Life (KDQoL)		F vs. NF				
		Physical health	33.7 vs. 40.7	ESRD CKD stage 5D (conventional hemodialysis)	Fried Phenotypes	151	27
		Kidney disease effects	51.6 vs. 66.8				
Independence	Functional Independence Measure (FIM)			Elderly, ≥65y/o	Edmonton Frail Scale (EFS)	137	11
	Spearman's correlation coefficient	Frailty diagnosis with global FIM	-0.703 (p<0.001)				
		Frailty diagnosis with motor FIM	-0.714 (p<0.001)				
		Frailty diagnosis with cognitive FIM	-0.575 (p<0.001)				
		EFS scores with gross FIM	-0.53 (p<0.01)				
Health-care utilization	Hospitalization		90% vs. 53% (p = 0.04)	≥ 65 yo, predialysis, eGFR < 20 mL/min	Groningen frailty indicator (GFI)	65	19
		Cumulative number of inpatient health-care visits	↑	CKD stage 1-5	Edmonton Frail Scale (EFS)	41	14
		Cumulative number of					

		emergency health-care visits					
		Cumulative number of total health-care visits					
		Admissions/year	0.77727 vs. 0.2838	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	¹⁰

Complications

	Effect (descriptions)	Risk Difference	CKD Severity	Frailty Assessment	Sample Size	Reference
Biological						
Cardiovascular	Cerebrovascular Accident	OR 1.55 (1.05-2.99)	CKD stage 5D (hemodialysis)	Fried Phenotypes	2275	⁴
	QRS duration	β coefficient = -0.29, t = -2.03 (p = 0.048)	CKD stage 5D (chronic hemodialysis)	Edmonton frailty scale (EFS)	41	²⁹
		β coefficient = -0.27, t = -1.84 (p = 0.05)		FRAIL scale		
	Vascular Access failure	HR 2.63 (1.03-6.71)	ESRD (CKD stage 5D)	Self-reported FRAIL scale	51	³⁰
Renal Function Decline	Risk for death or dialysis therapy	2.5 (1.4-4.4)-fold greater	CKD stages 1-4	Fried Phenotypes	336	⁷
Immunological	Mycophenolate mofetil (MMF) dose reduction (MDR)	HR 1.29 (1.01-1.66)	CKD stage 5T	Fried Phenotypes	525	¹³
Cognitive	Modified Mini-Mental State (3MS)	-2.37 to -2.80 (1 year) (p=0.03)	ESRD	Fried Phenotypes	324	⁹
	Declined, 1-4 years post-KT (points/week)	Slope = -0.04 vs. 0.005	CKD stage 5T	Fried physical frailty phenotypes (PFP)	665	³¹
	At 4 year post-KT (points)	-5.5 (87.4 vs.				

			92.9)				
Functional status	Disability		F vs. Prefrail vs. NF				
		Need for activities of daily livings (ADL) assistance	OR 11.32 (5.49-23.32) vs. 1.93 (1.01-3.68) vs. 1.00	CKD stage 5D (hemodialysis)	Fried Phenotypes	742	³²
Diabetes	Diabetes		OR 1.35 (1.10-1.65)	CKD stage 5D	Fried Phenotypes	2275	⁴
Body composition	Bones						
		Bone Mineral Density (BMD)	One year follow-up, with frailty	ESRD CKD stage 5D (chronic hemodialysis)	FRAIL Scale	43	³³
		L1	$\beta = -0.4$, $t = -2.18$, $p=0.04$				
		L4	$\beta = -0.39$, $t = -2.1$, $p=0.046$				
		Femur Neck (FN)	$\beta = -0.5$, $t = -2.96$, $p<0.01$				
			$\beta = -4$, $t = -3.17$, $p = 0.004$				
		Total	$\beta = -0.53$, $t = -3.27$, $p < 0.01$	ESRD CKD stage 5D (chronic	FRAIL Scale	43	³³

					hemodialysis)			
			Areas	One year follow-up, with frailty	ESRD CKD stage 5D (chronic hemodialysis)	FRAIL Scale	43	33
			Average L-spine areas	$\beta = -0.48$, $t = -2.84$, $p < 0.01$				
			Changes of average L-spine areas	$\beta = -0.5$, $t = -3.02$, $p < 0.01$				
			Z-score	One year follow-up, with frailty				
			Percentage change of L1 Z-score	$\beta = -0.45$, $t = -2.11$, $p = 0.049$				
			Vertebral Compression Fracture (VCF)	OR 1.8 ($p = 0.01$)	ESRD CKD stage 5D (chronic hemodialysis)	FRAIL Scale	43	34
			Quantitative ultrasound (QUS) parameters of calcaneus					
			Speed of sound (SOS)	Standardized β (range, p value)	CKD stage 5D (maintenance hemodialysis)	Fried Phenotypes	214	35
			Female (Five frailty criteria)	Negative (-0.253 to -0.439, $p \leq 0.034$)				

			Male (All criteria significant except weight loss)	Negative (-0.277 to -0.402, $p \leq$ 0.003)				
			Broadband ultrasound attenuation (BUA)	Standardized β (range, p value)				
			Female (All criteria significant except weakness and weight loss)	Negative (-0.209 to -0.354, $p \leq$ 0.045)				
			Male (All criteria significant except weight loss)	Negative (-0.171 to -0.371, $p \leq$ 0.045)				
			Stiffness index	Standardized β (range, p value)				
			Female (All criteria significant except weight loss)	Negative (-0.271 to -0.461, $p \leq$ 0.018)				
			Male (Five frailty criteria)	Negative (-0.183 to -0.461, $p \leq$ 0.048)				
			Muscles					
			Quadriceps muscle area (magnitude of association	Multivariable coefficient -30.3		Performance-based frailty (PbF)	80	36

		with PbF vs. 10 years of age)	cm ² (p = 0.02) vs. -6.6 cm ² (p = 0.0001)				
Laboratory data	Serum Albumin Concentrations (g/dL)						
		<3.2 vs. ≥3.9	OR 1.89 (1.30-2.59)	CKD stage 5D	Fried Phenotypes	2275	⁴
		Hypoalbuminemia	Negative association (p = 0.01)	CKD stage 5D (maintenance hemodialysis) (ESRD)	Frail Scale	46	³⁷
Psychological							
	Delirium	Post-KT delirium	OR 2.05 (1.02-4.13)	CKD stage 5T	Fried Phenotypes	893	²³
	Distress	Distress Thermometer	β = 0.35, t = 3.0 (95% CL = 0.12-0.58) (p = 0.003)	CKD stage 5D (hemodialysis)	Canadian frailty score	382	³⁸
Quality of Life	HRQoL						
		Fair/Poor HRQOL at follow-up (median 9.4 mo)	aOR 2.79 (1.32-5.90)	ESRD CKD stage 5T	Fried Phenotypes	233	³⁹
		Worsening HRQOL at follow-up (median 9.4 mo)	aRR 2.91 (1.08-7.80)				
		SF-36					
		Hierarchical regression	29% (p<0.001)	CKD stage 2-4	Fried Phenotypes	168	⁴⁰

		R ² change (effects of frailty on HRQoL) in Physical Component Summary (PCS)					
		Hierarchical regression R ² change (effects of frailty on HRQoL) in Mental Component Summary (MCS)	21.3% (p<0.001)				
		Physical components	Simple linear regression coefficient = -1.12 (-1.47 to -0.76) (p < 0.001)	CKD stages 3-5 (predialysis treatment)	Fried Phenotypes	61	41
		Mental components	Simple linear regression coefficient = -0.75 (-1.40 to -.016)				
		SF-12					
		MCS	Effect estimate 0.94 (0.91-0.97) (p<0.01)	CKD stage 5D (peritoneal dialysis, n=129; hemodialysis,	The Canadian Study of Health and Aging Clinical Frailty Scale (CFS)	251	42
		PCS	Effect estimate				

			0.88 (0.84-0.91) (p<0.01)	n=122)			
		KDQOL-SF scores within 3 months post-KT	F vs. NF				
		At KT	↓	CKD stage 5T	Fried Phenotypes	443	43
		Physical HRQoL	-6.31 points (95% CI -8.16 to -4.46)				
		Kidney disease-specific HRQoL	-6.53 points (95% CI -9.17 to -3.89)				
		Post-KT	Greater improvement				
		Physical HRQoL	1.35 points/month (0.65 to 2.05) vs. 0.34 points/month (-0.17 to 0.85)				
		Kidney disease-specific HRQoL	3.75 points/month (2.89 to 4.60) vs. 2.41 points/month				

					(1.78 to 3.04)				
				Constituent domains	Greater improvement				
				General health	4.93 points/month (3.51 to 6.35) vs. 2.87 points/month (1.82 to 3.92)				
				Effects of ESRD on daily living	7.10 points/month (5.68 to 8.51) vs. 4.01 points/month (2.99 to 5.03)				
				Cognitive function	2.88 points/month (1.80 to 3.96) vs. 1.28 points/month (0.50 to 2.07)				
				Social interaction	1.18 points/month (-0.06 to 2.43) vs. -				

					0.57 points/month (- 1.47 to 0.33)				
				Illness Intrusiveness Rating Scale	Effect estimate 1.14 (1.09-1.20)	CKD stage 5D (peritoneal dialysis, n=129; hemodialysis, n=122)	The Canadian Study of Health and Aging Clinical Frailty Scale (CFS)	251	42
				Barthel Index	Effect estimate 0.89 (0.86-.093)				
				Symptom score	Effect estimate 1.23 (1.13-1.34)				
				Hospital Anxiety and Depression Scale	Effect estimate 1.21 (1.11-1.31)				
				Falls	HR 2.1 (1.21- 3.92)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	1
					OR 2.39 (1.22- 4.71)	CKD stage 5D (maintenance hemodialysis)	Fried frailty index	762	44
				Higher numbers of falls	HR 3.09 (1.38- 6.90)	CKD stage 5D (hemodialysis)		95	45
				Time to first fall or fracture requiring medical attention	HR 1.60 (1.16- 2.20)	CKD stage 5D (maintenance hemodialysis)	Modified Fried Phenotypes by Bao Y ⁴⁶ .	1646	47
Graft Loss				Death-censored graft loss					

		F vs. NF (in patients with depressive symptoms)	aHR 6.20 (1.67, 22.95) vs. 3.16 (0.90, 11.04)	CKD stage 5T	Fried Phenotypes	773	⁴⁸
Health-care utilization	Hospitalization/Death		HR 1.56 (1.36-1.79)	CKD stage 5D	Fried Phenotypes	2275	⁴
	Hospitalization		HR 2.06 (1.18-3.58)	CKD stage 5D (hemodialysis)	Fried Phenotypes	205	¹
			aHR 1.80 (1.4-2.3)	CKD stage 5D (maintenance hemodialysis & peritoneal dialysis)	Adopted	1658	⁴⁹
			Relative risk = 1.43 (1.00-2.03)	CKD stage 5D (hemodialysis)	Fried Phenotypes	146	⁵⁰
		Number of hospitalizations for all causes	beta = 0.29 (p < 0.0001)	CKD stage 5D (peritoneal dialysis)	Chinese questionnaire	193	²⁶
		Number of hospitalizations related to cardiovascular events	beta = 0.37 (p < 0.0001)				
		Non-vascular access-related hospitalizations	aHR 1.98 (1.41-1.87) (內文應該寫錯，無勘誤)	CKD stage 5D	Fried Phenotypes	2275	⁴
	Time to first hospitalization		HR 1.26 (1.09-1.45)	CKD stage 5D (maintenance dialysis)	Earlier modification of Fried Phenotypes by Johansen et al ⁴ , but	1576	⁴⁶

				without weight loss.		
	Early Hospital Readmission (EHR)	aRR 1.61 (1.81-2.19) (p=0.002)	CKD stage 5T	Fried Phenotypes	383	51
	Duration of hospitalization					
	Hospital stay (days per year of follow up) (frail with depression vs. frail without depression vs. nonfrail)	26.62 (IQR 10.65-61.18) vs. 14.05 (IQR 3.57-37.27) vs. 8.04 (IQR 0.91-19.42) (p<0.0001)	CKD stage 5D (peritoneal dialysis)	In-house Chinese questionnaire	178	17
	Total length of hospital stay	beta = 0.34 (p < 0.0001)	CKD stage 5D (peritoneal dialysis)	Chinese questionnaire	193	26
	Longer Length of Stay (LOS)					
	with delayed graft function (DGF), LOS	Relative Risk 1.15 (1.03-1.29)	CKD stage 5T	Fried Phenotypes	589	52
	With DGF, LOS ≥2 weeks	OR 1.57 (1.06-2.33)				
	≥2 weeks		CKD stage 5 to 5T	Fried Phenotypes	569	53
	Change in 3 categories (more frail)	OR 2.02 (1.20-3.40)				
	Change in frailty scores (more frail)	OR 1.92 (1.13-3.25)				
	With depressive symptoms (aRR difference between F	aRR 1.88 (1.70-2.08) vs. 1.38	CKD stage 5T	Fried Phenotypes	773	48

		and NF)	(1.27-1.52)				
		CES-D score (10-point increase) (aRR increase between F and NF)	aRR 1.23 (1.16-1.31) vs. 1.17 (1.08-1.27)				
Mortality	Mortality		2.17 fold	CKD stage 5T	Fried Phenotypes	537	⁵⁴
			HR 1.57 (1.25-1.97)	CKD stage 5D (maintenance dialysis)	Earlier modification of Fried Phenotypes by Johansen et al ⁴ , but without weight loss.	1576	⁴⁶
			HR 2.24 (1.60-3.15)	CKD stage 5D	Fried Phenotypes	2275	⁴
			HR 1.22 (1.04-1.43)	CKD stage 5D (incident chronic dialysis)	CFS	390	⁵⁵
			HR 4.28 (1.22-14.98)	Predialysis (eGFR ≤ 25 mL)	PRISMA questionnaire & Timed up and Go test	104	⁵⁶
			aHR 9.83 (1.80-53.7)	CKD stage 5D (peritoneal dialysis)	CFS	119	¹²
			HR 2.60 (1.04-6.49)	CKD stage 5D (hemodialysis)	Fried Phenotypes	146	⁵⁰
			HR 2.37 (1.11-5.02)	CKD stage 5D (maintenance hemodialysis & peritoneal dialysis)	Adopted	1658	⁴⁹

		20.45% vs. 12.36% (p<0.005)	CKD stage 5D (hemodialysis)	Fried Phenotypes	320	¹⁰
	Performance-based frailty	HR 2.16 (1.41- 3.29)	CKD stage 5D (hemodialysis)	Fried Phenotypes & a definition that substitutes self-reported measures available on the Medical Outcomes Study 36-Item Short Form (SF-36) for the physical performance and exhaustion criteria.	771	⁵⁷
	Self-reported function- based frailty	HR 1.93 (1.24- 3.00)				
	Patients who met both performance- and self- reported function-based frailty	HR 2.46 (1.51- 4.01)				
	F vs. NF (in patients with depressive symptoms)	aHR 2.62 (1.03, 6.70) vs. 1.92 (0.68, 5.38)	CKD stage 5T	Fried Phenotypes	773	⁴⁸
	At 24-month follow up, frail with depression vs. frail without depression vs. nonfrail	62.5% vs. 71.4% vs 86.6% (p=0.001)	CKD stage 5D (peritoneal dialysis)	In-house Chinese questionnaire	178	¹⁷
	Prediction ability of comorbidities in F vs. NF	HR 0.75 (0.44- 1.29) vs. 1.66 (1.17-2.35)	CKD stage 5T (KT candidates, on waitlist)	Fried Phenotypes	2086	⁵⁸
	Out of 10 deceased within 1 year of initiation (percentage of F vs. NF)	30% vs. 9%	≥ 65 yo, predialysis, eGFR < 20 mL/min	Groningen frailty indicator (GFI)	65	¹⁹
	Risk for death or dialysis	2.5 (1.4-4.4)-fold	CKD stages 1-4	Fried Phenotypes	336	⁷

	therapy	greater				
	All-cause mortality					
	Adjusted	HR 1.66 (1.03-2.67)	CKD stage 5D (incident chronic dialysis)	Fried Phenotypes	370	59
	Among BMI ≥ 30 kg/m ²	HR 3.77 (1.10-12.92)				
	Above median Waist-Hip Ratio (WHR)	HR 2.38 (1.17-4.82)				
	Anti-neutrophil cytoplasmic antibody (ANCA)-associated vasculitides (AAV) patients' mortality	HR 2.43 (1.48-3.99)	CKD stage 5D to 5T (RRT [hemodialysis, peritoneal dialysis, transplantation])	Inability to walk without help	425	60
	Mortality of patients with monoclonal gammopathy and ESRD caused by myeloma cast nephropathy (MCN), immunoglobulin light chain amyloidosis (ALA), or light-chain deposition disease (LCDD)	HR, 1.93 (1.58-2.36)	CKD stage 5D to 5T (RRT [hemodialysis, peritoneal dialysis, transplantation])	Inability to walk without help	1462	61
	Post-KT mortality					
	Change in 3 categories (more frail)	HR 2.27 (1.11-4.65)	CKD stage 5 to 5T	Fried Phenotypes	569	53
	Change in frailty scores	HR 2.36 (1.12-				

	(more frail)	4.99)				
Composite	Composite outcomes of all-cause death or cardiovascular hospitalization	HR 23.58 (1.61-346.03)	CKD stage 5D (hemodialysis)	Multidimensional frailty score based on comprehensive geriatric assessment (CGA) protocol	46	⁶²
	30-day postoperative (KT) complications according to Comprehensive Complication Index (CCI)	β =13.31, 95% CI 5.72-20.89 (p = 0.0007)	CKD stage 5T	Groningen Frailty Indicator	150	⁶³

Bibliography

1. Yadla, M., John, J. &Mummadi, M. A study of clinical assessment of frailty in patients on maintenance hemodialysis supported by cashless government scheme. *Saudi J. Kidney Dis. Transplant.* (2017). doi:10.4103/1319-2442.198102
2. Johansen, K. L. *et al.* Factors Associated with Frailty and Its Trajectory among Patients on Hemodialysis. *Clin J Am Soc Nephrol* **12**, 1100–1108 (2017).
3. Mansur, H. N. Fragilidade na doença renal crônica: prevalência e fatores associados. (2012).
4. Johansen, K. L., Chertow, G. M., Jin, C. &Kutner, N. G. Significance of frailty among dialysis patients. *J. Am. Soc. Nephrol.* **18**, 2960–2967 (2007).
5. Wilhelm-Leen, E. R., Hall, Y. N., M, K. T. &Chertow, G. M. Frailty and chronic kidney disease: the Third National Health and Nutrition Evaluation Survey. *Am J Med* **122**, 664–71 e2 (2009).
6. Shlipak, M. G. *et al.* The Presence of Frailty in Elderly Persons with Chronic Renal Insufficiency. *Am. J. Kidney Dis.* **43**, 861–867 (2004).
7. Roshanravan, B. *et al.* A prospective study of frailty in nephrology-referred patients with CKD. *Am J Kidney Dis* **60**, 912–921 (2012).
8. Chiang, J. M. *et al.* Low testosterone is associated with frailty, muscle wasting and physical dysfunction among men receiving hemodialysis: a longitudinal analysis. *Nephrol. Dial. Transplant.* (2018). doi:10.1093/ndt/gfy252
9. McAdams-Demarco, M. A. *et al.* Frailty and cognitive function in incident hemodialysis patients. *Clin. J. Am. Soc. Nephrol.* **10**, 2181–2189 (2015).
10. Bancu, I. *et al.* Frail Patient in Hemodialysis: A New Challenge in Nephrology-Incidence in Our Area, Barcelones Nord and Maresme. *J. Aging Res.* **2017**, 7624139 (2017).
11. Fabrício-Wehbe, S. C. C. *et al.* Cross-cultural adaptation and validity of the ‘Edmonton Frail Scale - EFS’ in a Brazilian elderly sample. *Rev. Lat. Am. Enfermagem* **17**, 1043–1049 (2009).
12. Kamijo, Y., Kanda, E., Ishibashi, Y. &Yoshida, M. Sarcopenia and Frailty in PD: Impact on Mortality, Malnutrition, and

- Inflammation. *Perit. Dial. Int.* **38**, 447–454 (2018).
13. McAdams-Demarco, M. A. *et al.* Frailty, mycophenolate reduction, and graft loss in kidney transplant recipients. *Transplantation* **99**, 805–810 (2015).
 14. Adame Perez, S. I., Senior, P. A., Field, C. J., Jindal, K. & Mager, D. R. Frailty, Health-Related Quality of Life, Cognition, Depression, Vitamin D and Health-Care Utilization in an Ambulatory Adult Population with Type 1 or Type 2 Diabetes Mellitus and Chronic Kidney Disease: A Cross-Sectional Analysis. *Can. J. Diabetes* (2018). doi:10.1016/j.jcjd.2018.06.001
 15. Chao, C.-T., Chan, D.-C. & Huang, J.-W. Frail Phenotype Might Be Associated With Higher Appendicular but Not Truncal Fat Among End-Stage Renal Disease Patients. *Journal of pain and symptom management* **53**, e1–e4 (2017).
 16. Demircioglu, D. The Association of Vitamin D Levels and the Frailty Phenotype Among Non-geriatric Dialysis Patients: A Cross-sectional Study. *Clinics* (2018). doi:10.6061/clinics/2018/e116
 17. Szeto, C.-C. *et al.* Depression and Physical Frailty Have Additive Effect on the Nutritional Status and Clinical Outcome of Chinese Peritoneal Dialysis. *Kidney Blood Press. Res.* **43**, 914–923 (2018).
 18. Chao, C.-T., Lai, H.-J., Tsai, H.-B., Yang, S.-Y. & Huang, J.-W. Frail phenotype is associated with distinct quantitative electroencephalographic findings among end-stage renal disease patients: an observational study. *BMC Geriatr.* **17**, 277 (2017).
 19. Meulendijks, F. G. *et al.* Groningen frailty indicator in older patients with end-stage renal disease. *Ren. Fail.* **37**, 1419–1424 (2015).
 20. Margiotta, E. *et al.* SuO004GUT MICROBIOTA COMPOSITION AND FRAILITY IN ELDERLY PATIENTS WITH CHRONIC KIDNEY DISEASE. *Nephrol. Dial. Transplant.* **33**, i618–i618 (2018).
 21. DeSouza Orlandi, F. & Gesualdo, G. D. Assessment of the frailty level of elderly people with chronic kidney disease undergoing hemodialysis. *ACTA Paul. Enferm.* **27**, 29–34 (2014).
 22. Sales, C. *et al.* SP651ANXIETY AND DEPRESSION IN END STAGE RENAL DISEASE PATIENTS AND ITS ASSOCIATION WITH CLINICAL AND LABORATORIAL DATA. *Nephrol. Dial. Transplant.* **32**, iii355–iii355 (2017).
 23. Haugen, C. E. *et al.* Incidence, Risk Factors, and Sequelae of Post-kidney Transplant Delirium. *J. Am. Soc. Nephrol.* **29**, 1752–1759 (2018).
 24. Moffatt, H., Moorhouse, P., Mallery, L., Landry, D. & Tennankore, K. Using the Frailty Assessment for Care Planning Tool (FACT) to

- screen elderly chronic kidney disease patients for frailty: the nurse experience. *Clin. Interv. Aging* **13**, 843 (2018).
25. Johansen, K. L., Painter, P., Delgado, C. & Doyle, J. Characterization of physical activity and sitting time among patients on hemodialysis using a new physical activity instrument. *J. Ren. Nutr.* **25**, 25–30 (2015).
 26. Ng, J. K.-C. *et al.* Frailty in Chinese Peritoneal Dialysis Patients: Prevalence and Prognostic Significance. *Kidney Blood Press. Res.* **41**, 736–745 (2016).
 27. Noori, N., Sharma Parpia, A., Lakhani, R., Janes, S. & Goldstein, M. B. Frailty and the Quality of Life in Hemodialysis Patients: The Importance of Waist Circumference. *J. Ren. Nutr.* **28**, 101–109 (2018).
 28. deSousa Meira, A. *et al.* Frailty in elderly patients with chronic kidney disease under conservative treatment. *Rev. da Rede Enferm. do Nord.* **17**, 386–392 (2016).
 29. Chao, C.-T. & Huang, J.-W. Frailty severity is significantly associated with electrocardiographic QRS duration in chronic dialysis patients. *PeerJ* **3**, e1354 (2015).
 30. Chao, C.-T., Chiang, C.-K., Huang, J.-W. & Hung, K.-Y. Self-reported frailty among end-stage renal disease patients: A potential predictor of dialysis access outcomes. *Nephrology (Carlton, Vic.)* **22**, 333–334 (2017).
 31. Chu, N. M. *et al.* Frailty and Changes in Cognitive Function after Kidney Transplantation. *J. Am. Soc. Nephrol.* **30**, 336–345 (2019).
 32. Kutner, N. G., Zhang, R., Allman, R. M. & Bowling, C. B. Correlates of ADL difficulty in a large hemodialysis cohort. *Hemodial Int* **18**, 70–77 (2014).
 33. Chao, C.-T., Huang, J.-W. & Chan, D.-C. Frail phenotype might herald bone health worsening among end-stage renal disease patients. *PeerJ* **5**, e3542 (2017).
 34. Chao, C.-T., Chiang, C.-K., Huang, J.-W. & Chan, D.-C. Effect of Frail Phenotype on Bone Mass and Vertebral Compression Fracture in Individuals Undergoing Dialysis. *Journal of the American Geriatrics Society* **64**, e19-21 (2016).
 35. Yoneki, K. *et al.* Association between frailty and bone loss in patients undergoing maintenance hemodialysis. *J. Bone Miner. Metab.* (2018). doi:10.1007/s00774-017-0898-4
 36. Delgado, C., Doyle, J. W. & Johansen, K. L. Association of frailty with body composition among patients on hemodialysis. *J. Ren. Nutr.* **23**, 356–362 (2013).
 37. Chao, C.-T. *et al.* Simple self-report FRAIL scale might be more closely associated with dialysis complications than other frailty

- screening instruments in rural chronic dialysis patients. *Nephrology (Carlton)*. **20**, 321–328 (2015).
38. Camilleri, S., Chong, S., Tangvoraphonkchai, K., Yoowannakul, S. & Davenport, A. Effect of Self-Reported Distress Thermometer Score on the Maximal Handgrip and Pinch Strength Measurements in Hemodialysis Patients. *Nutr. Clin. Pract.* **32**, 682–686 (2017).
 39. McAdams-DeMarco, M. A. *et al.* Frailty and Health-Related Quality of Life in End Stage Renal Disease Patients of All Ages. *J. frailty aging* **5**, 174–179 (2016).
 40. Lee, S. J., Son, H. & Shin, S. K. Influence of frailty on health-related quality of life in pre-dialysis patients with chronic kidney disease in Korea: a cross-sectional study. *Health Qual. Life Outcomes* **13**, 70 (2015).
 41. Mansur, H. N., Colugnati, F. A., Grincenkov, F. R. & Bastos, M. G. Frailty and quality of life: a cross-sectional study of Brazilian patients with pre-dialysis chronic kidney disease. *Heal. Qual Life Outcomes* **12**, 27 (2014).
 42. Iyasere, O. U. *et al.* Quality of Life and Physical Function in Older Patients on Dialysis: A Comparison of Assisted Peritoneal Dialysis with Hemodialysis. *Clin. J. Am. Soc. Nephrol.* **11**, 423–430 (2016).
 43. McAdams-DeMarco, M. A. *et al.* Frailty and Postkidney Transplant Health-Related Quality of Life. *Transplantation* **102**, 291–299 (2018).
 44. Kutner, N. G., Zhang, R., Huang, Y. & Wasse, H. Falls among hemodialysis patients: potential opportunities for prevention? (2014). doi:10.1093/ckj/sfu034
 45. McAdams-DeMarco, M. A. *et al.* Frailty and falls among adult patients undergoing chronic hemodialysis: a prospective cohort study. *BMC Nephrol* **14**, 224 (2013).
 46. Bao, Y., Dalrymple, L., Chertow, G. M., Kaysen, G. A. & Johansen, K. L. Frailty, dialysis initiation, and mortality in end-stage renal disease. *Arch Intern Med* **172**, 1071–1077 (2012).
 47. Delgado, C. *et al.* Association of Self-Reported Frailty with Falls and Fractures among Patients New to Dialysis. *Am. J. Nephrol.* **42**, 134–140 (2015).
 48. Konel, J. M. *et al.* Depressive symptoms, frailty, and adverse outcomes among kidney transplant recipients. *Clin. Transplant.* **32**, e13391 (2018).
 49. Lee, S.-Y. *et al.* The prevalence, association, and clinical outcomes of frailty in maintenance dialysis patients. *J. Ren. Nutr.* **27**,

- 106–112 (2017).
50. McAdams-DeMarco, M. A. *et al.* Frailty as a novel predictor of mortality and hospitalization in individuals of all ages undergoing hemodialysis. *J Am Geriatr Soc* **61**, 896–901 (2013).
 51. McAdams-DeMarco, M. A. *et al.* Frailty and Early Hospital Readmission After Kidney Transplantation. *Am. J. Transplant.* **13**, 2091–2095 (2013).
 52. McAdams-DeMarco, M. A. *et al.* Frailty, Length of Stay, and Mortality in Kidney Transplant Recipients: A National Registry and Prospective Cohort Study. *Ann. Surg.* **266**, 1084–1090 (2017).
 53. Chu, N. M. *et al.* Dynamic Frailty Before Kidney Transplantation-Time of Measurement Matters. *Transplantation* (2019). doi:10.1097/TP.0000000000002563
 54. McAdams-DeMarco, M. A. *et al.* Frailty and mortality in kidney transplant recipients. *Am. J. Transplant* **15**, 149–154 (2015).
 55. Alfaadhel, T. A. *et al.* Frailty and mortality in dialysis: evaluation of a clinical frailty scale. *Clin. J. Am. Soc. Nephrol.* **10**, 832–840 (2015).
 56. Ali, H., Abdelaziz, T., Abdelaal, F. & Baharani, J. Assessment of prevalence and clinical outcome of frailty in an elderly predialysis cohort using simple tools. *Saudi J. Kidney Dis. Transpl.* **29**, 63–70 (2018).
 57. Johansen, K. L. *et al.* Association of Performance-Based and Self-Reported Function-Based Definitions of Frailty with Mortality among Patients Receiving Hemodialysis. *Clin. J. Am. Soc. Nephrol.* **11**, 626–632 (2016).
 58. Pérez Fernández, M. *et al.* Comorbidity, Frailty, and Waitlist Mortality among Kidney Transplant Candidates of All Ages. *Am. J. Nephrol.* **49**, 103–110 (2019).
 59. Fitzpatrick, J. *et al.* Frailty, body composition and the risk of mortality in incident hemodialysis patients: the Predictors of Arrhythmic and Cardiovascular Risk in End Stage Renal Disease study. *Nephrol. Dial. Transplant* **34**, 346–354 (2019).
 60. Romeu, M. *et al.* Survival of patients with ANCA-associated vasculitis on chronic dialysis: data from the French REIN registry from 2002 to 2011. *QJM* **107**, 545–555 (2014).
 61. Decourt, A. *et al.* Trends in Survival and Renal Recovery in Patients with Multiple Myeloma or Light-Chain Amyloidosis on Chronic Dialysis. *Clin. J. Am. Soc. Nephrol.* **11**, 431–441 (2016).
 62. Lee, S. W. *et al.* Is Frailty a Modifiable Risk Factor of Future Adverse Outcomes in Elderly Patients with Incident End-Stage Renal

Disease? *J. Korean Med. Sci.* **32**, 1800–1806 (2017).

63. Schopmeyer, L. *et al.* Frailty has a significant influence on postoperative complications after kidney transplantation-a prospective study on short-term outcomes. *Transpl. Int.* **32**, 66–74 (2019).